

involve

a journal of mathematics

A generalization of Eulerian numbers
via rook placements

Esther Banaian, Steve Butler, Christopher Cox,
Jeffrey Davis, Jacob Landgraf and Scarlitte Ponce



A generalization of Eulerian numbers via rook placements

Esther Banaian, Steve Butler, Christopher Cox,
Jeffrey Davis, Jacob Landgraf and Scarlitte Ponce

(Communicated by Jim Haglund)

We consider a generalization of Eulerian numbers which count the number of placements of cn rooks on an $n \times n$ chessboard where there are exactly c rooks in each row and each column, and exactly k rooks below the main diagonal. The standard Eulerian numbers correspond to the case $c = 1$. We show that for any c the resulting numbers are symmetric and give generating functions of these numbers for small values of k .

1. Introduction

Rook placements on boards have a wonderful and rich history in combinatorics (see, e.g., [Butler et al. 2011]). Traditionally the rooks are placed in a nonattacking fashion (i.e., at most one rook in each row and column) and the combinatorial aspects come from considering variations on the board shapes.

Instead of varying the board, we could also change the restrictions on how many rooks are allowed in each row and each column. If we have a square board and the number of rooks in each column and row is fixed, then this corresponds to counting nonnegative matrices with fixed row and column sums; see A000681, A001500, A257493, etc., in the [On-Line Encyclopedia of Integer Sequences](#) (OEIS).

In this paper, we will look at this latter case of placing multiple rooks in each row and column more closely. We begin in [Section 2](#) by exploring the connections between these rook placements and juggling patterns. In [Section 3](#), we look at Eulerian numbers (which correspond to the number of nonattacking rook placements on an $n \times n$ board with a fixed number of rooks below the main diagonal) and in [Section 4](#) generalize to the case in which c rooks are placed in each row and each column. In [Section 5](#), we provide generating functions for special cases of these generalized Eulerian numbers. We end with concluding remarks and open problems in [Section 6](#).

MSC2010: 05A15.

Keywords: Eulerian numbers, juggling, recursion, multiplex.

2. Minimal juggling patterns and rook placements

Juggling patterns can be described by a *siteswap* sequence listing the throws that the pattern requires, i.e., $t_1 t_2 \cdots t_n$ where at time $s \equiv i \pmod n$ we throw the ball so that it will land t_i beats in the future. A sequence of throws can be juggled if and only if there are no collisions, i.e., two balls landing at the same time, which is equivalent to $1 + t_1, 2 + t_2, \dots, n + t_n$ being distinct modulo n . One well-known property of siteswap sequences is that the average of the throws is the number of balls needed to juggle the pattern (see [Buhler et al. 1994; Polster 2003]).

A *minimal juggling pattern* is a valid juggling pattern $t_1 t_2 \cdots t_n$ with $0 \leq t_i \leq n - 1$. These form the basic building blocks of juggling patterns since all juggling patterns of period n arise by starting from some minimal juggling pattern and adding multiples of n to the various throws (such additions do not affect modular conditions). More about this approach was found by Buhler, Eisenbud, Graham and Wright [1994].

This naturally leads to the problem of enumerating minimal juggling patterns. This is done by relating such patterns to rook placements on a square board. In particular we will consider the $n \times n$ board \mathcal{B}_n , with labels on each cell (i, j) given by the rule

$$\begin{cases} j - i & \text{if } j \geq i, \\ n + j - i & \text{if } j < i. \end{cases}$$

We can interpret the rows of \mathcal{B}_n as the *throwing* times (modulo n) and the columns of \mathcal{B}_n as the *landing* times (modulo n). The label of the cell (i, j) is then the smallest possible throw required to throw at time i and land at time j .

Given a minimal juggling pattern $t_1 t_2 \cdots t_n$, we form a rook placement by placing a rook in row i on the cell labeled t_i for $1 \leq i \leq n$ (note that this forces the rook to be placed in the column corresponding to the landing time modulo n). Since landing times are unique modulo n , no two rooks will be in the same column, so this forms a nonattacking rook placement with n rooks. Conversely, given a nonattacking rook placement with n rooks we can form a minimal juggling pattern by reading off the cell labels of the covered square starting at the first row and reading down. This establishes the bijective relationship between minimal juggling patterns and nonattacking rook patterns on \mathcal{B}_n . An example of this is shown in Figure 1 for the minimal juggling pattern 24234.

We can extract information about the minimal juggling pattern by properties of the rook placements, including, for example, the number of balls.

Proposition 1. *The number of rooks below the main diagonal in a nonattacking rook placement on \mathcal{B}_n is the same as the number of balls necessary to juggle the corresponding minimal juggling pattern.*

0	1	2	3	4
4	0	1	2	3
3	4	0	1	2
2	3	4	0	1
1	2	3	4	0

Figure 1. A nonattacking rook placement on \mathcal{B}_5 corresponding to the minimal juggling pattern 24234.

Proof. Suppose there are k rooks below the main diagonal in a placement of n nonattacking rooks on \mathcal{B}_n . Then when we sum the labels of all the cells covered by a rook, i.e., we sum the throw heights for the juggling sequence, we have

$$\sum_{\ell=1}^n t_\ell = kn + \sum_{j=1}^n j - \sum_{i=1}^n i = kn.$$

Since the average of the throws is the number of balls needed for the sequence, the claim follows. □

Note that in [Figure 1](#) there are three rooks below the main diagonal and that the juggling pattern 24234 requires three balls to juggle.

2.1. Multiplex juggling and c -rook placements. A natural variation in juggling is to allow multiple balls to be caught and thrown at a time. This is known as *multiplex juggling*, and we will see that many of the basic ideas generalize well to this setting.

We will let c denote a hand capacity; i.e., at each beat we make c throws (allowing some of the throws to be 0, which happens when the number of actual balls thrown is less than c). Siteswap sequences of period n now correspond to a sequence of n sets, $T_1 T_2 \cdots T_n$, where each T_i is a (multi)set of the form $\{t_{i,1}, t_{i,2}, \dots, t_{i,c}\}$, denoted in shorthand notation as $[t_{i,1} t_{i,2} \cdots t_{i,c}]$. A multiplex juggling sequence is valid if and only if the *juggling modular condition* is satisfied. Namely, every $1 \leq \ell \leq n$ appears exactly c times in the multiset

$$\{t_{i,j} + i \pmod{n}\}_{\substack{1 \leq i \leq n \\ 1 \leq j \leq c}}$$

In other words, no more than c balls land at each time.¹ As in standard juggling patterns, the number of balls b needed to juggle the pattern relates to an average. In particular,

$$\frac{1}{n} \sum_{i=1}^n \sum_{j=1}^c t_{i,j} = b.$$

¹A 0 throw indicates a ball is not landing.

0	1	2	3	4
4	0	1	2	3
3	4	0	1	2
2	3	4	0	1
1	2	3	4	0

Figure 2. A 2-rook placement on \mathcal{B}_5 corresponding to the minimal multiplex juggling pattern [24][02][14][22][03].

We say a multiplex juggling sequence is a *minimal multiplex juggling sequence* if and only if $0 \leq t_{i,j} \leq n - 1$ for all throws $t_{i,j}$.

There is a relationship between period n , hand capacity c , multiplex juggling sequences and placements of “rooks” on \mathcal{B}_n . This is done by generalizing from nonattacking rook placements to *c-rook placements*, placements of cn rooks with exactly c rooks in every row and column, where multiple rooks are allowed in cells.

There is a bijection between minimal multiplex juggling patterns of period n with hand capacity c and *c-rook placements* on \mathcal{B}_n . In particular, for each i we place c rooks in the i -th row corresponding to $t_{i,1}, \dots, t_{i,c}$. Conversely, given a *c-rook placement* we can form a minimal multiplex juggling sequence by letting T_i denote the cells covered by the rooks in row i (with appropriate multiplicity). An example of this is shown in [Figure 2](#) for the minimal multiplex juggling pattern [24][02][14][22][03].

By the same argument used for [Proposition 1](#) we have the following.

Proposition 2. *The number of rooks below the main diagonal in a c-rook placement on \mathcal{B}_n is the same as the number of balls necessary to juggle the corresponding minimal multiplex juggling pattern.*

For example, the multiplex juggling pattern in [Figure 2](#) requires four balls to juggle.

3. Eulerian numbers

The Eulerian numbers, denoted $\langle n \atop k \rangle$, are usually defined as the number of permutations of $[n]$, $\pi = \pi_1\pi_2 \cdots \pi_n$, with k ascents ($\pi_i < \pi_{i+1}$), or equivalently the number of permutations with k descents ($\pi_i > \pi_{i+1}$). There is a bijection between permutations of $[n]$ with k descents and permutations with k drops ($i > \pi_i$), so that $\langle n \atop k \rangle$ also counts permutations of $[n]$ with k drops (see [\[Buhler et al. 1994\]](#)). Given an $n \times n$ board, with rows and columns labeled $1, 2, \dots, n$, we can use our permutation to form a nonattacking rook placement by placing rooks at positions (i, π_i) . A drop

$n \downarrow k \rightarrow$	0	1	2	3	4	5	6
1	1						
2	1	1					
3	1	4	1				
4	1	11	11	1			
5	1	26	66	26	1		
6	1	57	302	302	57	1	
7	1	120	1191	2416	1191	120	1

Table 1. The Eulerian numbers $\langle \begin{smallmatrix} n \\ k \end{smallmatrix} \rangle$ for $1 \leq n \leq 7$.

in the permutation corresponds to a rook below the main diagonal, so we will call any rook below the main diagonal a *drop*.

By Proposition 1, the number of drops in a nonattacking rook placement equals the number of balls necessary for the corresponding juggling pattern. Therefore, $\langle \begin{smallmatrix} n \\ k \end{smallmatrix} \rangle$ also counts the number minimal juggling patterns of period n using k balls.

The Eulerian numbers have many nice properties, some of which can be seen in Table 1. For example, they are symmetric, i.e., $\langle \begin{smallmatrix} n \\ k \end{smallmatrix} \rangle = \langle \begin{smallmatrix} n \\ n-k-1 \end{smallmatrix} \rangle$. This can be shown by noting if we start with a permutation with k ascents and reverse the permutation, we now have $n - 1 - k$ ascents (i.e., ascents go to descents and vice-versa; and there are $n - 1$ consecutive pairs). We will give a different proof of this symmetry in the next section using rook placements.

Another well-known property of the Eulerian numbers is a recurrence relation.

Proposition 3. *The Eulerian numbers satisfy $\langle \begin{smallmatrix} n \\ k \end{smallmatrix} \rangle = (n - k)\langle \begin{smallmatrix} n-1 \\ k-1 \end{smallmatrix} \rangle + (k + 1)\langle \begin{smallmatrix} n-1 \\ k \end{smallmatrix} \rangle$.*

This recurrence is again proven using permutations and ascents. Here, we provide an alternate proof using rook placements and drops.

Proof. Start by considering a nonattacking rook placement on an $(n - 1) \times (n - 1)$ board with $k - 1$ drops. Add an n -th row and n -th column, and place a rook in position (n, n) . The newly added rook is not below the diagonal and so we have not created any new drops. We can now create one additional drop by taking any rook (other than the one just added) which is *on or above* the main diagonal, say in position (i, j) , move that rook to position (n, j) and move the rook in position (n, n) to position (i, n) . This moves the rook in the j -th column below the main diagonal creating a new drop. Since no other rook moves, we now have precisely k drops and a nonattacking rook placement. Note that there are $(n - 1) - (k - 1) = n - k$ ways we could have chosen which rook to move, so that in total this gives $(n - k)\langle \begin{smallmatrix} n-1 \\ k-1 \end{smallmatrix} \rangle$ boards of size $n \times n$ with k drops.

Now, consider a nonattacking rook placement on an $(n - 1) \times (n - 1)$ board with k drops. Add an n -th row and n -th column, and place a rook in position (n, n) .

As before we switch, but now only switch with a rook which is *below* the main diagonal (i.e., a drop). This will not change the number of drops, so the result is a nonattacking rook placement on an $n \times n$ board with k drops. There are k rooks we can choose to switch with, or alternatively, we can leave the n -th rook in position (n, n) ; thus, there are $k + 1$ ways to build the desired rook placement, so that in total this gives $(k + 1)\binom{n-1}{k}$ boards of size $n \times n$ with k drops.

Finally, we note that each $n \times n$ board with k drops is formed uniquely from one of these operations. This can be seen by taking such a board and then noting the location of the rook(s) in the last row and in the last column. Suppose these are in positions (i, n) and (n, j) , respectively. We then move these rooks to positions (i, j) and (n, n) . This can at most decrease the number of drops by one (i.e., moving the rook in the last column does not affect the number of drops). Now removing the last row and column gives an $(n - 1) \times (n - 1)$ board having a nonattacking rook placement with either k or $k - 1$ drops. □

4. Generalized Eulerian numbers

The *generalized Eulerian numbers*, denoted $\binom{n}{k}_c$, are the number of c -rook placements on the $n \times n$ board with k drops. Just as the Eulerian numbers count the number of minimal juggling patterns of period n with k balls, the generalized Eulerian numbers count the number of minimal multiplex juggling patterns of period n with k balls and hand capacity c . Notice that the generalized Eulerian numbers reduce to the Eulerian numbers when $c = 1$. In [Table 2](#) we give some of the generalized Eulerian numbers for $c = 2$ and 3.

$n \downarrow k \rightarrow$	0	1	2	3	4	5	6	7	8	
1	1									
2	1	1	1							
3	1	4	11	4	1					
4	1	11	72	114	72	11	1			
5	1	26	367	1492	2438	1492	367	26	1	

$n \downarrow k \rightarrow$	0	1	2	3	4	5	6	7	8	9	
1	1										
2	1	1	1	1							
3	1	4	11	23	11	4	1				
4	1	11	72	325	595	595	325	72	11	1	

Table 2. Small values of the generalized Eulerian $\binom{n}{k}_c$ numbers for $c = 2$ (top) and 3 (bottom).

These numbers appear to satisfy a symmetry property similar to Eulerian numbers. We will give two proofs of this symmetry, one in terms of rook placements and the other using minimal multiplex juggling patterns.

Theorem 4. *Let n, k and c be nonnegative integers. Then $\langle n \rangle_k^c = \langle c(n-1)-k \rangle_c^n$.*

Proof. We construct a bijection between the rook placements with k rooks below the main diagonal and those with $c(n - 1) - k$ rooks below the diagonal. Consider a rook placement with c rooks in every row and column, and k rooks below the diagonal. Now, shift every rook one space to the right cyclically. Let us consider the number of rooks which are *strictly* above the main diagonal.

- All c rooks in the last column were shifted to the first column. So, none of these rooks are above the main diagonal.
- All of the k rooks that were initially below the main diagonal are now either on or still below the main diagonal.
- All other rooks will be above the diagonal.

Since there are cn rooks on the board total, there are $cn - c - k = c(n - 1) - k$ rooks above the diagonal after this shift. Finally, we switch the rows and columns of the board. This flips the rook placement across the main diagonal. After this transformation, there are now $c(n - 1) - k$ rooks *below* the main diagonal. This composition of transformations is invertible by switching rows and columns then shifting every rook left one space. Thus, the transformation gives a bijection, completing the proof. □

Before we can give the second proof, we must first establish some basic properties of (multiplex) juggling sequences.

Lemma 5. *If the sequence $T_1 T_2 \cdots T_n$ satisfies the juggling modular conditions with hand capacity c , and $\alpha \in \mathbb{Z}_n$ with $\gcd(\alpha, n) = 1$, then $(\alpha T_{1\alpha^{-1}})(\alpha T_{2\alpha^{-1}}) \cdots (\alpha T_{n\alpha^{-1}})$, where*

$$\alpha T_i := \{\alpha t_{i,1}, \dots, \alpha t_{i,c}\},$$

and the subscripts are taken modulo n , also satisfies the juggling modular conditions.

Proof. We have

$$A = \{\alpha t_{i\alpha^{-1},j} + i\}_{\substack{1 \leq i \leq n \\ 1 \leq j \leq c}} = \{\alpha(t_{i\alpha^{-1},j} + i\alpha^{-1})\}_{\substack{1 \leq i \leq n \\ 1 \leq j \leq c}} = \{\alpha(t_{i',j} + i')\}_{\substack{1 \leq i' \leq n \\ 1 \leq j \leq c}},$$

where we use that $\gcd(\alpha, n) = 1$ so that α is invertible modulo n and as i ranges between 1 and n , then so does $i' := i\alpha^{-1}$. Since $\{t_{i,j} + i\}_{1 \leq i \leq n, 1 \leq j \leq c}$ has c occurrences each of 1 through n , then scaling by α and taking terms modulo n we also have that A will have c occurrences each of 1 through n . □

Lemma 6. *If the sequence $T_1 T_2 \cdots T_n$ satisfies the juggling modular conditions of hand capacity c , and $\beta \in \mathbb{Z}$, then $(T_1 + \beta)(T_2 + \beta) \cdots (T_n + \beta)$, where*

$$T_i + \beta := [t_{i,1} + \beta, t_{i,2} + \beta, \dots, t_{i,c} + \beta],$$

still satisfies the juggling modular conditions.

Proof. The multiset

$$A = \{(t_{i,j} + \beta) + i\}_{\substack{1 \leq i \leq n \\ 1 \leq j \leq c}}$$

is found by taking $\{t_{i,j} + i\}_{1 \leq i \leq n, 1 \leq j \leq c}$ and shifting each element by β . Since $T_1 T_2 \cdots T_n$ satisfy the juggling modular conditions then so also must A . \square

Juggling proof of Theorem 4. We show there is a bijection between the minimal multiplex juggling sequences using k balls and those using $c(n - 1) - k$ balls for a fixed length n and hand capacity c . So let $T_1 T_2 \cdots T_n$ be a valid minimal multiplex juggling sequence with k balls and hand capacity c . By Lemma 5 and Lemma 6, if we scale each T_i by -1 (reversing the indexing) and add $n - 1$ then the resulting sets still satisfy the modular juggling conditions. In particular we have that the following satisfies the modular juggling condition:

$$(n - 1 - T_n)(n - 1 - T_{n-1}) \cdots (n - 1 - T_1).$$

We also note the resulting throws all lie between 0 and $n - 1$ so that this is indeed a minimal juggling pattern.

The number of balls in the new juggling sequence is

$$\frac{1}{n} \sum_{i=1}^n \sum_{j=1}^c ((n - 1) - t_{i,j}) = \frac{1}{n} \left(cn(n - 1) - \sum_{i=1}^n \sum_{j=1}^c t_{i,j} \right) = c(n - 1) - k.$$

Finally, we note that this operation is its own inverse, and thus gives the desired bijection. \square

5. Generalized Eulerian numbers for small k

We now look at determining the values of the generalized Eulerian numbers $\langle n \rangle_c^k$ for small k . This depends of course on both n and c . However, for a fixed k there are only finitely many c that need to be considered. This is a consequence of the following lemma.

Lemma 7. *For $c \geq k$ we have $\langle n \rangle_c^k = \langle n \rangle_k^k$.*

Proof. It will suffice to establish the following claim.

Claim. Every c -rook placement with k drops has at least $c - k$ rooks in every entry on the main diagonal.

We proceed to establish this by using induction on $k + c$. For $k + c = 1$, the only possible case is $k = 0$ and $c = 1$ for which there is only one placement, namely one rook in each cell on the main diagonal.

Now assume that we have established the claim for all k, c with $k + c < \ell$, and let $k + c = \ell$. Let S be a c -rook placement with k drops. We can interpret the rook placement as an incidence relationship of a regular bipartite graph. By Hall's marriage theorem, we know we can find a perfect matching in this bipartite graph which corresponds to T , a 1-rook placement contained in S . Suppose there are i drops in T . Then, $S - T$ is a $(c - 1)$ -rook placement with $k - i$ drops. Since $(c - 1) + (k - i) < c + k = \ell$, by our induction hypothesis, there are at least $c - k + i - 1$ rooks on each entry on the main diagonal in $S - T$, and hence also in S . If $i \geq 1$, we are done. If $i = 0$, then T is again the unique 1-rook placement where every rook is on the main diagonal, so S still has at least $c - k$ rooks on each entry on the main diagonal. \square

This can also be established in terms of minimal multiplex juggling patterns.

Juggling proof of Lemma 7. If there are k balls, then at each step we can throw at most k balls, i.e., each T_i has at least $c - k$ entries of 0. It follows that in the corresponding c -rook placement each row has at least $c - k$ rooks on the diagonal. \square

We will be looking at the generalized Eulerian numbers $\langle n \rangle_k^c$ for $k = 1, 2, 3$. By Lemma 7 this reduces down to only six cases to consider, namely, $\langle n \rangle_1^1, \langle n \rangle_2^1, \langle n \rangle_2^2, \langle n \rangle_3^1, \langle n \rangle_3^2$ and $\langle n \rangle_3^3$. Since $\langle n \rangle_k^1 = \langle n \rangle_k$, the cases $\langle n \rangle_1^1, \langle n \rangle_2^1$ and $\langle n \rangle_3^1$ have been previously determined (see A000295, A000460 and A000498, respectively, in the OEIS). So that leaves $\langle n \rangle_2^2, \langle n \rangle_3^2$ and $\langle n \rangle_3^3$ and in Table 3 we give the generating function for these three sequences. In the remainder of this section we will demonstrate the techniques used to determine the generating functions by working through the case for $\langle n \rangle_2^2$.

5.1. Placing rooks in a generic rook placement. We break the problem of counting c -rook placements into several subproblems according to the way the rooks below the main diagonal are placed relative to one another (i.e., relative placements instead of absolute placements). Given some generic placement of the k rooks below the main diagonal we can determine the number of ways to place the remaining rooks on or above the main diagonal. We then combine the results over all possible generic placements.

We will carefully work through the rook placement shown in Figure 3, which consists of two rooks below the main diagonal and where both rooks are in the same column and different rows. Here a, b, c and d are the number of rows between the various transition points (a transition point to passing a rook, or rooks, in a row or a column as we move along the main diagonal).

We place the remaining rooks one row at a time starting from the bottom and going to the top. For each new row, the way we place rooks will depend on all of

$$\sum_{n \geq 0} \left\langle \begin{matrix} n \\ 2 \end{matrix} \right\rangle_2 x^n = x^2 + 11x^3 + 72x^4 + 367x^5 + 1630x^6 + 6680x^7 + 26082x^8 + \dots$$

$$= \frac{x^2 - x^3 - x^4 - 3x^5 + 5x^6}{(1-x)^3(1-2x)^2(1-5x+5x^2)}$$

$$\sum_{n \geq 0} \left\langle \begin{matrix} n \\ 3 \end{matrix} \right\rangle_2 x^n = 4x^3 + 114x^4 + 1492x^5 + 13992x^6 + 109538x^7 + 769632x^8 + \dots$$

$$= \frac{4x^3 + 2x^4 - 300x^5 + 1748x^6 - 4676x^7 + 7058x^8 - 6648x^9 + 4397x^{10} - 2206x^{11} + 625x^{12}}{(1-x)^4(1-2x)^3(1-5x+5x^2)^2(1-8x+13x^2)}$$

$$\sum_{n \geq 0} \left\langle \begin{matrix} n \\ 3 \end{matrix} \right\rangle_3 x^n = x^2 + 23x^3 + 325x^4 + 3368x^5 + 28819x^6 + 218788x^7 + \dots$$

$$= \frac{x^2 - 7x^3 + 39x^4 - 336x^5 + 1844x^6 - 5545x^7 + 9697x^8 - 10404x^9 + 7532x^{10} - 4558x^{11} + 2435x^{12} - 700x^{13}}{(1-x)^4(1-2x)^3(1-5x+5x^2)^2(1-10x+27x^2-20x^3)}$$

Table 3. Generating functions for some of the generalized Eulerian numbers.




Figure 3. A 2-rook placement with two rooks below the main diagonal where both rooks are in the same column and different rows.

the choices we have made previously. However, it suffices to know only what is happening locally. In particular, we only need to know how many columns can have rooks placed into them, as well as the respective numbers that can go into those columns. We can represent these by a partition of what we will call the *excess* (the total number of rooks that can still be placed in the columns *after* the row has had its rooks placed). As we move one row up the board we will gain a new column (from the diagonal) and the excess will change in one of several ways.


- There are no rooks below or to the left of the new diagonal cell. Initially we now have a new column that can take up to c rooks, and we place c rooks in the row. The excess remains unchanged.
- There are τ rooks below the new diagonal cell. Initially we have the new column, but that can only take up to $c - \tau$ rooks (i.e., τ rooks have already

gone into the column), and we still have to place c rooks in the row. The excess decreases by τ .

- There are σ rooks to the left of the new diagonal cell. Initially we have the new column that can take up to c rooks, and we place $c - \sigma$ rooks in the row (i.e., σ rooks have already gone into the row). The excess increases by σ .

We note that it is possible for the last two situations to occur simultaneously.

In going from row to row we will transition from partitions of the old excess to partitions of the new excess. We illustrate this with an example in which case the excesses are both 2. We indicate a column which can still have r rooks placed into it by \boxed{r} , then underneath look at all possible ways we can place 2 rooks into those columns, and finally note the resulting set of columns contributing to the new excess:



This can be modeled by a transition matrix where the *columns* of the transition matrix correspond to the excess of the original row and the *rows* of the transition matrix correspond to the partitions of the excess of the new row:

$$\begin{matrix} & & \boxed{2} & \boxed{1} & \boxed{1} \\ & \boxed{2} & & & \\ \boxed{1} & \boxed{1} & \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix} & & \end{matrix}$$

Repeating this for all possible situations that might arise for transitioning between excesses 0, 1, or 2, we get the transition matrices in the following table:

		transition from			
		\emptyset	$\boxed{1}$	$\boxed{2}$	$\boxed{1} \boxed{1}$
transition to	\emptyset	(1)	(1)	(1)	(1)
	$\boxed{1}$	(1)	(2)	(2)	(3)
	$\boxed{2}$	(1)	(1)	(2)	(1)
	$\boxed{1} \boxed{1}$	(0)	(1)	(1)	(3)

We now start below the bottom row (in one possible way) and we move up from row to row and multiply on the *left* by the transition that we perform between the two rows. At any point we stop, the resulting vector will denote the number of ways to fill up the board to that row with a particular excess. In particular, if we

carry this procedure all the way to the top we will get a 1×1 matrix whose entry is the number of ways to fill in the rooks on and above the main diagonal.

For [Figure 3](#), where we have of runs of a , b , c and d rows as well as three other transitions to make, the resulting product that gives our count is

$$(1)^d(1\ 1) \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}^c \begin{pmatrix} 1 \\ 1 \end{pmatrix} (2)^b(1)(1)^a.$$

Finally, for this generic rook placement we sum over all possible choices of a , b , c and d that gives an $n \times n$ board, i.e.,

$$\sum_{a+b+c+d=n-3} (1)^d(1\ 1) \begin{pmatrix} 2 & 1 \\ 1 & 3 \end{pmatrix}^c \begin{pmatrix} 1 \\ 1 \end{pmatrix} (2)^b(1)(1)^a.$$

In order to help evaluate this sum, we will add in an extra parameter x that keeps track of how many of each transition we made, or viewed another way, the power of x corresponds to the number of rows we have. Therefore when counting the number of placements on an $n \times n$ board, we are interested in the coefficient of x^n of the expression

$$\sum_{a,b,c,d \geq 0} (x)^d x(1\ 1) \begin{pmatrix} 2x & x \\ x & 3x \end{pmatrix}^c x \begin{pmatrix} 1 \\ 1 \end{pmatrix} (2x)^b x(x)^a.$$

This sum can be decomposed as a combination of geometric sums giving

$$\begin{aligned} & \sum_{a,b,c,d \geq 0} (x)^d x(1\ 1) \begin{pmatrix} 2x & x \\ x & 3x \end{pmatrix}^c x \begin{pmatrix} 1 \\ 1 \end{pmatrix} (2x)^b x(x)^a \\ &= x^3 \left(\sum_{d \geq 0} x^d \right) (1\ 1) \left(\sum_{c \geq 0} \begin{pmatrix} 2x & x \\ x & 3x \end{pmatrix}^c \right) \begin{pmatrix} 1 \\ 1 \end{pmatrix} \left(\sum_{b \geq 0} (2x)^b \right) \left(\sum_{a \geq 0} x^a \right) \\ &= x^3 \cdot \frac{1}{1-x} \cdot (1\ 1) \left(I - \begin{pmatrix} 2x & x \\ x & 3x \end{pmatrix} \right)^{-1} \begin{pmatrix} 1 \\ 1 \end{pmatrix} \cdot \frac{1}{1-2x} \cdot \frac{1}{1-x} \\ &= \frac{x^3}{(1-x)^2(1-2x)} \cdot (1\ 1) \left(\frac{1}{1-5x+5x^2} \begin{pmatrix} 1-3x & x \\ x & 1-2x \end{pmatrix} \right) \begin{pmatrix} 1 \\ 1 \end{pmatrix} \\ &= \frac{x^3(2-3x)}{(1-x)^2(1-2x)(1-5x+5x^2)}. \end{aligned}$$

This is the generating function for one of the generic ways to place rooks. We can now repeat this procedure for every way in which we can place rooks below the main diagonal and add the individual generating functions together. All the seven generic cases, with their corresponding generating functions, are shown in [Figure 4](#). Adding the individual generating functions together then gives us the overall generating function that was given in [Table 3](#).




Figure 4. All generic 2-rook placements and corresponding generating functions.

This same process works for determining the generating function of $\langle n \rangle_k^c$ for any fixed k and c . The main challenge lies in that the number of generic cases that have to be considered grows drastically as we increase c and k . This is demonstrated in Table 4. It is possible to automate this process, which was used for determining the generating functions for $k = 3$ given in Table 3.

$k \downarrow c \rightarrow$	1	2	3	4	5	6	7
1	1						
2	4	7					
3	26	68	75				
4	236	940	1090	1105			
5	2752	16645	20360	20790	20821		
6	39208	360081	464111	477242	478376	478439	
7	660032	9202170	12492277	12933423	12974826	12977688	12977815

Table 4. The number of generic c -rook placements with k rooks below the main diagonal.

6. Conclusion

The generalized Eulerian numbers are a natural extension of the Eulerian numbers, at least in regards to the interpretation coming from rook placements. We have also seen that these numbers exhibit a symmetry similar to that of the Eulerian numbers. It would be interesting to know which other properties and relationships involving Eulerian numbers generalize. Some natural candidates to try and generalize include the following:

- Is there a generalization of the recurrence in [Proposition 3](#) for Eulerian numbers to generalized Eulerian numbers? Related to this, is there a simple generating function for the generalized Eulerian numbers?
- Is there a generalization of Worpitzky’s identity, $x^n = \sum_k \binom{n}{k} \langle x+k \rangle_n$, to generalized Eulerian numbers? Worpitzky’s identity is used in counting the number of juggling patterns (see [\[Buhler et al. 1994\]](#)), so a generalization might be useful in counting multiplex juggling patterns.
- Is there a generalization of the identity of Chung, Graham and Knuth [\[2010\]](#),

$$\sum_k \binom{a+b}{k} \langle k \rangle_{a-1} = \sum_k \binom{a+b}{k} \langle k \rangle_{b-1}?$$

Note that this uses the convention $\binom{0}{0} = 0$.

More about the Eulerian numbers and various identities and relationships that could be considered are given by Graham, Knuth and Patashnik [\[1994, §6.2\]](#).

We also note the original motivation for investigating these numbers was looking into the mathematics of multiplex juggling. There is a close connection between the mathematics of juggling and the mathematics of rook placements. We hope to see this relationship strengthened in future work.

Acknowledgment

The research was conducted at the 2015 REU program held at Iowa State University, which was supported by NSF DMS 1457443.

References

- [Buhler et al. 1994] J. Buhler, D. Eisenbud, R. Graham, and C. Wright, “Juggling drops and descents”, *Amer. Math. Monthly* **101**:6 (1994), 507–519. [MR](#) [Zbl](#)
- [Butler et al. 2011] F. Butler, M. Can, J. Haglund, and J. Remmel, “Rook theory notes”, 2011, available at <http://www.math.ucsd.edu/~remmel/files/Book.pdf>.
- [Chung et al. 2010] F. Chung, R. Graham, and D. Knuth, “A symmetrical Eulerian identity”, *J. Comb.* **1**:1 (2010), 29–38. [MR](#) [Zbl](#)
- [Graham et al. 1994] R. L. Graham, D. E. Knuth, and O. Patashnik, *Concrete mathematics: a foundation for computer science*, 2nd ed., Addison-Wesley, Reading, MA, 1994. [MR](#) [Zbl](#)
- [Polster 2003] B. Polster, *The mathematics of juggling*, Springer, 2003. [MR](#) [Zbl](#)

Received: 2016-04-26

Accepted: 2016-07-11

embanaian@csbsju.edu

College of St. Benedict, Collegeville, MN 56321, United States

butler@iastate.edu

*Department of Mathematics, Iowa State University,
Ames, IA 50011, United States*

cocox@andrew.cmu.edu

*Department of Mathematical Sciences, Carnegie Mellon
University, Pittsburgh, PA 15213, United States*

jeffrey.laylon.davis@emory.edu

*Department of Mathematics and Computer Science,
Emory University, Atlanta, GA 30322, United States*

jlandgr1@nd.edu

*Department of Mathematics, University of Notre Dame,
Notre Dame, IN 46556, United States*

sponce@iastate.edu

*Department of Mathematics, Iowa State University,
Ames, IA 50011, United States*

INVOLVE YOUR STUDENTS IN RESEARCH

Involve showcases and encourages high-quality mathematical research involving students from all academic levels. The editorial board consists of mathematical scientists committed to nurturing student participation in research. Bridging the gap between the extremes of purely undergraduate research journals and mainstream research journals, *Involve* provides a venue to mathematicians wishing to encourage the creative involvement of students.

MANAGING EDITOR

Kenneth S. Berenhaut Wake Forest University, USA

BOARD OF EDITORS

Colin Adams	Williams College, USA	Suzanne Lenhart	University of Tennessee, USA
John V. Baxley	Wake Forest University, NC, USA	Chi-Kwong Li	College of William and Mary, USA
Arthur T. Benjamin	Harvey Mudd College, USA	Robert B. Lund	Clemson University, USA
Martin Bohner	Missouri U of Science and Technology, USA	Gaven J. Martin	Massey University, New Zealand
Nigel Boston	University of Wisconsin, USA	Mary Meyer	Colorado State University, USA
Amarjit S. Budhiraja	U of North Carolina, Chapel Hill, USA	Emil Minchev	Ruse, Bulgaria
Pietro Cerone	La Trobe University, Australia	Frank Morgan	Williams College, USA
Scott Chapman	Sam Houston State University, USA	Mohammad Sal Moslehian	Ferdowsi University of Mashhad, Iran
Joshua N. Cooper	University of South Carolina, USA	Zuhair Nashed	University of Central Florida, USA
Jem N. Corcoran	University of Colorado, USA	Ken Ono	Emory University, USA
Toka Diagana	Howard University, USA	Timothy E. O'Brien	Loyola University Chicago, USA
Michael Dorff	Brigham Young University, USA	Joseph O'Rourke	Smith College, USA
Sever S. Dragomir	Victoria University, Australia	Yuval Peres	Microsoft Research, USA
Behrouz Emamizadeh	The Petroleum Institute, UAE	Y.-F. S. Pétermann	Université de Genève, Switzerland
Joel Foisy	SUNY Potsdam, USA	Robert J. Plemmons	Wake Forest University, USA
Erin W. Fulp	Wake Forest University, USA	Carl B. Pomerance	Dartmouth College, USA
Joseph Gallian	University of Minnesota Duluth, USA	Vadim Ponomarenko	San Diego State University, USA
Stephan R. Garcia	Pomona College, USA	Bjorn Poonen	UC Berkeley, USA
Anant Godbole	East Tennessee State University, USA	James Propp	U Mass Lowell, USA
Ron Gould	Emory University, USA	József H. Przytycki	George Washington University, USA
Andrew Granville	Université Montréal, Canada	Richard Rebarber	University of Nebraska, USA
Jerrold Griggs	University of South Carolina, USA	Robert W. Robinson	University of Georgia, USA
Sat Gupta	U of North Carolina, Greensboro, USA	Filip Saidak	U of North Carolina, Greensboro, USA
Jim Haglund	University of Pennsylvania, USA	James A. Sellers	Penn State University, USA
Johnny Henderson	Baylor University, USA	Andrew J. Sterge	Honorary Editor
Jim Hoste	Pitzer College, USA	Ann Trenk	Wellesley College, USA
Natalia Hritonenko	Prairie View A&M University, USA	Ravi Vakil	Stanford University, USA
Glenn H. Hurlbert	Arizona State University, USA	Antonia Vecchio	Consiglio Nazionale delle Ricerche, Italy
Charles R. Johnson	College of William and Mary, USA	Ram U. Verma	University of Toledo, USA
K. B. Kulasekera	Clemson University, USA	John C. Wierman	Johns Hopkins University, USA
Gerry Ladas	University of Rhode Island, USA	Michael E. Zieve	University of Michigan, USA

PRODUCTION

Silvio Levy, Scientific Editor

Cover: Alex Scorpan

See inside back cover or msp.org/involve for submission instructions. The subscription price for 2017 is US \$175/year for the electronic version, and \$235/year (+\$35, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues and changes of subscriber address should be sent to MSP.

Involve (ISSN 1944-4184 electronic, 1944-4176 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

Involve peer review and production are managed by EditFLOW® from Mathematical Sciences Publishers.

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

involve

2017

vol. 10

no. 4

New algorithms for modular inversion and representation by the form $x^2 + 3xy + y^2$	541
CHRISTINA DORAN, SHEN LU AND BARRY R. SMITH	
New approximations for the area of the Mandelbrot set	555
DANIEL BITTNER, LONG CHEONG, DANTE GATES AND HIEU D. NGUYEN	
Bases for the global Weyl modules of \mathfrak{sl}_n of highest weight $m\omega_1$	573
SAMUEL CHAMBERLIN AND AMANDA CROAN	
Leverage centrality of knight's graphs and Cartesian products of regular graphs and path powers	583
ROGER VARGAS, JR., ABIGAIL WALDRON, ANIKA SHARMA, RIGOBERTO FLÓREZ AND DARREN A. NARAYAN	
Equivalence classes of $GL(p, \mathbb{C}) \times GL(q, \mathbb{C})$ orbits in the flag variety of $gl(p + q, \mathbb{C})$	593
LETICIA BARCHINI AND NINA WILLIAMS	
Global sensitivity analysis in a mathematical model of the renal interstitium	625
MARIEL BEDELL, CLAIRE YILIN LIN, EMMIE ROMÁN-MELÉNDEZ AND IOANNIS SGOURALIS	
Sums of squares in quaternion rings	651
ANNA COOKE, SPENCER HAMBLÉN AND SAM WHITFIELD	
On the structure of symmetric spaces of semidihedral groups	665
JENNIFER SCHAEFER AND KATHRYN SCHLECHTWEG	
Spectrum of the Laplacian on graphs of radial functions	677
RODRIGO MATOS AND FABIO MONTENEGRO	
A generalization of Eulerian numbers via rook placements	691
ESTHER BANAÏAN, STEVE BUTLER, CHRISTOPHER COX, JEFFREY DAVIS, JACOB LANDGRAF AND SCARLITTE PONCE	
The H -linked degree-sum parameter for special graph families	707
LYDIA EAST KENNEY AND JEFFREY SCOTT POWELL	