

NISSUNA UMANA INVESTIGAZIONE SI PUO DIMANDARE VERA SCIENZA
S'ESSA NON PASSA PER LE MATEMATICHE DIMOSTRAZIONI
LEONARDO DA VINCI

vol. 1

no. 2

2013

MATHEMATICS AND MECHANICS
of
Complex Systems

FEDOR L. BAKHAREV, SERGEY A. NAZAROV AND GUIDO H. SWEERS

**A SUFFICIENT CONDITION FOR A DISCRETE SPECTRUM OF THE
KIRCHHOFF PLATE WITH AN INFINITE PEAK**



A SUFFICIENT CONDITION FOR A DISCRETE SPECTRUM OF THE KIRCHHOFF PLATE WITH AN INFINITE PEAK

FEDOR L. BAKHAREV, SERGEY A. NAZAROV AND GUIDO H. SWEERS

Sufficient conditions for a discrete spectrum of the biharmonic equation in a two-dimensional peak-shaped domain are established. Different boundary conditions from Kirchhoff's plate theory are imposed on the boundary and the results depend both on the type of boundary conditions and the sharpness exponent of the peak.

1. Motivation

Elliptic boundary value problems on domains which have a Lipschitz boundary and a compact closure, in particular when they generate positive self-adjoint operators, have fully discrete spectra. However, if the domain loses the Lipschitz property or compactness, other situations may occur. It is well-known that for the Dirichlet case boundedness is sufficient but not necessary for having discrete spectrum. See the famous paper [Rellich 1948] or the more recent [Rozenbljum 1972; van den Berg 1984]. On the other hand, for the Neumann problem of the Laplace operator there exist numerous examples of bounded domains such that the spectrum gets a nonempty continuous component (see e.g. [Courant and Hilbert 1953; Maz'ya and Poborchii 2006; 1997; Simon 1992; Hempel et al. 1991]).

The literature on the spectra for the Laplace operator with various boundary conditions on special domains is focused on domains that have a cusp, a finite or infinite peak or horn [van den Berg 1984; Hempel et al. 1991; Jakšić et al. 1992; Davies and Simon 1992; Jakšić 1993; Ivrii 1999; Boyarchenko and Levendorskii 2000; van den Berg and Lianantonakis 2001; Kovařík 2011] or even a rolled horn [Simon 1992].

The criteria in [Adams and Fournier 2003] and [Evans and Harris 1987] for the embedding $H^1(\Omega) \subset L_2(\Omega)$ to be compact show that the Neumann–Laplace

This work is supported by the Russian Foundation for Basic Research (project 12-01-00348). Bakharev is also supported by the Chebyshev Laboratory (Department of Mathematics and Mechanics, Saint Petersburg State University) under the grant 11.G34.31.0026 of the Government of the Russian Federation and grant 6.38.64.2012 of Saint Petersburg State University.

MSC2010: 35P05, 47A10, 74K20.

Keywords: Kirchhoff plate, cusp, peak, discrete and continuous spectra.

problem on a domain Ω with the infinite peak

$$\Pi_R = \{x = (x_1, x_2) \in \mathbb{R}^2 : x_1 > R, -H(x_1) < x_2 < H(x_1)\}, \quad (1)$$

where the function $H > 0$ is smooth and monotone decreasing, has discrete spectrum if and only if

$$\lim_{y \rightarrow +\infty} \int_y^{+\infty} \frac{H(\eta)}{H(y)} d\eta = 0 \quad \left(\iff \lim_{y \rightarrow +\infty} \frac{H(y + \epsilon)}{H(y)} = 0 \text{ for any } \epsilon > 0 \right). \quad (2)$$

The function H is assumed to have a first derivative that tends to zero and a bounded second derivative. It will be convenient to use the notation $\Upsilon(y) = (-H(y), H(y))$. Here is an image of such a domain:



The simplest boundary irregularity violating the Lipschitz condition is just the (finite) peak

$$\varpi_R = \{x : 0 < x_1 < R, -h(x_1) < x_2 < h(x_1)\}, \quad (3)$$

where $h(x_1) = h_0 x_1^{1+\alpha}$, $h_0 > 0$ and $\alpha > 0$. Nevertheless, the spectrum of the Neumann problem in the domain with this peak stays discrete. See [Remark 5.1](#).

A criterion ([\[Nazarov 2009\]](#)) for having essential spectrum in the Neumann problem for elliptic systems of second order differential equations with a polynomial property is derived in [\[Nazarov 1999\]](#). In particular it shows that the continuous spectrum of an elastic body with $\alpha \geq 1$ for the peak (3) is nonempty (see [\[Nazarov 2008; Bakharev and Nazarov 2009\]](#)). This phenomenon of generating wave processes in a finite volume, is known experimentally and used in the engineering practice to construct wave dampers, “black holes”, for elastic oscillations (see [\[Mironov 1988; Krylov and Tilman 2004\]](#), etc.).

In this paper we study the spectra of boundary value problems for the Kirchhoff model of a thin elastic plate described by the biharmonic operator Δ^2 . The boundary conditions that we consider model the three mechanically most reasonable cases, namely where the lateral sides of the peak are supplied with one of the following three types of the boundary conditions: clamped edge (*Dirichlet*), traction-free edge (*Neumann*) and hinged edge (*Mixed*). In all these cases the spectrum of the problem in a bounded domain with the peak as in (3) is discrete. We derive sufficient conditions for the spectrum to be discrete for the boundary value problem on an unbounded domain with a peak as in (1).

If a sufficient number of Dirichlet conditions are imposed on the lateral sides of the peak (the cases *D-N*, *M-M*, *D-M*, and *D-D*; see formulas (5)–(7) and

(11), (12)), then the proof that the spectrum is discrete becomes rather simple (Theorem 5). Indeed, it suffices to apply the weighted Friedrich's inequality (13) and to take into account the decay of the quantity $H(y)$ as $y \rightarrow +\infty$.

Our main interest concerns the cases $M-N$ and $N-N$. By applying weighted inequalities of Hardy type (Lemmas 7 and 8) we obtain a sufficient condition for the case $N-N$ to have discrete spectrum (Proposition 4). Indeed, as shown in [Adams and Fournier 2003], the second condition in (2) implies a criterion for the compact embedding $H^m(\Omega) \subset L_2(\Omega)$ for all m (we just need $m = 2$). One can also use that approach (Theorem 12) for the case $M-N$. This approach differs from the one used in [Adams and Fournier 2003; Evans and Harris 1987]. The different argument allows to obtain a condition for having discrete spectrum if one of the peak's edges is traction-free and the other one is hinged (the case $N-M$; see Theorem 13).

The obtained results essentially differ from each other: under the conditions (11) and also under (12) any decay of H is enough, the case $M-N$ needs a power decay rate with the exponent $\alpha > 1$, while the case $N-N$ needs a superexponential decay rate. See Remark 12.1 and 13.1.

2. The Kirchhoff plate model

Assumption 1. Let Ω be a domain in the plane \mathbb{R}^2 with a smooth (of class C^∞) boundary Γ such that, for some $R > 0$ and some monotone decreasing function $H : [R, \infty) \rightarrow \mathbb{R}^+$ with $\lim_{t \rightarrow \infty} H(t) = 0$,

- (1) $\{(x_1, x_2) \in \Omega; x_1 > R\} = \{(x_1, x_2); x_1 > R \text{ and } |x_2| < H(x_1)\}$ and
- (2) $\{(x_1, x_2) \in \Omega; x_1 < R\}$ is bounded.

We regard Ω as the projection of a thin isotropic homogeneous plate and apply the Kirchhoff theory (see [Mikhlin 1970, §30], [Nazarov 2002, Chapter 7], and so on). So we arrive at the fourth-order differential equation

$$\Delta^2 u(x) = \lambda u(x), \quad x \in \Omega, \quad (4)$$

which describes transverse oscillations of the plate. Here, $u(x)$ is the plate deflection, and λ a spectral parameter proportional to the square of the oscillation frequency.

The following sets of boundary conditions have a clear physical interpretation (see [Mikhlin 1970, §30], [Gazzola et al. 2010, §1.1], and so on):

(D): Dirichlet for a clamped edge:

$$u(x) = \partial_n u(x) = 0, \quad x \in \Gamma_D. \quad (5)$$

(N): Neumann for a traction-free edge:

$$\begin{cases} \partial_n \Delta u(x) - (1 - \nu)(\partial_s \kappa(x) \partial_s u(x) - \partial_s^2 \partial_n u(x)) = 0, \\ \Delta u(x) - (1 - \nu)(\partial_s^2 u(x) + \kappa(x) \partial_n u(x)) = 0, \end{cases} \quad x \in \Gamma_N. \quad (6)$$

(M): Mixed for a hinged edge:

$$u = \Delta u(x) - (1 - \nu)\kappa(x) \partial_n u(x) = 0, \quad x \in \Gamma_M. \quad (7)$$

Here, ∂_n and ∂_s stand for the normal and tangential derivatives, $\kappa(x)$ is the signed curvature of the contour Γ at the point $x \in \Gamma$ positive for convex boundary parts, and $\nu \in [0, 1/2)$ is the Poisson ratio. Finally, Γ_D , Γ_N , and Γ_M are the unions of finite families of open curves and $\Gamma = \overline{\Gamma_D} \cup \overline{\Gamma_N} \cup \overline{\Gamma_M}$, two of which may be empty. In what follows it is convenient to use the notation $y = x_1$ and $z = x_2$.

The general properties of the spectra depend on which of the boundary conditions (5)–(7) are imposed on the upper (+) and lower (–) sides,

$$\Sigma_R^\pm = \{x : y > R, z = \pm H(y)\},$$

of the peak. Let us give a precise statement. We define a symmetric bilinear form on $H^2(\Omega)$ by

$$a(u, u) = \int_{\Omega} \left(\left| \frac{\partial^2 u}{\partial x_1^2} \right|^2 + \left| \frac{\partial^2 u}{i} \partial x_2^2 \right|^2 + 2(1 - \nu) \left| \frac{\partial^2 u}{\partial x_1 \partial x_2} \right|^2 + 2\nu \frac{\partial^2 u}{\partial x_1^2} \frac{\partial^2 u}{\partial x_2^2} \right) dx \quad (8)$$

and $a(u, v) = \frac{1}{4}a(u + v, u + v) - \frac{1}{4}a(u - v, u - v)$. Then $\frac{1}{2}a(u, u)$ is the elastic energy stored in the plate. Since one directly verifies that

$$a(u, u) \geq (1 - \nu) \sum_{j,k=1}^2 \int_{\Omega} \left| \frac{\partial^2 u}{\partial x_j \partial x_k} \right|^2 dx, \quad (9)$$

the bilinear form $a(\cdot, \cdot)$ is nonnegative.

Definition 2. Let \mathbb{H} be the subspace of functions $u \in H^2(\Omega)$, satisfying the conditions (5) on Γ_D and $u = 0$ on Γ_M in the sense of traces.

By [Birman and Solomyak 1980, §10.1] \mathbb{H} is a Hilbert space with scalar product $a(\cdot, \cdot) + (\cdot, \cdot)_{\Omega}$, where $(\cdot, \cdot)_{\Omega}$ is the standard scalar product in the Lebesgue space $L_2(\Omega)$. Moreover, Theorem 2 of [Birman and Solomyak 1980, §10.1] implies that there exists a unique (unbounded) self-adjoint operator $A : D(A) \subset L_2(\Omega) \rightarrow L_2(\Omega)$ with $D(A) \subset D(A^{1/2}) = \mathbb{H}$ such that

$$(Au, v)_{\Omega} = a(u, v) \quad \text{for all } v \in \mathbb{H}. \quad (10)$$

Note that the eventually remaining boundary conditions in (5)–(7) appear in $D(A) \subset H^4(\Omega)$ as intrinsic natural boundary conditions from (9)–(8), see again [Mikhlin 1970, §30], [Gazzola et al. 2010, §1.1] etc.

Definition 3. By the spectrum for (4)–(7) we will mean $\sigma(A)$ with A defined in (10).

Since a is nonnegative, the spectrum $\sigma(A)$ belongs to $[0, \infty)$.

As a direct consequence of known results we may state the following.

Proposition 4. *If (2) holds, then the spectrum of the Equation (4), with either of the above boundary condition on the sides of the peak, is discrete.*

Proof. By [Birman and Solomyak 1980, §10.1 Theorem 5] the spectrum is discrete if and only if the embedding $\mathbb{H} \hookrightarrow L_2(\Omega)$ is compact. By [Adams and Fournier 2003; Evans and Harris 1987] one knows that $H^2(\Omega) \hookrightarrow L_2(\Omega)$ is compact whenever (2) holds true and $\mathbb{H} \subset H^2(\Omega)$. □

3. Simple cases: $D-N$, $M-M$, $D-M$ and $D-D$,

Theorem 5. *Suppose Ω is as in Assumption 1 and suppose that the boundary conditions for problem (4), as given in (5)–(7) contain one of the cases (11) or (12). Then the spectrum is discrete.*

Remark 5.1. By a similar reasoning, we may conclude that in the bounded domain ω , with the peak as in (3), Equation (4) has discrete spectrum for any set of conditions (5)–(7) on the arc $\partial\omega \setminus \mathbb{C}$. This fact follows from the inequality (see [Nazarov and Taskinen 2008]):

$$\| |x|^{-1} u; L_2(\omega) \|^2 \leq c (\| \nabla u; L_2(\omega) \|^2 + \| u; L_2(\omega \setminus \varpi_R) \|^2).$$

Proof. By assumption the boundary conditions provide at least one of the following two groups of relations:

$$u = 0 \quad \text{on } \Sigma_R^+ \cup \Sigma_R^-; \tag{11}$$

$$u = \partial_n u = 0 \quad \text{on } \Sigma_R^+ \text{ or on } \Sigma_R^-. \tag{12}$$

In both cases (11) and (12) the following version of Friedrich’s inequality is valid:

$$\int_{\Upsilon(y)} \left| \frac{\partial^2 u}{\partial z^2}(y, z) \right|^2 dz \geq \frac{c}{H(y)^4} \int_{\Upsilon(y)} |u(y, z)|^2 dz. \tag{13}$$

Therefore,

$$a(u, u) \geq c \int_{\Omega} H(y)^{-4} |u(x)|^2 dx. \tag{14}$$

The embedding operator $\gamma : \mathbb{H} \rightarrow L_2(\Omega)$ can be represented as the sum $\gamma_0 + \gamma_\rho$, where $\rho \geq R$ is large and positive, $\gamma_0 = \gamma - \gamma_\rho$, and γ_ρ contains the operator of multiplication by the characteristic function of Π_ρ . The operator γ_0 is compact, and the norm of γ_ρ , in view of (14), does not exceed $c \max \{H(x_1)^{-2}; x_1 \geq \rho\}$. Since the function H decays, this quantity goes to zero when $\rho \rightarrow +\infty$, i.e., the operator γ can be approximated by compact operators in the operator norm. Thus γ is compact and the result is proved. \square

4. Auxiliary inequalities

First of all we prove some one-dimensional weighted inequalities, two of which are of Hardy type involving a weight function h as follows.

Assumption 6. *Let h be a positive weight function of class C^2 on $[0, +\infty)$ such that*

- $\int_0^\infty h(s) ds < \infty$ and
- for some large T , we have $h'(t) < 0$ and $h''(t) > 0$ for $t \in (T, \infty)$.

Throughout this section h is supposed to satisfy this assumption.

Lemma 7. *If U is differentiable for $y \geq R$ and $U(R) = 0$, then*

$$\int_R^{+\infty} h(y) |U(y)|^2 dy \leq \int_R^{+\infty} F_h(y) |\partial_y U(y)|^2 dy,$$

where

$$F_h(y) = \frac{4}{h(y)} \left(\int_y^{+\infty} h(\tau) d\tau \right)^2.$$

Proof. Using the Cauchy–Bunyakovsky–Schwarz inequality, we have

$$\begin{aligned} & \int_R^{+\infty} h(y) |U(y)|^2 dy \\ &= 2 \int_R^{+\infty} h(y) \int_R^y \partial_t U(t) U(t) dt dy \\ &\leq 2 \int_R^{+\infty} \int_t^{+\infty} h(y) |\partial_t U(t) U(t)| dy dt \\ &\leq 2 \left(\int_R^{+\infty} h(t) |U(t)|^2 dt \right)^{1/2} \left(\int_R^{+\infty} h(t)^{-1} \left(\int_t^{+\infty} h(y) dy \right)^2 |\partial_t U(t)|^2 dt \right)^{1/2}, \end{aligned}$$

and the result follows through division by a common factor. \square

Lemma 8. *If U is differentiable for $y \geq R$ and $U(R) = 0$, then*

$$\int_R^{+\infty} h(y) |\partial_y U(y)|^2 dy \geq \int_R^{+\infty} G_h(y) |U(y)|^2 dy,$$

where

$$G_{h,R}(y) = \frac{1}{4h(y)} \left(\int_R^y h(\tau)^{-1} d\tau \right)^{-2}.$$

Proof. For functions v with $v(0) = 0$ the Hardy inequality tells us that

$$\int_0^{+\infty} t^{-2} |v(t)|^2 dt \leq 4 \int_0^{+\infty} |\partial_t v(t)|^2 dt.$$

We make the change $t \mapsto y \in [R, +\infty)$ where $t = \int_R^y h(\tau)^{-1} d\tau$, and set $U(y) = v(t)$. Then $\partial_t v(t) = h(y) \partial_y U(y)$ leads to the desired estimate. \square

Corollary 9. *If the function U is twice differentiable for $y \geq R$ and $U(R) = U'(R) = 0$, then*

$$\int_R^{+\infty} h(t) |\partial_t^2 U(t)|^2 dt \geq W_h(R) \int_R^{+\infty} h(t) |U(t)|^2 dt \quad (15)$$

is valid with

$$\begin{aligned} W_h(R) &:= \inf_{t \in [R, +\infty)} \frac{G_{h,R}(t)}{F_h(t)} \\ &= \inf_{t \in [R, +\infty)} \frac{1}{16} \left(\int_R^t h(\tau)^{-1} d\tau \right)^{-2} \left(\int_t^\infty h(\tau) d\tau \right)^{-2}. \end{aligned} \quad (16)$$

Lemma 10. *We have*

$$\inf_{t \in [R, +\infty)} \frac{G_{h^3,R}(t)}{h(t)h'(t)^2} \geq 1. \quad (17)$$

Suppose moreover that

$$\lim_{t \rightarrow +\infty} \partial_t (\log h(t)) = -\infty. \quad (18)$$

Then

$$W_h(R) \rightarrow +\infty \quad \text{and} \quad W_{h^3}(R) \rightarrow +\infty \quad \text{for} \quad R \rightarrow \infty.$$

Proof. Since $-h'(\tau) \geq -h'(t) > 0$ for $\tau < t$, we find

$$\begin{aligned} \frac{G_{h^3,R}(t)}{h(t)h'(t)^2} &= \frac{1}{4h'(t)^2 h(t)^4} \left(\int_R^t h(\tau)^{-3} d\tau \right)^{-2} \geq \frac{1}{4h(t)^4} \left(- \int_R^t h'(\tau) h(\tau)^{-3} d\tau \right)^{-2} \\ &= \frac{1}{4h(t)^4} \left(\frac{1}{2h(t)^2} - \frac{1}{2h(R)^2} \right)^{-2} \geq 1. \end{aligned}$$

Since (18) equals $h'(t)/h(t) \rightarrow -\infty$ for $t \rightarrow \infty$, we find that for $t \rightarrow \infty$ both

$$h(t) \left(\int_t^\infty h(\tau) d\tau \right)^{-1} \rightarrow \infty \quad \text{and} \quad \frac{1}{h(t)} \left(\int_R^t \frac{1}{h(\tau)} d\tau \right)^{-1} \rightarrow \infty.$$

Hence

$$\frac{G_{h,R}(t)}{F_h(t)} \rightarrow \infty \quad \text{as } t \rightarrow \infty,$$

and since the quotient also goes to infinity for $t \downarrow R$, it has a minimum in some $t_R \in (R, \infty)$. Calculating $(G_{h,R}(t)/F_h(t))' = 0$ we find

$$\frac{1}{h(t)} \int_t^\infty h(\tau) d\tau - h(t) \int_R^t h(\tau)^{-1} d\tau = 0.$$

Hence

$$\inf_{t \in [R, \infty)} \left(\int_R^t h(\tau)^{-1} d\tau \int_t^\infty h(\tau) d\tau \right)^{-1} = h(t_R)^2 \left(\int_{t_R}^\infty h(\tau) d\tau \right)^{-2},$$

which goes to infinity for $R \rightarrow \infty$ since $t_R > R$. The claim for $W_h(R)$ follows. The same argument holds true for $W_{h^3}(R)$. \square

5. Estimates for a traction-free boundary

We assume that the Neumann boundary conditions (6) are imposed at the both sides of the peak (1). Let us describe for $\rho \rightarrow +\infty$ the behavior of the multiplier $K(\rho)$ in the inequality

$$K(\rho) \int_{\Pi_\rho} |u(y, z)|^2 dy \leq \|u; H^2(\Omega)\|^2, \quad u \in H^2(\Omega). \quad (19)$$

If $K(\rho)$ increases unboundedly as $\rho \rightarrow +\infty$ then, as above, Theorem 10.1.5 of [Birman and Solomyak 1980] ensures that the spectrum of the equation (4)–(7) stays discrete even in the case both sides of the peak are supplied with the traction-free boundary conditions (N) and, moreover, for any other boundary conditions from the list (5)–(7).

Proposition 11. *Suppose that Assumption 6 is satisfied for $h = H$ and that*

$$\lim_{t \rightarrow \infty} \partial_t (\log H(t)) = -\infty.$$

Then for ρ sufficiently large (19) holds true with

$$K(\rho) = c \min\{H^{-4}(\rho), W_H(\rho), W_{H^3}(\rho)\}. \quad (20)$$

Proof. It is sufficient to check the inequality (19) for smooth functions which vanish for $y < \rho$. We use the representation

$$u(x) = u(y, z) = u_0(y) + zu_1(y) + u^\perp(y, z)$$

where, for $y > \rho$, the component u^\perp is subject to the orthogonality conditions

$$\begin{aligned} \int_{\Upsilon(y)} u^\perp(y, z) dz &= 0, \\ \int_{\Upsilon(y)} \partial_z u^\perp(y, z) dz &= u^\perp(y, H(y)) - u^\perp(y, -H(y)) = 0. \end{aligned} \tag{21}$$

Let us process the integrals on the right-hand side of

$$\int_{\Pi_\rho} |\nabla_x^2 u(x)|^2 dx = I_1 + 4I_2 + I_3, \tag{22}$$

where

$$I_1 := \int_{\Pi_\rho} |\partial_z^2 u(x)|^2 dx, \quad I_2 := \int_{\Pi_\rho} |\partial_y \partial_z u(x)|^2 dx, \quad I_3 := \int_{\Pi_\rho} |\partial_y^2 u(x)|^2 dx.$$

Since $I_1 = \int_\rho^{+\infty} \int_{\Upsilon(y)} |\partial_z^2 u^\perp(y, z)|^2 dz dy$ and since, by the orthogonality conditions in (21), inequality (13) holds here also, we find that

$$I_1 \geq c \int_{\Pi_\rho} H(y)^{-4} |u^\perp(x)|^2 dx. \tag{23}$$

For the last term in (22) we have

$$I_3 = \int_{\Pi_\rho} |\partial_y^2 u_0(y) + z\partial_y^2 u_1(y) + \partial_y^2 u^\perp(y, z)|^2 dx \geq J_1 + J_2 + 2J_3 + 2J_4, \tag{24}$$

where

$$\begin{aligned} J_1 &= g \int_{\Pi_\rho} |\partial_y^2 u_0(y)|^2 dx, & J_3 &= g \int_{\Pi_\rho} \partial_y^2 u_0(y) \partial_y^2 u^\perp(y, z) dx, \\ J_2 &= g \int_{\Pi_\rho} |z\partial_y^2 u_1(y)|^2 dx, & J_4 &= g \int_{\Pi_\rho} z\partial_y^2 u_1(y) \partial_y^2 u^\perp(y, z) dx. \end{aligned}$$

We readily notice that according to the inequality (15) the estimates

$$J_1 \geq W_H(\rho) \int_\rho^{+\infty} 2H(y) |u_0(y)|^2 dy = W_H(\rho) \int_{\Pi_\rho} |u_0(y)|^2 dx \tag{25}$$

and

$$\begin{aligned} J_2 &= \frac{2}{3} \int_{\rho}^{+\infty} H^3(y) |\partial_y^2 u_1(y)|^2 dy \geq \frac{2}{3} W_{H^3}(\rho) \int_{\rho}^{+\infty} H^3(y) |u_1(y)|^2 dy \\ &= W_{H^3}(\rho) \int_{\Pi_{\rho}} |zu_1(y)|^2 dx \end{aligned} \quad (26)$$

are fulfilled. For our purpose we need $W_H(\rho) \rightarrow +\infty$ and $W_{H^3}(\rho) \rightarrow +\infty$ for $\rho \rightarrow \infty$ and this we will assume.

Besides, by the Cauchy–Bunyakovsky–Schwarz inequality, we have

$$|J_3| \leq J_1^{1/2} \left(\int_{\rho}^{+\infty} \frac{1}{2H(y)} \left(\int_{\Upsilon(y)} \partial_y^2 u^{\pm}(y, z) dz \right)^2 dy \right)^{1/2}.$$

We now deal with the inner integral in z in the last expression. To this end, we take into account the orthogonality conditions (21) and the trace inequality. We differentiate the first equality in (21) twice with respect to y and obtain

$$\begin{aligned} \sum_{\pm} (2\partial_y u^{\pm}(y, \pm H(y)) \partial_y H(y) + u^{\pm}(y, \pm H(y)) \partial_y^2 H(y) \pm \partial_z u^{\pm}(y, \pm H(y)) (\partial_y H(y))^2) \\ + \int_{\Upsilon(y)} \partial_y^2 u^{\pm}(y, z) dz = 0. \end{aligned}$$

Thus,

$$\begin{aligned} \left(\int_{\Upsilon(y)} \partial_y^2 u^{\pm}(y, z) dz \right)^2 \\ \leq c \sum_{\pm} (|\partial_z u^{\pm}(y, \pm H(y))|^2 |\partial_y H(y)|^4 + |u^{\pm}(y, \pm H(y))|^2 |\partial_y^2 H(y)|^2 \\ + |\partial_y u^{\pm}(y, \pm H(y))|^2 |\partial_y H(y)|^2). \end{aligned}$$

For the first two terms between the brackets we use the trace inequality

$$|\partial_z u^{\pm}(y, \pm H(y))|^2 |H(y)|^2 + |u^{\pm}(y, \pm H(y))|^2 \leq c |H(y)|^3 \int_{\Upsilon(y)} |\partial_z^2 u^{\pm}(y, z)|^2 dz.$$

For the third term, we write down the chain of inequalities

$$\begin{aligned} |\partial_y u^{\pm}(y, \pm H(y))|^2 \\ \leq c H(y) \int_{\Upsilon(y)} |\partial_y \partial_z u^{\pm}(y, z)|^2 dz + c |H(y)|^{-2} \left(\int_{\Upsilon(y)} \partial_y u^{\pm}(y, z) dz \right)^2 \\ \leq c H(y) \int_{\Upsilon(y)} |\partial_y \partial_z u^{\pm}(y, z)|^2 dz \\ + 2c \left(\frac{\partial_y H(y)}{H(y)} \right)^2 (|u^{\pm}(y, H(y))|^2 + |u^{\pm}(y, -H(y))|^2). \end{aligned}$$

As a result, we find that

$$\left| \int_{\Upsilon(y)} \partial_y^2 u^\perp(y, z) dz \right|^2 \leq c(|\partial_y H(y)|^4 H(y) + |\partial_y^2 H(y)|^2 |H(y)|^3) \int_{\Upsilon(y)} |\partial_z^2 u^\perp(y, z)|^2 dz + c|\partial_y H(y)|^2 |H(y)| \int_{\Upsilon(y)} |\partial_y \partial_z u^\perp(y, z)|^2 dz.$$

The final inequality for the integral J_3 takes the form

$$|J_3| \leq c_1(\rho) J_1^{1/2} I_1^{1/2} + c_2(\rho) J_1^{1/2} K_1^{1/2} \tag{27}$$

where $K_1 = \|\partial_{yz}^2 u^\perp; L_2(\Pi_\rho)\|^2$ and

$$c_1(\rho) = c \sup_{y \in [\rho, +\infty)} (|\partial_y H(y)|^2 + |\partial_y^2 H(y)| |H(y)|),$$

$$c_2(\rho) = c \sup_{y \in [\rho, +\infty)} |\partial_y H(y)|.$$

Both suprema tend to 0 for $\rho \rightarrow +\infty$. A similar argument shows that

$$|J_4| \leq c_1(\rho) J_2^{1/2} I_1^{1/2} + c_2(\rho) J_2^{1/2} K_1^{1/2}. \tag{28}$$

It remains to process the second term in (22), that is,

$$I_2 = \int_{\Pi_\rho} |z \partial_y u_1(y) + \partial_y \partial_z u^\perp(y, z)|^2 dx$$

$$= \int_{\Pi_\rho} |\partial_y \partial_z u^\perp(y, z)|^2 dx + \int_{\Pi_\rho} |\partial_y u_1(y)|^2 dx + 2 \int_{\Pi_\rho} \partial_y u_1(y) \partial_y \partial_z u^\perp(y, z) dx.$$

$$\qquad \qquad \qquad \cong K_1 \qquad \qquad \qquad \cong K_2 \qquad \qquad \qquad \cong 2K_3$$

So it follows that

$$K_1 = I_2 - K_2 - 2K_3 \leq I_2 + 2|K_3|. \tag{29}$$

We continue by estimating the integral K_3 :

$$|K_3| = \left| \int_\rho^{+\infty} \partial_y u_1(y) \int_{\Upsilon(y)} \partial_y \partial_z u^\perp(y, z) dz dy \right|$$

$$\leq \left(\int_\rho^{+\infty} G_{H^3, \rho}(y) |\partial_y u_1(y)|^2 dy \right)^{\frac{1}{2}} \left(\int_\rho^{+\infty} G_{H^3, \rho}(y)^{-1} \left| \int_{\Upsilon(y)} \partial_y \partial_z u^\perp(y, z) dz \right|^2 dy \right)^{\frac{1}{2}}$$

$$\leq c J_2^{1/2} \left(\int_\rho^{+\infty} G_{H^3, \rho}(y)^{-1} \left| \int_{\Upsilon(y)} \partial_y \partial_z u^\perp(y, z) dz \right|^2 dy \right)^{1/2}.$$

Differentiating the second formula (21) with respect to y yields

$$\int_{\Upsilon(y)} \partial_y \partial_z u^\pm(y, z) dz + \sum_{\pm} \partial_z u^\pm(y, \pm H(y)) \partial_y H(y) = 0.$$

By the trace inequality we find that

$$\begin{aligned} \left| \int_{\Upsilon(y)} \partial_y \partial_z u^\pm(y, z) dz \right|^2 &= \left(\sum_{\pm} \partial_z u^\pm(y, \pm H(y)) \right)^2 |\partial_y H(y)|^2 \\ &\leq c |H(y)| |\partial_y H(y)|^2 \int_{\Upsilon(y)} |\partial_z^2 u^\pm(y, z)|^2 dz. \end{aligned}$$

Thus, from the relation (18), which implies (17), we get

$$|K_3| \leq c \sup_{y \in [\rho, +\infty)} \{ |G_{H^3, \rho}(y)|^{-1/2} |\partial_y H(y)| |H(y)|^{1/2} \} J_2^{1/2} I_1^{1/2} \leq c J_2^{1/2} I_1^{1/2}. \quad (30)$$

We find by combining (29) and (30) that

$$K_1 \leq I_2 + c J_2^{1/2} I_1^{1/2}$$

and so (27) and (28) yield, respectively,

$$|J_3| \leq c_1(\rho) J_1^{1/2} I_1^{1/2} + c_2(\rho) J_1^{1/2} (I_2 + c J_2^{1/2} I_1^{1/2})^{1/2}, \quad (31)$$

$$|J_4| \leq c_1(\rho) J_2^{1/2} I_1^{1/2} + c_2(\rho) J_2^{1/2} (I_2 + c J_2^{1/2} I_1^{1/2})^{1/2}. \quad (32)$$

Using first (22) and (24), next (31) and (32) for ρ large enough, and finally (23), (25) and (26) we conclude that indeed

$$\begin{aligned} \|\nabla_x^2 u; L_2(\Pi_\rho)\|^2 &= I_1 + 4I_2 + I_3 \\ &\geq I_1 + 4I_2 + J_1 + J_2 + 2J_3 + 2J_4 \\ &\geq \frac{1}{2}(I_1 + 4I_2 + J_1 + J_2) \geq \frac{1}{2}(I_1 + J_1 + J_2) \\ &\geq c \min\{H^{-4}(\rho), W_H(\rho), W_{H^3}(\rho)\} \|u; L_2(\Pi_\rho)\|^2, \end{aligned}$$

whenever ρ is large enough. □

6. Traction-free boundaries: N - N

Theorem 12. *Suppose that H satisfies Assumption 6 with $h = H$ and that*

$$\lim_{t \rightarrow \infty} \partial_t (\log H(t)) = -\infty. \quad (33)$$

Then the embedding $H^2(\Omega) \hookrightarrow L_2(\Omega)$ is compact and the spectrum of the problem (4) with the Neumann boundary conditions (6) on both sides of the peak is discrete.

Proof. By [Proposition 11](#), $K(\rho)$ can be estimated as in [\(20\)](#). [Assumption 6](#) implies that [Lemma 10](#) holds true and hence $K(\rho) \rightarrow +\infty$ for $\rho \rightarrow +\infty$. One concludes as in the proof of [Theorem 5](#) through an approximation by compact operators. \square

Remark 12.1. Note that the functions $H(y) = y^{-\alpha}$ and $H(y) = \exp(-\alpha y)$, $\alpha > 0$, do not satisfy the requirement in [\(2\)](#) or [\(33\)](#). The functions $H(y) = \exp(-y^{1+\alpha})$ with $\alpha > 0$ however do.

7. An incomplete Dirichlet condition: $M-N$

In this section the boundary conditions only contain a single stable condition

$$u(x) = 0, \quad x \in \Sigma_\rho^+ \quad (\text{or } x \in \Sigma_\rho^-). \quad (34)$$

Theorem 13. *Suppose that $\lim_{y \rightarrow \infty} H(y)^{-3} G_H(y) = +\infty$. Then the problem in [\(4\)](#) with the boundary condition as in [\(34\)](#) has discrete spectrum.*

Remark 13.1. The functions $H(y) = y^{-1-\alpha}$ with $\alpha > 0$ meet the condition in [Theorem 13](#).

Proof. By [\(34\)](#), Friedrich's inequality holds on the section $\Upsilon(y)$ and, consequently,

$$\|\mathcal{A} \partial_z u; L_2(\Pi_\rho)\|^2 \geq c \|\mathcal{A} H^{-2} u; L_2(\Pi_\rho)\|^2$$

for every positive weight function $y \mapsto \mathcal{A}(y)$. The function $v = \partial_z u$ can be represented as the sum $v(x) = v_0(y) + v^\perp(x)$ where, for $y > \rho$, the component v^\perp satisfies the first condition in [\(21\)](#). Therefore,

$$\begin{aligned} & \int_{\Pi_\rho} |\nabla_x^2 u(x)|^2 dx \\ & \geq \int_{\Pi_\rho} |\nabla_x v(x)|^2 dx \\ & \geq \int_\rho^{+\infty} 2H(y) |\partial_y v_0(y)|^2 dy + \int_{\Pi_\rho} |\partial_z v^\perp(y, z)|^2 dx + 2 \int_{\Pi_\rho} \partial_y v_0(y) \partial_y v^\perp(y, z) dx \\ & =: I_4 + I_5 + 2I_6. \end{aligned}$$

Setting $Z_H(y) = H(y)^{-1} G_H(y)$, we get

$$\begin{aligned} I_4 & \geq \int_\rho^{+\infty} 2G_H(y) |v_0(y)|^2 dy \geq \int_\rho^{+\infty} 2Z_H(y) H(y) |v_0(y)|^2 dy \\ & = \|Z_H v_0; L_2(\Pi_\rho)\|^2. \end{aligned}$$

Friedrich's inequality implies

$$I_5 = \|\partial_z v^\perp; L_2(\Pi_\rho)\| \geq c \|H^{-2} v^\perp; L_2(\Pi_\rho)\|^2.$$

Furthermore,

$$\begin{aligned} I_6 &= \int_{\Pi_\rho} \partial_y v_0(y) \partial_y v^\perp(y, z) dx \leq I_5^{1/2} \left(\int_\rho^{+\infty} \frac{1}{2H(y)} \left| \int_{\Upsilon(y)} \partial_y v^\perp(y, z) dz \right|^2 \right)^{1/2} \\ &\leq I_5^{1/2} \left(\int_\rho^{+\infty} \frac{1}{2H(y)} |v^\perp(y, H(y)) \partial_y H(y) - v^\perp(y, -H(y)) \partial_y H(y)|^2 \right)^{1/2}. \end{aligned}$$

Thus $I_6 \leq c |\partial_y H(\rho)| I_5^{1/2} I_6^{1/2}$ holds and hence

$$\|\nabla_x^2 u; L_2(\Pi_\rho)\|^2 \geq c \|\min\{H^{-4}; H^{-3} G_H\} u; L_2(\Pi_\rho)\|^2.$$

Compactness and hence the discrete spectrum follow from the assumption on H . □

References

- [Adams and Fournier 2003] R. A. Adams and J. J. F. Fournier, *Sobolev spaces*, 2nd ed., Pure and Applied Mathematics **140**, Elsevier/Academic Press, Amsterdam, 2003.
- [Bakharev and Nazarov 2009] F. L. Bakharev and S. A. Nazarov, “On the structure of the spectrum of a problem in the theory of elasticity for a body with a supersharp peak”, *Sibirsk. Mat. Zh.* **50:4** (2009), 746–756. In Russian; translated in *Sib. Math. J.* **50:4** (2009), 587–595.
- [van den Berg 1984] M. van den Berg, “On the spectrum of the Dirichlet Laplacian for horn-shaped regions in \mathbf{R}^n with infinite volume”, *J. Funct. Anal.* **58:2** (1984), 150–156.
- [van den Berg and Lianantonakis 2001] M. van den Berg and M. Lianantonakis, “Asymptotics for the spectrum of the Dirichlet Laplacian on horn-shaped regions”, *Indiana Univ. Math. J.* **50:1** (2001), 299–333.
- [Birman and Solomyak 1980] M. S. Birman and M. Z. Solomyak, *Спектральная теория самосопряженных операторов в гильбертовом пространстве*, Leningrad Univ., 1980. Translated as *Spectral theory of selfadjoint operators in Hilbert space*, Reidel, Dordrecht, 1987.
- [Boyarchenko and Levendorskii 2000] S. I. Boyarchenko and S. Z. Levendorskii, “Spectral asymptotics of the Neumann Laplacian on oscillating horns”, *Russ. J. Math. Phys.* **7:1** (2000), 1–14.
- [Courant and Hilbert 1953] R. Courant and D. Hilbert, *Methods of mathematical physics*, vol. 1, Interscience, New York, 1953.
- [Davies and Simon 1992] E. B. Davies and B. Simon, “Spectral properties of Neumann Laplacian of horns”, *Geom. Funct. Anal.* **2:1** (1992), 105–117.
- [Evans and Harris 1987] W. D. Evans and D. J. Harris, “Sobolev embeddings for generalized ridged domains”, *Proc. London Math. Soc.* (3) **54:1** (1987), 141–175.
- [Gazzola et al. 2010] F. Gazzola, H.-C. Grunau, and G. Sweers, *Polyharmonic boundary value problems: Positivity preserving and nonlinear higher order elliptic equations in bounded domains*, Lecture Notes in Mathematics **1991**, Springer, Berlin, 2010.
- [Hempel et al. 1991] R. Hempel, L. A. Seco, and B. Simon, “The essential spectrum of Neumann Laplacians on some bounded singular domains”, *J. Funct. Anal.* **102:2** (1991), 448–483.
- [Ivrii 1999] V. Ivrii, “Precise spectral asymptotics for Neumann Laplacian in domains with cusps”, *Appl. Anal.* **71:1-4** (1999), 139–147.
- [Jakšić 1993] V. Jakšić, “On the spectrum of the Neumann Laplacian of long-range horns: a note on the Davies–Simon theorem”, *Proc. Amer. Math. Soc.* **119:2** (1993), 663–669.

- [Jakšić et al. 1992] V. Jakšić, S. Molčanov, and B. Simon, “Eigenvalue asymptotics of the Neumann Laplacian of regions and manifolds with cusps”, *J. Funct. Anal.* **106**:1 (1992), 59–79.
- [Kovařík 2011] H. Kovařík, “Eigenvalue asymptotic of Robin Laplace operators on two-dimensional domains with cusps”, *J. Lond. Math. Soc.* (2) **83**:1 (2011), 256–271.
- [Krylov and Tilman 2004] V. V. Krylov and F. J. B. S. Tilman, “Acoustic ‘black holes’ for flexural waves as effective vibration dampers”, *J. Sound Vib.* **274** (2004), 605–619.
- [Maz’ya and Poborchii 1997] V. G. Maz’ya and S. V. Poborchii, *Differentiable functions on bad domains*, World Scientific, River Edge, NJ, 1997.
- [Maz’ya and Poborchii 2006] V. G. Maz’ya and S. V. Poborchii, “Theorems for embedding Sobolev spaces on domains with a peak and on Hölder domains”, *Algebra i Analiz* **18**:4 (2006), 95–126.
- [Mikhlin 1970] S. G. Mikhlin, Вариационные методы в математической физике, 2nd ed., Nauka, Moscow, 1970. First edition translated as *Variational methods in mathematical physics*, Pergamon Press, Oxford, 1964.
- [Mironov 1988] M. A. Mironov, “Propagation of a flexural wave in a plate whose thickness decreases smoothly to zero in a finite interval”, *Soviet Phys. Acoust.* **34** (1988), 318–319.
- [Nazarov 1999] S. A. Nazarov, “Polynomial property of selfadjoint elliptic boundary value problems, and the algebraic description of their attributes”, *Uspekhi Mat. Nauk* **54**:5(329) (1999), 77–142. In Russian; translated in *Russian Math. Surveys* **54**:5 (1999), 947–1014.
- [Nazarov 2002] S. A. Nazarov, *Asymptotic theory of thin plates and rods*, vol. 1: *Dimension reduction and integral estimates*, Nauchnaya Kniga, Novosibirsk, 2002.
- [Nazarov 2008] S. A. Nazarov, “On the spectrum of a problem in the theory of elasticity for a spiked body”, *Sibirsk. Mat. Zh.* **49**:5 (2008), 1105–1127.
- [Nazarov 2009] S. A. Nazarov, “On the essential spectrum of boundary value problems for systems of differential equations in a bounded domain with a cusp”, *Funktional. Anal. i Prilozhen.* **43**:1 (2009), 55–67. In Russian; translated in *Funct. Anal. Appl.* **43**:1 (2009), 44–54.
- [Nazarov and Taskinen 2008] S. A. Nazarov and J. Taskinen, “On the spectrum of the Steklov problem in a domain with a peak”, *Vestnik Sankt-Pet. Univ. Ser. I Mat. Mekh. Astr.* **2008**:1 (2008), 56–65. In Russian; translated in *Vestnik St. Petersburg Univ. Math.* **41**:1 (2008), 45–52.
- [Rellich 1948] F. Rellich, “Das Eigenwertproblem von $\Delta u + \lambda u = 0$ in Halbröhren”, pp. 329–344 in *Studies and essays presented to R. Courant on his 60th birthday*, Interscience, New York, 1948.
- [Rozenbljum 1972] G. V. Rozenbljum, “The eigenvalues of the first boundary value problem in unbounded domains”, *Mat. Sb. (N.S.)* **89**:(131) (1972), 234–247. In Russian; translated in *Math. USSR Sb.* **18** (1972), 235–248 (1973).
- [Simon 1992] B. Simon, “The Neumann Laplacian of a jelly roll”, *Proc. Amer. Math. Soc.* **114**:3 (1992), 783–785.

Received 7 Mar 2012. Revised 21 Jul 2012. Accepted 28 Aug 2012.

FEDOR L. BAKHAREV: fbakharev@yandex.ru

Mathematics and Mechanics Faculty, St. Petersburg State University, 198904 St. Petersburg, Russia

SERGEY A. NAZAROV: sergnazarov@yahoo.co.uk

Institute of Problems of Mechanical Engineering, Russian Academy of Sciences, 199178 St. Petersburg, Russia

Current address: Mathematics and Mechanics Faculty, St. Petersburg State University, Universitetsky pr., 28, 198504, Stary Peterhof, Russia

GUIDO H. SWEERS: gsweers@math.uni-koeln.de

Mathematisches Institut, Universität zu Köln, D-50931 Cologne, Germany



MATHEMATICS AND MECHANICS OF COMPLEX SYSTEMS

msp.org/memocs

EDITORIAL BOARD

ANTONIO CARCATERRA	Università di Roma "La Sapienza", Italia
ERIC A. CARLEN	Rutgers University, USA
FRANCESCO DELL'ISOLA	(CO-CHAIR) Università di Roma "La Sapienza", Italia
RAFFAELE ESPOSITO	(TREASURER) Università dell'Aquila, Italia
ALBERT FANNJIANG	University of California at Davis, USA
GILLES A. FRANCFORT	(CO-CHAIR) Université Paris-Nord, France
PIERANGELO MARCATI	Università dell'Aquila, Italy
JEAN-JACQUES MARIGO	École Polytechnique, France
PETER A. MARKOWICH	DAMTP Cambridge, UK, and University of Vienna, Austria
MARTIN OSTOJA-STARZEWSKI	(CHAIR MANAGING EDITOR) Univ. of Illinois at Urbana-Champaign, USA
PIERRE SEPPECHER	Université du Sud Toulon-Var, France
DAVID J. STEIGMANN	University of California at Berkeley, USA
PAUL STEINMANN	Universität Erlangen-Nürnberg, Germany
PIERRE M. SUQUET	LMA CNRS Marseille, France

MANAGING EDITORS

MICOL AMAR	Università di Roma "La Sapienza", Italia
CORRADO LATTANZIO	Università dell'Aquila, Italy
ANGELA MADEO	Université de Lyon-INSA (Institut National des Sciences Appliquées), France
MARTIN OSTOJA-STARZEWSKI	(CHAIR MANAGING EDITOR) Univ. of Illinois at Urbana-Champaign, USA

ADVISORY BOARD

ADNAN AKAY	Carnegie Mellon University, USA, and Bilkent University, Turkey
HOLM ALTENBACH	Otto-von-Guericke-Universität Magdeburg, Germany
MICOL AMAR	Università di Roma "La Sapienza", Italia
HARM ASKES	University of Sheffield, UK
TEODOR ATANACKOVIĆ	University of Novi Sad, Serbia
VICTOR BERDICHEVSKY	Wayne State University, USA
GUY BOUCHITTÉ	Université du Sud Toulon-Var, France
ANDREA BRAIDES	Università di Roma Tor Vergata, Italia
ROBERTO CAMASSA	University of North Carolina at Chapel Hill, USA
MAURO CARFORE	Università di Pavia, Italia
ERIC DARVE	Stanford University, USA
FELIX DARVE	Institut Polytechnique de Grenoble, France
ANNA DE MASI	Università dell'Aquila, Italia
GIANPIETRO DEL PIERO	Università di Ferrara and International Research Center MEMOCS, Italia
EMMANUELE DI BENEDETTO	Vanderbilt University, USA
BERNOLD FIEDLER	Freie Universität Berlin, Germany
IRENE M. GAMBA	University of Texas at Austin, USA
SERGEY GAVRILYUK	Université Aix-Marseille, France
TIMOTHY J. HEALEY	Cornell University, USA
DOMINIQUE JEULIN	École des Mines, France
ROGER E. KHAYAT	University of Western Ontario, Canada
CORRADO LATTANZIO	Università dell'Aquila, Italy
ROBERT P. LIPTON	Louisiana State University, USA
ANGELO LUONGO	Università dell'Aquila, Italia
ANGELA MADEO	Université de Lyon-INSA (Institut National des Sciences Appliquées), France
JUAN J. MANFREDI	University of Pittsburgh, USA
CARLO MARCHIORO	Università di Roma "La Sapienza", Italia
GÉRARD A. MAUGIN	Université Paris VI, France
ROBERTO NATALINI	Istituto per le Applicazioni del Calcolo "M. Picone", Italy
PATRIZIO NEFF	Universität Duisburg-Essen, Germany
ANDREY PIATNITSKI	Narvik University College, Norway, Russia
ERRICO PRESUTTI	Università di Roma Tor Vergata, Italy
MARIO PULVIRENTI	Università di Roma "La Sapienza", Italia
LUCIO RUSSO	Università di Roma "Tor Vergata", Italia
MIGUEL A. F. SANJUAN	Universidad Rey Juan Carlos, Madrid, Spain
PATRICK SELVADURAI	McGill University, Canada
ALEXANDER P. SEYRANIAN	Moscow State Lomonosov University, Russia
MIROSLAV ŠILHAVÝ	Academy of Sciences of the Czech Republic
GUIDO SWEERS	Universität zu Köln, Germany
ANTOINETTE TORDESILLAS	University of Melbourne, Australia
LEV TRUSKINOVSKY	École Polytechnique, France
JUAN J. L. VELÁZQUEZ	Bonn University, Germany
VINCENZO VESPRI	Università di Firenze, Italia
ANGELO VULPIANI	Università di Roma La Sapienza, Italia

MEMOCS (ISSN 2325-3444 electronic, 2326-7186 printed) is a journal of the International Research Center for the Mathematics and Mechanics of Complex Systems at the Università dell'Aquila, Italy.

Cover image: "Tangle" by © John Horigan; produced using the *Context Free* program (contextfreeart.org).

PUBLISHED BY

 **mathematical sciences publishers**
nonprofit scientific publishing
<http://msp.org/>

© 2013 Mathematical Sciences Publishers

Self-organized stochastic tipping in slow-fast dynamical systems	129
Mathias Linkerhand and Claudius Gros	
Well-posedness for dislocation-based gradient viscoplasticity, II: general nonassociative monotone plastic flows	149
Sergiy Nesenenko and Patrizio Neff	
Symmetry classes for even-order tensors	177
Marc Olive and Nicolas Auffray	
A direct approach to nonlinear shells with application to surface-substrate interactions	211
Miroslav Šilhavý	
A sufficient condition for a discrete spectrum of the Kirchhoff plate with an infinite peak	233
Fedor L. Bakharev, Sergey A. Nazarov and Guido H. Sweets	

MEMOCS is a journal of the International Research Center for the Mathematics and Mechanics of Complex Systems at the Università dell'Aquila, Italy.

