

Pacific Journal of Mathematics

A CHARACTERIZATION OF THE SUBGROUPS OF THE ADDITIVE RATIONALS

ROSS A. BEAUMONT AND HERBERT S. ZUCKERMAN

A CHARACTERIZATION OF THE SUBGROUPS OF THE ADDITIVE RATIONALS

ROSS A. BEAUMONT AND H. S. ZUCKERMAN

1. Introduction. In the class of abelian groups every element of which (except the identity) has infinite order, the subgroups of the additive group of rational numbers have the simplest structure. These rational groups are the groups of rank one, or generalized cyclic groups, an abelian group G being said to have rank one if for any pair of elements, $a \neq 0, b \neq 0$, in G , there exist integers m, n , such that $ma = nb \neq 0$. Although many of the properties of these groups are known [1], it seems worthwhile to give a simple characterization from which their properties can easily be derived. This characterization is given in Theorems 1 and 2 of §2, and the properties of the rational groups are obtained as corollaries of these theorems in §3. In §4, all rings which have a rational group as additive group are determined.

Let $p_1, p_2, \dots, p_j, \dots$ be an enumeration of the primes in their natural order; and associate with each p_j an exponent k_j , where k_j is a nonnegative integer or the symbol ∞ . We consider sequences $i; k_1, k_2, \dots, k_j, \dots$, where i is any positive integer such that $(i, p_j) = 1$ if $k_j > 0$, and define $(i; k_1, k_2, \dots, k_j, \dots) = (i; k_j)$ to be the set of all rational numbers of the form ai/b , where a is any integer and b is an integer such that $b = \prod'_{p_j} p_j^{n_j}$ with $n_j \leq k_j$. Then each sequence determines a well-defined set of rational numbers. The symbol \prod' designates a product over an arbitrary subset of the primes that satisfy whatever conditions are put on them; \prod designates a product over all primes that satisfy the given conditions.

2. Characterization of the rational groups. We show that the nontrivial subgroups of R are exactly the subsets $(i; k_j)$ defined in the introduction.

THEOREM 1. *The set $(i; k_j)$ is a subgroup of R^+ , the additive group of rational numbers. We have $(i; k_j) = (i'; k'_j)$ if and only if $i = i', k_j = k'_j$ for all j .*

Received October 11, 1950. Presented to the American Mathematical Society, June 17, 1950.

Pacific J. Math. 1 (1951), 169-177.

Proof. If $ai/b \in (i; k_j)$, $ci/d \in (i; k_j)$, then $b = \prod_{p_j} p_j^{n_j}$, $d = \prod_{p_j} p_j^{m_j}$, and $[b, d] = \prod_{p_j} p_j^{s_j}$, where $s_j = \max(n_j, m_j) \leq k_j$. Writing $[b, d] = bb' = dd'$, we have

$$\frac{ai}{b} - \frac{ci}{d} = \frac{b'ai}{[b,d]} - \frac{d'ci}{[b,d]} = \frac{(b'a - d'c)i}{[b,d]} \in (i; k_j).$$

It is clear that different sequences determine different subgroups.

In the sequel we need the following properties of a subgroup $G \neq 0$ of R^+ .

(1) Every $\zeta \in G$ has the form $\zeta = ai/b$, $(ai, b) = 1$, where i is the least positive integer in G .

For every ζ we have $\zeta = m/b$, where $(m, b) = 1$; and if i is the least positive integer in G , then $m = ai + r$ and $m - ai \in G$ imply $r = 0$.

(2) If $ai/b \in G$, $i \in G$, and $(a, b) = 1$, then $i/b \in G$.

For there exist integers k, l such that $ka + lb = 1$ and

$$\frac{i}{b} = \frac{(ka + lb)i}{b} = \frac{kai}{b} + li \in G.$$

(3) If $ai/b \in G$ where i is the least positive integer in G , and $(a, b) = 1$, then $(i, b) = 1$.

By (2), $i/b \in G$; and if $(i, b) \neq 1$, $h/b' \in G$ with $h < i$. Then $b'(h/b') = h \in G$.

We assume in the proof of the remaining properties that the elements of G are written in the canonical form ai/b with $(ai, b) = 1$ and i the least positive integer in G .

(4) If $ai/bc \in G$, then $i/b \in G$.

For $cai/bc = ai/b \in G$ and $i/b \in G$ by (2).

(5) If $ai/b, ci/d \in G$, and if $(b, d) = 1$, then $i/bd \in G$. For by (2) we have For by (2) we have

$$\frac{i}{bd} = \frac{(kb + ld)i}{bd} = \frac{ki}{d} + \frac{li}{b} \in G.$$

THEOREM 2. *If $G \neq 0$ is a subgroup of R^+ , then there exists a sequence $(i; k_1, k_2, \dots, k_j, \dots)$ such that $G = (i; k_j)$.*

Proof. By (1), every $\zeta \in G$ has the form $\zeta = ai/b$, $(ai, b) = 1$, where i is the

least positive integer in G . We write all elements of G in this form. If, for every l , there exist $ai/b \in G$ such that $p_j^l | b$, let $k_j = \infty$. If not, let $k_j = \max k$ such that $p_j^k | b$ for some $ai/b \in G$. Since $(ai, b) = 1$, we have $(i, p_j) = 1$ if $k_j > 0$. By the definition of i and k_j , G is contained in $(i; k_j)$. Now every element of $(i; k_j)$ has the form $ai/(p_1^{n_1} \cdots p_r^{n_r})$, where $n_j \leq k_j$ and $(a, p_1^{n_1} \cdots p_r^{n_r}) = 1$. By (4) and the definition of k_j , G contains every $i/p_j^{n_j}$ with $n_j \leq k_j$, and by (5), G contains $ai/(p_1^{n_1} \cdots p_r^{n_r})$. Hence $G = (i; k_j)$.

3. Properties of the rational groups. In this section, properties of the rational groups are obtained as corollaries of the theorems of §1.

COROLLARY 1. *The group $(i; k_j)$ is a subgroup of $(i'; k'_j)$ if and only if $k_j \leq k'_j$ and $i = mi'$.*

COROLLARY 2. *The group $(i; k_j)$ is cyclic if and only if $k_j < \infty$ for all j and $k_j = 0$ for almost all j .*

Proof. If $(i; k_j)$ is cyclic, it is generated by ai/b with $(ai, b) = 1$. Since every element of $(i; k_j)$ has the form nai/b , we have $a = 1$ and $b = \prod_{k_j > 0} p_j^{k_j}$. Conversely $(i; k_j)$ contains $i/\prod_{k_j > 0} p_j^{k_j}$, and this element generates $(i; k_j)$.

COROLLARY 3. *We have $(i; k_j) \cong (i'; k'_j)$ if and only if both $k_j = k'_j$ for almost all j , and, whenever $k_j \neq k'_j$, both are finite. Every isomorphism between $(i; k_j)$ and $(i'; k'_j)$ is given by*

$$\frac{ai}{b} \longleftrightarrow \frac{mai'}{nb},$$

where

$$m = \left(\prod'_{k_j = k'_j = \infty} p_j^{a_j} \right) \left(\prod_{\substack{k_j \geq k'_j \\ k_j \text{ finite}}} p_j^{k_j - k'_j} \right),$$

$$n = \left(\prod'_{k_h = k'_h = \infty} p_h^{b_h} \right) \left(\prod_{\substack{k'_h \geq k_h \\ k'_h \text{ finite}}} p_h^{k'_h - k_h} \right).$$

Proof. If $(i; k_j) \cong (i'; k'_j)$, then $i \rightarrow mi'/n$ with $(mi', n) = 1$. If $\eta \rightarrow i'$, then $m\eta \rightarrow mi'$ and $ni \rightarrow mi'$, so that $m\eta = ni$, or $\eta = ni/m$.

Hence $ni/m \rightarrow i'$. We write

$$m = p_{\alpha_1}^{a_1} \cdots p_{\alpha_r}^{a_r}, \quad a_l > 0, \quad n = p_{\beta_1}^{b_1} \cdots p_{\beta_s}^{b_s}, \quad b_m > 0;$$

then for $n_j \leq k_j$ we have

$$\frac{i}{p_j^{n_j}} \rightarrow \frac{p_{\alpha_1}^{a_1} \cdots p_{\alpha_r}^{a_r} i'}{p_j^{n_j} p_{\beta_1}^{b_1} \cdots p_{\beta_s}^{b_s}};$$

while for $n'_j \leq k'_j$ we have

$$\frac{p_{\beta_1}^{b_1} \cdots p_{\beta_s}^{b_s} i}{p_j^{n'_j} p_{\alpha_1}^{a_1} \cdots p_{\alpha_r}^{a_r}} \rightarrow \frac{i'}{p_j^{n'_j}}.$$

We have the following alternatives with consequences which follow from (3):

- I. $j = \alpha_l \quad : \quad n_j - k'_j \leq a_l \leq k_j - n'_j$
 II. $j = \beta_m \quad : \quad n'_j - k_j \leq b_m \leq k'_j - n_j$
 III. $\left. \begin{array}{l} j \neq \alpha_l \\ j \neq \beta_m \end{array} \right\} : n_j \leq k'_j, \quad n'_j \leq k_j$

It follows that $k_j = \infty$ implies $k'_j = \infty$ and conversely. With both k_j and k'_j finite we may choose $n_j = k_j$ and $n'_j = k'_j$ and we have:

- I. $j = \alpha_l \quad : \quad a_l = k_j - k'_j$
 II. $j = \beta_m \quad : \quad b_m = k'_j - k_j$
 III. $\left. \begin{array}{l} j \neq \alpha_l \\ j \neq \beta_m \end{array} \right\} : k_j = k'_j$

We have $k_j = k'_j$ if and only if $j \neq \alpha_l, j \neq \beta_m$. In particular, we have $k_j = k'_j$ for almost all j . If $k_j > k'_j$, then $j = \alpha_l$ and $a_l = k_j - k'_j$. If $k'_j > k_j$, then $j = \beta_m$ and $b_m = k'_j - k_j$.

Now $i \rightarrow mi'/n$ implies $ai/b \rightarrow ami'/bn$, so that the only isomorphisms between $(i; k_j)$ and $(i'; k'_j)$ are those described in the corollary. Incidentally, we

have derived necessary conditions for the relation $(i; k_j) \cong (i'; k'_j)$.

With the necessary conditions satisfied, we check that the given correspondence actually is an isomorphism. These conditions imply that the correspondence is single-valued with a single-valued inverse from $(i; k_j)$ onto $(i'; k'_j)$. It is clear that addition is preserved.

COROLLARY 4. *The group $(i; k_j)$ admits only the identity automorphism if and only if k_j is finite for all j .*

Proof. If k_j is finite for all j , we have by Corollary 3, with $k_j = k'_j$ for all j , that $m = n = 1$. Conversely, if any $k_j = \infty$, then the correspondence of Corollary 3 gives us nontrivial automorphisms.

The multiplicative group of the field of rational numbers, R^\times , is a direct product of the infinite cyclic subgroups of R^\times generated by the prime numbers p_k for all k . Such a subgroup consists of the elements $p_k, p_k^2, \dots, 1, 1/p_k, 1/p_k^2, \dots$.

COROLLARY 5. *The group of automorphisms of $(i; k_j)$ is isomorphic to the direct product of all of the infinite cyclic subgroups of R^\times generated by those primes p_k for which $k_j = \infty$.*

Proof. By Corollary 3, there is a (1-1) correspondence between the automorphisms of $(i; k_j)$ and the rational numbers M/N with $(M, N) = 1$, where M and N are arbitrary products of those primes for which $k_j = \infty$. This correspondence clearly preserves multiplication and the set of all rationals M/N has the stated structure as a group with respect to multiplication.

COROLLARY 6. *For any two subgroups $(i; k_j)$ and $(i'; k'_j)$ of R^+ , the set T consisting of all ordinary products of an element of $(i; k_j)$ with an element of $(i'; k'_j)$ is again a subgroup of R^+ .*

Proof. We have $T = (I; K_j)$, where

$$I = \frac{ii'}{\prod p_j p_j^{s_j}}, \quad K_j = k_j + k'_j - s_j$$

with $s_j = \min(\alpha_j, k'_j) + \min(\alpha'_j, k_j)$, where α_j is the highest power of p_j that divides i , and α'_j the highest power of p_j that divides i' .

COROLLARY 7. *If $(i; k_j) \geq (i'; k'_j)$ and $p_j^{l_j}$ is the maximum power of p_j such that $p_j^{l_j}$ divides i'/i , then the difference group $(i; k_j) - (i'; k'_j)$ is a direct sum*

whence $ac(i \times i)$

$$\begin{aligned}
 &= \left[\frac{ai}{b} \times \frac{ci}{d} \right] + \dots + \left[\frac{ai}{b} \times \frac{ci}{d} \right] \quad (\text{by the distributive laws in } S) \\
 &\qquad\qquad\qquad bd \text{ summands} \\
 &= (bd) \left[\frac{ai}{b} \times \frac{ci}{d} \right].
 \end{aligned}$$

THEOREM 3. *If there is an infinite number of k_j such that $0 < k_j < \infty$, then the only ring S with $(i; k_j)$ as additive group is the null ring. If $0 < k_j < \infty$ for only a finite number of k_j , then S is a ring with additive group $(i; k_j)$ if and only if multiplication in S is defined by*

$$\frac{ai}{b} \times \frac{ci}{d} = \frac{acA' \left(\prod_{0 < k_j < \infty} p_j^{k_j} \right) i}{bd \prod'_{k_j = \infty} p_j^{n_j}},$$

where A' and n_j are arbitrary.

Proof. If S is a ring with additive group $(i; k_j)$, then $i \times i = Ai/B \in (i; k_j)$, where $(Ai, B) = 1$, $B = \prod' p_j^{n_j}$, $n_j \leq k_j$. By the lemma, we have

$$\frac{ai}{b} \times \frac{ci}{d} = \frac{acAi}{bdB}.$$

If $0 < k_r < \infty$, this yields in particular

$$\frac{i}{p_r^{k_r}} \times \frac{i}{p_r^{k_r}} = \frac{Ai}{p_r^{2k_r} B}.$$

Therefore $(p_r, B) = 1$, for otherwise we would have $2k_r + n_r \leq k_r$, which is impossible. Hence, $B = \prod' p_j^{n_j}$ is a product of primes for which $k_j = \infty$, and it is necessary that $p_r^{k_r} | A$. If there is an infinite number of primes p_j with $0 < k_j < \infty$, then $A = 0$ and $(ai/b) \times (ci/d) = 0$. This proves the first statement in the theorem.

If $0 < k_j < \infty$ for only a finite number of primes p_j , then

$$A = A' \prod_{0 < k_j < \infty} p_j^{k_j}.$$

Together with what has been proved above, this gives

$$\frac{ai}{b} \times \frac{ci}{d} = \frac{acA' \left(\prod_{0 < k_j < \infty} p_j^{k_j} \right) i}{bd \prod'_{k_j = \infty} p_j^{n_j}},$$

where A' and $n_j > 0$ are arbitrary integers.

Conversely, this definition of multiplication always makes $(i; k_j)$ a ring. Closure with respect to \times is insured by providing $p_j^{k_j}$ in the numerator when $0 < k_j < \infty$, and the associative and distributive laws are readily verified.

COROLLARY 1. *The set $(i; k_j)$ is a subring of R if and only if there is no k_j such that $0 < k_j < \infty$.*

Proof. Let $(i; k_j)$ be a subring of R and assume that for at least one k_j we have $0 < k_j < \infty$. If $0 < k_j < \infty$ for infinitely many k_j , then $(i; k_j)$ is not a subring of R , since by Theorem 3 it is the null ring. If $0 < k_j < \infty$ for a finite number of k_j , then multiplication in any ring with $(i; k_j)$ as additive group is given by the formula of the theorem. Hence this must reduce to ordinary multiplication for some choice of A' and n_j ; that is,

$$\frac{A' \prod_{0 < k_j < \infty} p_j^{k_j}}{\prod'_{k_j = \infty} p_j^{n_j}} = i; \quad A' \prod_{0 < k_j < \infty} p_j^{k_j} = i \prod'_{k_j = \infty} p_j^{n_j}.$$

By hypothesis, at least one p_j with $k_j > 0$ appears in the left member of the above equality. Since no prime appears in both products, we have $p_j | i$. This contradicts $(i, p_j) = 1$ for $k_j > 0$.

Conversely, let every k_j be either 0 or ∞ . By the theorem, we have

$$\frac{ai}{b} \times \frac{ci}{d} = \frac{acA' i}{bd \prod'_{k_j = \infty} p_j^{n_j}},$$

and we may select $A' = i, \prod'_{k_j = \infty} p_j^{n_j} = 1$, yielding ordinary multiplication.

COROLLARY 2. *If $(i; k_j)$ is a subring of R , then $(i; k_j)$ is a ring under the*

multiplication

$$\frac{ai}{b} \times \frac{ci}{d} = \frac{ac}{bd} \left(\frac{ei}{f} \right)$$

for arbitrary $ei/f \in (i; k_j)$.

Proof. By Corollary 1, we have $k_j = 0$ or $k_j = \infty$, so that every element of $(i; k_j)$ has the form

$$\frac{A' i}{\prod'_{k_j=\infty} p_j^{n_j}} ;$$

and by the theorem these are just the multipliers which are used to define multiplication.

COROLLARY 3. *If S is a ring with additive group $(i; k_j)$, then either S is a null ring or S is isomorphic to a subring of R .*

Proof. If S is not null, the correspondence

$$\frac{ai}{b} \rightarrow \frac{aA}{bB}, \quad \text{where} \quad \frac{ai}{b} \times \frac{ci}{d} = \frac{acAi}{bdB},$$

is (1-1) from S on a subset of R , and

$$\frac{ai}{b} + \frac{ci}{d} = \frac{(da + bc)i}{bd} \rightarrow \frac{(da + bc)A}{bdB} = \frac{aA}{bB} + \frac{cA}{dB},$$

$$\frac{ai}{b} \times \frac{ci}{d} = \frac{acAi}{bdB} \rightarrow \frac{acA^2}{bdB^2} = \frac{aA}{bB} \frac{cA}{dB}.$$

COROLLARY 4. *All rings with additive group R^+ are isomorphic to R .*

Proof. The correspondence of Corollary 3 clearly exhausts R .

REFERENCE

1. R. Baer, *Abelian groups without elements of finite order*, Duke Math. J. 3 (1937), 68-122.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

HERBERT BUSEMANN
University of Southern California
Los Angeles 7, California

R. M. ROBINSON
University of California
Berkeley 4, California

E. F. BECKENBACH, Managing Editor
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

R. P. DILWORTH	P. R. HALMOS	BØRGE JESSEN	J. J. STOKER
HERBERT FEDERER	HEINZ HOPF	PAUL LÉVY	E. G. STRAUS
MARSHALL HALL	R. D. JAMES	GEORGE PÓLYA	KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA	UNIVERSITY OF SOUTHERN CALIFORNIA
CALIFORNIA INSTITUTE OF TECHNOLOGY	STANFORD UNIVERSITY
UNIVERSITY OF CALIFORNIA, BERKELEY	WASHINGTON STATE COLLEGE
UNIVERSITY OF CALIFORNIA, DAVIS	UNIVERSITY OF WASHINGTON
UNIVERSITY OF CALIFORNIA, LOS ANGELES	* * *
UNIVERSITY OF CALIFORNIA, SANTA BARBARA	AMERICAN MATHEMATICAL SOCIETY
OREGON STATE COLLEGE	NATIONAL BUREAU OF STANDARDS,
UNIVERSITY OF OREGON	INSTITUTE FOR NUMERICAL ANALYSIS

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1951 BY PACIFIC JOURNAL OF MATHEMATICS

Pacific Journal of Mathematics

Vol. 1, No. 2

December, 1951

Tom M. (Mike) Apostol, <i>On the Lerch zeta function</i>	161
Ross A. Beaumont and Herbert S. Zuckerman, <i>A characterization of the subgroups of the additive rationals</i>	169
Richard Bellman and Theodore Edward Harris, <i>Recurrence times for the Ehrenfest model</i>	179
Stephen P.L. Diliberto and Ernst Gabor Straus, <i>On the approximation of a function of several variables by the sum of functions of fewer variables</i>	195
Isidore Isaac Hirschman, Jr. and D. V. Widder, <i>Convolution transforms with complex kernels</i>	211
Irving Kaplansky, <i>A theorem on rings of operators</i>	227
W. Karush, <i>An iterative method for finding characteristic vectors of a symmetric matrix</i>	233
Henry B. Mann, <i>On the number of integers in the sum of two sets of positive integers</i>	249
William H. Mills, <i>A theorem on the representation theory of Jordan algebras</i>	255
Tibor Radó, <i>An approach to singular homology theory</i>	265
Otto Szász, <i>On some trigonometric transforms</i>	291
James G. Wendel, <i>On isometric isomorphism of group algebras</i>	305
George Milton Wing, <i>On the L^p theory of Hankel transforms</i>	313