

# Pacific Journal of Mathematics

**RECURRENCE TIMES FOR THE EHRENFEST MODEL**

RICHARD BELLMAN AND THEODORE EDWARD HARRIS

# RECURRENCE TIMES FOR THE EHRENFEST MODEL

RICHARD BELLMAN AND THEODORE HARRIS

1. Introduction and summary. In 1907, P. and T. Ehrenfest [1] used a simple urn scheme as a pedagogic device to elucidate some apparent paradoxes in thermodynamic theory. Their model undergoes fluctuations intuitively related to fluctuations about equilibrium of certain thermodynamic systems. In view of an apparent discord among physicists [6, pp.139-145] we shall not try to force an analogy with entropy.

The original Ehrenfest scheme was defined as follows. Initially,  $2N$  balls are divided in an arbitrary manner between two urns, 1 and 2, the balls being numbered from 1 to  $2N$ . An integer between 1 and  $2N$  is selected at random, each such integer having probability  $(2N)^{-1}$ , and the ball with the number selected is transferred from one urn to the other. The process is repeated any number of times. If  $n_1$  and  $n_2$  are the numbers of balls in urns 1 and 2 respectively before a transfer, it is clear that the probability is  $n_1/(2N)$  that the transfer is from urn 1 to urn 2 and  $n_2/(2N)$  that it is in the contrary direction.

Let  $x'(n)$  be the number of balls in urn 1 after  $n$  transfers, and let  $L'_{j,k}$  be the smallest integer  $m$  such that  $x'(m) = k$ , given that  $x'(0) = j$ . If  $k = j$ , we call  $L'_{k,k}$  the *recurrence time* for the state  $k$ . If  $k \neq j$ , we call  $L'_{j,k}$  the *first-passage time* from  $j$  to  $k$ . The distribution of  $x'(n)$ , known classically, was derived by Kac [5] as an example of the use of matrix methods. Kac then found the mean and variance of  $L'_{j,k}$ , attributing some of his methods to Uhlenbeck. Friedman [4] found the moment-generating function for  $x'(n)$  (for the Ehrenfest and more general models) by solving a difference-differential equation.

Instead of the original Ehrenfest model, we shall discuss a modified scheme with a continuous time parameter, which was apparently first suggested by A. J. F. Siegert [9]. In this scheme there are two urns and  $2N$  balls initially divided between them arbitrarily. Each ball acts, independently of all the others, as follows: there is a probability of  $(1/2) dt + o(dt)$  that the ball changes urns between  $t$  and  $t + dt$ , and a probability of  $1 - [(1/2) dt + o(dt)]$  that the ball remains

---

Received August 5, 1950, and, in revised form, November 8, 1950.

*Pacific J. Math.* 1 (1951), 179-193.

in place between  $t$  and  $t + dt$ . Standard reasoning then shows that the total probability of a change by some ball between  $t$  and  $t + dt$  is  $Ndt + o(dt)$ , and that consequently the probability density for the length of time between transfers is  $Ne^{-Nt} dt$ . When a transfer occurs, it is readily seen that the probabilities that it is from urn 1 to urn 2 or from urn 2 to urn 1, respectively, depend on the relative number of balls in the two urns exactly as for the original Ehrenfest model. Thus we see that the present scheme is essentially the original Ehrenfest scheme where the drawings are made at random times. As we shall see, the time-continuous scheme is easier to handle analytically.

Let  $x(t)$  be the number of balls in urn 1 at time  $t$ ; we shall sometimes speak of this number as the *state* of the system. Then  $x(t)$  is a random function which can take integer values from 0 to  $2N$ ;  $x(t)$  executes a random walk—with a “restoring force”—about the equilibrium value  $N$ . It is clear that the random walk is a Markov process.

Let  $L_{j,k}$ ,  $j \neq k$ , be the *first-passage time* from state  $j$  to state  $k$ ; that is,  $L_{j,k}$  is the infimum of  $t$  such that  $x(t) = k$ , given that  $x(0) = j$ . Let  $L_{k,k}$  be the *recurrence time* for the state  $k$ ; that is,  $L_{k,k}$  is the infimum of  $t$  such that  $x(t) = k$  and  $x(\tau) \neq k$  for  $0 < \tau < t$ , given that  $x(0) = k$ . We shall discuss the probability distributions of  $L_{j,k}$  and  $L_{k,k}$ .

The probability distribution of  $L_{j,k}$  depends, of course, on the size of the model (that is, on the number  $N$ ). When it is necessary to emphasize this dependence we shall sometimes employ the notation  $L_{j,k}^{(N)}$  in place of  $L_{j,k}$ .

We shall use the notation  $P(A)$  for the probability of the event  $A$ ;  $P(A | B)$  for the conditional probability of  $A$ , given  $B$ ;  $E(X)$  for the mean, or expected value, of the random variable  $X$ . By the *distribution* of a random variable  $X$  we mean the function (of say  $u$ ) given by  $P(X \leq u)$ . The statement that a sequence of distributions converges to a distribution  $F(u)$  will mean convergence at all continuity points of  $F(u)$ .

There are two limiting situations in which the distribution of  $L_{j,k}$  is of interest.

(a) Consider a simple thermodynamic system such as an ideal gas in a container. Let us think of the container as consisting of two halves which, however, are not separated by a partition. Suppose that initially the molecules are spread in a rather uniform manner through the two halves of the container. According to classical kinetic theory, if we wait long enough, a time will come, in general, when all the molecules are in one half of the container. Such events, where the fraction of molecules in one half of the container is appreciably different from

(1/2), are evidently enormously rare if the number of molecules is large. Correspondingly, we should like to show that the random variable  $L_{N,k}$ , where  $|k - N|$  is of the order of magnitude of  $N$ , is very large with high probability when  $N$  is large. Now the mean of  $L_{N,k}$  is extremely large when  $N$  is large. However, as Kac has observed, the standard deviation is of the same order of magnitude as the mean. Thus we cannot conclude from the values of the first two moments that  $L_{N,k}$  is large with high probability. We shall show, however, that *the distribution of  $L_{N,k}/E(L_{N,k})$  converges to  $1 - e^{-u}$  as  $N \rightarrow \infty$  provided  $k/N$  remains less than some fixed number  $\lambda_1 < 1$  (Theorem 1).*

The situation with respect to  $L_{k,k}$ , where again  $k/N < \lambda_1 < 1$ , is somewhat different. If  $k/N$  is appreciably different from 0, a very short recurrence time is not improbable. The distribution of  $L_{k,k}/E(L_{k,k})$  has for large  $N$  a "lump" of probability of magnitude  $k/N$  concentrated near 0, the remainder of the distribution being exponential (Theorem 2).

(b) In the theory of the Brownian motion and elsewhere in physics and statistics an important role is played by the stationary Gaussian Markov process  $z(t)$  which we scale so that

$$E[z(t)] = 0, \quad E[z(t)]^2 = 1/2.$$

This process is defined by the requirement that the joint distribution of  $z(t_1), \dots, z(t_m)$  for any distinct numbers  $t_1, \dots, t_m$  is Gaussian and dependent only on the differences  $t_i - t_j$  and that the autocorrelation function is given by

$$E[z(s) z(t + s)] = (1/2) e^{-|t|}.$$

If  $N$  is large, the  $z(t)$  process, under the conditional hypothesis that  $z(0)$  has an appropriate value, is approximated by the process

$$z_N(t) = \frac{x(t) - N}{N^{1/2}}$$

in a sense described in Section 6. (It should be remembered that  $x(t)$  depends on  $N$ .) By considering the distribution of  $L_{N,k}$  where  $(k - N)/N^{1/2} \rightarrow -\xi_0 < 0$ ,  $N \rightarrow \infty$ , we obtain in Theorem 3 the Laplace transform of the distribution of  $L$ , the first time at which  $z(t) = -\xi_0$ , given  $z(0) = 0$ . This result is not new, having been obtained by Siegert [10] and by Darling (unpublished). However, the present method of derivation seems instructive.

Results similar to those given under (a) and (b) are obtained for the random

variable  $L_{N,k}^*$ , the first time  $|x(t) - N| = N - k$ , given  $x(0) = N$ .

2. **The mathematical model.** Suppose that there are initially  $j$  balls in urn 1 and  $2N - j$  in urn 2. Associate with the  $i$ th ball a random function  $x_i(t)$  defined as follows:  $x_i(t)$  is 1 if the  $i$ th ball is in urn 1 at time  $t$ , and 0 otherwise. From the elementary theory of Markov processes (see, for example, Kolmogorov [7]), we have

$$(1) \quad P[x_i(t) = x_i(0)] = \frac{1 + e^{-t}}{2},$$

$$P[x_i(t) = 1 - x_i(0)] = \frac{1 - e^{-t}}{2}.$$

We may define the generating function of  $x_i(t)$  by

$$P[x_i(t) = 0] + sP[x_i(t) = 1].$$

Then the generating function of  $x_i(t)$  is, from (1),

$$(1/2)[1 + e^{-t} + (1 - e^{-t})s],$$

or

$$(1/2)[1 - e^{-t} + (1 + e^{-t})s],$$

according as  $x_i(0)$  is 0 or 1. Since the quantities  $x_i(t)$ ,  $i = 1, \dots, 2N$ , are independent, the generating function for  $x(t) = \sum x_i(t)$  is

$$(2) \quad \sum_{k=0}^{2N} P[x(t) = k | x(0) = j] s^k$$

$$= 2^{-2N} [1 - e^{-t} + (1 + e^{-t})s]^j [1 + e^{-t} + (1 - e^{-t})s]^{2N-j}$$

$$= \sum_{k=0}^{2N} Q_{j,k}(t) s^k,$$

where we have introduced the notation  $Q_{j,k}(t)$  for  $P[x(t) = k | x(0) = j]$ . Formula (2) was given by Siebert [9; 11].

Because of the simple nature of the process under consideration it is easy to show that  $L_{j,k}$  and  $L_{k,k}$  are (measurable) random variables with absolutely continuous distributions. We omit the proof. We let  $P_{j,k}(u)$  be the probability density

of  $L_{j,k}$ ,

$$(3) \quad \int_0^u P_{j,k}(y) dy = P[L_{j,k} \leq u].$$

Define

$$(4) \quad Q_k = \lim_{t \rightarrow \infty} Q_{j,k}(t) = \binom{2N}{k} / 2^{2N},$$

$$(5) \quad m_k = 1 / [(N - k)Q_k], \quad k < N.$$

It is convenient to notice that, as  $N \rightarrow \infty$ ,

$$(6) \quad m_k = \frac{[\lambda^\lambda (2 - \lambda)^{2 - \lambda}]^N \left[ \frac{\pi \lambda (2 - \lambda)}{N} \right]^{1/2} \left[ 1 + \frac{1}{\lambda} O\left(\frac{1}{N}\right) \right]}{1 - \lambda}, \quad k \neq 0,$$

where we have put  $\lambda = k/N$ , and  $O(1/N)$  is independent of  $\lambda$ .

The quantities  $L_{j,k}$ ,  $Q_k$ , and so on, depend on the size of the model; when it is necessary to emphasize this dependence we shall write  $L_{j,k}^{(N)}$ ,  $Q_k^{(N)}$ , and so on.

**3. Distribution of  $L_{j,k}$ ,  $j \neq k$ .** In this section we consider the distribution of the first-passage time from state  $j$  to state  $k$  for large  $N$ , where  $|j - k|$  is of the order of magnitude of  $N$ . As far as the limiting distributions are concerned, we can restrict ourselves without loss of generality to consideration of  $L_{N,k}$ ,  $k < N$ . For example, if  $j > N > k$  then we can write

$$L_{j,k} = L_{j,N} + L_{N,k}.$$

The first-passage time from  $j$  to  $N$ , representing movement toward equilibrium, is negligible relative to  $L_{N,k}$  and does not affect the asymptotic result. On the other hand if  $N > j > k$ , we have

$$L_{j,k} = L_{N,k} - L_{N,j}$$

and it is not difficult to show that  $L_{N,j}$  is negligible compared with  $L_{N,k}$ .

If the first passage to the state  $k$  occurs at time  $\tau$ , the probability that the state at time  $t$  is again  $k$  is  $Q_{k,k}(t - \tau)$ . We have therefore

$$(7) \quad Q_{j,k}(t) = \int_0^t P_{j,k}(\tau) Q_{k,k}(t - \tau) d\tau, \quad j \neq k.$$

Formula (7) is the continuous counterpart of a formula long used for discrete

processes and recently exploited by Feller [2]. Taking Laplace transforms of both sides of (7) we have

$$(8) \quad \int_0^\infty P_{j,k}(t) e^{-\sigma t} dt = \frac{\int_0^\infty Q_{j,k}(t) e^{-\sigma t} dt}{\int_0^\infty Q_{k,k}(t) e^{-\sigma t} dt}, \quad \Re(\sigma) > 0,$$

Since the quantities  $Q_{j,k}(t)$  are polynomials in  $e^{-\lambda t}$ , as we observe from (2), both the numerator and the denominator in (8) have a simple pole at  $\sigma = 0$ , and their quotient is therefore analytic in the circle  $|\sigma| < 1$ .

For simplicity denote  $L_{N,k}^{(N)}$  by  $L_k^{(N)}$ . We have the following result.

**THEOREM 1.** *The distribution function of  $L_k^{(N)}/m_k^{(N)}$  converges to  $1 - e^{-u}$ ,  $u \geq 0$ , as  $N \rightarrow \infty$ , provided  $k/N \leq \lambda_1 < 1$ , the convergence being uniform in  $k$  and  $u$ ,*

The proof will bring out the fact that likewise

$$(9) \quad E[L_k^{(N)}]/m_k^{(N)} \rightarrow 1, \quad N \rightarrow \infty, \quad k/N \leq \lambda_1.$$

Theorem 1 will follow from this lemma:

**LEMMA 1.** *For the complex variable  $\sigma$ , let*

$$\phi_k^{(N)}(\sigma) = \int_0^\infty P_{N,k}^{(N)}(u) \exp[-\sigma u/m_k^{(N)}] du.$$

*Further, let  $\{k(N)\}$  be a sequence of nonnegative integers such that  $k(N)/N \rightarrow \lambda_0 \leq \lambda_1 < 1$  as  $N \rightarrow \infty$ . Then (the convergence being bounded and uniform provided  $|\sigma| < \sigma_0 < 1$ ),*

$$(10) \quad \lim_{N \rightarrow \infty} \phi_{k(N)}^{(N)}(\sigma) = \frac{1}{1 + \sigma}, \quad |\sigma| < 1.$$

*Proof of Theorem 1.* The function  $\phi_k^{(N)}(-\sigma)$  is the moment-generating function" of the quantity  $L_k^{(N)}/m_k^{(N)}$ . Lemma 1 then implies, as is well known, that

$$(11) \quad \lim_{N \rightarrow \infty} P[L_{k(N)}^{(N)}/m_{k(N)}^{(N)} \leq u] = 1 - e^{-u}$$

uniformly for  $u \geq 0$  provided  $k(N)/N \rightarrow \lambda_0$ . Lemma 1 also implies, since we have convergence in a complex neighborhood of  $\sigma = 0$ , that

$$\lim_{N \rightarrow \infty} E[L_{k(N)}^{(N)}] / m_{k(N)} = 1,$$

so that (11) is still true if we replace  $m_{k(N)}^{(N)}$  by  $E[L_{k(N)}^{(N)}]$ .

Now if Theorem 1 were not true then an  $\epsilon > 0$  and a sequence  $\{h(N)\}, h(N)/N \leq \lambda_1$ , would exist such that for infinitely many integers  $N$  we would have

$$\sup_{0 \leq u < \infty} |1 - e^{-u} - P[L_{h(N)}^{(N)} / m_{h(N)} \leq u]| > \epsilon.$$

Extracting a convergent subsequence from  $\{h(N)/N\}$ , we are led to a contradiction of (11).

*Proof of Lemma 1.* The proof of Lemma 1, which is somewhat indirect, proceeds as follows. We can obtain an expression for  $\phi_k^{(N)}(\sigma)$  by substituting  $\sigma/m_k^{(N)}$  for  $\sigma$  in (8), obtaining

$$(12) \quad \phi_k^{(N)}(\sigma) = \frac{\int_0^\infty Q_{N,k}(t) e^{-\sigma t/m_k} dt}{\int_0^\infty Q_{k,k}(t) e^{-\sigma t/m_k} dt} = \frac{J_1}{J_2}.$$

We can obtain an asymptotic estimate of  $J_1$  as we shall see later. However, a direct estimate of  $J_2$  appears difficult to obtain. We shall therefore resort to another expression for  $\phi_k^{(N)}(\sigma)$  which is easier to estimate. Having estimates for  $\phi_k^{(N)}(\sigma)$  and for  $J_1$ , we can get an estimate of  $J_2$ , which will be necessary for Theorem 2.

Since a direct proof of Lemma 1 is easy if all terms in the sequence  $\{k(N)\}$  are 0 we can suppose  $k > 0$ . If  $0 < k < N$  we have, from elementary reasoning, the important relation

$$(13) \quad L_{N,0} = L_{N,k} + L_{k,0}.$$

On account of the Markovian nature of the process,  $L_{N,k}$  and  $L_{k,0}$  are independent random variables and the Laplace transform of the distribution of their sum is the product of the Laplace transforms of their individual distributions. Therefore, using (8) and (13), we have

$$(14) \quad E(e^{-sL_{N,0}}) = E(e^{-sL_{N,k}}) E(e^{-sL_{k,0}}),$$



or

$$\begin{aligned}
 (15) \quad \phi_k^{(N)}(\sigma) &= \int_0^\infty P_{N,k}(t) e^{-\sigma t/m_k} dt \\
 &= \frac{\int_0^\infty P_{N,0}(t) e^{-\sigma t/m_k} dt}{\int_0^\infty P_{k,0}(t) e^{-\sigma t/m_k} dt} = \frac{\int_0^\infty Q_{N,0}(t) e^{-\sigma t/m_k} dt}{\int_0^\infty Q_{k,0}(t) e^{-\sigma t/m_k} dt} \\
 &= \frac{\int_0^\infty (1 - e^{-2t})^N e^{-\sigma t/m_k} dt}{\int_0^\infty (1 - e^{-t})^k (1 + e^{-t})^{2N-k} e^{-\sigma t/m_k} dt}
 \end{aligned}$$

The advantage of (15) over (12) is that  $Q_{k,0}(t)$  is a simpler function than  $Q_{k,k}(t)$ .

The numerator of the last fraction in (15) is  $(1/2)B [N + 1, (1/2)\sigma/m_k]$ . The denominator, with the substitution  $e^{-t} = y$ , becomes

$$(16) \quad I = \int_0^1 (1 - y)^k (1 + y)^{2N-k} y^{(\sigma/m_k)-1} dy .$$

We now have to estimate  $I$  as  $N \rightarrow \infty$  under the hypothesis  $k/N \rightarrow \lambda_0 < 1$ . [We shall write simply  $k$  for  $k(N)$ .] We shall restrict  $\sigma$  to the circumference of a circle, say  $|\sigma| = (1/2)$ , since it is clearly sufficient to prove Lemma 1 for such a circle. Write

$$I = \int_0^\epsilon + \int_\epsilon^1 = I_1 + I_2 , \quad 0 < \epsilon < 1 - \lambda_0 \leq 1 .$$

Making use of (6) and the fact that  $(1 - y)^k (1 + y)^{2N-k}$  increases to a maximum at  $y = 1 - k/N$  and then decreases,  $1 - k/N$  being larger than  $\epsilon$  for sufficiently large  $N$ , we have

$$\begin{aligned}
 (17) \quad I_1 &= \int_0^{1/N^2} + \int_{1/N^2}^\epsilon \\
 &= \frac{mk}{\sigma} [1 + o(1)] + \int_{1/N^2}^\epsilon (1 - y)^k (1 + y)^{2N-k} y^{(\sigma/m_k)-1} dy
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{m_k}{\sigma} [1 + o(1)] + O[(1 - \epsilon)^k (1 + \epsilon)^{2N-k} \log N] \\
 &= \frac{m_k}{\sigma} [1 + o(1)] + o(m_k),
 \end{aligned}$$

where  $o(\ )$  is independent of  $\sigma$  for  $|\sigma| = 1/2$ .

To estimate  $I_2$  we distinguish the cases  $\lambda_0 > 0$  and  $\lambda_0 = 0$ . If  $\lambda_0 > 0$ , then  $I_2$  can be estimated using the method of Laplace; see [8, p. 77]. We obtain then, setting  $k/N = \lambda$ , (see (6)),

$$(18) \quad I_2 = [\lambda^\lambda (2 - \lambda)^{2-\lambda}]^N \left[ \frac{\pi \lambda (2 - \lambda)}{N(1 - \lambda)} \right]^{1/2} [1 + o(1)] = m_k [1 + o(1)].$$

Going back to (15), we obtain (10) from (17) and (18), since

$$(1/2)B[N + 1, (1/2)\sigma/m_k] = (m_k/\sigma)[1 + o(1)].$$

This completes the proof of Lemma 1 for the case  $\lambda_0 > 0$ . If  $\lambda_0 = 0$ , the integral  $I_2$  can be estimated by making a change of the variable of integration which shows the integral to be asymptotically equivalent to a Beta function. We need not enter into details.

**4. Distribution of  $L_{k,k}$ .** We shall establish the following result.

**THEOREM 2.** Assume  $\lambda = k/N \leq \lambda_1 < 1$ , and put

$$F_\lambda(u) = \lambda + (1 - \lambda) [1 - e^{-(1-\lambda)u}], \quad u \geq 0.$$

Then for every  $b > 0$  we have

$$\lim_{N \rightarrow \infty} \sup_{b \leq u < \infty} |P[L_{k,k}^{(N)} / E(L_{k,k}^{(N)}) \leq u] - F_\lambda(u)| = 0$$

uniformly in  $k$ .

*Proof.* As in the case of Theorem 1, it is sufficient to prove that the Laplace transform of the distribution of  $NQ_k^{(N)} L_{k,k}^{(N)}$  approaches

$$\lambda_0 + (1 - \lambda_0)^2 / (1 + \sigma - \lambda_0)$$

provided  $k/N = \lambda \rightarrow \lambda_0 < 1$ . (We know from the general theory of Markov processes that

$$EL_{k,k}^{(N)} = 1 / (NQ_k^{(N)}),$$

see Feller, [3, p. 325].)

The relation which replaces (7) when  $j = k$  is

$$(19) \quad Q_{k,k}(t) = e^{-Nt} + \int_0^t P_{k,k}(\tau) Q_{k,k}(t-\tau) d\tau,$$

the term  $e^{-Nt}$  in (19) being the probability that the system remains in state  $k$  the entire time from 0 to  $t$ . From (19), we have

$$(20) \quad \int_0^\infty P_{k,k}(t) e^{-\sigma Q_k N t} dt \\ = 1 - \frac{1}{N(1 + \sigma Q_k) \int_0^\infty Q_{k,k}(t) e^{-\sigma Q_k N t} dt}, \quad |\sigma| = \frac{1}{2}.$$

If we equate the right side of (8), with  $j = N$  and with  $\sigma$  replaced by  $\sigma N Q_k$ , to the right side of (15) with  $\sigma/m_k$  replaced by  $\sigma N Q_k$ , we obtain

$$\frac{\int_0^\infty Q_{N,k}(t) e^{-\sigma N Q_k t} dt}{\int_0^\infty Q_{k,k}(t) e^{-\sigma N Q_k t} dt} = \phi_k^{(N)}(\sigma N Q_k m_k) = \phi_k^{(N)}\left(\frac{\sigma}{1-\lambda}\right),$$

or

$$(21) \quad \int_0^\infty Q_{k,k}(t) e^{-\sigma N Q_k t} dt = \frac{\int_0^\infty Q_{N,k}(t) e^{-\sigma N Q_k t} dt}{\phi_k^{(N)} \sigma / (1-\lambda)} \\ = \frac{I_3}{\phi_k^{(N)} \sigma / (1-\lambda)}.$$

To estimate  $I_3$ , which is the numerator of (12) with  $\sigma$  replaced by  $\sigma/(1-\lambda)$ , we need two lemmas.

LEMMA 2. Given  $\epsilon > 0$ , let

$$t_N(\epsilon) = \sup \left\{ t: \max_{0 \leq r \leq 2N} \frac{|Q_{N,r}^{(N)}(t) - Q_r^{(N)}|}{Q_r^{(N)}} \geq \epsilon \right\}.$$

Then

$$(22) \quad t_N(\epsilon) = O(\log N), \quad N \rightarrow \infty.$$

*Proof.* By (2),  $Q_{N,r}(t)$  is the coefficient of  $s^r$  in

$$(1 - e^{-2t})^N (1 + zs + s^2)^N / 2^{2N},$$

where  $z = z(t) = 2(1 + e^{-2t}) / (1 - e^{-2t})$ . Since for large  $N$  the root of the equation

$$(1 - e^{-2t})^N = 1 - \delta, \quad 0 < \delta < 1, \quad \delta \text{ fixed},$$

is approximately  $t \sim (1/2) \log N$ , it suffices to prove Lemma 2 for the quantities

$$t_N^*(\epsilon) = \sup \left\{ t : \max_{0 \leq r \leq 2N} \frac{|c_r^{(N)}(t) - c_r^{(N)}|}{c_r^{(N)}} \geq \epsilon \right\},$$

where we have set

$$(23) \quad (1 + zs + s^2)^N = \sum c_r^{(N)}(t) s^r,$$

$$c_r^{(N)} = c_r^{(N)}(\infty) = 2^{2N} Q_r^{(N)}.$$

Suppose  $\epsilon > 0$  is given. Choose an arbitrary  $\alpha > 1$ . Let  $\epsilon_1 < \epsilon$  be a positive number and define

$$(24) \quad \epsilon_{N+1} = \epsilon_N (1 + 1/N^\alpha), \quad N = 1, 2, \dots$$

Note that  $\{\epsilon_N\}$  is a bounded increasing sequence. We select  $\epsilon_1$  small enough so that  $\epsilon_N < \epsilon$ , for all  $N$ . Now define a sequence  $\bar{t}_1 \leq \bar{t}_2 \leq \dots$  as follows:  $\bar{t}_1 = t_1^*(\epsilon_1)$ ;  $\bar{t}_{N+1}$  for  $N \geq 1$  is the maximum of  $\bar{t}_N$  and the positive root of

$$(25) \quad (1/3)[z(t) - 2](1 + \epsilon_N)/\epsilon_N = 1/N^\alpha.$$

[Note that  $z(t)$  is monotone decreasing.] It is then clear from (25) that

$$(26) \quad \bar{t}_N \sim (1/2)\alpha \log N, \quad N \rightarrow \infty.$$

We now wish to show inductively that

$$(27) \quad \frac{|c_r^{(N)}(t) - c_r^{(N)}|}{c_r^{(N)}} \leq \epsilon_N \quad \text{for } t \geq \bar{t}_N, \quad N = 1, 2, \dots$$

Clearly (27) holds for  $N = 1$ , since  $\epsilon_1 < \epsilon$  and  $\bar{t}_1 = t_1^*$ . Suppose that (27) is true for a general  $N$ . From (23), we have

$$(28) \quad \begin{aligned} c_r^{(N+1)}(t) &= c_r^{(N)}(t) + zc_{r-1}^{(N)}(t) + c_{r-2}^{(N)}(t), \\ c_r^{(N+1)} &= c_r^{(N)} + 2c_{r-1}^{(N)} + c_{r-2}^{(N)}. \end{aligned}$$

Using (27), (28), and the fact that  $c_{r-1}^{(N)}/c_r^{(N+1)} \leq 1/3$ , we have for  $t \geq \bar{t}_N$ ,

$$(29) \quad \frac{|c_r^{(N+1)}(t) - c_r^{(N+1)}|}{c_r^{(N+1)}} \leq \epsilon_N \left[ 1 + \frac{1}{3} |z(t) - 2| \left( \frac{1 + \epsilon_N}{\epsilon_N} \right) \right].$$

From the definition of  $\epsilon_N$  it is then clear that (27) holds with  $N$  replaced by  $N + 1$ . Then  $t > \bar{t}_N$  implies that the left side of (27) is less than  $\epsilon$ . Use of (26) now completes the proof of Lemma 2.

LEMMA 3. Assume  $k/N \leq \lambda_1$ . Then

$$\max_{0 \leq t \leq \infty} Q_{N,k}^{(N)}(t) < \exp[-3(1 - \lambda_1)^2 N/5].$$

*Proof.* Lemma 3 is an immediate consequence of a result of S. Bernstein on sums of independent random variables; see Uspensky [12, p. 205]. To apply Bernstein's result, we consider the  $2N$  balls as consisting of  $N$  pairs, each pair having initially one ball in urn 1 and one in urn 2, letting Uspensky's random variable  $x_i$  be the number of balls from the  $i$ th pair in urn 1, minus 1, at time  $t$ . Now

$$Q_{N,k}^{(N)}(t) = P \left[ \sum_{i=1}^N x_i = k - N \right] < P \left[ \sum_{i=1}^N x_i \leq k - N \right],$$

and the applicability of Bernstein's result is obvious.

We now return to the proof of Theorem 2. To estimate the integral  $I_3$  defined in (21), write

$$(30) \quad I_3 = \int_0^\infty (Q_{N,k}(t) - Q_k) e^{-\sigma N Q_k t} dt + 1/(N\sigma).$$

Write the integral on the right side of (30) as

$$\int_0^{t_N(\epsilon)} + \int_{t_N(\epsilon)}^\infty = I'_3 + I''_3$$

for an arbitrary  $\epsilon > 0$ , where  $t_N(\epsilon)$  is defined in Lemma 2. Using Lemmas 2 and 3, we have

$$(31) \quad |I_3''| < \epsilon Q_k \int_{t_N(\epsilon)}^{\infty} e^{-\sigma N Q_k t} dt < \epsilon / (N\sigma),$$

$$|I_3'| = O\{\log N \exp[-3(1 - \lambda_1)^2 N / 5]\}.$$

Thus  $I_3 \sim 1/(N\sigma)$ . Putting this estimate in (21) and recalling from Theorem 1 that

$$\phi_k^{(N)}\left(\frac{\sigma}{1 - \lambda}\right) \rightarrow \frac{1}{1 + \sigma / (1 - \lambda_0)},$$

we get the desired result from (20).

**5. Intuitive interpretation.** Theorem 1 means intuitively that if we take  $m_k$  as our time unit, the attainment of the state  $k$  is an occurrence of the "chance" type; that is, the probability of attaining  $k$  during a given time interval is almost independent of the past history of the process. This interpretation suggests that Theorem 1 should be true for more general types of processes with a central tendency.

Theorem 2 seems to mean that if the initial state is  $k$  there is a probability  $\lambda$  of returning to  $k$  before leaving its immediate neighborhood; there is a probability  $1 - \lambda$  of getting completely away from the neighborhood before the first return; in this case the first return has the distribution of first passage times given in Theorem 1.

**6. Application to stationary Gaussian Markov processes.** In Theorems 1 and 2 we considered rare or microscopic fluctuations of  $x(t)$ . But if  $N$  is large  $x(t)$  will for the most part deviate little from its mean value  $N$ , and to consider the ordinary fluctuations of  $x(t)$  we consider

$$z_N(t) = [x(t) - N] / N^{1/2}.$$

Let  $t_1, \dots, t_m$  be a fixed set of nonnegative numbers. The joint distribution of  $z_N(t_1), \dots, z_N(t_m)$ , given  $z_N(0) = 0$ , approaches, as  $N \rightarrow \infty$ , the joint distribution of  $z(t_1), \dots, z(t_m)$ , given  $z(0) = 0$ , where  $z(t)$  is the stationary Gaussian Markov process with

$$E[z(t)] = 0, \quad E[z(s) z(s + t)] = (1/2) e^{-|t|}.$$

Define the random variable  $L$  to be the smallest value of  $t$  for which  $z(t) = -\xi_0 < 0$ , given  $z(0) = 0$ . It is intuitively clear that the distribution of  $L$  is given by the limiting distribution of  $L_{N,k}$  as  $N \rightarrow \infty$  provided we let

$$(32) \quad (k - N)/N^{1/2} \rightarrow -\xi_0.$$

A rigorous proof of this statement is not difficult but we omit it.

To find the limiting Laplace transform for the distribution of  $L_{N,k}$  under the hypothesis (32), we consider (15) with  $\sigma > 0$  in place of  $\sigma/m_k$ , and let  $k = N - \xi N^{1/2}$ . The substitution  $e^{-t} = y/N^{1/2}$  puts the denominator in the form

$$(33) \quad \frac{1}{N^{\sigma/2}} \left[ \int_0^{N^\alpha} + \int_{N^\alpha}^{N^{1/2}} \left( 1 - \frac{y}{N^{1/2}} \right)^{N - \xi N^{1/2}} \times \left( 1 + \frac{y}{N^{1/2}} \right)^{N + \xi N^{1/2}} y^{\sigma-1} dy \right],$$

where  $\alpha$  is an arbitrary number between 0 and  $1/6$ . If  $0 < y < N^\alpha$ , then

$$\left( 1 - \frac{y}{N^{1/2}} \right)^{N - \xi N^{1/2}} \left( 1 + \frac{y}{N^{1/2}} \right)^{N + \xi N^{1/2}} = e^{-y^2 + 2\xi y} [1 + O(N^{-1/2 + 3\alpha})].$$

Hence,

$$\int_0^{N^\alpha} \rightarrow \int_0^\infty e^{-y^2 + 2\xi_0 y} y^{\sigma-1} dy.$$

The second integral inside the bracket in (33) goes to 0 as  $N \rightarrow \infty$ .

The numerator of (15), with  $\sigma$  in place of  $\sigma/m_k$ , is

$$(1/2)B(N + 1, \sigma/2) N^{\sigma/2}.$$

We thus have the following result.

**THEOREM 3.** *The Laplace transform of the distribution of  $L$  is given by*

$$(34) \quad \frac{(1/2)\Gamma(\sigma/2)}{\int_0^\infty e^{-y^2 + 2\xi_0 y} y^{\sigma-1} dy}.$$

Formula (34) was obtained by Siebert and by Darling through direct consideration of the  $z(t)$  process. It is interesting to notice that the present procedure utilizes (13) which has no counterpart for the  $z(t)$  process.

7. **Two-sided limits.** Let  $L_{N,k}^*$ ,  $N \geq k$ , be the first time  $|x(t) - N| = N - k$ , given  $x(0) = N$ . Let  $L^*$  be the first time  $|z(t)| = \xi_0 > 0$ , given  $z(0) = 0$ . Arguments similar to those used for Theorems 1 and 3 give the following two results.

**THEOREM 1a.** *Under the conditions of Theorem 1 the limiting distribution of  $L_{N,k}^*/m_k$  is  $1 - e^{-2u}$ ,  $u \geq 0$ .*

**THEOREM 3a.** *The distribution of  $L^*$  has the Laplace transform*

$$\frac{(1/2)\Gamma(\sigma/2)}{\int_0^\infty e^{-y^2} y^{\sigma-1} \cosh(2\xi_0 y) dy} .$$

8. **Added in proof.** An argument has been found which rigorizes the remarks of Section 5 and gives a proof of Theorems 1 and 2 for more general processes.

REFERENCES

1. P. and T. Ehrenfest, *Über zwei bekannte Einwände gegen das Boltzmannsche H-Theorem*, Phys. Z. 8 (1907), 311.
2. W. Feller, *Fluctuation theory of recurrent events*, Trans. Amer. Math. Soc. 67 (1949), 98-119.
3. ———, *Introduction to Probability*, Wiley, New York, 1950.
4. Bernard Friedman, *A simple urn model*, Comm. Pure Appl. Math. 11 (1949), 59.
5. Mark Kac, *Random walk and the theory of Brownian motion*, Amer. Math. Monthly 54 (1947), 369.
6. A. I. Khinchin, *Statistical Mechanics*, Dover, New York, 1949, pp.139-145.
7. A. Kolmogorov, *Analytischen methoden in der Wahrscheinlichkeitsrechnung*, Math. Ann. 104 (1931), 432.
8. G. Pólya and G. Szegő, *Aufgaben u. Lehrsätze I*, Dover, New York, 1945.
9. A. J. F. Siegert, *Note on the Ehrenfest problem*, Los Alamos Scientific Laboratory, MDDC-1406 (LADC-438).
10. ———, *On the first passage time problem (abstract)*, Physical Rev. 70 (1946), 449.
11. ———, *On the approach to statistical equilibrium*, Physical Rev. 76 (1949), 1708-1714.
12. J. V. Uspensky, *Introduction to Mathematical Probability*, McGraw-Hill, New York, 1937.

STANFORD UNIVERSITY  
THE RAND CORPORATION





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

HERBERT BUSEMANN  
University of Southern California  
Los Angeles 7, California

R. M. ROBINSON  
University of California  
Berkeley 4, California

E. F. BECKENBACH, Managing Editor  
University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

R. P. DILWORTH	P. R. HALMOS	BØRGE JESSEN	J. J. STOKER
HERBERT FEDERER	HEINZ HOPF	PAUL LÉVY	E. G. STRAUS
MARSHALL HALL	R. D. JAMES	GEORGE PÓLYA	KÔSAKU YOSIDA

## SPONSORS

UNIVERSITY OF BRITISH COLUMBIA	UNIVERSITY OF SOUTHERN CALIFORNIA
CALIFORNIA INSTITUTE OF TECHNOLOGY	STANFORD UNIVERSITY
UNIVERSITY OF CALIFORNIA, BERKELEY	WASHINGTON STATE COLLEGE
UNIVERSITY OF CALIFORNIA, DAVIS	UNIVERSITY OF WASHINGTON
UNIVERSITY OF CALIFORNIA, LOS ANGELES	* * *
UNIVERSITY OF CALIFORNIA, SANTA BARBARA	AMERICAN MATHEMATICAL SOCIETY
OREGON STATE COLLEGE	NATIONAL BUREAU OF STANDARDS,
UNIVERSITY OF OREGON	INSTITUTE FOR NUMERICAL ANALYSIS

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Spécial price to individual faculty members of supporting institutions and to members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1951 BY PACIFIC JOURNAL OF MATHEMATICS

Tom M. (Mike) Apostol, <i>On the Lerch zeta function</i> .....	161
Ross A. Beaumont and Herbert S. Zuckerman, <i>A characterization of the subgroups of the additive rationals</i> .....	169
Richard Bellman and Theodore Edward Harris, <i>Recurrence times for the Ehrenfest model</i> .....	179
Stephen P.L. Diliberto and Ernst Gabor Straus, <i>On the approximation of a function of several variables by the sum of functions of fewer variables</i> .....	195
Isidore Isaac Hirschman, Jr. and D. V. Widder, <i>Convolution transforms with complex kernels</i> .....	211
Irving Kaplansky, <i>A theorem on rings of operators</i> .....	227
W. Karush, <i>An iterative method for finding characteristic vectors of a symmetric matrix</i> .....	233
Henry B. Mann, <i>On the number of integers in the sum of two sets of positive integers</i> .....	249
William H. Mills, <i>A theorem on the representation theory of Jordan algebras</i> .....	255
Tibor Radó, <i>An approach to singular homology theory</i> .....	265
Otto Szász, <i>On some trigonometric transforms</i> .....	291
James G. Wendel, <i>On isometric isomorphism of group algebras</i> .....	305
George Milton Wing, <i>On the <math>L^p</math> theory of Hankel transforms</i> .....	313