

# Pacific Journal of Mathematics

**REMARKS ON THE SPACE  $H^p$**

STANLEY SIMON WALTERS

# REMARKS ON THE SPACE $H^p$

STANLEY S. WALTERS

**1. Introduction.** The space  $H^p$  is the collection of all single-valued complex functions  $f$  which are regular on the interior of the unit circle in the complex plane, and for which

$$\sup_{0 \leq r < 1} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta < \infty .$$

In [6] it was shown that  $H^p$ ,  $0 < p < 1$ , is a linear topological space in which the metric is  $\|f - g\|^p$ , where we define

$$\|f\| = \sup_{0 \leq r < 1} \left( \frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta})|^p d\theta \right)^{1/p} .$$

It was moreover shown that  $(H^p)^*$ , the conjugate of  $H^p$ , has sufficiently many elements (linear functionals on  $H^p$ ) so as to distinguish elements in  $H^p$ , in the sense that if  $f \neq 0$  is in  $H^p$ , then there is a  $\gamma \in (H^p)^*$  such that  $\gamma(f) \neq 0$ .

In the present paper it will be shown that if  $\gamma$  is in  $(H^p)^*$ ,  $0 < p < 1$ , then there exists a unique function  $G$  which is regular in the open unit circle, continuous on the closed circle,<sup>1</sup> and such that

$$\gamma(f) = \lim_{r=1} \frac{1}{2\pi} \int_0^{2\pi} f(\rho e^{i\theta}) G\left(\frac{r}{\rho} e^{-i\theta}\right) d\theta, \quad r < \rho < 1 ,$$

for every  $f$  in  $H^p$ . It is further shown that the following is true of  $G$ :

---

Received November 29, 1950.

The author is indebted to Professor Angus E. Taylor for his valuable assistance in the preparation of this paper.

<sup>1</sup>The author wishes to acknowledge the fact that the referee suggested the plausibility that  $G$  is continuous on the closed unit circle.

*Pacific J. Math.* 1 (1951), 455-471.

(a) if  $0 < p < 1/n$ ,  $n = 2, 3, \dots$ , then  $[d^{n-1} G(z)]/dz^{n-1}$  is continuous on the closure of the unit circle;

(b) if  $0 < p < 1/2n$ ,  $n = 1, 2, \dots$ , then  $G(e^{it})$  has a continuous  $n$ th derivative with respect to  $t$ ; and

(c) if  $0 < p < 1/2$ , then the power series for  $G$  converges absolutely on the boundary of the unit circle.

It is moreover shown that if  $G$  is regular on the open unit circle and is such that

$$\lim_{r=1} \frac{1}{2\pi} \int_0^{2\pi} f(\rho e^{i\theta}) G\left(\frac{r}{\rho} e^{-i\theta}\right) d\theta, \quad r < \rho < 1,$$

exists for every  $f$  in  $H^p$ , then the functional so defined is in  $(H^p)^*$ . Thus  $(H^p)^*$  is equivalent to a subspace of the functions which are regular on the open unit circle and continuous on the closed unit circle when  $0 < p < 1$ ; and indeed, as  $p$  tends toward zero, the spaces  $(H^p)^*$  are equivalent to subspaces of spaces whose members have far stronger properties than merely the property of being continuous on the closure of the unit circle.

A generalization of a theorem by Khintchine and Ostrowski [1, p.157], which is a sort of generalization of Vitali's theorem, will also be presented; namely, it will be shown that a bounded sequence in  $H^p$ ,  $0 < p < \infty$ , whose boundary values converge on a set of positive measure, converges uniformly on all compact subsets of the unit circle. Khintchine and Ostrowski proved this theorem in the case that the sequence consists of uniformly bounded elements.

It is worth remarking that under the present "norm"  $\|\cdot\|$ ,  $H^p$ ,  $0 < p < 1$ , is definitely *not* a normed linear space, this being due to the complete failure of Minkowski's inequality for index smaller than unity. As a result, it is conjectured by the author that  $H^p$ ,  $0 < p < 1$ , is not a normed linear space at all (and hence contains no bounded convex neighborhood). If this conjecture is true, then  $H^p$ ,  $0 < p < 1$ , offers an interesting example of a linear topological space which is not locally convex (since  $H^p$  is clearly locally bounded) and whose conjugate space has sufficiently many members so as to distinguish the elements in  $H^p$ .

**2. Representation of linear functionals on  $H^p$ ,  $0 < p < 1$ .** In this section we shall suppose always that  $0 < p < 1$ . We let  $\Delta$  be the set of all  $z$  such that  $|z| < 1$ , and  $\mathfrak{U}$  the class of all single-valued complex functions which are regular

on  $\Delta$ . We shall first make some definitions and prove several lemmas before proving the representation theorem.

For many of the topological terms used in the ensuing, see [3]. By a *complete linear topological space*, we shall mean a space in which  $f_n - f_m \rightarrow 0$  implies  $\lim_{n \rightarrow \infty} f_n$  exists in the space. *Locally bounded linear topological space* and *normed linear space* will be abbreviated LBLTS and NLS respectively. By  $F^*$ , where  $F$  is a linear topological space, we shall mean the conjugate of  $F$ , that is, the space of linear functionals on  $F$ .

If  $F$  is a LBLTS, it is easy to show that  $F^*$  is a complete NLS (Banach space) in which

$$\|\gamma\| = \sup_{f \in U} |\gamma(f)| ,$$

where  $\gamma \in F^*$ , and  $U$  is a fixed bounded neighborhood of the origin. Moreover, the topology so introduced into  $F^*$  is independent of  $U$ . With respect to  $H^P$ , we let  $U$  be the unit sphere, so that

$$\|\gamma\| = \sup_{\|f\|=1} |\gamma(f)| .$$

It is then simple to prove the following theorem, merely by modeling the proof exactly after that given in the theory of NLS's.

LEMMA 1. *If  $F$  is a complete LBLTS, and  $\Gamma$  is a subset of  $F^*$  having the property that, for each fixed  $f$  in  $F$ ,  $\gamma(f)$  is bounded as  $\gamma$  varies over  $\Gamma$ , then  $\Gamma$  is a bounded set.*

We remind ourselves that  $H^P$  is locally bounded, and is moreover complete by [6]. We make the following definitions, where  $f$  and  $g$  are any elements in  $\mathcal{U}$ :

$$(i) \quad \gamma_n(f) = f^{(n)}(0)/n! , \quad n = 0, 1, \dots ,$$

$$(ii) \quad T_w f : T_w f(z) = f(wz) , \quad w \in \Delta , z \in \Delta ,$$

$$(iii) \quad u_n : u_n(z) = z^n , \quad z \in \Delta , n = 0, 1, \dots ,$$

$$(iv) \quad B(f, g; z) = \sum_{n=0}^{\infty} \gamma_n(f) \gamma_n(g) z^n , \quad z \in \Delta .$$

It is easily verified that

$$B(f, g; z) = \frac{1}{2\pi} \int_0^{2\pi} f(z_1 e^{i\theta}) g(z_2 e^{-i\theta}) d\theta ,$$

where  $z_1 z_2 = z$ , and  $z_1$  and  $z_2$  are in  $\Delta$ . The proof is made by expansion of the integrand above in a Taylor series about the origin and then term-by-term integration. In particular,

$$B(f, g; r) = \frac{1}{2\pi} \int_0^{2\pi} f(\rho e^{i\theta}) g\left(\frac{r}{\rho} e^{-i\theta}\right) d\theta , \quad r < \rho < 1 .$$

LEMMA 2. *If  $f$  is in  $H^P$ , then  $T_w f$  is in  $H^P$ , and moreover*

$$T_w f = \sum_{n=0}^{\infty} \gamma_n(f) w^n u_n .$$

*Proof.* Let  $g = \sum_{n=0}^{\infty} \gamma_n(f) w^n u_n$ . We first show that this series converges. Note that  $\|u_n\| = 1$ , and

$$|\gamma_n(f)| \leq \left(\frac{pn + 1}{pn}\right) (pn + 1)^{1/p} \cdot \|f\| .$$

The last inequality appears in [6, Theorem 6]. Thus

$$\left\| \sum_{n=l}^m \gamma_n(f) w^n u_n \right\|^P \leq \sum_{n=l}^m \|\gamma_n(f) w^n u_n\|^P \rightarrow 0 \quad \text{as } l, m \rightarrow \infty ,$$

whence  $\sum_{n=0}^{\infty} \gamma_n(f) w^n u_n$  converges, by the completeness of  $H^P$ . Then, noting [6, Theorem 8], which tells us that a convergent sequence in  $H^P$  converges point-wise to its limit, we have

$$g(z) = \sum_{n=0}^{\infty} \gamma_n(f) w^n u_n(z) = \sum_{n=0}^{\infty} \gamma_n(f) (wz)^n$$

But  $T_w f(z) = \sum_{n=0}^{\infty} \gamma_n(f) (wz)^n$ . This completes the proof.

We note that it was obvious that  $T_w f$  was in  $H^P$  in the first place, merely from the definition of  $H^P$ ; but the form for  $T_w f$ , which was obtained above, will be

needed later.

**THEOREM 1.** *If  $G \in \mathfrak{A}$  such that  $\lim_{r=1} B(f, G; r) = \gamma(f)$  (that is, we assume that this limit exists) for all  $f$  in  $H^p$ , then  $\gamma$  is in  $(H^p)^*$ . Conversely, if  $\gamma$  is in  $(H^p)^*$ , then there exists a unique  $G$  in  $\mathfrak{A}$  such that  $\gamma(f) = \lim_{r=1} B(f, G; r)$  for all  $f$  in  $H^p$ .*

*Proof.* To prove the first part of our theorem, let  $\gamma_r(f) = B(f, G; r)$ . Clearly  $\gamma_r(f)$  is distributive in  $f$ . Suppose  $\|f\| = 1$  and  $r < p < 1$ . Then

$$|\gamma_r(f)| \leq \sum_{n=0}^{\infty} |\gamma_n(f)| \cdot |\gamma_n(G)| r^n \leq \sum_{n=0}^{\infty} |\gamma_n(G)| \left( \frac{pn+1}{pn} \right) (pn+1)^{1/p} r^n.$$

Thus,  $\gamma_r(f)$  is bounded in  $f$  for  $\|f\| = 1$ ,  $r$  being fixed. It is then clear that  $\gamma_r$  is in  $(H^p)^*$ . Since  $\lim_{r=1} \gamma_r(f)$  exists, it follows that  $\gamma_r(f)$  is continuous on  $0 \leq r \leq 1$  for each fixed  $f$  in  $H^p$ . Thus  $\{\gamma_r(f)\}$  is bounded for  $0 \leq r < 1$ . As a result of Lemma 1, we may conclude that  $\{\|\gamma_r\|\}$  is bounded for  $0 \leq r < 1$ ; that is, there exists an  $M$  such that  $\|\gamma_r\| \leq M$  for  $0 \leq r < 1$ . Let  $\|f\| = 1$ . Then  $|\gamma_r(f)| \leq M$ , whence  $|\gamma(f)| \leq M$ . Thus  $\gamma$  is necessarily in  $(H^p)^*$  since it is bounded on the unit sphere in  $H^p$ .

We now prove the second part of Theorem 1. We note that if  $\lim_{r=1} B(f, G; r) = \gamma(f)$  for some  $G$  and all  $f$ , then

$$\gamma(u_n) = \lim_{r=1} B(u_n, G; r) = \lim_{r=1} \gamma_n(G) r^n = \gamma_n(G);$$

that is,  $\gamma_n(G) = \gamma(u_n)$  for all  $n$ , or merely  $G(z) = \sum_{n=0}^{\infty} \gamma(u_n) z^n$ . We note that  $\sum_{n=0}^{\infty} \gamma(u_n) z^n$  converges, for  $|\gamma(u_n)| \leq \|\gamma\| \cdot \|u_n\| = \|\gamma\|$ . Let us now verify that  $G$ , as defined, has the desired property. We see that

$$B(f, G; r) = \sum_{n=0}^{\infty} \gamma_n(f) \gamma(u_n) r^n = \gamma \left\{ \sum_{n=0}^{\infty} \gamma_n(f) r^n u_n \right\} = \gamma(T_r f).$$

But  $\|T_r f - f\| \rightarrow 0$ ; see [5] for this result; note that

$$\|T_r f - f\| = \left( \frac{1}{2\pi} \int_0^{2\pi} |f(re^{i\theta}) - f(e^{i\theta})|^p d\theta \right)^{1/p},$$

where  $f(e^{i\theta})$  is the boundary function for  $f(z)$ . Thus  $\gamma(T_r f) \rightarrow \gamma(f)$ , or  $B(f, G; r) \rightarrow \gamma(f)$ . Our proof is thus complete.

**THEOREM 2.** *The function  $G$  in Theorem 1 is continuous on the closure of  $\Delta$ .*

*Proof.* We first verify that  $f_t(z) = (1 - ze^{it})^{-1}$  is in  $H^p$  for every real  $t$ . It suffices to show that  $f_0$  is in  $H^p$ . We see that

$$|1 - re^{i\theta}|^{-2} = [(1 - re^{i\theta})(1 - re^{-i\theta})]^{-1} = (1 - 2r \cos \theta + r^2)^{-1},$$

whence

$$|1 - re^{i\theta}|^{-p} = (1 - 2r \cos \theta + r^2)^{-p/2}.$$

From the character of  $(1 - 2r \cos \theta + r^2)$ , we see that it suffices to show that  $\int_0^\delta (1 - 2r \cos \theta + r^2)^{-p/2} d\theta$  is bounded in  $r$ , where  $\delta$  is any positive number. We note that the following is true for  $0 \leq \theta \leq \delta$  (where  $\delta$  is some sufficiently small positive number) and for all  $r$  such that  $1/2 \leq r < 1$ :

$$\begin{aligned} 1 - 2r \cos \theta + r^2 &\geq 1 - 2r \left(1 - \frac{\theta^2}{2} + \frac{\theta^4}{24}\right) + r^2 = (1 - 2r + r^2) + r\theta^2 \left(1 - \frac{\theta^2}{12}\right) \\ &= (1 - r)^2 + r\theta^2 \left(1 - \frac{\theta^2}{12}\right) \\ &\geq \frac{r\theta^2}{2} \geq \frac{\theta^2}{4}. \end{aligned}$$

Thus,  $(1 - 2r \cos \theta + r^2)^{-p/2} \leq 4^{p/2} \theta^{-p}$ . Since  $\theta^{-p}$  is integrable on  $[0, \delta]$ , our statement is proved.

We remind ourselves that we are trying to show that  $G$  is continuous on the assumption that

$$\gamma(f) = \lim_{r \rightarrow 1} \frac{1}{2\pi} \int_0^{2\pi} f(\rho e^{i\theta}) G\left(\frac{r}{\rho} e^{-i\theta}\right) d\theta, \quad r < \rho < 1,$$

exists for each  $f$  in  $H^p$ . Let  $\gamma_r$  be defined as in the proof of Theorem 1. Then

$$\begin{aligned}
 \gamma_r(f_t) &= \frac{1}{2\pi} \int_0^{2\pi} \frac{1}{1 - \rho e^{i(\theta+t)}} \cdot G\left(\frac{r}{\rho} e^{-i\theta}\right) d\theta \\
 &= \frac{1}{2\pi} \int_0^{2\pi} \frac{G\left(\frac{r}{\rho} e^{i\theta}\right)}{1 - \rho e^{i(t-\theta)}} d\theta \\
 &= \frac{1}{2\pi} \int_0^{2\pi} \frac{G\left(\frac{r}{\rho} e^{i\theta}\right) \frac{r}{\rho} e^{i\theta}}{\frac{r}{\rho} e^{i\theta} - r e^{it}} d\theta \\
 &= G(re^{it}), \qquad r < \rho < 1.
 \end{aligned}$$

The last equality is true by virtue of Cauchy's integral formula. We then have shown that  $G(re^{it}) = \gamma_r(f_t)$ . Consequently, since  $\lim_{r=1} \gamma_r(f_t)$  exists by hypothesis,  $\lim_{r=1} G(re^{it})$  exists for all  $t$ , and in fact

$$\gamma(f_t) = G(e^{it}),$$

where we define  $G(e^{it})$  to be the boundary function  $\lim_{r=1} G(re^{it})$ .

We now show that  $\lim_{t=t_0} f_t = f_{t_0}$  in the topology of  $H^p$ . Now, for any  $g$  in  $H^p$ , letting  $g(e^{i\theta})$  be its boundary function, we know that

$$\sup_{0 \leq r < 1} \int_0^{2\pi} |g(re^{i\theta})|^p d\theta = \int_0^{2\pi} |g(e^{i\theta})|^p d\theta.$$

It is easily verified that (see, for example, [4, Theorem 7, p. 29])

$$\lim_{t=t_0} \int_0^{2\pi} |g(e^{i(\theta+t)}) - g(e^{i(\theta+t_0)})|^p d\theta = 0.$$

Clearly  $f_t(e^{i\theta}) = (1 - e^{i(\theta+t)})^{-1}$ , whence  $f_t(e^{i\theta}) = f_0(e^{i(\theta+t)})$ . Thus  $\lim_{t=t_0} f_t = f_{t_0}$ , in the topology of  $H^p$ .



Now, by Theorem 1,  $\gamma$  is continuous, whence  $\lim_{t \rightarrow t_0} \gamma(f_t) = \gamma(f_{t_0})$ ; hence  $\lim_{t \rightarrow t_0} G(e^{it}) = G(e^{it_0})$ . We have now shown that  $G(e^{it})$  is continuous.

We remember that, in the course of proving Theorem 1, we showed that  $\{\gamma_r\}$  is bounded in  $r$  as a subset of  $(H^p)^*$ . Obviously  $\{f_t\}$  is a bounded subset of  $H^p$ , all of the elements having the same norm. Thus  $\gamma_r(f_t)$  is bounded in both  $r$  and  $t$ . In other words,  $G(re^{it})$  is bounded in  $r$  and  $t$ , or equivalently  $G$  is uniformly bounded on  $\Delta$ . We then know that

$$G(re^{it}) = \frac{1}{2\pi} \int_0^{2\pi} G(e^{it}) P_r(\theta - t) d\theta,$$

where  $P_r(\theta)$  is the Poisson kernel. But, since  $G(e^{it})$  is continuous, the right side above is necessarily a continuous function on the closed unit circle. Our proof is now complete.

It will now be shown that even more can be said of  $G$  when  $0 < p < 1/2$ .

**THEOREM 3.** *If  $0 < p < 1/2$ , then  $G(e^{it})$  satisfies the Lipschitz condition of order one.*

*Proof.* It suffices to show that

$$\|f_{t+h} - f_t\| = \|f_h - f_0\| \leq A \cdot |1 - e^{ih}|$$

for some fixed constant  $A$ . We have

$$\begin{aligned} \|f_h - f_0\| &= \left( \frac{1}{2\pi} \int_0^{2\pi} \left| \frac{1}{1 - e^{i(\theta+h)}} - \frac{1}{1 - e^{i\theta}} \right|^p d\theta \right)^{1/p} \\ &= \frac{|1 - e^{ih}|}{(2\pi)^{1/p}} \left\{ \int_0^{2\pi} |(1 - e^{i(\theta+h)})(1 - e^{i\theta})|^{-p} d\theta \right\}^{1/p}, \end{aligned}$$

The proof will then be complete after we have shown that

$$\int_0^{2\pi} |(1 - e^{i(\theta+h)})(1 - e^{i\theta})|^{-p} d\theta$$

is bounded for all sufficiently small  $h$ . It is evident that

$$|(1 - e^{i\theta})(1 - e^{i(\theta+h)})|^2 = 4(1 - \cos \theta)[1 - \cos(\theta + h)],$$

and hence

$$|(1 - e^{i\theta})(1 - e^{i(\theta+h)})|^p = 4^{p/2} (1 - \cos \theta)^{p/2} (1 - \cos (\theta + h))^{p/2}.$$

We now must show that

$$\int_0^{2\pi} (1 - \cos \theta)^{-p/2} (1 - \cos (\theta + h))^{-p/2} d\theta$$

is bounded in  $h$  for all sufficiently small  $h$ . We note that the following is true for all sufficiently small  $\theta$  and  $h$ :

$$1 - \cos \theta \geq \frac{\theta^2}{2} \left(1 - \frac{\theta^2}{12}\right) \geq \frac{\theta^2}{4},$$

$$1 - \cos (\theta + h) \geq \frac{(\theta + h)^2}{4}.$$

Thus we have

$$(1 - \cos \theta)^{-p/2} [1 - \cos (\theta + h)]^{-p/2} \leq 4^p \theta^{-p} (\theta + h)^{-p}$$

for all sufficiently small  $\theta$  and  $h$ . Since  $\theta^{-2p}$  is integrable on the interval  $[0, 2\pi]$ , it is then rather easy to show that

$$\int_0^{2\pi} (1 - \cos \theta)^{-p/2} [1 - \cos (\theta + h)]^{-p/2} d\theta$$

is bounded in  $h$  for all sufficiently small  $h$ .

We now have the rather interesting result:

**COROLLARY.** *If  $0 < p < 1/2$ , then  $\sum_{n=0}^{\infty} |\gamma_n(G)| < \infty$ .*

*Proof.* Since  $G(e^{it})$  is of bounded variation, it follows that  $G(z)$  is a power series of bounded variation according to [7, § 7.5]. Hence the conclusion is obtained by [7, (i), p. 158].

We shall now show that even more may be said of  $G$  when  $0 < p < 1/2$ .

**THEOREM 4.** *If  $0 < p < 1/2$ , then  $(d/dz)G(z)$  is continuous on the closure of*

$\Delta$ , and moreover  $(d/dt)G(e^{it})$  is continuous on  $[0, 2\pi]$ .

*Proof.* By Cauchy's integral formulas (where  $(d/dz)G(z) = G'(z)$ ):

$$\begin{aligned} G'(re^{it}) &= \frac{1}{2\pi} \int_0^{2\pi} \frac{G\left(\frac{r}{\rho} e^{i\theta}\right) \frac{r}{\rho} e^{i\theta}}{\left(\frac{r}{\rho} e^{i\theta} - re^{it}\right)^2} d\theta = \frac{1}{2\pi} \int_0^{2\pi} \frac{G\left(\frac{r}{\rho} e^{-i\theta}\right) \frac{\rho}{r} e^{i\theta}}{(1 - \rho e^{i(\theta+t)})^2} d\theta \\ &= \frac{1}{2\pi r} \int_0^{2\pi} [f_t^2(\rho e^{i\theta}) \cdot \rho e^{i\theta}] \cdot G\left(\frac{r}{\rho} e^{-i\theta}\right) d\theta, \quad r < \rho < 1. \end{aligned}$$

Thus  $G'(re^{it}) = (1/r)\gamma_r(f_t^2 u_1)$ . We note that since  $0 < p < 1/2$ , we have  $f_t^2 \in H^p$ , whence  $f_t^2 \cdot u_1 \in H^p$ , since  $u_1$  is bounded. Thus we show exactly as in Theorem 2 that

$$\gamma(f_t^2 u_1) = G'(e^{it}),$$

$G'(e^{it})$  is continuous in  $t$ ,

$G'(z)$  is uniformly bounded on  $\Delta$ ,

$$G'(re^{it}) = \frac{1}{2\pi} \int_0^{2\pi} G'(e^{i\theta}) P_r(\theta - t) d\theta,$$

where we define  $G'(e^{it})$  to be the boundary value of  $G'(z)$ . Let us now consider

$$F(re^{it}) = \frac{1}{2\pi} \int_0^{2\pi} \left[ -ie^{-i\theta} \frac{d}{d\theta} G(e^{i\theta}) \right] P_r(\theta - t) d\theta.$$

We note that  $G(e^{i\theta})$  is absolutely continuous by virtue of Theorem 2, whence  $(d/d\theta)G(e^{i\theta})$  is integrable. We also note that

$$G(re^{it}) = \frac{1}{2\pi} \int_0^{2\pi} G(e^{i\theta}) P_r(\theta - t) d\theta = \sum_{n=0}^{\infty} C_n r^n e^{int},$$

where

$$C_n = \frac{1}{2\pi} \int_0^{2\pi} G(e^{i\theta}) e^{-in\theta} d\theta .$$

Moreover, it is not at all difficult to verify that the real and imaginary parts of  $-ie^{i\theta}(d/d\theta)G(e^{i\theta})$  are conjugate, whence

$$F(re^{it}) = \sum_{n=0}^{\infty} d_n r^n e^{int} ,$$

where

$$d_n = \frac{1}{2\pi} \int_0^{2\pi} \left[ -ie^{-i\theta} \frac{d}{d\theta} G(e^{i\theta}) \right] e^{-in\theta} d\theta .$$

Integration by parts readily yields

$$d_n = (n+1) C_n ;$$

that is,

$$F(re^{it}) = \sum_{n=0}^{\infty} (n+1) C_{n+1} r^n e^{int} ,$$

and hence  $F(z) = G'(z)$ . Thus, we necessarily have

$$G'(e^{i\theta}) = -ie^{-i\theta} \frac{d}{d\theta} G(e^{i\theta})$$

almost everywhere. Since  $G'(e^{i\theta})$  is continuous, it follows that  $G(e^{i\theta})$  necessarily has a continuous derivative, and in fact

$$\frac{d}{d\theta} G(e^{i\theta}) = ie^{i\theta} G'(e^{i\theta}) .$$

This completes the proof of the theorem.

We sum up by presenting the following theorem, which is readily proved by induction, the proof being modeled after that given for Theorem 4.

**THEOREM 5.** *If  $0 < p < 1/n, n = 2, 3, \dots$ , then  $(d^{n-1}/dz^{n-1}) G(z)$  is continuous on the closure of  $\Delta$ . Moreover, if  $0 < p < 1/2n, n = 1, 2, \dots$ , then  $G(e^{it})$  has a continuous  $n$ th derivative with respect to  $t$ .*

**3. Generalization of Vitali's Theorem.** In this section we assume merely that  $p$  is any positive real number. We here need the following:

**LEMMA 3.** *If  $\{f_n\}$  is a bounded sequence in  $H^p$ , and if  $\lim_{n=\infty} f_n(z)$  exists on a set having at least one limit point in  $\Delta$ , then  $\lim_{n=\infty} f_n(z)$  exists uniformly on all compact subsets of  $\Delta$ .*

*Proof.* The proof is a simple consequence of the following inequalities:

$$|f(z)| \leq \frac{\|f\|}{(1-|z|)^{1/p}}, \quad \text{when } 0 < p \leq 1,$$

and

$$|f(z)| \leq \frac{\|f\|}{1-|z|} \quad \text{when } 1 \leq p < \infty.$$

The first of the above inequalities appears in [6, Theorem 2]. The second is easily obtained as follows. By Cauchy's integral formula,

$$f(z) = \frac{1}{2\pi} \int_0^{2\pi} \frac{f(\rho e^{i\theta}) \rho e^{i\theta}}{\rho e^{i\theta} - z} d\theta,$$

and hence, by Hölder's inequality,

$$\begin{aligned} |f(z)| &\leq \frac{\rho}{\rho - |z|} \frac{1}{2\pi} \int_0^{2\pi} |f(\rho e^{i\theta})| d\theta \\ &\leq \frac{\rho}{\rho - |z|} \left( \frac{1}{2\pi} \int_0^{2\pi} |f(\rho e^{i\theta})|^p d\theta \right)^{1/p}, \end{aligned}$$

whence

$$|f(z)| \leq \frac{\|f\|}{1-|z|}.$$

Let

$$N(r) = \begin{cases} \frac{1}{(1-r)^{1/p}}, & 0 < p \leq 1. \\ \frac{1}{1-r}, & 1 \leq p < \infty. \end{cases}$$

It is then clear that  $|f_n(z)| \leq N(r) \cdot M$  when  $|z| \leq r < 1$ , where  $\|f_n\| \leq M$  for all  $n$ . We choose  $r$  so large that the set  $|z| < r$  includes a set having a limit point in  $|z| < r$  and such that  $\lim_{n \rightarrow \infty} f_n(z)$  exists on this set. Then, by Vitali's theorem,  $\lim_{n \rightarrow \infty} f_n(z)$  exists uniformly on all compact subsets of  $|z| < r$ , and hence on all compact subsets of  $\Delta$ . This completes the proof.

**THEOREM 6.** *Suppose  $\{f_n\}$  is a bounded sequence in  $H^p$ . Further, suppose  $\lim_{n \rightarrow \infty} f_n(e^{i\theta})$  exists on a set of positive measure in the interval  $[0, 2\pi]$ . Then  $\lim_{n \rightarrow \infty} f_n(z)$  exists uniformly on all compact subsets of  $\Delta$ .*

*Proof.* It suffices, by the preceding lemma, to show that  $\lim_{n \rightarrow \infty} f_n(z)$  exists on some neighborhood of the origin. Thus, we shall show that this is the case whenever  $|z| < 1/9$ . Let  $|z_0| < 1/9$ , and suppose  $\lim_{n \rightarrow \infty} f_n(z_0)$  does not exist. Then we may find a positive number  $\alpha$  and subsequences  $\{f_{n_k}\}$  and  $\{f_{m_k}\}$  of  $\{f_n\}$  which have the property that  $|f_{n_k}(z_0) - f_{m_k}(z_0)| > \alpha$  for all  $k$ . We then define  $q_k = f_{n_k} - f_{m_k}$ . It is clear that  $\{q_k\}$  is a bounded sequence in  $H^p$ . We then write  $q_k = g_k \cdot h_k$ , by virtue of F. Riesz's decomposition theorem [5], where  $g_k$  and  $h_k$  are such that

- (i)  $g_k \in H^p$  and  $g_k(z) \neq 0$  for all  $z$  in  $\Delta$ ,
- (ii)  $|h_k(z)| \leq 1$  on  $\Delta$  and  $|h_k(e^{i\theta})| = 1$  almost everywhere,
- (iii)  $\|g_k\| = \|q_k\|$ .

We note that  $l_k(z) = [g_k(z)]^{p/2}$  is in  $H^2$ , and in fact  $\{l_k\}$  is a bounded sequence in  $H^2$ . Since  $\lim_{k \rightarrow \infty} [f_{n_k}(e^{i\theta}) - f_{m_k}(e^{i\theta})] = 0$  on a set of positive measure, it follows that  $\lim_{k \rightarrow \infty} l_k(e^{i\theta}) = 0$  on a set  $E$  of measure  $\mu > 0$ . We next shall show that  $\lim_{k \rightarrow \infty} l_k(z_0) = 0$ , which will in turn imply that  $\lim_{k \rightarrow \infty} g_k(z_0) = 0$ , and hence imply  $\lim_{k \rightarrow \infty} q_k(z_0) = 0$ , a contradiction to  $|q_k(z_0)| > \alpha$  for all  $k$ .

Let  $A > 0$ , and define

$$\phi(\theta) = \begin{cases} A/\mu & \text{on } E \\ A/(\mu - 2\pi) & \text{on } CE, \end{cases}$$

where  $CE$  is the set  $[0, 2\pi] - E$ . There is no loss in supposing that  $\mu \leq \pi$ . Define

$$u_0(re^{i\theta}) = \frac{1}{2\pi} \int_0^{2\pi} \phi(t) P_r(\theta - t) dt,$$

where  $P_r(\theta)$  is the Poisson kernel. Then  $u_0$  is harmonic in  $\Delta$  and  $\lim_{r \rightarrow 1} u_0(re^{i\theta}) = \phi(\theta)$  a.e., by virtue of Fatou's theorem; see [7, § 3.442]. Let

$$u(re^{i\theta}) = u_0(re^{i\theta}) - u_0(z_0).$$

We note that

$$u_0(re^{i\theta}) = \frac{a_0}{2} + \sum_{n=1}^{\infty} r^n (a_n \cos n\theta + b_n \sin n\theta),$$

where  $\{a_n, b_n\}$  are the Fourier coefficients of  $\phi(\theta)$ . Since  $u_0(0) = 0$ , this being due to the fact that  $\int_0^{2\pi} \phi(t) dt = 0$  and  $P_0(\theta - t) = 1$ , we then have  $a_0$  equal to zero, or

$$u_0(re^{i\theta}) = \sum_{n=1}^{\infty} r^n (a_n \cos n\theta + b_n \sin n\theta).$$

We note that  $|a_n| \leq 2A/\pi$  as well as  $|b_n| \leq 2A/\pi$ , whence

$$|u_0(re^{i\theta})| \leq \frac{4A}{\pi} \sum_{n=1}^{\infty} r^n = \frac{4A}{\pi} \frac{r}{1-r} < \frac{A}{2\pi} \leq \frac{A}{2(2\pi - \mu)}$$

provided  $0 \leq r < 1/9$ .

Let  $v(z)$  be the harmonic conjugate of  $u(z)$  which vanishes at  $z_0$ , and define  $g(z) = e^{u(z)+iv(z)}$ . Then  $g \in \mathfrak{U}$ , and  $g(z_0) = 1$ . Moreover, since  $|g(z)| = e^{u(z)}$ , we have  $\lim_{r \rightarrow 1} |g(re^{i\theta})| = e^{\phi(\theta) - u_0(z_0)}$ . By Cauchy's integral formula

we have

$$l_k(z_0) = \frac{1}{2\pi} \int_0^{2\pi} l_k(\rho e^{i\theta}) g(\rho e^{i\theta}) \frac{\rho e^{i\theta}}{\rho e^{i\theta} - z_0} d\theta, \quad |z_0| < \rho < 1.$$

This is true since  $l_k(z_0) = l_k(z_0) g(z_0)$ . We note that  $u(z)$  is bounded in  $\Delta$ , and hence so is  $g(z)$ . Since

$$\lim_{\rho=1} \int_0^{2\pi} |l_k(\rho e^{i\theta}) - l_k(e^{i\theta})|^2 d\theta = 0,$$

and since  $g(z)$  is bounded on  $\Delta$ , it is then evident that

$$\lim_{\rho=1} \int_0^{2\pi} |l_k(\rho e^{i\theta}) g(\rho e^{i\theta})| d\theta = \int_0^{2\pi} |l_k(e^{i\theta}) g(e^{i\theta})| d\theta.$$

Hence

$$|l_k(z_0)| \leq \frac{1}{2\pi} \frac{1}{1 - |z_0|} \int_0^{2\pi} |l_k(e^{i\theta})| e^{\phi(\theta) - u_0(z_0)} d\theta.$$

Consequently

$$\begin{aligned} |l_k(z_0)| &\leq e^{A/\mu - u_0(z_0)} \frac{1}{2\pi} \left( \frac{1}{1 - |z_0|} \right) \int_E |l_k(e^{i\theta})| d\theta \\ &\quad + e^{A/(\mu - 2\pi) - u_0(z_0)} \left( \frac{1}{2\pi} \right) \left( \frac{1}{1 - |z_0|} \right) \int_{CE} |l_k(e^{i\theta})| d\theta. \end{aligned}$$

Since

$$\frac{1}{2\pi} \int_{CE} |l_k(e^{i\theta})| d\theta \leq \frac{1}{2\pi} \int_0^{2\pi} |l_k(e^{i\theta})| d\theta \leq \left( \frac{1}{2\pi} \int_0^{2\pi} |l_k(e^{i\theta})|^2 d\theta \right)^{1/2}$$

and since  $\{l_k\}$  is a bounded subset of  $H^2$ , we see that

$$\frac{1}{2\pi} \int_{CE} |l_k(e^{i\theta})| d\theta$$



is bounded with respect to  $k$ . Moreover

$$\begin{aligned} \frac{A}{\mu - 2\pi} - u_0(z_0) &\leq \frac{A}{\mu - 2\pi} + |u_0(z_0)| \leq \frac{A}{\mu - 2\pi} + \frac{4A}{\pi} \left( \frac{|z_0|}{1 - |z_0|} \right) \\ &\leq \frac{A}{\mu - 2\pi} + \frac{A}{2(2\pi - \mu)} = \frac{A}{2(\mu - 2\pi)}. \end{aligned}$$

By virtue of Schwarz's inequality, where  $\xi$  is an arbitrary measurable subset of  $[0, 2\pi]$ , we have

$$\int_{\xi} |l_k(e^{i\theta})| d\theta \leq [m(\xi)]^{1/2} \left( \int_{\xi} |l_k(e^{i\theta})|^2 d\theta \right)^{1/2}.$$

Hence, by a convergence theorem of Lebesgue (see [2, p. 190]), we have

$$\lim_{k \rightarrow \infty} \int_E |l_k(e^{i\theta})| d\theta = 0,$$

since  $\lim_{k \rightarrow \infty} l_k(e^{i\theta}) = 0$  on  $E$ . Now, for arbitrary  $\epsilon > 0$ , we choose  $A$  so large that

$$e^{A/[2(\mu - 2\pi)]} \cdot \left( \frac{1}{2\pi} \right) \left( \frac{1}{1 - |z_0|} \right) \int_{CE} |l_k(e^{i\theta})| d\theta < \epsilon/2.$$

and hence we obtain, from the foregoing,

$$e^{A/[\mu - 2\pi]} - u_0(z_0) \left( \frac{1}{2\pi} \right) \left( \frac{1}{1 - |z_0|} \right) \int_{CE} |l_k(e^{i\theta})| d\theta < \epsilon/2.$$

Having so chosen  $A$ , choose  $K$  so large that  $k > K$  implies

$$e^{A/\mu - u_0(z_0)} \left( \frac{1}{2\pi} \right) \left( \frac{1}{1 - |z_0|} \right) \int_E |l_k(e^{i\theta})| d\theta < \epsilon/2.$$

Hence,  $k > K$  implies  $|l_k(z_0)| < \epsilon/2 + \epsilon/2 = \epsilon$ . This completes the proof of the theorem.

## REFERENCES

1. L. Bieberbach, *Lehrbuch der Funktionentheorie*, vol. II, Leipzig, 1931.
2. L. M. Graves, *Theory of functions of real variables*, New York, 1946.
3. D. H. Hyers, *Linear topological spaces*, Bull. Amer. Math. Soc. **51** (1945), 1-24.
4. J. E. Littlewood, *Theory of functions*, Cambridge, England, 1944.
5. F. Riesz, *Über die Randwerte einer analytischen Funktion*, Math. Zeit. **18** (1923), 87-95.
6. S. S. Walters, *The space  $H^p$  with  $0 < p < 1$* , Proc. Amer. Math. Soc. **1** (1950), 800-805.
7. A. Zygmund, *Trigonometrical series*, Warsaw, 1935.

UNIVERSITY OF CALIFORNIA, LOS ANGELES



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

HERBERT BUSEMANN  
University of Southern California  
Los Angeles 7, California

R. M. ROBINSON  
University of California  
Berkeley 4, California

E. F. BECKENBACH, Managing Editor  
University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

R. P. DILWORTH	P. R. HALMOS	BØRGE JESSEN	J. J. STOKER
HERBERT FEDERER	HEINZ HOPF	PAUL LÉVY	E. G. STRAUS
MARSHALL HALL	R. D. JAMES	GEORGE PÓLYA	KÔSAKU YOSIDA

## SPONSORS

UNIVERSITY OF BRITISH COLUMBIA	UNIVERSITY OF SOUTHERN CALIFORNIA
CALIFORNIA INSTITUTE OF TECHNOLOGY	STANFORD UNIVERSITY
UNIVERSITY OF CALIFORNIA, BERKELEY	WASHINGTON STATE COLLEGE
UNIVERSITY OF CALIFORNIA, DAVIS	UNIVERSITY OF WASHINGTON
UNIVERSITY OF CALIFORNIA, LOS ANGELES	* * *
UNIVERSITY OF CALIFORNIA, SANTA BARBARA	AMERICAN MATHEMATICAL SOCIETY
OREGON STATE COLLEGE	NATIONAL BUREAU OF STANDARDS,
UNIVERSITY OF OREGON	INSTITUTE FOR NUMERICAL ANALYSIS

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Spécial price to individual faculty members of supporting institutions and to members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1951 BY PACIFIC JOURNAL OF MATHEMATICS

R. P. Boas, <i>Completeness of sets of translated cosines</i> .....	321
J. L. Brenner, <i>Matrices of quaternions</i> .....	329
Edmond Darrell Cashwell, <i>The asymptotic solutions of an ordinary differential equation in which the coefficient of the parameter is singular</i> .....	337
James Dugundji, <i>An extension of Tietze's theorem</i> .....	353
John G. Herriot, <i>The polarization of a lens</i> .....	369
J. D. Hill, <i>The Borel property of summability methods</i> .....	399
G. G. Lorentz, <i>On the theory of spaces <math>\Lambda</math></i> .....	411
J. H. Roberts and W. R. Mann, <i>On a certain nonlinear integral equation of the Volterra type</i> .....	431
W. R. Utz, <i>A note on unrestricted regular transformations</i> .....	447
Stanley Simon Walters, <i>Remarks on the space <math>H^p</math></i> .....	455
Hsien Chung Wang, <i>Two theorems on metric spaces</i> .....	473