

Pacific Journal of Mathematics

**ON THE REALIZABILITY OF HOMOTOPY GROUPS AND
THEIR OPERATIONS**

SZE-TSEN HU

ON THE REALIZABILITY OF HOMOTOPY GROUPS AND THEIR OPERATIONS

SZE-TSEN HU

1. Introduction. Let B be a given arcwise connected topological space and b_0 a basic point of B . Then we obtain a sequence of homotopy groups

$$\pi_1(B), \pi_2(B), \dots, \pi_n(B), \dots.$$

The fundamental group $\pi_1(B)$ is in general non-abelian and written multiplicatively. All higher homotopy groups $\pi_n(B)$, $n \geq 2$, are abelian and written additively. The group $\pi_1(B)$ operates on the left of every higher homotopy group $\pi_n(B)$, $n \geq 2$; that is to say, for every $w \in \pi_1(B)$ and every $a \in \pi_n(B)$, a unique element $wa \in \pi_n(B)$ is determined, and

$$w(a_1 + a_2) = wa_1 + wa_2, \quad w_1(w_2a) = (w_1w_2)a, \quad 1a = a.$$

For arbitrary elements $a \in \pi_m(B)$ and $b \in \pi_n(B)$, $m \geq 2$, $n \geq 2$, a Whitehead product $a \circ b$ is defined [10, p. 411], which is an element of $\pi_{m+n-1}(B)$. The Whitehead product is known to be bilinear; namely,

$$(a_1 + a_2) \circ b = a_1 \circ b + a_2 \circ b, \quad a \circ (b_1 + b_2) = a \circ b_1 + a \circ b_2.$$

Roughly speaking, the realizability problem is whether these homotopy groups and mutual operations described above are otherwise completely arbitrary. It can be formulated precisely as follows. Let

$$\pi_1, \pi_2, \dots, \pi_n, \dots$$

be a given sequence of abstract groups. All groups except the first one are abelian and additive, while π_1 is written multiplicatively. There are given two kinds of operations between these groups. First, the group π_1 operates on the left of every group π_n with $n \geq 2$. Secondly, for arbitrary elements $\alpha \in \pi_m$, $\beta \in \pi_n$, $m \geq 2$, $n \geq 2$, a bilinear product $\alpha \circ \beta$ is defined and is an element of the group π_{m+n-1} .

Received October 8, 1950, and in revised form February 28, 1951. Presented to the American Mathematical Society, October 28, 1950.

Pacific J. Math. 1 (1951), 583-602.

The realizability problem is to construct an arcwise connected topological space B and a basic point $b_0 \in B$ satisfying the following conditions:

(1.1) *There exists, for each integer $n \geq 1$, an isomorphism $h_n: \pi_n(B) \approx \pi_n$ of $\pi_n(B)$ onto π_n .*

(1.2) *For arbitrary elements $w \in \pi_1(B)$ and $a \in \pi_n(B)$, $n \geq 2$, we have $h_n(wa) = h_1(w)h_n(a)$.*

(1.3) *For arbitrary elements $a \in \pi_m(B)$ and $b \in \pi_n(B)$, $m \geq 2$, $n \geq 2$, we have $h_{m+n-1}(a \circ b) = h_m(a) \circ h_n(b)$.*

This general problem has not yet been solved. The first partial solution was given by J. H. C. Whitehead [12]. By means of an inductive construction based on his previous contributions, he succeeded to give an infinite polytope B which satisfies the conditions (1.1) and (1.2). However, he gave no explicit information as to the Whitehead products of the higher homotopy groups of the space he constructed.

The object of the present work is to give a synthetic and algebraic construction of an arcwise connected topological space B with a basic point b_0 and prove the following:

REALIZABILITY THEOREM. *There exists an arcwise connected topological space B and a basic point $b_0 \in B$ satisfying the conditions (1.1), (1.2), and*

(1.4) *For arbitrary elements $a \in \pi_m(B)$ and $b \in \pi_n(B)$, $m \geq 2$, $n \geq 2$, we have $a \circ b = 0$.*

Our principal construction is motivated by the following observations:

(a) Let π be a given group and n a positive integer (we assume π to be abelian if $n > 1$). Then we can construct an arcwise connected space $P(\pi, n)$ such that:

$$(1.5) \quad \pi_n(P(\pi, n)) \approx \pi, \quad \pi_i(P(\pi, n)) = 0 \quad (i \neq n);$$

(1.6) If $n > 1$, there is a correspondence which associates with each endomorphism $h: \pi \rightarrow \pi$ a continuous map $h^\#: P(\pi, n) \rightarrow P(\pi, n)$ such that $(h_1 h_2)^\# = h_1^\# h_2^\#$, and $h^\#$ is the identity if h is the identity.

(b) Let π_2, π_3, \dots , be a sequence of abelian groups, and let Y denote the topological product of all the spaces $P(\pi_n, n)$, $n = 2, 3, \dots$. Then Y is simply

connected and $\pi_i(Y) \approx \pi_i$ for $i \geq 2$; moreover, all the Whitehead products in Y vanish. This is a consequence of J. H. C. Whitehead [13, p.289].

(c) Let G be a group of homeomorphisms of Y , and let $X = P(\pi_1, 1)$ where π_1 is any given group. Let $\chi: \pi_1 \rightarrow G$ be a homomorphism. Let \tilde{X} denote the universal covering space of X . It is well known that \tilde{X} is a bundle space over X with discrete fiber π_1 and discrete structural group π_1 . The homomorphism $\chi: \pi_1 \rightarrow G$ induces a bundle space B over X with fiber Y and structural group G which is weakly associated with \tilde{X} . Then the operations of π_1 on $\pi_n = \pi_n(Y)$ are given by $w \rightarrow \chi_*(w)$, where $\chi_*(w)$ is the automorphism of $\pi_n(Y)$ induced by the map $\chi(w): Y \rightarrow Y$. By suitable choice of the homomorphism $\chi: \pi_1 \rightarrow G$, the bundle space B has the properties described in the Realizability Theorem.

As an application, we are able to show that Whitehead products of the higher homotopy groups of a given topological space are essential invariants of the space; that is, they are not completely determined by the homotopy groups and the operations of the fundamental group upon the higher homotopy groups.

2. Semi-simplicial polytope. First of all, let us recall the definition of semi-simplicial complexes of S. Eilenberg and J. A. Zilber [2] as what follows.

A *semi-simplicial complex* K is a collection of elements $\{\sigma\}$ called *simplexes* together with two functions. The first function associates with each simplex σ an integer $q \geq 0$ called the dimension of σ ; we then say that σ is a q -simplex. The second function associates with each q -simplex σ ($q > 0$) of K and with each i ($0 \leq i \leq q$) a $(q-1)$ -simplex $\sigma^{(i)}$ called the i -th *face* of σ , subject to the condition

$$(2.1) \quad [\sigma^{(j)}]^{(i)} = [\sigma^{(i)}]^{(j-1)}$$

for $q > 1$ and $i < j$. We may pass to lower dimensional faces of σ by iteration. If $0 \leq i_1 < \dots < i_n \leq q$ then we define inductively

$$\sigma^{(i_1, \dots, i_n)} = [\sigma^{(i_2, \dots, i_n)}]^{(i_1)}.$$

This is a $(q-n)$ -simplex. If $0 \leq j_0 < \dots < j_{q-n} \leq q$ is the set complementary to $\{i_1, \dots, i_n\}$ then we also write

$$\sigma^{(i_1, \dots, i_n)} = \sigma_{(j_0, \dots, j_{q-n})}.$$

In particular, $\sigma_{(i)}$ for $0 \leq i \leq q$ is a 0-simplex called the i -th *vertex* of σ . We shall

also refer to $\sigma_{(0)}$ as the *leading vertex* and $\sigma_{(0,1)}$ as the *leading edge*. For any two simplexes σ and τ of K , we shall write

$$\tau < \sigma$$

if either $\tau = \sigma$ or $\tau = \sigma^{(i_1, \dots, i_n)}$ for some set (i_1, \dots, i_n) of integers $0 \leq i_1 < \dots < i_n \leq q$. A subcomplex L of K is a subcollection of simplexes of K with the property that $\sigma \in L$ and $\tau < \sigma$ imply $\tau \in L$. Obviously, every semi-simplicial complex K is a closure finite abstract complex [5, p.91] with its incidence numbers defined by means of the bounding relation

$$\partial\sigma = \sum_{i=0}^q (-1)^i \sigma^{(i)} .$$

Now, let K be a given semi-simplicial complex. We shall construct a topological space $P(K)$, called the *semi-simplicial polytope associated with K* .

For every integer $q \geq 0$, to every q -simplex σ of K let us associate an open geometric q -cell w_σ , called the *open q -cell corresponding to σ* , which is the interior of some ordered geometric q -simplex s_σ ; that is,

$$w_\sigma = \text{Int } s_\sigma, \quad s_\sigma = \langle v_0, \dots, v_q \rangle .$$

If s_σ is 0-dimensional, we define $\text{Int } s_\sigma = s_\sigma$. We assume that no two of these open cells $\{w_\sigma \mid \sigma \in K\}$ have a point in common. Let each open cell w_σ have the euclidean topology and the affine geometry of the geometric simplex s_σ .

Let σ be an arbitrary q -simplex of K and s_σ be the ordered geometric q -simplex associated with σ as above. We define the *closed q -cell $\text{Cl } w_\sigma$* as a set by taking

$$\text{Cl } w_\sigma = \bigcup_{\tau < \sigma} w_\tau .$$

There is a natural transformation

$$\mu_\sigma : s_\sigma \rightarrow \text{Cl } w_\sigma$$

of s_σ onto $\text{Cl } w_\sigma$ defined as follows. For each n -dimensional face $(0 \leq n \leq q)$,

$$s' = \langle v_{j_0}, \dots, v_{j_n} \rangle, \quad 0 \leq j_0 < \dots < j_n \leq q ,$$

of s_σ , we define μ_σ on the interior $\text{Int } s'$ of s' to be the unique barycentric map of $\text{Int } s'$ onto w_τ , $\tau = \sigma_{(j_0, \dots, j_n)}$, which preserves the order of vertices. Give

$\text{Cl}w_\sigma$ the identification topology determined by μ_σ ; that is to say, a set $M \subset \text{Cl}w_\sigma$ is called *open* if and only if its inverse image $\mu_\sigma^{-1}(M) \subset s_\sigma$ is open.

Let us denote by $P(K)$ the union of all open cells w_σ corresponding to the simplexes σ of K . We define a topology of $P(K)$ as follows: A set M of $P(K)$ is said to be *open* if $M \cap \text{Cl}w_\sigma$ is an open set of $\text{Cl}w_\sigma$ for every closed cell $\text{Cl}w_\sigma$. The topological space $P(K)$ thus obtained is the semi-simplicial polytope associated with K . It is a polyhedral realization of the semi-simplicial complex K .

We remark that, for each simplex σ of K , the natural transformation

$$\mu_\sigma: s_\sigma \rightarrow \text{Cl}w_\sigma \subset P(K)$$

is a continuous map of s_σ onto $\text{Cl}w_\sigma$ and $\mu_\sigma|_{w_\sigma}$ is the identity. Following J.H.C. Whitehead [11, p.221], we shall call it the *characteristic map* for the open cell w_σ of $P(K)$.

Obviously, $P(K)$ is a CW-complex in the sense of J.H.C. Whitehead [11, p.223]. Hence we have the following assertions.

(2.2) $P(K)$ is a normal Hausdorff space.

(2.3) A transformation $f: P(K) \rightarrow R$ of $P(K)$ into an arbitrary topological space R is a continuous map if and only if the partial transformation $f|_{\text{Cl}w_\sigma}$ is continuous for each closed cell $\text{Cl}w_\sigma$ of $P(K)$.

3. Simplicial maps. We recall the definition of simplicial maps of semi-simplicial complexes [2, p.500]. A *simplicial map* $T: K_1 \rightarrow K_2$ of a semi-simplicial complex K_1 into another such complex K_2 is a function which to each q -simplex σ of K_1 assigns a q -simplex $\tau = T(\sigma)$ of K_2 in such a fashion that

$$\tau^{(i)} = T(\sigma^{(i)}) \quad (i = 0, \dots, q).$$

(3.1) A simplicial map $T: K_1 \rightarrow K_2$ induces a unique continuous map $f_T: P(K_1) \rightarrow P(K_2)$, which maps w_σ of $P(K_1)$ barycentrically onto w_τ of $P(K_2)$ with $\tau = T(\sigma)$.

Proof. For each integer $q \geq 0$, let K_1^q denote the q -dimensional skeleton of K_1 ; that is, K_1^q is the set of simplexes in K_1 with dimensions not exceeding q . Then $P(K_1^q)$ can be chosen as a subpolytope of $P(K_1)$ and will be called the q -dimensional skeleton of $P(K_1)$. Define a map

$$\phi_0: P(K_1^0) \rightarrow P(K_2)$$

as follows: For each simplex σ in K_1^0 , $\text{Cl}w_\sigma = w_\sigma$ is a single point of $P(K_1^0)$. Since $\tau = T(\sigma)$ is of dimension zero, $\text{Cl}w_\tau = w_\tau$ is a point of $P(K_2)$. Then ϕ_0 is defined by taking $\phi_0(w_\sigma) = w_\tau$. It is clear that ϕ_0 is uniquely determined by T . Since $P(K_1^0)$ is discrete, ϕ_0 is a continuous map.

Now assume that there exists a unique continuous map.

$$\phi_{q-1} : P(K_1^{q-1}) \rightarrow P(K_2)$$

which maps w_σ of $P(K_1^{q-1})$ barycentrically onto w_τ of $P(K_2)$ with $\tau = T(\sigma)$ for a certain integer $q > 0$. We are going to construct a map

$$\phi_q : P(K_1^q) \rightarrow P(K_2)$$

as follows: Let σ be an arbitrary q -simplex of K_1^q and $\tau = T(\sigma)$; w_σ is the interior of the ordered geometric simplex s_σ , and w_τ that of s_τ . Denote by $B_\sigma : s_\sigma \rightarrow s_\tau$ the unique barycentric map of s_σ onto s_τ preserving the order of vertices. Then ϕ_q is defined by taking

$$\phi_q(x) = \begin{cases} \phi_{q-1}(x) & (x \in P(K_1^{q-1})), \\ B_\sigma(x) & (x \in w_\sigma, \sigma \in K_1^q). \end{cases}$$

Now ϕ_q is uniquely determined by T and maps w_σ barycentrically onto w_τ with $\tau = T(\sigma)$ for each simplex σ of K_1^q . To prove the continuity of ϕ_q , it is sufficient to prove that of the partial map $\psi_\sigma = \phi_q|_{\text{Cl}w_\sigma}$ for each q -simplex σ of K_1^q . By means of the property of T and that of ϕ_n ($n \leq q$), it is easily seen that in the following diagram

$$\begin{array}{ccc} s_\sigma & \xrightarrow{B_\sigma} & s_\tau \\ \downarrow \mu_\sigma & & \downarrow \mu_\tau \\ \text{Cl} w_\sigma & \xrightarrow{\psi_\sigma} & \text{Cl} w_\tau \end{array}$$

commutativity holds; that is, $\mu_\tau B_\sigma = \psi_\sigma \mu_\sigma$, where $\tau = T(\sigma)$ and μ_σ, μ_τ are characteristic maps. Let U be an arbitrary open set of $\text{Cl}w_\tau$ and $V = \psi_\sigma^{-1}(U)$ in $\text{Cl}w_\sigma$. Since $\mu_\tau B_\sigma$ is continuous, $\mu_\sigma^{-1}(V) = (\mu_\tau B_\sigma)^{-1}(U)$ is an open set of s_σ . By the definition of the topology of $\text{Cl}w_\sigma$, V is open. Hence ψ_σ is continuous. This proves the continuity of ϕ_q . Hence we have completed the inductive construction of a sequence of continuous maps $\{\phi_q\}$, uniquely determined by T , such that

$$\phi_q|P(K_1^{q-1}) = \phi_{q-1}$$

for every $q > 0$, and ϕ_q maps w_σ barycentrically onto w_τ , $\tau = T(\sigma)$, for each simplex $\sigma \in K_1^q$.

The required continuous map $f_T: P(K_1) \rightarrow P(K_2)$ is defined by taking

$$f_T|P(K_1^q) = \phi_q \quad (q = 0, 1, \dots).$$

This completes the proof of (3.1).

4. The singular polytope $P(X)$. Let X be a given topological space. The singular complex $S(X)$ [2, p.502] is a typical semi-simplicial complex. The semi-simplicial polytope associated with $S(X)$ and constructed in §2 is essentially the *singular polytope* of J.B. Giever [4, p.182], which will be denoted simply by $P(X)$.

For the remainder of the present section, we shall assume that X is arcwise connected and that $x_0 \in X$ is a given point. Following S. Eilenberg, we denote by $S_n(X)$ the subcomplex of $S(X)$ consisting of all singular simplexes σ such that all faces of σ of dimensions less than n are collapsed at x_0 . The associated semi-simplicial polytope of $S_n(X)$ can be chosen naturally as a subpolytope of $P(X)$ and will be denoted by $P_n(X)$.

Now let M be a minimal subcomplex of $S(X)$ [2, p.502]. We can choose the associated polytope $P(M)$ as a subpolytope of $P(X)$. The following assertion is an immediate consequence of a corollary of Eilenberg and Zilber [2, p.503].

(4.1) *If the homotopy groups $\pi_i(X)$ vanish for each $i < n$, then $P(M)$ is a subpolytope of $P_n(X)$.*

Let Δ_q be a given ordered geometric q -simplex and let I be the closed unit interval of real numbers. The topological product $\Delta_q \times I$ has a *standard triangulation* into ordered simplexes without the introduction of new vertices. By means of this standard triangulation and the arguments analogous to those used in §3 and those used by Eilenberg and Zilber [2, p.504], it is not difficult to construct a homotopy

$$\delta_t: P(X) \rightarrow P(X) \quad (0 \leq t \leq 1).$$

subject to the following conditions:

- (i) δ_0 is the identity map;

- (ii) δ_1 maps $P(X)$ into $P(M)$;
- (iii) $\delta_t | P(M)$ is the identity map for all $t \in I$.

Then the main result of Eilenberg and Zilber [2] can be stated as follows.

(4.2) *The polytope $P(M)$ is a deformation retract of $P(X)$.*

Note that δ_t is not simplicial if $0 < t < 1$. The family of simplicial maps

$$f_{\phi_t} : P(X) \longrightarrow P(X) \quad (0 \leq t \leq 1),$$

induced according to (3.1) by the family $\phi_t : S(X) \longrightarrow S(X)$ ($0 \leq t \leq 1$), of Eilenberg and Zilber [2, p.504] is not continuous in t because of our topology introduced in $P(X)$.

The following assertion is a direct consequence of (4.2) and a theorem of J.B.Giever [4, Theorem VI].

(4.3) *The homotopy groups of $P(M)$ are isomorphic with those of X ; that is,*

$$\pi_i(P(M)) \approx \pi_i(X) \quad (i \geq 1).$$

5. **The polytope $P(\pi, n)$.** Throughout the present section, let π be a (discrete) group and n a positive integer. If $n = 1$, we make no assumption on π and write it multiplicatively; otherwise, we assume π abelian and written additively. Eilenberg and MacLane [3, p.517] define a semi-simplicial complex $K(\pi, n)$ which is very useful in the relations between homology and homotopy groups. A q -simplex σ of $K(\pi, n)$ is a function σ with values in π defined over all sets of arguments $0 \leq a_0 \leq \dots \leq a_n \leq q$ and subject to two specified conditions, namely the conditions (2.2) and (2.3) of [3]. We denote by $P(\pi, n)$ the semi-simplicial polytope associated with $K(\pi, n)$. Since $K(\pi, n)$ has only one 0-simplex, $P(\pi, n)$ is arcwise connected. We shall use the unique vertex p_0 of $P(\pi, n)$ as the base point for the homotopy groups.

THEOREM 1. *The homotopy groups of the polytope $P(\pi, n)$ are given below:*¹

$$\begin{aligned} \pi_n(P(\pi, n)) &\approx \pi; \\ \pi_i(P(\pi, n)) &= 0 \quad (i \neq n). \end{aligned}$$

¹This theorem is known to S. Eilenberg. He mentioned this fact in his address delivered before the Topology Conference of the International Congress of Mathematicians, 1950.

Proof. According to Realizability Theorem of J.H.C.Whitehead [12, p.261], there exists an arcwise connected topological space X with

$$(5.1) \quad \pi_n(X) \approx \pi, \quad \pi_i(X) = 0 \quad (i \neq n).$$

Choose a minimal complex M of the singular complex $S(X)$ [2, p.502], and consider their associated polytopes $P(X)$ and $P(M) \subset P(X)$. It follows from (4.3) and (5.1) that

$$(5.2) \quad \pi_n(P(M)) \approx \pi, \quad \pi_i(P(M)) = 0 \quad (i \neq n).$$

Since $\pi_i(X) = 0$ for each $i < n$, (4.1) tells us that $P(M)$ is a subpolytope of $P_n(X)$.

According to Eilenberg and MacLane [3, p.517], there is a natural simplicial map

$$\kappa: M \rightarrow K(\pi, n).$$

Since $\pi_i(X) = 0$ for each $i > n$, a result of Eilenberg and MacLane [3, p.519] gives a simplicial map

$$\bar{\kappa}: K(\pi, n) \rightarrow M$$

such that $\kappa\bar{\kappa}$ is the identity on $K(\pi, n)$. It follows also from the construction of κ and $\bar{\kappa}$ given by Eilenberg and MacLane [3] that $\bar{\kappa}\kappa(\sigma) = \sigma$ for every n -simplex of M . Now let

$$f: P(M) \rightarrow P(\pi, n), \quad \bar{f}: P(\pi, n) \rightarrow P(M)$$

be the continuous maps induced respectively by κ and $\bar{\kappa}$ according to (3.1). Denoting the n -dimensional skeleton of $P(M)$ by $P^n(M)$, we obtain the result that $f\bar{f}$ is the identity map on $P(\pi, n)$, and $\bar{f}f \upharpoonright P^n(M)$ is that on $P^n(M)$.

Since $\pi_i(P(M)) = 0$ for each $i > n$, it follows from a standard obstruction method that $\bar{f}f: P(M) \rightarrow P(M)$ is homotopic with the identity map on $P(M)$. Since $f\bar{f}$ is the identity map on $P(\pi, n)$, this proves that $P(M)$ and $P(\pi, n)$ are of the same homotopy type. Hence (5.2) implies Theorem 1.

Let π_* be a subgroup of π . Then $K(\pi_*, n)$ is the subcomplex of $K(\pi, n)$ consisting of all simplexes σ of $K(\pi, n)$ such that

$$\sigma(a_0, \dots, a_n) \in \pi_*$$

for all sets of arguments $0 \leq a_0 \leq \dots \leq a_n \leq \dim \sigma$. We can imbed $P(\pi_*, n)$ as

a subpolytope of $P(\pi, n)$ in an obvious way. If π_* is the subgroup consisting of a single element, then we use the notation $P_0(\pi, n)$ for this $P(\pi_*, n)$. It follows from Theorem 1 that

$$(5.3) \quad \pi_i(P_0(\pi, n)) = 0$$

for all integers $i \geq 1$. Now (5.3) and the exactness of the homotopy sequence imply that the identity map

$$j : (P(\pi, n), p_0) \longrightarrow (P(\pi, n), P_0(\pi, n))$$

induces the onto isomorphisms:

$$(5.4) \quad j_* : \pi_i(P(\pi, n)) \approx \pi_i(P(\pi, n), P_0(\pi, n)) \quad (i \geq 2).$$

For the remainder of the present section, we shall assume $n \geq 2$. There is a natural homomorphism

$$k_* : \pi \longrightarrow \pi_n(P(\pi, n), P_0(\pi, n))$$

described as follows. For an arbitrary element ω of π , there is one and only one n -simplex σ of $K(\pi, n)$ such that $\sigma(0, \dots, n) = \omega$. The open n -cell w_σ of $P(\pi, n)$ is the interior of a geometric n -simplex s_σ with ordered vertices. The order of the vertices determines an orientation of the pair $(s_\sigma, \partial s_\sigma)$. The characteristic map

$$\mu_\sigma : s_\sigma \longrightarrow \text{Cl } w_\sigma$$

carries the pair $(s_\sigma, \partial s_\sigma)$ into the pair $(P(\pi, n), P_0(\pi, n))$ and maps each vertex of s_σ into p_0 . Hence μ_σ determines an element $[\mu_\sigma]$ of the group $\pi_n(P(\pi, n), P_0(\pi, n))$. The homomorphism k_* is defined by setting $k_*(\omega) = [\mu_\sigma]$.

By a careful examination of the proof of Theorem 1, it is not difficult to see that the homomorphism $j_*^{-1}k_*$ is an isomorphism of π onto $\pi_n(P(\pi, n))$; that is,

$$(5.5) \quad \lambda_n = j_*^{-1}k_* : \pi \approx \pi_n(P(\pi, n)).$$

Hence k_* is also an isomorphism onto.

Now let $h : \pi \longrightarrow \pi$ be a given endomorphism of the group π . Then h induces a simplicial map

$$\eta : K(\pi, n) \longrightarrow K(\pi, n)$$

described as follows: For each simplex $\sigma \in K(\pi, n)$, $\eta(\sigma)$ is the simplex of $K(\pi, n)$ such that $\dim \eta(\sigma) = \dim \sigma$ and

$$\eta(\sigma)(a_0, \dots, a_n) = h(\sigma(a_0, \dots, a_n))$$

for every set of arguments $0 \leq a_0 \leq \dots \leq a_n \leq \dim \sigma$. Let us denote by

$$h^\# = f_\eta : P(\pi, n) \longrightarrow P(\pi, n)$$

the continuous map induced by the simplicial map η according to (3.1). The following properties of the correspondence $h \longrightarrow h^\#$ are immediate.

(5.6) For any two endomorphisms $h_1, h_2 : \pi \longrightarrow \pi$ of the group π , we have $(h_1 h_2)^\# = h_1^\# h_2^\#$.

(5.7) If $h : \pi \longrightarrow \pi$ is the identity endomorphism of the group π , then $h^\#$ is the identity map of $P(\pi, n)$.

Since $h^\#(p_0) = p_0$, $h^\#$ induces a homomorphism

$$h_* : \pi_n(P(\pi, n)) \longrightarrow \pi_n(P(\pi, n)) .$$

THEOREM 2. In the following rectangle of homomorphisms

$$\begin{array}{ccc} \pi & \xrightarrow{\lambda_n} & \pi_n(P(\pi, n)) \\ \downarrow h & & \downarrow h_* \\ \pi & \xrightarrow{\lambda_n} & \pi_n(P(\pi, n)) \end{array}$$

the commutativity relation $h_* \lambda_n = \lambda_n h$ holds.

Proof. It is obvious that the partial map $h^\#|_{P_0(\pi, n)}$ coincides with the identity map on $P_0(\pi, n)$. Hence $h^\#$ induces an endomorphism h_0 of the relative homotopy group $\pi_n(P(\pi, n), P_0(\pi, n))$. Since $\lambda_n = j_*^{-1} k_*$, the above rectangle can be decomposed into the following two:

$$\begin{array}{ccccc} \pi & \xrightarrow{k_*} & \pi_n(P(\pi, n), P_0(\pi, n)) & \xleftarrow{j_*} & \pi_n(P(\pi, n)) \\ \downarrow h & & \downarrow h_0 & & \downarrow h_* \\ \pi & \xrightarrow{k_*} & \pi_n(P(\pi, n), P_0(\pi, n)) & \xleftarrow{j_*} & \pi_n(P(\pi, n)) . \end{array}$$

The commutativity of the left rectangle, $k_* h = h_0 k_*$, is a direct consequence of the definitions of k_* and $h^\#$. The commutativity of the right rectangle, $h_0 j_* = j_* h_*$, is a property of the induced homomorphisms of the homotopy sequence. Since j_* is

an isomorphism onto, we have $j_*^{-1}h_0 = h_*j_*^{-1}$. Hence we obtain

$$h_*\lambda_n = h_*j_*^{-1}k_* = j_*^{-1}h_0k_* = j_*^{-1}k_*h = \lambda_n h.$$

This completes the proof.

(5.8) COROLLARY. *If we identify the groups π and $\pi_n(P(\pi, n))$ by means of the isomorphism λ_n , then the endomorphisms h and h_* coincide.*

6. Existence of the space B . Throughout the present section and the following one, let

$$\{\pi_n\} = \pi_1, \quad \pi_2, \dots, \quad \pi_n, \dots$$

be a given sequence of groups, where π_1 is a (multiplicative) group and π_n ($n > 1$) is an (additive) abelian group admitting π_1 as a given group of left operators; that is, for every $\xi \in \pi_1$ and every $\alpha \in \pi_n$, the element $\xi\alpha \in \pi_n$ is defined and

$$\xi(\alpha + \beta) = \xi\alpha + \xi\beta, \quad \xi(\eta\alpha) = (\xi\eta)\alpha, \quad 1\alpha = \alpha.$$

For each integer $n \geq 1$, let $P_n = P(\pi_n, n)$ denote the polytope associated with the complex $K(\pi_n, n)$. We shall use the following notations:

$$X = P_1, \quad Y = P_2 \times P_3 \times \dots \times P_n \times \dots.$$

Both X and Y are arcwise connected Hausdorff spaces. Let $\theta_n: Y \rightarrow P_n$ ($n = 2, 3, \dots$) denote the projection of Y onto the factor space P_n . The following properties of the space Y are immediate consequences of results in a note due to J.H.C. Whitehead, [13, p.289]:

(6.1) Y is 1-connected; that is, $\pi_1(Y) = 0$.

(6.2) $\pi_n(Y) \approx \pi_n$ for every integer $n \geq 2$.

(6.3) The Whitehead products in Y are all trivial; that is, for any two elements $a \in \pi_m(Y)$ and $b \in \pi_n(Y)$, we have $a \circ b = 0$.

Each element $\xi \in \pi_1$ determines, for every $n = 2, 3, \dots$, an automorphism

$$\xi_n: \pi_n \approx \pi_n$$

defined by $\xi_n(\omega) = \xi\alpha \in \pi_n$ for any $\alpha \in \pi_n$. According to §5, ξ_n induces a homeomorphism

$$\xi_n^\# : P_n \longrightarrow P_n \quad (n = 2, 3, \dots)$$

of P_n onto P_n . Define a homeomorphism

$$\xi^\# : Y \longrightarrow Y \quad (\xi \in \pi_1)$$

of Y onto itself by taking

$$(6.4) \quad \theta_n \xi^\#(y) = \xi_n^\# \theta_n(y) \quad (y \in Y; n = 2, 3, \dots).$$

The association $\xi \longrightarrow \xi^\#$ clearly determines a homomorphism

$$\rho : \pi_1 \longrightarrow \text{Hom}(Y)$$

of the discrete group π_1 into the discrete group $\text{Hom}(Y)$ of all homeomorphisms of Y onto itself. Let

$$G = \rho(\pi_1) \subset \text{Hom}(Y);$$

then G is a *topological transformation group* of Y and is isomorphic with the quotient group of π_1 over the kernel of the homomorphism ρ .

Remembering the isomorphism $\lambda_1 : \pi_1 \approx \pi_1(X)$ defined by (5.5), we shall call

$$\chi = \rho \lambda_1^{-1} : \pi_1(X) \longrightarrow G$$

the *characteristic homomorphism*.

Now let us consider the universal covering space \tilde{X} of X . It is well known that \tilde{X} is a bundle space over X with discrete fiber π_1 and structural group π_1 . Then the characteristic homomorphism χ induces a *weakly associated bundle space* B over X with Y as fiber and G as structural group [7]. The bundle space B is uniquely determined up to an equivalence in the sense of fiber bundles. In the following sections, we shall give an explicit construction of the bundle space B .

7. Barycentric subdivisions of semi-simplicial polytopes. Let K be a given semi-simplicial complex and $P(K)$ its associated polytope. We are going to define barycentric subdivisions of $P(K)$.

For each simplex $\sigma \in K$, let us denote by s'_σ the *barycentric first derived* [6, p. 3] of the ordered geometric simplex s_σ associated with σ .

Since the characteristic map

$$\mu_\sigma : s_\sigma \longrightarrow \text{Cl } w_\sigma$$

reduces to the identity map if it is restricted within the interior w_σ of s_σ , μ_σ induces a simplicial subdivision of w_σ into $\mu_\sigma(\text{Int } s'_\sigma)$, named the *barycentric first derived* w'_σ of w_σ , which is a finite set of open geometric simplexes. If we replace each open cell w_σ of $P(K)$ by its barycentric first derived w'_σ , we obtain a subdivision of $P(K)$, called the *first barycentric subdivision* $P'(K)$ of $P(K)$.

More generally, let us denote by $s_\sigma^{(n)}$ the *barycentric n -th derived* [6, p. 3] of s_σ . Then the characteristic map μ_σ induces a simplicial subdivision of w_σ into $\mu_\sigma(\text{Int } s_\sigma^{(n)})$, called the *barycentric n -th derived* $w_\sigma^{(n)}$ of w_σ . If we replace each open cell w_σ by its barycentric n -th derived $w_\sigma^{(n)}$, we obtain the *n -th barycentric subdivision* $P^{(n)}(K)$ of $P(K)$. It is clear that the characteristic map μ_σ carries each open simplex of $s_\sigma^{(n)}$ barycentrically onto some open simplex of $P^{(n)}(K)$.

Let v be an arbitrary vertex (that is, an open 0-simplex) of $P^{(n)}(K)$, where $n \geq 1$. The *star* of v , denoted by $\text{St}(v)$, is defined to be the union of all open simplexes ξ of $P^{(n)}(K)$ such that $\text{Cl } \xi$ contains v . The following assertion can easily be proved.

(7.1) *The star of each vertex of $P^{(n)}(K)$ ($n \geq 1$) is contractible (in itself) to a point.*

By a *simplicial polytope* P , we understand the union of a collection of closed geometric simplexes $\{s_\alpha\}$, where the index α runs over a certain abstract set A , such that (i) every face of an arbitrary simplex s_α of the collection belongs to the collection and (ii) the intersection $s_\alpha \cap s_\beta$ of any two simplexes of the collection is either vacuous or a face on both of them, with the topology defined as follows: A set $M \subset P$ is said to be open if and only if, for each closed geometric simplex s_α of the collection, $M \cap s_\alpha$ is an open set s_α in its euclidean topology. Simplicial polytopes are called topological polyhedra by J. H. C. Whitehead [9, p. 316]. The following assertion can easily be proved.

(7.2) *For each $n \geq 2$, $P^{(n)}(K)$ is a simplicial polytope.*

8. Explicit construction of the bundle. Let us return to the notations of §6. The vertices of the first barycentric subdivision of X are barycenters $\{x_\sigma\}$ of the open cells $\{w_\sigma\}$ of X , where σ runs over all simplexes of the semi-simplicial complex $K = K(\pi_1, 1)$. In particular, we shall denote by x_0 the vertex which corresponds to the unique 0-simplex of K . Hence x_0 is the basic point of the fundamental group $\pi_1(X)$ of X . Let V_σ denote the star of the vertex x_σ in the first barycentric subdivision X . Then we obtain an open covering $\Omega = \{V_\sigma\}$ of X ,

indexed by $\sigma \in K$. According to (7.1), each member V_σ of the covering Ω is contractible (in itself) to the point x_σ .

For each $\sigma \in K$, let $s_\sigma = \langle v_0, \dots, v_q \rangle$ be the associated geometric simplex. Denote by $c_\sigma : I \rightarrow s_\sigma$ the unique linear map such that $c_\sigma(0) = v_0$ and $c_\sigma(1) = x_\sigma$. Define a path

$$C_\sigma : I \rightarrow X$$

joining x_0 to x_σ by taking $C_\sigma = \mu_\sigma c_\sigma$, where $\mu_\sigma : s_\sigma \rightarrow Clw_\sigma$ is the characteristic map.

Let σ and τ be any two simplexes of K such that $V_\sigma \cap V_\tau$ is nonvoid. Take a point $x \in V_\sigma \cap V_\tau$. Choose a path $D : I \rightarrow V_\sigma$ joining x_σ to x and a path $E : I \rightarrow V_\tau$ joining x_τ to x . Then the closed path $C_\tau E D^{-1} C_\sigma^{-1}$ represents an element $\xi_{\tau\sigma}$ of $\pi_1(X)$ which clearly does not depend on the choice of the point x and the paths D and E . Let σ, τ, θ be any three simplexes of K such that $V_\sigma \cap V_\tau \cap V_\theta$ is nonvoid; then it is easy to see that $\xi_{\theta\tau} \xi_{\tau\sigma} = \xi_{\theta\sigma}$. Call

$$g_{\tau\sigma} = \chi(\xi_{\tau\sigma}) \in G.$$

Then, as constant maps on $V_\sigma \cap V_\tau$ into G , the collection $\{g_{\tau\sigma}\}$ together with the covering $\Omega = \{V_\sigma\}$ form¹ a system of coordinate transformations in X with values in G [7, §3.1].

The construction of N.E.Steenrod [7, §3.2] gives a fiber bundle (with coordinate system)

$$F = \{B, X, p, Y, G, V_\sigma, \phi_\sigma\}$$

with base space X , fiber Y , group G , and the coordinate transformations $\{g_{\tau\sigma}\}$. To clarify the precise situation, we shall briefly describe the entities of F as follows.

Let us regard the indexing complex K as a topological space with the discrete topology; that is to say, every simplex σ of K is considered as a point which is an open set of K . Let T be the subset of $X \times Y \times K$ consisting of these triples (x, y, σ) such that $x \in V_\sigma$. Define in T an equivalence relation:

$$(x, y, \sigma) \sim (x', y', \tau) \quad \text{if} \quad x = x', \quad g_{\tau\sigma} \cdot y = y'.$$

¹The author has the advantage of reading the book [7] in manuscript. The system $\{g_{\tau\sigma}\}$ of coordinate transformations constructed here is essentially a particular case of that constructed by N.E.Steenrod [7, Sec.13.8]. The sketch given here is to clarify the precise situation.

Define the *bundle space* B to be the totality of these equivalence classes in T . Let

$$\omega : T \rightarrow B$$

assign to each (x, y, σ) its equivalence class $[x, y, \sigma]$. Give B the identification topology determined by ω ; namely, a set U in B is called open if $\omega^{-1}(U)$ is an open set of T . Then B is a Hausdorff space and ω a continuous open map. The *projection*

$$p : B \rightarrow X$$

is defined by $p([x, y, \sigma]) = x$. The *coordinate functions*

$$\phi_\sigma : V_\sigma \times Y \rightarrow p^{-1}(V_\sigma) \quad (\sigma \in K)$$

are defined by $\phi_\sigma(x, y) = [x, y, \sigma]$ for each $x \in V_\sigma$ and $y \in Y$ with σ running over K .

Let y_0 denote the point of Y such that $\theta_n(y_0)$ is the unique vertex of P_n for each $n = 2, 3, \dots$. Denote by 0 the unique 0 -simplex of K and call $b_0 = [x_0, y_0, 0] \in B$. Then we have $p(b_0) = x_0$ and $\phi_0(x_0, y_0) = b_0$. We shall understand that x_0, y_0, b_0 are respectively the basic points of the various homotopy groups of the spaces X, Y, B studied in the next section.

9. The homotopy groups of the bundle space B . In the present section, we shall study in details the homotopy groups of the bundle space B , constructed in the foregoing section, and their mutual operations. The realizability theorem, stated in the introduction, follows as an immediate consequence of these investigations.

First of all, let us recall the (*exact*) homotopy sequence [7, §17.3],

$$\begin{aligned} \dots & \xrightarrow{P_{n+1}^*} \pi_{n+1}(X) \xrightarrow{\Delta_{n+1}} \pi_n(Y) \xrightarrow{i_n^*} \pi_n(B) \xrightarrow{P_n^*} \pi_n(X) \xrightarrow{\Delta_n} \dots \\ \dots & \xrightarrow{P_2^*} \pi_2(X) \xrightarrow{\Delta_2} \pi_1(Y) \xrightarrow{i_1^*} \pi_1(B) \xrightarrow{P_1^*} \pi_1(X), \end{aligned}$$

of the fiber bundle $F = \{B, X, p, Y, G, V_\sigma, \phi_\sigma\}$, with x_0, y_0, b_0 as the basic points of the homotopy groups of the spaces X, Y, B , respectively. Here, the homomorphisms p_n^* ($n \geq 1$) are those induced by the projection $p : B \rightarrow X$, and i_n^* ($n \geq 1$) those induced by the map $i : Y \rightarrow B$ defined by

$$(9.1) \quad i(y) = \phi_0(x_0, y_0) = [x_0, y_0, 0] \quad (y \in Y).$$

If we identify λ with $p^{-1}(x_0)$, then i is the injection of $p^{-1}(x_0)$ into B .

THEOREM 3. *For each integer $n \geq 1$, there is a natural isomorphism $h_n: \pi_n(B) \approx \pi_n$ of $\pi_n(B)$ onto π_n .*

Proof. First, let us prove that p_1^* is an isomorphism onto. By means of the arcwise connectedness of the fiber $p^{-1}(x_0)$, it can be shown by a standard argument that p_1^* maps $\pi_1(B)$ onto $\pi_1(X)$. According to (6.1), we have $\pi_1(Y) = 0$. An application of the exactness of the homotopy sequence gives that the kernel of p_1^* is $i_1^*(\pi_1(Y)) = 0$. Hence p_1^* is an isomorphism onto. We define

$$(9.2) \quad h_1 = \lambda_1^{-1} p_1^* : \pi_1(B) \approx \pi_1,$$

where $\lambda_1: \pi_1 \approx \pi_1(X)$ is the isomorphism defined by (5.5) for $n = 1$.

Next let $n \geq 2$. Since $X = P_1$, we have

$$\pi_{n+1}(X) = 0 = \pi_n(X).$$

Then it follows from the exactness of the homotopy sequence that i_n^* is an isomorphism onto. The projection $\theta_n: Y \rightarrow P_n$ induces an isomorphism onto:

$$\theta_n^* : \pi_n(Y) \approx \pi_n(P_n).$$

We define

$$(9.3) \quad h_n = \lambda_n^{-1} \theta_n^* i_n^{*-1} : \pi_n(B) \approx \pi_n \quad (n \geq 2),$$

where $\lambda_n: \pi_n \approx \pi_n(P_n)$ is the isomorphism defined by (5.5). This completes the proof of the theorem.

According to S. Eilenberg [1], the fundamental group $\pi_1(B)$ operates on the left of $\pi_n(B)$ for each $n \geq 2$. Let $w \in \pi_1(B)$ and $a \in \pi_n(B)$ be arbitrarily given elements. Choose a path $C: I \rightarrow B$ with $C(0) = b_0 = C(1)$ which represents w , and a map $f: I^n \rightarrow B$ with $f(\partial I^n) = b_0$ which represents a , where I denotes the closed unit interval of real numbers and I^n the closed unit n -cube of the euclidean n -space with ∂I^n denoting its boundary. Let $f_t: I^n \rightarrow B$ ($0 \leq t \leq 1$) be any homotopy such that $f_1 = f$ and $f_t(\partial I^n) = C(t)$ for all $0 \leq t \leq 1$; then the element $wa \in \pi_n(B)$ is represented by the map f_0 . We remind that π_1 operates on the left π_n for each $n \geq 2$.

THEOREM 4. *For arbitrary elements $w \in \pi_1(B)$ and $a \in \pi_n(B)$ ($n \geq 2$) we have*

$$h_n(wa) = h_1(w)h_n(a).$$

Proof. Choose a path $C: I \rightarrow X$ with $C(0) = x_0 = C(1)$ which represents the element $p_1^*(w)$ of the fundamental group $\pi_1(X)$. Denote by Y_0 the fiber over x_0 ; that is, $Y_0 = p^{-1}(x_0) = i(Y)$. Let $j: Y_0 \rightarrow Y$ denote the inverse of i . According to N. E. Steenrod [7, §3.1], there is a homotopy $H_t: Y_0 \rightarrow B$ ($0 \leq t \leq 1$) such that H_1 is the identity and $pH_t(Y_0) = C(t)$ for each $0 \leq t \leq 1$. More precisely, we choose H_t to be a translation of Y_0 along C^{-1} into itself [7, §13.1]. Call

$$\xi = h_1(w) = \lambda_1^{-1}p_1^*(w) \in \pi_1.$$

It follows from Steenrod's proof in his construction [7, §13.8] of the system $\{g_{\tau\sigma}\}$ of the coordinate transformations of the fiber bundle F that the homeomorphism

$$\psi = jH_0i : Y \rightarrow Y$$

is in G , and

$$\psi = \lambda p_1^*(w) = \rho h_1(w) = \rho(\xi) = \xi^\#,$$

where $\rho: \pi_1 \rightarrow \text{Hom}(Y)$ is the homomorphism which maps ξ into $\xi^\#$ defined by (7.4). Hence H_0 is the homeomorphism $i\xi^\#j$ of Y_0 onto itself and maps b_0 into itself. Define a path $\tilde{C}: I \rightarrow B$ by taking $\tilde{C}(t) = H_t(b_0)$ for each $t \in I$. Since $p\tilde{C} = C$, and p_1^* is an isomorphism, \tilde{C} represents the element w of $\pi_1(B)$.

Choose a map $g: I^n \rightarrow Y$ with $g(\partial I^n) = y_0$ which represents the element $i_n^{*-1}(a)$ of $\pi_n(Y)$. Then the map $f = ig: I^n \rightarrow B$ is a representative of the element $a \in \pi_n(B)$ and maps I^n into Y_0 . Define a homotopy $f_t: I^n \rightarrow B$ ($0 \leq t \leq 1$) by taking $f_t = H_t f$ for each $0 \leq t \leq 1$. Then we have $f_1 = f$ and $f_t(\partial I^n) = \tilde{C}(t)$ for every $0 \leq t \leq 1$. Hence, by definition, the map $f_0 = H_0 f = i\xi^\#g$ represents the element $wa \in \pi_n(B)$. It follows that $\xi^\#g$ is a representative of the element $i_n^{*-1}(wa)$ of $\pi_n(Y)$.

The homeomorphism $\xi_n^\#: P_n \rightarrow P_n$ induces an automorphism

$$\xi_* : \pi_n(P_n) \approx \pi_n(P_n).$$

By (7.4), we have $\theta_n \xi^\#g = \xi_n^\# \theta_n g$. Hence

$$\theta_n^* i_n^{*-1}(wa) = \xi_* \theta_n^* i_n^{*-1}(a).$$

According to Theorem 2, we have $\lambda_n^{-1} \xi_* = \xi \lambda_n^{-1}$. So we deduce that

$$h_n(wa) = \lambda_n^{-1} \theta_n^* i_n^{*-1}(wa) = \lambda_n^{-1} \xi_* \theta_n^* i_n^{*-1}(a)$$

$$= \xi \lambda_n^{-1} \theta_n^* i_n^{*-1}(a) = \xi h_n(a) = h_1(w) h_n(a).$$

This completes the proof.

For arbitrarily given elements $a \in \pi_m(B)$ and $b \in \pi_n(B)$ ($m \geq 2, n \geq 2$), let us choose representative maps $f: I^m \rightarrow B$ and $g: I^n \rightarrow B$ with $f(\partial I^m) = b_0 = g(\partial I^n)$. Since $I^{m+n} = I^m \times I^n$, we have

$$(9.4) \quad \partial I^{m+n} = (I^m \times \partial I^n) \cup (\partial I^m \times I^n).$$

The Whitehead product [10, p. 411] of the elements a and b is an element $a \circ b$ of $\pi_{m+n-1}(B)$ determined by the map $h: \partial I^{m+n} \rightarrow B$ which is defined as follows:

$$(9.5) \quad h(x, y) = \begin{cases} f(x) & (x \in I^m, y \in \partial I^n), \\ g(y) & (x \in \partial I^m, y \in I^n). \end{cases}$$

THEOREM 5. *For arbitrary elements $a \in \pi_m(B)$ and $b \in \pi_n(B)$ ($m \geq 2, n \geq 2$), we have $a \circ b = 0$.*

Proof. Let $\alpha = i_m^{*-1}(a) \in \pi_m(Y)$ and $\beta = i_n^{*-1}(b) \in \pi_n(Y)$. Then we have $i_{m+n-1}^*(\alpha \circ \beta) = a \circ b$. Hence Theorem 5 is an immediate consequence of (6.3).

According to Theorems 3-5, our bundle space B constructed in §8 satisfies all the conditions in the Realizability Theorem stated in the introduction. This completes the proof of the Realizability Theorem.

10. An application. Take an even sphere S^{2r} and let

$$\pi_n = \pi_n(S^{2r}) \quad (n = 1, 2, \dots).$$

The foregoing construction gives an arcwise connected topological space B with

$$\pi_n(B) \approx \pi_n(S^{2r}) \quad (n = 1, 2, \dots).$$

Since $\pi_1(B) = 0 = \pi_1(S^{2r})$, the operations of the fundamental groups on the higher homotopy groups are all trivial for both B and S^{2r} . However, the Whitehead products of the higher homotopy groups are essentially different for the spaces B and S^{2r} . In fact, if e is a generator of the group $\pi_{2r}(S^{2r})$, then the Whitehead product $e \circ e$ is nonzero because it has Hopf invariant ± 2 [8, p. 205]; but all the Whitehead products for the space B are zero. This proves that the Whitehead products of a topological space are essential invariants of the space and that they are not determined by the homotopy groups together with the operations of the fundamental

group upon the higher homotopy groups.

REFERENCES

1. S. Eilenberg, *On the relation between the fundamental group of a space and the higher homotopy groups*, Fund. Math. **32** (1939), 167-175.
2. S. Eilenberg and J. A. Zilber, *Semi-simplicial complexes and singular homology*, Ann. of Math. **51** (1950), 499-513.
3. S. Eilenberg and S. MacLane, *Relations between homology and homotopy groups of spaces, II*, Ann. of Math. **51** (1950), 514-533.
4. J. B. Giever, *On the equivalence of the two singular homology theories*, Ann. of Math. **51** (1950), 178-191.
5. S. Lefschetz, *Algebraic topology*, Amer. Math. Soc. Colloquium Publications, Vol. 27, New York, 1942.
6. ———, *Topics in topology*, Ann. of Math., Studies No. 10, Princeton University Press, Princeton, 1942.
7. N. E. Steenrod, *The topology of fiber bundles*, Princeton Math. Series, No. 14, Princeton University Press, Princeton, 1951.
8. G. W. Whitehead, *A generalization of the Hopf invariant*, Ann. of Math. **51** (1950), 192-237.
9. J. H. C. Whitehead, *Simplicial spaces, nuclei and m -groups*, Proc. London Math. Soc. **45** (1939), 243-327.
10. ———, *On adding relations to homotopy groups*, Ann. of Math. **42** (1941), 409-428.
11. ———, *Combinatorial homotopy, I, II*, Bull. Amer. Math. Soc. **55** (1949), 213-245, 453-496.
12. ———, *On the realizability of homotopy groups*, Ann. of Math. **50** (1949), 261-263.
13. ———, *On the groups $\pi_r(V_{n,m})$ and sphere-bundles*, Proc. London Math. Soc., Series 2, **48** (1944), 243-291.

TULANE UNIVERSITY AND
INSTITUTE FOR ADVANCED STUDY

ERRATA, VOLUME I

- Richard Bellman and Theodore Harris, *Recurrence times for the Ehrenfest model*, p. 188, formula (21): read $\phi_k^{(N)}[\sigma/(1-\lambda)]$ for $\phi_k^{(N)}\sigma/(1-\lambda)$ (twice).
 p. 191, line 9 from bottom of page: read "macroscopic" for "microscopic."
- W. Karush, *An iterative method for finding characteristic vectors of a symmetric matrix*, p. 240, formula (9): read $s < r^i$ ($i = 0, 1, 2, \dots$).

ACKNOWLEDGMENT

The editors gratefully acknowledge the services of the following persons not members of the Editorial Staff who have been consulted concerning the preparation of the first volume of this Journal:

R. P. Agnew, S. Agmon, R. Arens, R. Baer, G. Birkhoff, R. P. Boas, R. H. Bruck, H. V. Craig, C. Davis, R. P. Dilworth, N. Dunford, H. P. Edmundson, S. Eilenberg, A. Erdélyi, J. W. Green, R. M. Hayes, G. Hedlund, A. E. Heins, M. R. Hestenes, J. D. Hill, V. Hlavaty, R. D. James, M. Kac, I. Kaplansky, D. H. Lehmer, J. Lehner, H. Lewy, G. Lorentz, C. C. MacDuffee, G. W. Mackey, E. J. McShane, H. B. Mann, E. J. Mickle, D. Montgomery, C. B. Morrey, A. P. Morse, I. Niven, L. J. Paige, W. T. Puckett, W. T. Reid, M. S. Robertson, J. B. Rosser, H. Samelson, I. E. Segal, A. Seidenberg, P. A. Smith, I. S. Sokolnikoff, R. H. Sorgenfrey, J. D. Swift, O. Szász, O. Taussky, A. E. Taylor, A. W. Tucker, S. E. Warschawski, W. R. Wasow, G. W. Whitehead, A. L. Whiteman, F. Wolf, J. W. T. Youngs, A. Zygmund.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

M.M. SCHIFFER*

Stanford University
Stanford, California

E. HEWITT

University of Washington
Seattle 5, Washington

R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

E.F. BECKENBACH**

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN

P. R. HALMOS

BØRGE JESSEN

J. J. STOKER

HERBERT FEDERER

HEINZ HOPF

PAUL LÉVY

E. G. STRAUS

MARSHALL HALL

R. D. JAMES

GEORGE PÓLYA

KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

HUGHES AIRCRAFT COMPANY

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. Manuscripts intended for the outgoing editors should be sent to their successors. All other communications to the editors should be addressed to the managing editor, E.G. Straus, at the University of California Los Angeles 24, California.

50 reprints of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50; back numbers (Volumes 1, 2, 3) are available at \$2.50 per copy. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

* To be succeeded in 1955, by H.L. Royden, Stanford University, Stanford, California.

** To be succeeded in 1955, by E.G. Straus, University of California, Los Angeles 24, Calif.

UNIVERSITY OF CALIFORNIA PRESS · BERKELEY AND LOS ANGELES

COPYRIGHT 1954 BY PACIFIC JOURNAL OF MATHEMATICS

Pacific Journal of Mathematics

Vol. 1, No. 4

, 1951

R. H. Bruck, <i>Loops with transitive automorphism groups</i>	481
Paul R. Garabedian, <i>A partial differential equation arising in conformal mapping</i>	485
Magnus R. Hestenes, <i>Applications of the theory of quadratic forms in Hilbert space to the calculus of variations</i>	525
Sze-Tsen Hu, <i>On the realizability of homotopy groups and their operations</i>	583