

Pacific Journal of Mathematics

ON A PAPER OF NIVEN AND ZUCKERMAN

J. W. S. CASSELS

ON A PAPER OF NIVEN AND ZUCKERMAN

J. W. S. CASSELS

1. **Introduction.** Let 'digit' mean an integer in the range $0 \leq a < 10$. For digits $a_1, a_2, \dots, a_r; b_1, b_2, \dots, b_s$ ($s \geq r$) and integer m , denote by

$$R_m(a_1, \dots, a_r, b_1, \dots, b_s)$$

the number of solutions of

$$b_n = a_1, b_{n+1} = a_2, \dots, b_{n+r-1} = a_r \quad (0 < n < n+r \leq s; n \equiv m \pmod{r}),$$

so that

$$(1) \quad 0 \leq R_m(a_1, \dots, a_r; b_1, \dots, b_s) \leq s - r + 1.$$

Suppose that

$$x_1, x_2, \dots$$

is an infinite sequence of digits. It has been shown [2] that if

$$(2) \quad \lim_{N \rightarrow \infty} \frac{1}{N} \sum_{m=1}^r R_m(a_1, \dots, a_r; x_1, \dots, x_N) = 10^{-r}$$

for all integers r and digits a_1, \dots, a_r , then

$$(3) \quad \lim_{N \rightarrow \infty} \frac{1}{N} R_m(a_1, \dots, a_r; x_1, \dots, x_N) = r^{-1} 10^{-r}$$

for all integers r, m , and digits a_1, \dots, a_r . A possibly simpler proof is as follows.

2. **Proof.** Let $\epsilon > 0$ and digits a_1, \dots, a_r be given. The simple argument of Hardy-Wright [1] shows that if the integer s is fixed large enough, then

Received April 10, 1952.

Pacific J. Math. 3 (1953), 555-557

$$(4) \quad \text{Max}_{\mu} \left| R_{\mu}(a_1, \dots, a_r; b_1, \dots, b_s) - \frac{s-r+1}{r 10^r} \right| < \epsilon(s-r+1)$$

except for at most $\epsilon 10^s$ sets of digits b_1, \dots, b_s . ('Exceptional' sets.) Thus, by (2) with b_1, \dots, b_s for a_1, \dots, a_r , the number of exceptional sets

$$(5) \quad x_t, x_{t+1}, \dots, x_{t+s-1} \quad (1 \leq t \leq N-s+1)$$

is at most $2 \in N$ for all large enough N .

On the other hand,

$$(6) \quad (s-r+1) R_m(a_1, \dots, a_r; x_1, \dots, x_N)$$

differs from

$$(7) \quad \sum_{t=1}^{N-s+1} R_{m-t+1}(a_1, \dots, a_r; x_t, \dots, x_{t+s-1})$$

by at most $2s^2$, since each solution of

$$a_1 = x_n, a_2 = x_{n+1}, \dots, a_r = x_{n+r-1} \quad (s \leq n \leq N-s; n \equiv m \pmod{r})$$

contributes exactly $s-r+1$ both to (6) and to (7). Hence, using the estimate (3) for the at most $2 \in N$ exceptional sets (5), and the estimate (4) for the others, we have

$$\begin{aligned} & \left| R_m(a_1, \dots, a_r; x_1, \dots, x_N) - \frac{N-s+1}{r 10^r} \right| \\ & \leq \frac{2s^2}{s-r+1} + \epsilon(N-s+1) + 2 \in N, \end{aligned}$$

and so

$$\limsup_N \left| \frac{1}{N} R_m(a_1, \dots, a_r; x_1, \dots, x_N) - r^{-1} 10^{-r} \right| \leq 3 \epsilon.$$

Since ϵ is arbitrarily small, this proves (3) as required.

REFERENCES

1. G. H. Hardy and E. M. Wright, *An introduction to the theory of numbers*, First Edition, Oxford, 1938, §9.13.
2. I. Niven and H. S. Zuckerman, *On the definition of normal numbers*, Pacific J. Math. 1 (1951), 103-110.

TRINITY COLLEGE,
CAMBRIDGE, ENGLAND

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

R. M. ROBINSON

University of California
Berkeley 4, California

*R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

E. F. BECKENBACH, Managing Editor

University of California
Los Angeles 24, California

*During the absence of Herbert Busemann in 1952.

ASSOCIATE EDITORS

R. P. DILWORTH

P. R. HALMOS

BØRGE JESSEN

J. J. STOKER

HERBERT FEDERER

HEINZ HOPF

PAUL LÉVY

E. G. STRAUS

MARSHALL HALL

R. D. JAMES

GEORGE PÓLYA

KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

• • •

AMERICAN MATHEMATICAL SOCIETY

NATIONAL BUREAU OF STANDARDS,

INSTITUTE FOR NUMERICAL ANALYSIS

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December, by the University of California, Berkeley 4, California. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

UNIVERSITY OF CALIFORNIA PRESS · BERKELEY AND LOS ANGELES

COPYRIGHT 1952 BY PACIFIC JOURNAL OF MATHEMATICS

| | |
|---|-----|
| Shmuel Agmon, <i>On the singularities of Taylor series with reciprocal coefficients</i> | 431 |
| Richard Arens, <i>A generalization of normed rings</i> | 455 |
| Iacopo Barsotti, <i>Intersection theory for cycles of an algebraic variety</i> | 473 |
| Leonard M. Blumenthal, <i>Two existence theorems for systems of linear inequalities</i> | 523 |
| Frank Herbert Brownell, III, <i>Translation invariant measure over separable Hilbert space and other translation spaces</i> | 531 |
| J. W. S. Cassels, <i>On a paper of Niven and Zuckerman</i> | 555 |
| Nelson Dunford, <i>Spectral theory. II. Resolutions of the identity</i> | 559 |
| Eugene Lukacs and Otto Szász, <i>On analytic characteristic functions</i> | 615 |
| W. A. Mersman, <i>Evaluation of an integral occurring in servomechanism theory</i> | 627 |
| Lawrence Edward Payne and Alexander Weinstein, <i>Capacity, virtual mass, and generalized symmetrization</i> | 633 |
| Choy-Tak Taam, <i>The boundedness of the solutions of a differential equation in the complex domain</i> | 643 |