

Pacific Journal of Mathematics

**GENERALIZED CONVEXITY AND SURFACES OF NEGATIVE
CURVATURE**

PAUL ARNOLD CLEMENT

GENERALIZED CONVEXITY AND SURFACES OF NEGATIVE CURVATURE

PAUL A. CLEMENT

Introduction. In a study [4] of surfaces whose Gaussian, or total, curvature K satisfies the relation $K \leq 0$, a number of functions having geometrical significance have been shown to be convex. In the present paper, a study of surfaces whose Gaussian curvature satisfies $K \leq K_0$, where K_0 is a negative constant, leads to the determination of a class of functions which are subfunctions (defined in §1.1) of a two-parameter family of functions determined by the bound K_0 . This is a natural generalization because the convexity property is equivalent to the subfunction property with respect to the particular two-parameter family (nonvertical straight lines) determined by the bound $K_0 = 0$.

A main objective will be to exhibit functions which have a geometrical significance and also have the subfunction property for surfaces with $K \leq K_0$. This property then implies certain inequality relations for functions associated with certain geometrical configurations on such surfaces.

I. SUBFUNCTIONS

1.1. Definitions. A real-valued function $g(x)$ of a single real variable x defined on an open interval (a, b) , with $-\infty \leq a < x < b \leq +\infty$, is said to be a *convex function* of x provided $g(x)$ satisfies the inequality

$$(1.11) \quad g[tx_1 + (1-t)x_2] \leq tg(x_1) + (1-t)g(x_2)$$

for all x_1, x_2 in (a, b) and for all t on the range $0 \leq t \leq 1$. If $g(x)$ is of class C^2 , it is convex if and only if $g''(x) \geq 0$ throughout the interval.

Geometrically, (1.11) indicates that no part of the graph of the curve $y=g(x)$ lies above the chord joining two points upon it within the interval (a, b) .

A generalization of the foregoing characteristic geometric property of convex

Received July 6, 1952. This paper is largely a condensation of a June, 1949, doctoral dissertation, University of California, Los Angeles; I gratefully acknowledge the guidance of Professor Beckenbach.

Pacific J. Math. 3 (1953), 333-368

functions leads to the theory of subfunctions [3]. Let $\{h_{\alpha\beta}(x)\}$ be a two-parameter family of continuous functions such that for all x_1, x_2 in (a, b) and every y_1, y_2 there exists a unique member $h_{\alpha_0\beta_0}(x)$ of the family such that $h_{\alpha_0\beta_0}(x_i) = y_i$ ($i = 1, 2$). Then a function $g(x)$ is said to be a *subfunction* of the given family on (a, b) provided¹ we have

$$(1.12) \quad g[tx_1 + (1-t)x_2] \leq h_{\alpha_1\beta_1}[tx_1 + (1-t)x_2]$$

for all x_1, x_2 in (a, b) and for all t on the range $0 \leq t \leq 1$, and where

$$h_{\alpha_1\beta_1}(x_i) = g(x_i) \quad (i = 1, 2).$$

Geometrically, (1.12) indicates that in the subinterval (x_1, x_2) no part of the graph of the curve $y = g(x)$ lies above the member of the parameter family joining the points $[x_1, g(x_1)]$ and $[x_2, g(x_2)]$. We note that if $g(x)$ is convex, it is a subfunction of the two-parameter family of nonvertical straight lines.

1.2. A fundamental theorem. Necessary and sufficient conditions that a function $g(x)$ be a subfunction of a certain type of two-parameter family have been obtained by Shniad [10]. The following lemma and theorem are due to him; proofs are included because of the fundamental use made of the theorem in subsequent developments.

LEMMA 1.1. *If $\phi(x)$ is a positive continuous function of x , and $\psi(x)$ is a strictly increasing continuous function of x , on $a < x < b$, then the condition that $g(x)$ be a subfunction of the family $A\phi + B\phi\psi$, where A and B are parameters of the family, is equivalent to the condition that g/ϕ be a convex function of ψ .*

Proof. The hypotheses on ϕ and ψ ensure that g/ϕ is a continuous function of ψ . To prove the existence of a unique member of the family through any two points (x_i, y_i) ($i = 1, 2$), with the x_i distinct and in the interval, it suffices to note that

$$\begin{vmatrix} \phi(x_1) & \phi(x_1)\psi(x_1) \\ \phi(x_2) & \phi(x_2)\psi(x_2) \end{vmatrix} = \phi(x_1)\phi(x_2)[\psi(x_2) - \psi(x_1)] \neq 0.$$

Let x_1 and x_2 satisfy $a < x_1 < x_2 < b$, and let

¹A more general definition of the subfunction property is given in [5]; in [3] it is shown that a function satisfying (1.12) necessarily is continuous on (a, b) .

$$h_{\alpha_1 \beta_1}(x) = A_1 \phi(x) + B_1 \phi(x) \psi(x)$$

with

$$h_{\alpha_1 \beta_1}(x_i) = g(x_i) \quad (i = 1, 2).$$

Then the condition

$$h_{\alpha_1 \beta_1}(x) \geq g(x) \quad \text{for } x_1 < x < x_2$$

is equivalent to the condition

$$A_1 + B_1 \psi(x) \geq \frac{g(x)}{\phi(x)} \quad \text{for } x_1 < x < x_2,$$

or that $g(x)/\phi(x)$ be a convex function of ψ on the range $\psi(a+) < \psi < \psi(b-)$.

THEOREM 1.2.² *Let $\phi(x)$, $\psi(x)$, and $g(x)$ be functions having the following properties on an interval $a < x < b$:*

- a) *the functions ϕ , ψ , and g have continuous second derivatives,*
- b) *the inequalities $\phi(x) > 0$ and $\psi'(x) > 0$ hold, and*
- c) *each of the functions $\phi(x)$ and $\phi(x)\psi(x)$ is a solution of the differential equation*

$$h'' + Ph' + Qh = 0,$$

where P and Q are continuous on the interval.

Then a necessary and sufficient condition that $g(x)$ be a subfunction of the family $A\phi + B\phi\psi$ on the given interval is that

$$g'' + Pg' + Qg \geq 0$$

on the interval.

Proof. From Lemma 1.1 it follows that g is a subfunction of the family if and only if g/ϕ is a convex function of ψ . Since g/ϕ has a continuous second derivative with respect to ψ , the latter condition is equivalent to

²The conclusion of this theorem is obtained in a more general setting in [9]. However, the proof is immediate for the theorem as stated here, and this form is sufficient for our purposes.

$$\frac{1}{(\psi')^2 \phi} \left\{ g'' + g' \left[-\frac{\psi''}{\psi'} - \frac{2\phi'}{\phi} \right] + g \left[\frac{\psi''\phi'}{\psi'\phi} - \frac{\phi''}{\phi} + 2 \left(\frac{\phi'}{\phi} \right)^2 \right] \right\} \geq 0.$$

From the Wronskian relation we easily verify that ϕ and $\phi\psi$ are linearly independent solutions of the differential equation. Then the theorem follows from uniqueness properties of linearly independent solutions of this type of differential equation.

1.3. Sub- K_0 functions. The differential equation we are to consider is

$$h'' + K_0 h = 0,$$

where K_0 is a negative constant, and the interval of definition is $0 \leq x < b \leq \infty$. The two-parameter family of solutions of the equation is given by

$$(1.31) \quad \{h_{\alpha\beta}(x)\} = \{\alpha \cosh(\sqrt{-K_0}x) + \beta \sinh(\sqrt{-K_0}x)\},$$

where α and β are the parameters. A property of this family is given in the following lemma; we omit the proof.

LEMMA 1.3. *If A: (x_1, y_1) and B: (x_2, y_2) are two points with $x_1 \neq x_2$, then there is one and only one curve of the family $\{h_{\alpha\beta}(x)\}$ passing through A and B. Thus, if $y_1 \geq 0$ and $y_2 \geq 0$, the curve $h_{\alpha_1\beta_1}(x)$ passing through A and B satisfies $h_{\alpha_1\beta_1}(x) \geq 0$ for $x_1 \leq x \leq x_2$.*

DEFINITION. A function $g(x)$ will be said to be a *sub- K_0 function* of x if it is a subfunction of the family $\{h_{\alpha\beta}(x)\}$ of (1.31) on the interval $0 \leq x < b \leq \infty$. Moreover, $g(x)$ will be said to be a *K_0 -function* if the sign of equality of its subfunction relation (1.12) holds throughout the interval; and it will be a *strictly sub- K_0 function* if the strict inequality holds throughout for $0 < t < 1$.

It is convenient to introduce a second-order differential operator \mathfrak{S} defined by

$$\mathfrak{S} \equiv D^2 + K_0,$$

where K_0 is a negative constant; we may write \mathfrak{S}_x to indicate the variable for differentiation.

REMARK. With the choices

$$\phi(x) \equiv \cosh(\sqrt{-K_0}x) \text{ and } \psi(x) \equiv \tanh(\sqrt{-K_0}x),$$

the family $\{\alpha\phi + \beta\psi\}$ coincides with the family (1.31), and these functions ϕ and ψ satisfy the hypotheses of Theorem 1.2. Hence a function $g(x)$ of class C^2 is a sub- K_0 function (K_0 -function) if and only if $\mathfrak{S}g(x) \geq 0$ ($\mathfrak{S}g(x) = 0$) on the interval.

Certain elementary properties of sub- K_0 functions are given in the following theorems. The proofs are omitted as they merely involve applying the foregoing remark to appropriate members of the family $\{h_{\alpha\beta}(x)\}$.

THEOREM 1.4. *Any linear combination of sub- K_0 functions with nonnegative coefficients is a sub- K_0 function.*

THEOREM 1.5. *Let $f(x)$ be a nonnegative sub- K_0 function, and let k be a constant ≥ 1 . Then $[f(x)]^k$ is a sub- K_0 function; in fact, $[f(x)]^k$ is a sub- kK_0 function.*

THEOREM 1.6. *Let $f_i(x)$ ($i = 1, 2, \dots, n$) be convex functions of x which are nonnegative and monotonic nondecreasing and at least one of which is a sub- K_0 function. Then the product function $f_1 f_2 \cdots f_n$ is a sub- K_0 function.*

II. SURFACES OF NEGATIVE CURVATURE

2.1. Geodesic parameters. Let an analytic surface S be represented by geodesic parameters [7, p. 174] (u, v) , so that

$$(2.11) \quad ds^2 = du^2 + \mu^2(u, v)dv^2 \quad (\mu \geq 0),$$

and

$$(2.12) \quad da = \mu(u, v)du dv,$$

where the curves $v = \text{constant}$ are the geodesics, and the curves $u = \text{constant}$ are the *geodesic parallels*. The surface S is said to be given in *geodesic representation*.

Singular points of the geodesic family are points where $\mu = 0$; other points, where $\mu > 0$, are *regular points*.

The Gaussian curvature K of S exists at all regular points. If S is given in geodesic representation, the Gaussian curvature is given [7, p. 181] by the formula

$$(2.13) \quad K = -\frac{1}{\mu} \frac{\partial^2 \mu}{\partial u^2}.$$

DEFINITION. An analytic surface S will be said to be a *sub- K_0 surface* if its Gaussian curvature is bounded from above by K_0 , a negative constant, at all regular points of S . Moreover, S will be said to be a *K_0 -surface* if its Gaussian curvature everywhere is K_0 . If S is a sub- K_0 surface which is not a K_0 -surface, it will be said to be a *strictly sub- K_0 surface*.

2.2. Geodesic parallels. We have the following lemma.

LEMMA 2.1. *If an analytic surface S is given in geodesic representation, then a necessary and sufficient condition that S be a sub- K_0 surface is that the function $\mu(u, v_0)$ be a sub- K_0 function of u for each line-segment $u_1 < u < u_2$, $v = v_0$ in the (u, v) -domain of definition.*

Proof. The result follows directly from (2.13) and Lemma 1.3 by an argument analogous to that in [4, p. 286]. The proof reveals that μ is a strictly sub- K_0 function of u if and only if S is a strictly sub- K_0 surface, and that μ is a K_0 -function of u if S is a K_0 -surface.

Let S be a sub- K_0 surface given in geodesic representation. Then we have the following results.

THEOREM 2.2. *Let the arcs $C(u)$ ($u_1 \leq u \leq u_2$), of length $l(u)$, be arcs of geodesic parallels between geodesics $v = v_1$ and $v = v_2$ ($v_1 < v_2$) on S . Then the length $l(u)$ is a sub- K_0 function of u (that is, of the geodesic length $u - u_1$); $l(u)$ is a strictly sub- K_0 function if S is a strictly sub- K_0 surface, and $l(u)$ is a K_0 -function if S is a K_0 -surface.*

Proof. A proof may be patterned on that of the related theorem in [4, p. 287], in which we substitute the appropriate member (which is of class C^2) of the family (1.31) in a subfunction inequality in place of the convexity inequality.

THEOREM 2.3. *Let the arcs $C(u)$ ($u_1 - W < u < u_1 + W$), of length $l(u)$, be arcs of geodesic parallels between geodesics $v = v_1$ and $v = v_2$ ($v_1 < v_2$) on S , and let $a(w)$ denote the area of the part of S enclosed by $v = v_1$, $C(u_1 + w)$, $v = v_2$, $C(u_1 - w)$ ($0 \leq w < W$). Then $a(w)$ is a sub- K_0 function of w ; $a(w)$ is a strictly sub- K_0 function if S is a strictly sub- K_0 surface, and is a K_0 -function of w if S is a K_0 -surface.*

Proof. The proof is similar to that in [4, p. 288] when we consider subfunction properties instead of convexity properties.

2.3. Geodesic polar coordinates. Let the analytic surface S be represented

in geodesic polar coordinates [7, p.181] (u, v) , that is, coordinates for which (2.11), (2.12), and

$$(2.31) \quad \mu(0, v) = 0, \quad \left[\frac{\partial \mu}{\partial u} \right]_{u=0} = 1$$

are satisfied. The curve $u = u_0$ is a *geodesic circle* with center at the *pole* P of the representation and *geodesic radius* u_0 .

We shall write r, θ for u, v , respectively.

Hereafter we indicate functions determined by, or calculated for, a K_0 -surface by a subscript zero. Some such functions can be determined explicitly.

LEMMA 2.4. *Let S_0 be a K_0 -surface, and let $l_0(r)$ and $a_0(r)$ denote the circumference and area, respectively, of the geodesic circle on S_0 with fixed center P and geodesic radius r . Then*

$$(2.32) \quad l_0(r) \equiv \frac{2\pi}{\sqrt{-K_0}} \sinh(\sqrt{-K_0}r)$$

and

$$(2.33) \quad a_0(r) \equiv \frac{2\pi}{-K_0} [\cosh(\sqrt{-K_0}r) - 1].$$

Moreover,

$$l_0(r) > 2\pi r \quad (r > 0 \text{ on } S),$$

and

$$a_0(r) > \pi r^2 \quad (r > 0 \text{ on } S).$$

Proof. Since $\mathfrak{S}_u \mu = 0$, we find that the function $\mu_0(r)$ of the family (1.31) satisfying (2.31) is

$$\mu_0(r) = \frac{1}{\sqrt{-K_0}} \sinh(\sqrt{-K_0}r).$$

When we evaluate (2.11) and (2.12) for a geodesic circle using this expression for μ , we obtain the formulas of the lemma. The inequalities are easily established; cf. [4, p.291-292].

We remark that the functions $l_0(r)$ and $a_0(r)$ will occur in formulas which refer to a sub- K_0 surface S ; in such cases, (2.32) and (2.33) provide definitions of these functions on S .

III. SUBFUNCTIONS FOR GEODESIC CIRCLES

3.1. Definition. Some functions of geometrical significance involving the geodesic radius have certain properties in common. We collect these properties in the following definition.

CONDITION C. For a given sub- K_0 surface S and for a given pole P of geodesic polar coordinates on S , a function $\phi(r)$ of the geodesic radius r satisfies Condition C provided: $\phi(0) = 0$; for $r \geq 0$ on S , $\phi(r)$ is a continuous, nondecreasing sub- K_0 function of r ; $\phi(r) \equiv 0$ if S is a K_0 -surface, but otherwise $\phi(r)$ is a strictly sub- K_0 function.

If we let $K_0 = 0$, the K_0 -surface becomes a developable surface, and the "sub- K_0 function of r " property becomes the usual "convex function of r " property. Thus our Condition C specializes to Condition A of [4, p.289] when $K_0 = 0$.

It follows from the theorems of § 1.3 that sums and products of functions which satisfy Condition C also satisfy Condition C.

3.2. The length function. Hereafter we assume that $\mu(r, \theta)$ is of class C^2 , which ensures the existence of the derivatives we write. We now consider a geodesic circle C_r on S with fixed center P and geodesic radius r .

LEMMA 3.1. *Let S be an analytic sub- K_0 surface, and let $l(r)$ denote the length of the circumference of C_r . Then $l(r)$ satisfies the differential relation*

$$(3.21) \quad \mathfrak{S}l(r) \equiv l''(r) + K_0 l(r) \geq 0 \quad (r \geq 0 \text{ on } S).$$

Proof. The result is immediate since $\mathfrak{S}_r \mu(r, \theta) \geq 0$ for $r \geq 0$ on S . We note that equality holds in (3.21) if and only if S is a K_0 -surface, that is, in our notation, if and only if we have $\mathfrak{S}l_0(r) = 0$, where $l_0(r)$ is given by (2.32).

LEMMA 3.2. *Let S be an analytic sub- K_0 surface, and let $a(r)$ denote the area of C_r . Then $a(r)$ satisfies the differential relation*

$$(3.22) \quad a''(r) + K_0 a(r) - 2\pi \equiv l'(r) + K_0 a(r) - 2\pi \geq 0 \quad (r \geq 0 \text{ on } S).$$

Proof. By differentiating the area function

$$a(r) = \int_0^r \int_0^{2\pi} \mu(\rho, \theta) d\rho d\theta,$$

we get

$$a''(r) = \int_0^{2\pi} \frac{\partial \mu}{\partial r} d\theta \equiv l'(r).$$

Since $a(0) = 0$, and $l'(0) = 2\pi$ by (2.31), we have equality in (3.22) for $r = 0$. The derivative of the function

$$l'(r) + K_0 a(r) - 2\pi$$

is $\mathcal{C}l(r)$, which is nonnegative by Lemma 3.1; hence the left member of (3.22) is monotonic nondecreasing, and (3.22) holds. It is readily seen that equality holds in (3.22) if and only if S is a K_0 -surface.

THEOREM 3.3. *Let S be an analytic sub- K_0 surface, and let $l(r)$ denote the length of the circumference of C_r . Then the function*

$$\phi_1(r) \equiv l(r) - l_0(r) \tag{r \ge 0 \text{ on } S},$$

satisfies Condition C.

Proof. The functions $\mu(r, \theta)$ and $\mu_0(r, \theta)$ associated with the surfaces S and S_0 , respectively, both satisfy (2.31), and are such that

$$\frac{\partial^2 \mu}{\partial r^2} + K\mu = 0 \tag{r \ge 0 \text{ on } S},$$

$$\frac{\partial^2 \mu_0}{\partial r^2} + K_0 \mu_0 = 0 \tag{r \ge 0 \text{ on } S},$$

where $K \leq K_0$. By Sturm's oscillation theorems [8, Chap. X], it follows that

$$\mu(r, \theta) - \mu_0(r, \theta) \geq 0 \tag{r \ge 0 \text{ on } S},$$

and

$$\frac{\partial \mu}{\partial r} - \frac{\partial \mu_0}{\partial r} \geq 0 \tag{r \ge 0 \text{ on } S}.$$

Hence we find that $\phi_1(0) = 0$, $\phi_1(r) \equiv 0$ on S if and only if S is a K_0 -surface, and

$$\phi_1'(r) \equiv l'(r) - l_0'(r) = \int_0^{2\pi} \left(\frac{\partial \mu}{\partial r} - \frac{\partial \mu_0}{\partial r} \right) d\theta \geq 0 \quad (r \geq 0 \text{ on } S).$$

Then calculation shows that

$$\mathfrak{S}\phi_1(r) = \mathfrak{S}l(r),$$

whence

$$\mathfrak{S}\phi_1(r) \geq 0 \quad (r \geq 0 \text{ on } S),$$

by Lemma 3.1. Thus, by Theorem 1.2, $\phi_1(r)$ satisfies Condition C.

COROLLARY 3.4. *If S is an analytic sub- K_0 surface, then $l(r)$ is a monotonic increasing sub- K_0 function of r and satisfies the inequality*

$$l(r) \geq l_0(r) \quad (r \geq 0 \text{ on } S);$$

$l(r)$ is a strictly sub- K_0 function if and only if S is not a K_0 -surface.

Proof. The inequality follows from Theorem 3.3 and the identity

$$l(r) = l_0(r) + \phi_1(r) \quad (r \geq 0 \text{ on } S).$$

REMARK. The function $\phi_1(r)$ may be modified to form a new function in the following way: replace the function $l_0(r)$ (Lemma 2.4) in $\phi_1(r)$ by its Maclaurin series expansion from which has been deleted any finite or infinite number of terms. The new function so obtained is a *nonnegative, monotonic increasing, sub- K_0 function* of r . In like manner, similar functions may be formed from subsequent ϕ functions which involve subtractive functions $l_0(r)$ and $a_0(r)$. We omit proofs.

3.3. The area function. On a surface where $K \leq K_0$, the area function $a(r)$ for a geodesic circle C_r has properties similar to those given for $l(r)$.

THEOREM 3.5. *Let S be an analytic sub- K_0 surface, and let $a(r)$ denote the area of C_r . Then the function*

$$\phi_2(r) \equiv a(r) - a_0(r) \quad (r \geq 0 \text{ on } S)$$

satisfies Condition C.

Proof. Verification is immediate by use of Lemma 3.2 and Theorem 1.2.

COROLLARY 3.6. *If S is an analytic sub- K_0 surface, then $a(r)$ is a monotonic increasing sub- K_0 function of r and satisfies the inequality*

$$a(r) \geq a_0(r) \quad (r \geq 0 \text{ on } S);$$

$a(r)$ is a strictly sub- K_0 function if and only if S is not a K_0 -surface.

Proof. The inequality follows from Theorem 3.5 and the identity

$$a(r) \equiv a_0(r) + \phi_2(r) \quad (r \geq 0 \text{ on } S).$$

We shall find additional theorems for the functions $a(r)$ and $\phi_2(r)$ showing certain subfunction properties of these functions when an additional assumption is made for the surface S . In the sequel we use the following lemma, which shows that certain conditions which clearly imply the sub- K_0 function property for a function also imply this property for its square root.

LEMMA 3.7. *If $g(r)$ is a nonnegative function for which $g'''(r)$ exists in the interval $\alpha \leq r < \beta$, and $g(r)$ satisfies*

$$(3.31) \quad h(\alpha) \equiv 2g(\alpha)g''(\alpha) - [g'(\alpha)]^2 + 4K_0[g(\alpha)]^2 \geq 0$$

and

$$g'''(r) + 4K_0g'(r) \geq 0 \quad (\alpha \leq r < \beta),$$

then $[g(r)]^{1/2}$ is a sub- K_0 function in $\alpha \leq r < \beta$ and is a strictly sub- K_0 function there provided

$$(3.32) \quad g'''(r) + 4K_0g'(r) > 0 \quad (\alpha < r < \beta).$$

Proof. If we let $f(r) \equiv [g(r)]^{1/2}$, then at points where $f(r) \neq 0$ we have

$$\mathfrak{S}f(r) = \frac{1}{4} [g(r)]^{-3/2} h(r).$$

Moreover,

$$(3.33) \quad h'(r) = 2g(g''' + 4K_0g'),$$

so that from the hypotheses we get

$$h(\alpha) \geq 0, \quad h'(r) \geq 0 \quad (\alpha \leq r < \beta),$$

whence $h(r) \geq 0$. Thus $\mathfrak{S}f(r) \geq 0$ at points where $f(r) \neq 0$. And, since the non-negative function $f(r)$ satisfies the subfunction inequality (1.12) for points where $f(r) = 0$, it follows that the continuous function $f(r)$ is a sub- K_0 function for $\alpha \leq r < \beta$.

With (3.31) and (3.32) the nonnegative sub- K_0 (and hence convex) function $g(r)$ can vanish at no more than one point of $\alpha \leq r < \beta$, whence, by (3.33), we have $h(r) > 0$ ($\alpha < r < \beta$). It follows that we have $\mathfrak{S}f(r) > 0$ except for at most one point of $\alpha < r < \beta$, so that $f(r)$ is a strictly sub- K_0 function for $\alpha \leq r < \beta$. This completes the proof of the lemma.

An additional assumption on the surface S causes certain functions immediately to satisfy (3.33) for $r \geq 0$ on S . Thus, if S satisfies $K \leq 4K_0$ for its Gaussian curvature, then a modification of the proof of Theorem 3.3 indicates that we have

$$a'''(r) + 4K_0 a'(r) \equiv l''(r) + 4K_0 l(r) \geq 0 \quad (r \geq 0 \text{ on } S),$$

with equality holding if and only if S is a $4K_0$ -surface. We now determine some functions which have certain subfunction properties in common; these properties are collected in:

CONDITION D. For a given sub- $4K_0$ surface S and for a given pole P of the geodesic polar coordinates on S , a function $\psi(r)$ of the geodesic radius r satisfies *Condition D* provided: $\psi(0) = 0$; for $r \geq 0$ on S , $\psi(r)$ is a continuous monotonic nondecreasing sub- K_0 function of r ; and $\psi(r)$ is a strictly sub- K_0 function except possibly when S is a $4K_0$ -surface.

THEOREM 3.8. *Let S be an analytic sub- $4K_0$ surface, and let $a(r)$ denote the area of the geodesic circle C_r . Then*

$$\psi_1(r) \equiv [a(r)]^{1/2}$$

and

$$\psi_2(r) \equiv [\phi_2(r)]^{1/2} \equiv [a(r) - a_0(r)]^{1/2}$$

satisfy *Condition D*, and $\psi_1(r)$ is a K_0 -function if S is a $4K_0$ -surface.

Proof. We have

$$a'(r) = l(r), \quad a''(r) = l'(r), \quad a'''(r) = l''(r);$$

hence, beside $a(0) = 0$, we have

$$a'(0) = 0, \quad a'''(r) \geq 0 \quad (r \geq 0 \text{ on } S),$$

with

$$a'''(r) + 4K_0 a'(r) > 0$$

for $r > 0$ on S unless $K \equiv 4K_0$. Then for $r \geq 0$ on S , $a(r)$ satisfies the hypotheses on $g(r)$ of Lemma 3.7, so that $\psi_1(r)$ satisfies Condition D for $r \geq 0$ on S , and is a strictly sub- K_0 function if S is not a $4K_0$ -surface. If S is a $4K_0$ -surface, then

$$[a(r)]^{1/2} \equiv \left[\frac{2\pi}{-4K_0} \{ \cosh(\sqrt{-4K_0}r) - 1 \} \right]^{1/2} \equiv \left(\frac{\pi}{-K_0} \right)^{1/2} \sinh(\sqrt{-K_0}r),$$

and thus it is a K_0 -function.

The proof for $\psi_2(r)$ is similar in method and is omitted.

We can find other functions which satisfy Condition D. Let $l_1(r)$ and $a_1(r)$ denote the length of circumference and area, respectively, of the geodesic circle C_r on a $4K_0$ -surface. Formulas for $l_1(r)$ and $a_1(r)$ can be written (see Lemma 2.4), and these expressions serve to define $l_1(r)$ and $a_1(r)$ for a surface S having arbitrary curvature. If S is a sub- $4K_0$ surface, then, by methods analogous to those of Lemma 3.1 and Lemma 3.2, we find the relations

$$l'(r) \geq 2\pi - 4K_0 a(r) \quad (r \geq 0 \text{ on } S),$$

and

$$l''(r) + 4K_0 l(r) \geq 0 \quad (r \geq 0 \text{ on } S),$$

with the equality sign holding for $r > 0$ on S if and only if S is a $4K_0$ -surface; that is,

$$(3.34) \quad l_1'(r) = 2\pi - 4K_0 a_1(r),$$

and

$$l_1''(r) + 4K_0 l_1(r) = 0.$$

THEOREM 3.9. *Let S be an analytic sub- $4K_0$ surface, and let $l(r)$ and $a(r)$ denote the circumference and area function, respectively, of C_r on S . Then the functions*

$$\psi_3(r) \equiv l(r) - l_1(r),$$

$$\psi_4(r) \equiv a(r) - a_1(r),$$

and

$$\psi_5(r) \equiv [a(r) - a_0(r)]^{1/2}$$

satisfy Condition D.

Proof. The method is that used in earlier theorems wherein now we apply the four relations which immediately precede Theorem 3.9.

REMARK. It was indicated earlier that our Condition C reduces to Condition A of [4, p. 289] if $K_0 = 0$. Now if $K_0 = 0$, the assumption that S satisfies $K \leq 4K_0$ imposes no new requirement upon the surface. In fact, our Condition D becomes Condition A if $K_0 = 0$ and if the function $\psi(r)$ is identically zero when the surface is developable.

The role played by the condition $K \leq 4K_0$, when $K_0 \neq 0$ for "square root" functions is indicated in the following theorem.

THEOREM 3.10. *Let S be an analytic sub- K_0 surface, and let $a(r)$ denote the area of C_r on S . Then in order that the function*

$$\psi_1(r) \equiv [a(r)]^{1/2}$$

be a sub- K_0 function of r for every possible pole P , it is necessary and sufficient that S be a sub- $4K_0$ surface.

Proof. The sufficiency has been established in Theorem 3.8.

Now let P_1 be a point of S where $K_1 > 4K_0$, and let P_1 be the pole of a geodesic polar coordinate system. Since S is analytic, there exists a neighborhood of P_1 in which $K > 4K_0$, and hence a value $r_1 > 0$ such that the geodesic circle of radius r_1 lies entirely within this neighborhood. In this coordinate system we have

$$\frac{\partial^2 \mu}{\partial r^2} + 4K_0 \mu < 0 \quad (0 < r \leq r_1 \text{ on } S),$$

and then it easily follows that

$$l''(r) + 4K_0 l(r) < 0 \quad (0 < r \leq r_1 \text{ on } S).$$

By calculation we get that

$$\mathfrak{S}\psi_1(r) = \frac{1}{4} a^{-3/2} [2al' - l^2 + 4K_0 a^2] \equiv \frac{1}{4} a^{-3/2} h(r),$$

where $h(r)$ is the bracketed expression. Then we have that $h(0) = 0$, and

$$h'(r) = 2a(l'' + 4K_0 l) < 0 \quad (0 < r \leq r_1 \text{ on } S);$$

hence $h(r) < 0$ for $0 < r \leq r_1$, and thus also

$$\mathfrak{S}\psi_1(r) < 0 \quad (0 < r \leq r_1 \text{ on } S).$$

Then by Theorem 1.2, $\psi_1(r)$, when evaluated in a coordinate system with such a pole, cannot be a sub- K_0 function.

IV. THE ISOPERIMETRIC INEQUALITY AND RELATED FUNCTIONS

4.1. The isoperimetric inequality. Let L and A denote the perimeter and area, respectively, of a simply connected region bounded by an analytic curve on a surface of nonpositive curvature. The isoperimetric inequality

$$(4.11) \quad \theta \equiv \frac{L^2}{4\pi} - A \geq 0$$

holds for such a region. In fact, the following theorem [6, p. 670-672] has been established:

For an analytic surface S , a necessary and sufficient condition that (4.11) hold for all simply connected regions bounded by analytic curves on S is that $K \leq 0$ on S . Further, if $K \leq 0$ but $K \neq 0$ on S , then the strict sign of inequality holds in (4.11); while if $K \equiv 0$ on S , then the sign of equality holds in (4.11) only for geodesic circles on S .

We shall study the function θ of (4.11) and some modifications of it for sub- K_0 function properties when S is assumed to be a sub- K_0 surface and the region is that determined by a geodesic circle. A well-known generalization of the function θ for geodesic circles on surfaces of constant negative curvature K_0 is the function

$$(4.12) \quad \phi_3(r) \equiv \frac{l^2(r)}{4\pi} + \frac{K_0 a^2(r)}{4\pi} - a(r),$$

which we shall call the *isoperimetric function*.

THEOREM 4.1. *Let S be an analytic sub- K_0 surface, and let $l(r)$ denote the length of the circumference, and $a(r)$ the area, of the geodesic circle C_r on S . Then the isoperimetric function $\phi_3(r)$ satisfies Condition C.*

Proof. Squaring the inequality (3.22) and using (3.21), we obtain

$$4\pi \mathfrak{C} \phi_3(r) \geq [l'^2 + K_0 l^2 - 4\pi^2] \equiv h(r),$$

where $h(r)$ is the function in brackets. Then we see that $h(0) = 0$, and that $h'(r) \geq 0$ for $r \geq 0$ by (3.21); hence $\mathfrak{C} \phi_3(r) \geq 0$ for $r \geq 0$ on S , and thus $\phi_3(r)$ is a sub- K_0 function by Theorem 1.2. The other requirements of Condition C are easily found to be satisfied by $\phi_3(r)$.

COROLLARY 4.2. *If S is an analytic sub- K_0 surface, and $l(r)$ and $a(r)$ as in the theorem, then the function*

$$\theta_1(r) \equiv \frac{l^2(r)}{4\pi} - a(r)$$

is a continuous monotonic nondecreasing sub- K_0 function of r .

Proof. It follows from the proof of Lemma 3.2 that $a(r)$ is a continuous monotonic nondecreasing sub- K_0 function of r , and then that $a^2(r)$ also has these properties by Theorem 1.5. Then, using the positive coefficient $-K_0/4\pi$, we may apply Theorem 1.4 and get

$$\theta_1(r) \equiv \phi_3(r) - \frac{K_0 a^2(r)}{4\pi},$$

so that $\theta_1(r)$ has the properties stated in Corollary 4.2.

COROLLARY 4.3. *If S is an analytic sub- K_0 surface, then*

$$(4.13) \quad \frac{l^2(r)}{4\pi} - a(r) > \frac{l^2(r)}{4\pi} + \frac{K_0 a^2(r)}{4\pi} - a(r) \geq 0 \quad (r > 0 \text{ on } S),$$

where the sign of equality holds for $r > 0$ on S if and only if S is a K_0 -surface.

Proof. This corollary is an immediate consequence of Theorem 4.1 and Corollary 4.2.

4.2. Modifications of the isoperimetric function. We shall consider modifications of the isoperimetric function, $\phi_3(r)$, which are produced by adding certain functions to it and/or by replacing $l(r)$ by $l_0(r)$ or $2\pi r$ and $a(r)$ by $a_0(r)$ or zero. For example, the function $\theta_1(r)$ may be considered a modification of $\phi_3(r)$ formed by replacing the $a^2(r)$ function in $\phi_3(r)$ by zero.

THEOREM 4.4. *Let S be an analytic sub- K_0 surface, and let $l(r)$, $l_0(r)$, and $a(r)$, $a_0(r)$ denote length and area functions associated with the geodesic circle C_r . Then the functions*

$$\phi_4(r) \equiv \frac{l_0(r)l(r)}{4\pi} + \frac{K_0 a_0^2(r)}{4\pi} - a(r)$$

and

$$\phi_5(r) \equiv \frac{l_0(r)l(r)}{4\pi} + \frac{K_0 a_0(r)a(r)}{4\pi} - \frac{a_0(r)}{2} - \frac{a(r)}{2}$$

satisfy Condition C.

Proof. We establish the result that

$$(4.21) \quad l_0 l' - ll_0' \equiv l_0 l' + K_0 a_0 l - 2\pi l \geq 0 \quad (r \geq 0 \text{ on } S).$$

The function on the left is zero when $r = 0$, and its derivative is the nonnegative (by Lemma 3.1) function $l_0(r) \oslash l(r)$; hence (4.21) holds.

Now $\phi_4(0) = 0$, and $\phi_4'(r) \geq 0$ by (4.21) and Theorem 3.3; thus $\phi_4(r)$ is monotonic nondecreasing. The calculation for $\oslash \phi_4(r)$ may be arranged so that

$$\begin{aligned} 4\pi \oslash \phi_4(r) \equiv & [l_0 \oslash l - K_0 l(l - l_0) - 2K_0 a_0(l' - l_0') \\ & - 4\pi K_0(a - a_0) + K_0(l_0^2 + K_0 a_0^2 - 4\pi a_0)]. \end{aligned}$$

Then we have $\mathfrak{S} \phi_4(r) \geq 0$ for $r \geq 0$ on S , since each parenthesis above is non-negative by previous results — the last one, in particular, being identically zero according to Corollary 4.3. Thus $\phi_4(r)$ is a sub- K_0 function by Theorem 1.2. Finally, $\phi_4(r)$ satisfies Condition C since the signs of equality hold in the relations above if and only if S is a K_0 -surface, and obviously $\phi_4(r) \equiv 0$ if S is a K_0 -surface.

For $\phi_5(r)$, we find that $\phi_5(0) = 0$ and that $\phi_5'(r) \geq 0$ by Lemma 3.2. We may arrange the calculation so that

$$\begin{aligned} 4\pi \mathfrak{S} \phi_5(r) &\equiv [l_0 \mathfrak{S} l + (l_0' l' + K_0 l_0 l - 2\pi l_0' - 2\pi K_0 a_0)] \\ &\geq l_0' l' + K_0 l_0 l - 4\pi^2 \equiv h(r), \end{aligned}$$

where $h(r)$ is the function after the inequality sign. Clearly $h(0) = 0$, and we find that

$$h'(r) \equiv l_0' \mathfrak{S} l \geq 0.$$

Hence, $h(r) \geq 0$ for $r \geq 0$ on S , and then Theorem 1.2 ensures that $\phi_5(r)$ is a sub- K_0 function. The other conditions to complete the proof are easily verified.

Theorem 4.4 then admits a corollary which is analogous to the isoperimetric inequality for the functions $\phi_4(r)$ and $\phi_5(r)$; we omit its statement, but we remark that the inequality for $\phi_5(r)$ is sharper than the isoperimetric inequality (4.13) in that it presents a better estimate (greater lower bound) for $l(r)$.

The next theorem presents another function determined by the modification process.

THEOREM 4.5. *For a surface and functions as in Theorem 4.4, the function*

$$\phi_6(r) \equiv \frac{rl(r)}{2} - a(r) - \frac{rl_0(r)}{2} + a_0(r) \equiv \frac{r}{2} [l(r) - l_0(r)] - [a(r) - a_0(r)]$$

satisfies Condition C.

We omit the computations and also the corollary stating the inequality satisfied by $\phi_6(r)$.

It may be noted that with $\phi_6(r)$ satisfying Condition C it readily follows that $\phi_4(r)$ does. For if the function

$$\frac{1}{4\pi} (l_0 - 2\pi r) (l - l_0),$$

which satisfies Condition C (in part by Theorem 1.6), is added to $\phi_6(r)$, we obtain $\phi_4(r)$, which then satisfies Condition C (in part by Theorem 1.4).

Theorem 4.5 suggests a consideration of the substitution of $l(r) - l_0(r)$ and $a(r) - a_0(r)$ for the functions $l(r)$ and $a(r)$. When this substitution is made in the isoperimetric function, we find that the new function does not satisfy our conditions. Nevertheless, in the next theorem we have a result of this procedure.

THEOREM 4.6. *For a surface and functions as in Theorem 4.4, the function*

$$\phi_7(r) \equiv \frac{1}{4\pi} [l(r) - l_0(r)]^2 + \frac{K_0}{4\pi} [a(r) - a_0(r)]^2$$

satisfies Condition C.

Proof. We find that $\phi_7(0) = 0$ and that $\phi_7'(r) \geq 0$ by Lemma 3.2 and Theorem 3.3; thus $\phi_7(r)$ is monotonic nondecreasing. By computation we find that

$$4\pi \zeta \phi_7(r) \geq [2(l' - l'_0)(l' + K_0 a - 2\pi) + K_0(l - l_0)^2 + K_0^2(a - a_0)^2] \equiv h(r),$$

where $h(r)$ is the bracketed expression. We see that $h(0) = 0$, and that its derivative satisfies

$$\begin{aligned} h'(r) &\geq 2[(l'' - l''_0)(l' + K_0 a - 2\pi) + K_0(l - l_0)(l' - l'_0) + K_0^2(a - a_0)(l - l_0)] \\ &= 2(l'' + K_0 l)(l' + K_0 a - 2\pi) \geq 0 \end{aligned}$$

by Lemmas 3.1 and 3.2. Hence $h(r) \geq 0$ for $r \geq 0$ on S , with equality holding if and only if S is a K_0 -surface. It follows that $\phi_7(r)$ satisfies Condition C since obviously $\phi_7(r) \equiv 0$ if S is a K_0 -surface.

Theorem 4.6 admits refinements of the inequalities which appear in Corollary 4.3 and Corollary 3.4.

COROLLARY 4.7. *Let an analytic sub- K_0 surface S be referred to a geodesic polar coordinate system with pole P . Then, for geodesic circles, the isoperimetric function $\phi_3(r)$ and the functions $\phi_5(r)$ and $\phi_7(r)$ satisfy the inequalities*

$$(4.22) \quad \phi_3(r) \geq 2\phi_5(r) \geq 0$$

and

$$\phi_3(r) \geq \phi_7(r) \geq 0,$$

where the signs of equality hold for $r > 0$ on S if and only if S is a K_0 -surface, in which case all functions are identically zero.

Proof. It is easily seen that

$$\phi_7(r) \equiv \phi_3(r) - 2\phi_5(r),$$

and the corollary then follows from Theorems 4.6 and 4.3.

COROLLARY 4.8. *Let an analytic sub- K_0 surface S be referred to a geodesic polar coordinate system with pole P . Then the length of the circumference of a geodesic circle of radius r satisfies the inequality*

$$(4.23) \quad l(r) \geq l_0(r) + \sqrt{-K_0} [a(r) - a_0(r)],$$

where the sign of equality holds for $r > 0$ on S if and only if S is a K_0 -surface.

Proof. Since $K_0 < 0$, $4\pi\phi_7(r)$ has real factors. The factor $\phi_8(r)$, where $\phi_8(r) \equiv l - l_0 + \sqrt{-K_0}(a - a_0)$, satisfies $\phi_8(r) \geq 0$ by Corollaries 3.4 and 3.6; hence so also does the other factor by Corollary 4.7. This other factor yields (4.23).

Less precise relations may be obtained from the isoperimetric function by using the theorems of § 1.3.

THEOREM 4.9. *Let S be an analytic sub- K_0 surface with length and area functions relating to geodesic circles on S as previously defined. Then the functions*

$$\phi_9(r) \equiv \frac{1}{4\pi} [l^2(r) - l_0^2(r)] - [a(r) - a_0(r)],$$

$$\phi_{10}(r) \equiv \frac{1}{4\pi} [l^2(r) - l_0^2(r)] + \frac{K_0}{4\pi} [a^2(r) - a_0^2(r)],$$

and

$$\phi_{11}(r) \equiv l^2(r) - \frac{a(r)}{a_0(r)} l_0^2(r) \equiv l^2(r) - 4\pi a(r) \cosh\left(\frac{\sqrt{-K_0} r}{2}\right)$$

satisfy Condition C.

Proof. We refer to § 1.3 and merely indicate the verification of the desired subfunction property of these functions. Thus, $\phi_9(r)$ results from adding the function $-(K_0/4\pi)[a^2(r) - a_0^2(r)]$, which satisfies Condition C, to the isoperimetric function $\phi_3(r)$. The function $\phi_{10}(r)$ is obtained by adding the function $\phi_2(r) \equiv a(r) - a_0(r)$ to $\phi_3(r)$. And the function $\phi_{11}(r)$ is obtained by adding $-K_0 a(r)[a(r) - a_0(r)]$, which satisfies Condition C, to $4\pi\phi_3(r)$.

4.3. Another kind of modification. The properties of the isoperimetric function and its modifications which we have developed now enable us to introduce new functions which satisfy our conditions. These new functions are produced by replacing each term of an expression by its square root.

THEOREM 4.10. *Let S be an analytic sub- $4K_0$ surface with length and area functions relating to a geodesic circle on S as previously defined. Then the functions*

$$\psi_6(r) \equiv \sqrt{a(r)} - \sqrt{a_0(r)}$$

and

$$\psi_7(r) \equiv \sqrt{a(r)} - \sqrt{a_1(r)}$$

satisfy Condition D.

Proof. We have $\psi_7(0) = 0$, and

$$\psi_7'(r) = \frac{1}{2} \left[\frac{l}{\sqrt{a}} - \frac{l_1}{\sqrt{a_1}} \right] \geq 0$$

for $r \geq 0$ on S by the properties of $\phi_{11}(r)$ of Theorem 4.9, since now $l_1(r)$ and $a_1(r)$ behave analogously to $l_0(r)$ and $a_0(r)$ of that theorem. Hence $\psi_7(r)$ is a monotonic nondecreasing function of r . Then using (3.34) and the isoperimetric identity satisfied by $l_1(r)$, we get

$$\ominus \psi_7(r) = \frac{a^{-3/2}}{4} [2al' - l^2 + 4K_0a^2] \geq 0$$

for $r \geq 0$ on S , since the function in brackets is identical with that which would occur for the function $\psi_1(r)$ of Theorem 3.8. Thus $\psi_7(r)$ satisfies Condition D.

The proof for $\psi_6(r)$ is similar to this for $\psi_7(r)$.

The next theorem presents a modification of the function $\phi_{11}(r)$ of Theorem 4.9.

THEOREM 4.11. *Let S be an analytic sub- K_0 surface with length and area functions relating to geodesic circles on S as previously defined. Then the function*

$$\phi_{12}(r) \equiv l(r)\sqrt{a_0(r)} - l_0(r)\sqrt{a(r)}$$

satisfies Condition C.

Proof. We first establish the inequality

$$(4.31) \quad 2a(r)l'(r) - l^2(r) + K_0 a^2(r) \geq 0 \quad (r \geq 0 \text{ on } S),$$

where the sign of equality holds for $r > 0$ on S if and only if S is a K_0 -surface. The result is immediate, since the function on the left in (4.31) is zero at $r = 0$, and its derivative is nonnegative for $r \geq 0$.

Clearly $\phi_{12}(0) = 0$, and $\phi_{12}(r) \geq 0$ for $r > 0$ on S since $\phi_{11}(r)$ satisfies Condition C. Then, by substituting for $l'(r)$ and $l'_0(r)$ from (4.31), we find that

$$\phi'_{12}(r) \geq \frac{1}{2} \left[\left(\frac{l}{a} + \frac{l_0}{a_0} \right) \left(l\sqrt{a_0} - l_0\sqrt{a} \right) - K_0\sqrt{aa_0} (\sqrt{a} - \sqrt{a_0}) \right] \geq 0.$$

Hence $\phi_{12}(r)$ is monotonic nondecreasing. Then using (3.21) we find that

$$2\mathfrak{S}\phi_{12}(r) \geq \left(\frac{l'l_0}{\sqrt{a_0}} - \frac{ll'_0}{\sqrt{a}} \right) + \left(\frac{l'_0 l}{\sqrt{a_0}} - \frac{l_0^2 l}{2a_0\sqrt{a_0}} - \frac{l'_0 l}{\sqrt{a}} \right) + \left(\frac{l'l_0}{\sqrt{a_0}} - \frac{l'l_0}{\sqrt{a}} + \frac{l^2 l_0}{2a\sqrt{a}} \right).$$

Now using (4.31) in the last two parentheses, we get

$$\begin{aligned}
 4\mathfrak{S}\phi_{12}(r) &\geq 2\left(\frac{l'l_0}{\sqrt{a_0}} - \frac{ll'_0}{\sqrt{a}}\right) + \frac{ll_0}{aa_0} (l\sqrt{a_0} - l_0\sqrt{a}) - K_0 l_0 \sqrt{\frac{a}{a_0}} (\sqrt{a} - \sqrt{a_0}) \\
 &\geq \frac{2}{\sqrt{a}} \left(l'l_0 \sqrt{\frac{a}{a_0}} - ll'_0 \right) \geq \frac{2}{\sqrt{a}} (l'l_0 - ll'_0).
 \end{aligned}$$

Hence, by (4.21), it follows that $\mathfrak{S}\phi_{12}(r) \geq 0$ for $r \geq 0$ on S . Thus, on citing Theorem 1.2 and the obvious fact that $\phi_{12}(r) \equiv 0$ if S is a K_0 -surface, we have shown that $\phi_{12}(r)$ satisfies Condition C.

V. EXTENSIONS AND GENERALIZATIONS

5.1. Geodesic circular sectors. The generalization from a basic configuration of geodesic circles to one of geodesic circular sectors is indicated in [4, p. 296], and its relations apply immediately to this study.

We state some representative results.

THEOREM 5.1. *Let S be an analytic sub- K_0 surface, and let $l(r; \theta_1, \theta_2)$ and $a(r; \theta_1, \theta_2)$ denote respectively the length of the bounding arc and the area of the geodesic circular sector on S with fixed pole P , fixed angle from θ_1 to θ_2 , $\theta_1 < \theta_2$, and geodesic radius r . Then the functions*

$$\phi_{13}(r) \equiv l(r; \theta_1, \theta_2) - l_0(r; \theta_1, \theta_2),$$

$$\phi_{14}(r) \equiv a(r; \theta_1, \theta_2) - a_0(r; \theta_1, \theta_2),$$

and

$$\phi_{15}(r) \equiv \frac{l^2(r; \theta_1, \theta_2)}{2(\theta_2 - \theta_1)} + \frac{K_0 a^2(r; \theta_1, \theta_2)}{2(\theta_2 - \theta_1)} - a(r; \theta_1, \theta_2),$$

$$(r \geq 0, \theta_1 \leq \theta \leq \theta_2 \text{ on } S),$$

satisfy Condition C.

The proof for each function is similar to the proof of the analogous result for the corresponding function for geodesic circles, and will not be given here.

Other functions which satisfy Condition C or Condition D for geodesic circular sectors (the analogues of those for geodesic circles) obviously could be

written. It is clear that, as corollaries, we then obtain certain inequality relations between the length and area functions for a suitably restricted surface.

5.2. Regular super- K_0 surfaces. The preceding results concerning sub- K_0 surfaces hold in the large and are unaffected by singular points. We now describe somewhat analogous results for surfaces whose Gaussian curvature satisfies $K \geq K_0$; such surfaces will be called super- K_0 surfaces. We still assume $K_0 < 0$, although some of the results hold, in the small, for K_0 any constant. In general, our results will hold only on parts of S where there are no singular points of the surface, or of the family of geodesics, other than at the pole of geodesic polar coordinates; and some of the results hold only in the small even where there are no singular points.

A function $f(x)$ is said to be a *super- K_0 function* provided $-f(x)$ is a sub- K_0 function.

A surface S given in geodesic coordinates, or in geodesic polar coordinates, will be said to be *regular* provided there are no singular points on S except, in the case of geodesic polar coordinates, at the pole P .

Lemma 2.1 holds if we add the restriction that S is regular, and replace "sub- K_0 " by "super- K_0 ." Theorems 2.2 and 2.3 hold with the same alterations, and the inequality relations given by (3.21), (3.22), and (4.21) hold with the inequality signs reversed.

THEOREM 5.2. *Let S be a regular analytic super- K_0 surface, and let $l(r)$ and $a(r)$ denote the length and area functions for a geodesic circle C_r . Then the functions $-\phi_j(r)$ ($j = 1, 2, 4, 5, 6, 8$) satisfy Condition C.*

Proof. The theorem follows in routine fashion by an examination of earlier calculations for these functions in relation to (3.21), (3.22), and (4.21) with the inequality signs reversed.

Now consider the isoperimetric function $\phi_3(r)$. We compute $\phi_3'(r)$, and find that $\phi_3(r)$ is *monotonic nonincreasing on any regular super- K_0 surface S , and is monotonic decreasing if S is not a K_0 -surface*. Actually, since $l'(0) = 2\pi$, it follows from a consideration of $\mathfrak{S}\phi_3(r)$ that there is an $r_0 = r_0(S, P)$ such that $\phi_3(r)$ is a super- K_0 function for $0 \leq r \leq r_0$.

From the properties of the functions $\phi_j(r)$ we obtain results for $l(r)$ and $a(r)$. We have

$$l(r) = l_0(r) + \phi_1(r),$$

$$l'(r) = l'_0(r) + \phi'_1(r),$$

and

$$\mathfrak{S}l(r) = \mathfrak{S}\phi_1(r).$$

Since the functions $-\phi_j(r)$ satisfy Condition C on regular super- K_0 surfaces, we have

$$\phi_1(r) \leq 0, \quad \mathfrak{S}\phi_1(r) \leq 0 \quad (r \geq 0 \text{ on } S).$$

It follows that on regular analytic super- K_0 surfaces the function $l(r)$ is a super- K_0 function and satisfies $l(r) \leq l_0(r)$; $l(r)$ is a strictly super- K_0 function and satisfies the strict inequality for $r > 0$ on S if S is not a K_0 -surface. Also, on these surfaces we have $\phi'_1(0) = 0$, so that, since $\phi_1(r)$ is a super- K_0 function, for a given regular analytic super- K_0 surface and for a given pole P on S , either $l(r)$ is monotonic increasing on S or there is an $r_0 = r_0(S, P) > 0$ such that $l(r)$ is monotonic increasing for $0 \leq r \leq r_0$ and monotonic decreasing for $r \geq r_0$ on S .

Again, we have

$$a(r) = a_0(r) + \phi_2(r),$$

$$a'(r) = l'_0(r) + \phi'_1(r) = l'(r),$$

and

$$\mathfrak{S}a(r) = 2\pi + \mathfrak{S}\phi_2(r).$$

On regular analytic super- K_0 surfaces we have

$$\phi_2(r) \leq 0, \quad \phi''_2(0) = 0, \quad \mathfrak{S}\phi_2(r) \leq 0 \quad (r \geq 0 \text{ on } S).$$

Hence on regular super- K_0 surfaces, $a(r)$ satisfies $a(r) \leq a_0(r)$; the strict inequality holds for $r > 0$ on S if S is not a K_0 -surface. Further, for a given regular analytic super- K_0 surface, and for a given pole P on S , either $a(r)$ is a strictly sub- K_0 function, or there is an $r_0 = r_0(S, P) > 0$ such that $a(r)$ is a strictly sub- K_0 function for $0 \leq r \leq r_0$ and a strictly super- K_0 function for $r \geq r_0$ on S . The interval $0 \leq r \leq r_0$ on which $a(r)$ is a sub- K_0 function coincides with the interval on which $l(r)$ is increasing.

From the properties of $\phi_4(r)$, $\phi_5(r)$ and $\phi_6(r)$ described in Theorem 5.2, we deduce some inequalities of interest. Thus, *on regular analytic super- K_0 surfaces we have the inequalities*

$$a(r) \geq \frac{1}{4\pi} [l_0(r)l(r) + K_0 a_0^2(r)],$$

$$a(r) \geq \frac{1}{l'_0(r)} [l_0(r)l(r) - 2\pi a_0(r)],$$

and

$$a(r) \geq a_0(r) - \frac{r}{2} [l_0(r) - l(r)],$$

associated with $\phi_4(r)$, $\phi_5(r)$, and $\phi_6(r)$ respectively, with the signs of equality holding for $r > 0$ on S if and only if S is a K_0 -surface.

When the proof in Theorem 4.6 is examined in light of the new basic inequalities for regular analytic super- K_0 surfaces, we find that $\phi_7(r)$ remains a monotonic, nondecreasing sub- K_0 function. The function $4\pi\phi_7(r)$ is factorable in such a way that $\phi_8(r)$ is a factor; then, by Theorem 5.2, the other factor satisfies the inequality

$$l(r) - l_0(r) - \sqrt{-K_0} [a(r) - a_0(r)] \leq 0.$$

Hence, using this last relation, *on regular analytic super- K_0 surfaces we have the inequalities*

$$l(r) \leq l_0(r) - \sqrt{-K_0} [a_0(r) - a(r)] \leq l_0(r),$$

and

$$a_0(r) \geq a(r) \geq a_0(r) - \frac{1}{\sqrt{-K_0}} [l_0(r) - l(r)],$$

with the signs of equality holding for $r > 0$ on S if and only if S is a K_0 -surface.

The ϕ functions related to geodesic circular sectors (see Theorem 5.1) have analogous properties on regular super- K_0 surfaces.

5.3. Surface characterization. Heretofore we have assumed S to be either

a sub- K_0 surface or a super- K_0 surface. In certain instances we have obtained, in the two cases, conclusions which are distinct except for the dividing class of K_0 -surfaces. Thus by logical exclusion we obtain several characterizations of the indicated classes of surfaces.

For example, a regular analytic surface S is a super- K_0 surface, but not a K_0 -surface, if and only if for each pole P on S we have

$$(5.31) \quad l(r) < l_0(r)$$

for all $r > 0$ on S .

Proof. In § 5.2 we have shown that the condition $K \geq K_0$, $K \neq K_0$, on S implies (5.31). Conversely, if we should have $K_1 < K_0$ at some P_1 on S , then we would have $K < K_0$ throughout some neighborhood of P_1 , and therefore, in the neighborhood, we would have $l(r) > l_0(r)$; also, if we should have $K \equiv K_0$ on S , then we would have $l(r) \equiv l_0(r)$; hence (5.31) implies $K \geq K_0$, $K \neq K_0$ on S .

In the same way we could establish similar results for each function in the following theorem.

THEOREM 5.3. *The regular analytic surface S is i) a sub- K_0 surface, but not a K_0 -surface, ii) a super- K_0 surface, but not a K_0 -surface, or iii) a K_0 -surface, if and only if we have*

$$(5.32) \quad \text{i) } \phi_j(r) > 0, \quad \text{ii) } \phi_j(r) < 0, \quad \text{or} \quad \text{iii) } \phi_j(r) \equiv 0, \\ (j = 1, 2, \dots, 6, 8, 9, \dots, 15),$$

respectively, for all poles P and all $r > 0$ on S .

By Theorem 1.2, it is evident that we might replace (5.32) with the differential conditions

$$\text{i) } \mathfrak{S} \phi_j(r) > 0, \quad \text{ii) } \mathfrak{S} \phi_j(r) < 0, \quad \text{or} \quad \text{iii) } \mathfrak{S} \phi_j(r) \equiv 0 \\ (j = 1, 2, 4, 5, 6, 8, 13, 14).$$

5.4. Geodesically similar curves. The preceding theory may be applied to more general configurations than geodesic circles and sectors. Thus we may study comparison functions which involve length and area functions relating to a class of curves upon an arbitrary surface S as compared to the corresponding curves upon a K_0 -surface or in the plane.

It is evident that r has heretofore played a dual role: it has served as the parameter for the family of geodesic circles (sectors) on S with centers at the pole P , and it also has been a variable of the geodesic polar coordinate system. We now rephrase the previous conditions in terms of the parameter of the family of curves to be considered.

CONDITION A(k). For a given surface S of nonpositive Gaussian curvature, and for a given one-parameter family of curves $C(k)$, a function $\lambda(k)$ of the parameter k satisfies Condition A(k) provided: $\lambda(0) = 0$; for $k \geq 0$, $\lambda(k)$ is a continuous monotonic nondecreasing convex function of k ; $\lambda(k) \equiv 0$ if S is a developable surface, but otherwise is monotonic increasing and strictly convex.

CONDITION C(k). For a given sub- K_0 surface S , and for a given one-parameter family of curves $C(k)$, a function $\tau(k)$ of the parameter k satisfies Condition C(k) provided: $\tau(0) = 0$; for $k \geq 0$, $\tau(k)$ is a continuous monotonic nondecreasing sub- K_0 function of k ; $\tau(k) \equiv 0$ if S is a K_0 -surface, but otherwise $\tau(k)$ is a strictly sub- K_0 function of k .

On a surface S referred to geodesic polar coordinates (r, θ) with a given pole P , we first consider the family of curves $C(k)$ of parameter k given by

$$(5.41) \quad r = kf(\theta), \quad k \geq 0,$$

where $f(\theta)$ admits a continuous derivative and $f(\theta) \geq 1$. We remark that the condition $f(\theta) \geq 1$ is merely a normalization; for, if $f(\theta_0) \leq 1$ and $f(\theta) \neq 0$ in a closed interval, $\alpha \leq \theta \leq \beta$, then $f(\theta)$ is bounded away from zero in (α, β) , say $f(\theta) \geq m > 0$ in (α, β) . Then a new parameter k_1 may be introduced by setting $k = mk_1$, so that

$$r = k_1 f_1(\theta) = k_1 [mf(\theta)],$$

and this representation satisfies our requirements. It may be noted that $f(\theta) \equiv 1$ presents the case of geodesic circles (sectors). The curves $C(k)$ of the family given by (5.41) are said to be *similarly situated* or *homothetic*, and we shall call them geodesically similar.

THEOREM 5.4. *Let S be an analytic surface of nonpositive Gaussian curvature referred to geodesic polar coordinates with given pole P_0 . Let $l_p(k; \alpha, \beta)$ and $l_s(k; \alpha, \beta)$ denote the lengths of the curve of the family $C(k)$ of (5.41) from $\theta = \alpha$ to $\theta = \beta$, ($\alpha < \beta$), for the parameter value k in the plane and on the surface S , respectively, and let $ap(k; \alpha, \beta)$ and $as(k; \alpha, \beta)$ denote the*

areas of the sectors formed by the curve of the family $C(k)$, $\theta = \alpha$, and $\theta = \beta$ ($\alpha < \beta$), for the parameter value k in the plane and on the surface S respectively. Then the functions

$$\lambda_1(k; \alpha, \beta) \equiv l_S(k; \alpha, \beta) - l_P(k; \alpha, \beta)$$

and

$$\lambda_2(k; \alpha, \beta) \equiv a_S(k; \alpha, \beta) - a_P(k; \alpha, \beta)$$

satisfy Condition A (k).

Proof (outlined). For θ_0 fixed and $\alpha < \theta_0 < \beta$, let

$$\lambda_0(k, \theta_0) \equiv \left[\mu^2 + \left(\frac{dr}{d\theta} \right)^2 \right]^{1/2} - \left[r^2 + \left(\frac{dr}{d\theta} \right)^2 \right]^{1/2},$$

where $r = r(\theta)$ is given by (5.41). We find that $\lambda_0(0, \theta_0) = 0$, and that $\partial \lambda_0 / \partial k$ and $\partial^2 \lambda_0 / \partial k^2$ are nonnegative since $\mu(r, \theta_0)$ is a convex function of r . On verifying the other requirements, we have that, for each fixed value θ_0 , $\lambda_0(k, \theta_0)$ satisfies Condition A (k).

Since

$$\lambda_1(k; \alpha, \beta) \equiv \int_{\alpha}^{\beta} \lambda_0(k, \theta) d\theta,$$

it follows (See [4, Theorem 1, p. 287].) that $\lambda_1(k; \alpha, \beta)$ satisfies Condition A (k). If the function $l_S(k; \alpha, \beta)$ alone is considered, then the relations used also indicate that $l_S(k; \alpha, \beta)$ is a convex function of k , that it is strictly convex if S is not a developable surface, and that it is linear (as a function of k) if S is developable.

Now, with (5.41),

$$\lambda_2(k; \alpha, \beta) \equiv \int_{\alpha}^{\beta} \int_0^r [\mu(\rho, \theta) - \rho] d\rho d\theta,$$

and its first and second derivatives are found to be nonnegative by use of the convexity of $\mu(r, \theta)$. The remainder of the argument is direct.

We find other results for the area functions:

THEOREM 5.5. *Let S be an analytic sub- K_0 surface referred to geodesic polar coordinates with given pole P_0 . Let $a_S(k; \alpha, \beta)$, $a_{S_0}(k; \alpha, \beta)$, and $a_P(k; \alpha, \beta)$ denote the areas of the sectors formed by the curve of the family $C(k)$ of (5.41) for the parameter value k , $\theta = \alpha$, and $\theta = \beta$ ($\alpha < \beta$) on the surface S , the K_0 -surface S_0 , and in the plane respectively. Then the function*

$$\lambda_2(k; \alpha, \beta) \equiv a_S(k; \alpha, \beta) - a_P(k; \alpha, \beta)$$

is a monotonic nondecreasing sub- K_0 function of k , and the function

$$\tau_1(k; \alpha, \beta) \equiv a_S(k; \alpha, \beta) - a_{S_0}(k; \alpha, \beta)$$

satisfies Condition C(k).

Proof. By Theorem 5.4, $\lambda_2(k; \alpha, \beta)$ is nonnegative and monotonic nondecreasing. By calculation,

$$\mathfrak{S}_k \lambda_2(k; \alpha, \beta) = \int_{\alpha}^{\beta} \int_0^r \left[\mathfrak{S}_{\rho} \mu + \frac{\partial^2 \mu}{\partial \rho^2} (f^2 - 1) - K_0 \rho \right] d\rho d\theta > 0$$

for $r > 0$ ($k > 0$) on S since $f(\theta) \geq 1$. Hence $\lambda_2(k; \alpha, \beta)$ is a sub- K_0 function of k .

For the other function, we find that

$$\frac{\partial \tau_1}{\partial k} = \int_{\alpha}^{\beta} [\mu(r, \theta) - \mu_0(r, \theta)] f(\theta) d\theta$$

and

$$\mathfrak{S}_k \tau_1(k; \alpha, \beta) = \int_{\alpha}^{\beta} \int_0^r [f^2 \mathfrak{S}_{\rho} (\mu - \mu_0) - K_0 (\mu - \mu_0) (f^2 - 1)] d\rho d\theta.$$

These are nonnegative by the proof of Theorem 3.3, and the rest of the argument is immediate.

5.5. The Steiner configuration. Let C be an arbitrary closed convex curve in the plane, of length L and area F , and let $C(\rho)$ be a curve parallel to C at a distance ρ from it, ρ being measured along the outward normal to C , of length $L(\rho)$ and area $F(\rho)$. The family of curves $C(\rho)$ will be called a *Steiner configuration*; it is a classical result of Steiner [2, p. 128] that

$$L(\rho) = L + 2\pi\rho$$

and

$$F(\rho) = F + \rho L + \pi\rho^2.$$

Generalizations of these formulas for curves lying on a curved surface have been given in [1; 2], and explicit formulas found in the case of surfaces of constant curvature. We shall establish the sub- K_0 function property of some functions which involve the $L(\rho)$ and $F(\rho)$ functions for the Steiner configuration associated with a suitable curve C on an arbitrary sub- K_0 surface, $K_0 < 0$. It is evident that our preceding theory for geodesic circles of center P_0 on S is obtained from a Steiner configuration on S , if the curve C is a geodesic circle of center P_0 on S .

Let the curve C be a simple, closed, bounding, and differentiable curve on the surface S . Introduce a geodesic representation with coordinates (u, v) in which $u = 0$ is the curve C , and $v = \text{constant}$ are the geodesics orthogonal to C ; further, let v be the arc length of C measured positively for motion on the curve which keeps the bounded area to the left, and let u be the arc length of geodesics normal to C . Sufficient conditions for the validity of such a coordinate system in a region of S have been given [1; 2]. We shall assume that our coordinate system is valid and term *admissible* those curves which satisfy the above conditions.

Then, for an admissible curve C of length L and area F , and for ρ fixed, the length $L(\rho)$ of $C(\rho)$ is given by

$$(5.51) \quad L(\rho) = \int_C \mu(\rho, v) dv,$$

and the area $F(\rho)$ of $C(\rho)$ is given by

$$(5.52) \quad F(\rho) = F + \int_C \int_0^\rho \mu(u, v) du dv.$$

For a K_0 -surface, $K_0 < 0$, Abascal [1, p. 843] has shown that these relations simplify to (in our notation)

$$(5.53) \quad L_0(\rho) = L + l_0(\rho) - \frac{K_0}{2\pi} [F l_0(\rho) + L a_0(\rho)],$$

and

$$(5.54) \quad F_0(\rho) = F + a_0(\rho) + \frac{L l_0(\rho)}{2\pi} - \frac{K_0 F a_0(\rho)}{2\pi},$$

where $l_0(\rho)$ and $a_0(\rho)$ are given in Lemma 2.4.

LEMMA 5.6. *The functions $L_0(\rho)$ and $F_0(\rho)$ satisfy the relations*

$$L_0'(\rho) + K_0 F_0(\rho) - 2\pi = 0,$$

$$L_0''(\rho) + K_0 L_0(\rho) = 0,$$

and

$$(5.55) \quad \frac{L_0^2(\rho)}{4\pi} + \frac{K_0 F_0^2(\rho)}{4\pi} - F_0(\rho) = M = \text{constant}$$

for $\rho \geq 0$ on the K_0 -surface S_0 .

Proof. The first two relations follow easily from (5.53), (5.54), and the properties of $l_0(\rho)$ and $a_0(\rho)$. The third relation is immediate since the derivative of its left member is zero.

THEOREM 5.7. *Let S be an analytic sub- K_0 surface, and let $C(\rho)$ denote the curves of the Steiner configuration for an admissible curve C on S . Then the length function $L(\rho)$ is a sub- K_0 function of ρ ; $L(\rho)$ is a strictly sub- K_0 function if S is a strictly sub- K_0 surface, and it is a K_0 -function of ρ if S is a K_0 -surface. Further, the area function $F(\rho)$ is a strictly sub- K_0 function.*

Proof. These properties of $L(\rho)$ were established in Theorem 2.2.

By calculation from (5.52) we get

$$(5.56) \quad \mathfrak{S}_\rho F(\rho) = \int_C \int_0^\rho \mathfrak{S}_u \mu \, du \, dv + \int_C \left(\frac{\partial \mu}{\partial u} \right)_{\rho=0} dv + K_0 F.$$

For our geodesic representation, it is known [7, p. 188] that

$$K_g(v) = \left[\frac{\partial \mu(u, v)}{\partial u} \right]_{u=0},$$

where $K_g(v)$ is the geodesic curvature of C . By the Gauss-Bonnet theorem [7, p. 191], noting that C has no exterior angles, we get

$$(5.57) \quad \int_C K_g(v) dv = 2\pi - \iint_F K \mu du dv \geq 2\pi - K_0 F.$$

since $K \leq K_0$. With $\mathfrak{S}_u \mu \geq 0$ and (5.57), it follows from (5.56) that

$$(5.58) \quad \mathfrak{S}_\rho F(\rho) \equiv L'(\rho) + K_0 F(\rho) \geq 2\pi,$$

and then Theorem 1.2 ensures the result of the theorem.

We shall now make comparison between the length and area functions for Steiner configurations on a sub- K_0 surface S and on the K_0 -surface. However, our expressions may be considered to be functions formed with respect to S alone because of (5.53), (5.54), and the known formulas for $l_0(\rho)$ and $a_0(\rho)$.

THEOREM 5.8. *Let S be an analytic sub- K_0 surface, and let $C(\rho)$ denote the curves of the Steiner configuration for an admissible curve C of length L and area F on S . Let $C_0(\rho)$ denote the curves of the Steiner configuration for any admissible curve C_0 of length $L_0 = L$ and area $F_0 = F$ on the K_0 -surface S_0 . Then the functions*

$$\tau_2(\rho) \equiv L(\rho) - L_0(\rho)$$

and

$$\tau_3(\rho) \equiv F(\rho) - F_0(\rho)$$

satisfy Condition C(ρ), where ρ is the parameter of the family.

Proof. There is equality in (5.57) if S is a K_0 -surface, and the proof using (5.51) and (5.52) is similar to those of Theorem 3.3 and Theorem 3.5.

Theorem 5.8 admits the corollary that *the functions $L(\rho)$ and $F(\rho)$ satisfy the inequalities*

$$L(\rho) \geq L_0(\rho) \equiv L + l_0(\rho) + \frac{|K_0|}{2\pi} [F l_0(\rho) + L a_0(\rho)]$$

and

$$F(\rho) \geq F_0(\rho) \equiv F + a_0(\rho) + \frac{L l_0(\rho)}{2\pi} + \frac{|K_0| F a_0(\rho)}{2\pi};$$

both functions are strictly sub- K_0 functions and satisfy the strict inequalities

for $\rho > 0$ on S if S is not a K_0 -surface, and they satisfy the equalities if S is a K_0 -surface. We remark that the conditions $L_0 = L$ and $F_0 = F$ were imposed to meet the requirements of Condition $C(\rho)$. The sub- K_0 function properties and inequality relations above would hold equally well for any admissible C_0 such that $L_0 \leq L$ and $F_0 \leq F$.

We shall now establish some results for functions involving $L(\rho)$ and $F(\rho)$ which are analogous to the isoperimetric function and to its modifications.

THEOREM 5.9. *Let $L(\rho)$ and $F(\rho)$ be the length and area functions, respectively, of the curves of a Steiner configuration on an analytic sub- K_0 surface S . Then the function*

$$\theta(\rho) \equiv \frac{L^2(\rho)}{4\pi} - F(\rho)$$

is a positive monotonic strictly increasing sub- K_0 function of ρ ; further, if C_0 on the K_0 -surface S_0 satisfies $L_0 = L$ and $F_0 = F$, then the function

$$\tau_4(\rho) \equiv \frac{L^2(\rho) - L_0^2(\rho)}{4\pi} - [F(\rho) - F_0(\rho)]$$

satisfies Condition $C(\rho)$.

Proof. It is known [6] that $\theta(\rho) > 0$ on sub- K_0 surfaces. With (5.58) and Theorem 5.7, routine computations show that $\theta'(\rho)$ and $\ominus\theta(\rho)$ are positive, establishing the properties of $\theta(\rho)$. The properties of $\tau_4(\rho)$ are established in routine manner by the use of (5.58) and Theorem 5.8.

THEOREM 5.10. *Let $L(\rho)$ and $F(\rho)$ be the length and area functions, respectively, of the curves of a Steiner configuration on a sub- K_0 surface S , and let $L_0(\rho)$ and $F_0(\rho)$ be the length and area functions, respectively, of the curves of a Steiner configuration on a K_0 -surface S_0 . Let the admissible curve C_0 on S_0 satisfy $L_0 = L$ and $F_0 = F$. Then the function*

$$\tau_5(\rho) \equiv \left[\frac{L^2(\rho)}{4\pi} + \frac{K_0 F^2(\rho)}{4\pi} - F(\rho) \right] - \left[\frac{L_0^2(\rho)}{4\pi} + \frac{K_0 F_0^2(\rho)}{4\pi} - F_0(\rho) \right]$$

satisfies Condition $C(\rho)$.

Proof. Obviously $\tau_5(0) = 0$, and using (5.55) we get $\tau_5'(\rho) \geq 0$ by (5.58).

By another calculation we find that

$$4\pi\mathfrak{S}\tau_5(\rho) \geq \{2L'(\rho)[L'(\rho) + K_0F(\rho) - 2\pi] + K_0L^2(\rho) + K_0^2F^2(\rho) - 4\pi K_0F(\rho) - 4\pi K_0M\},$$

where the constant M is given by (5.55). We then use (5.58) just as we used (3.22) in the proof of Theorem 4.1, and we get

$$(5.59) \quad 4\pi\mathfrak{S}\tau_5(\rho) \geq [L'^2(\rho) + K_0L^2(\rho) - 4\pi^2 - 4\pi K_0M] \equiv h(\rho)$$

where $h(\rho)$ is the function in brackets. By (5.53) and (5.54), we verify that

$$4\pi K_0M = L_0'^2(\rho) + K_0L_0^2(\rho) - 4\pi^2,$$

and when this is substituted in (5.59), it follows that $h(0) \geq 0$ since $L'(0) \geq L_0'(0)$ by (5.58). By computation and use of Theorem 5.7 we find that $h'(\rho) \geq 0$ for $\rho \geq 0$; hence $h(\rho) \geq 0$ for $\rho \geq 0$, and $\tau_5(\rho)$ is a sub- K_0 function. Since further considerations show that the signs of equality hold above if and only if S is a K_0 -surface, we have, with the final remark that $\tau_5(\rho) \equiv 0$ if S is a K_0 -surface, the result that $\tau_5(\rho)$ satisfies Condition C (ρ).

We remark that the last two theorems imply inequalities for the functions $L(\rho)$ and $F(\rho)$ somewhat similar to (4.13); we omit the formal statements.

Let the symbol $\Phi_k(\rho)$ denote the new functions produced from the functions $\phi_k(r)$ when the functions $r, a(r), a_0(r), l(r),$ and $l_0(r)$, associated with geodesic circles, are replaced by the functions $\rho, F(\rho), F_0(\rho), L(\rho),$ and $L_0(\rho)$, respectively, associated with the Steiner configuration of an admissible curve C . For example,

$$\tau_5(\rho) \equiv \Phi_3(\rho) - M,$$

where M is the constant of (5.55). It may then be verified (indeed almost solely by inspection of the proof that the corresponding $\phi(r)$ function satisfies Condition C) that *the functions*

$$\Phi_k(\rho) - M \qquad (k = 4, 5),$$

and

$$\Phi_k(\rho) \qquad (k = 6, 7, 8, 9, 10, 11),$$

satisfy Condition $C(\rho)$.

Again, we might formulate a Condition $D(\rho)$ which is analogous to Condition D in the way that Condition $C(\rho)$ corresponds to Condition C . Then large parts of the theory in § 3.3 on "square-root" functions are found to apply to similar functions associated with a Steiner configuration. Finally, it may be shown that much of the theory in § 5.2, 5.3 can be generalized to hold for appropriate functions associated with Steiner configurations.

REFERENCES

1. E. V. Abascal, *A generalization of Steiner's formulae*, Bull. Amer. Math. Soc. 53 (1947), 841-844.
2. C. B. Allendoerfer, *Steiner's formulae on a general S^{n+1}* , Bull. Amer. Math. Soc. 54 (1948), 128-135.
3. E. F. Beckenbach, *Generalized convex functions*, Bull. Amer. Math. Soc. 43 (1937), 363-371.
4. ———, *Some convexity properties of surfaces of negative curvature*, Amer. Math. Monthly 55 (1948), 285-301.
5. E. F. Beckenbach and R. H. Bing, *On generalized convex functions*, Trans. Amer. Math. Soc. 58 (1945), 220-230).
6. E. F. Beckenbach and T. Radó, *Subharmonic functions and surfaces of negative curvature*, Trans. Amer. Math. Soc. 35 (1933), 662-674.
7. L. P. Eisenhart, *An introduction to differential geometry*, Princeton, 1940.
8. E. L. Ince, *Ordinary differential equations*, New York, 1944.
9. M. M. Peixoto, *Generalized convex functions and second order differential inequalities*, Bull. Amer. Math. Soc. 55 (1949) 563-572.
10. H. Shniad, *Convexity properties of means of moduli of analytic functions*, Thesis, University of California, Los Angeles, 1948.

UNIVERSITY OF CALIFORNIA, LOS ANGELES, AND
STATE COLLEGE OF WASHINGTON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

R. M. ROBINSON

University of California
Berkeley 4, California

R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

E. HEWITT

University of Washington
Seattle 5, Washington

E. F. BECKENBACH

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN

HERBERT FEDERER

MARSHALL HALL

P. R. HALMOS

HEINZ HOPF

R. D. JAMES

BØRGE JESSEN

PAUL LÉVY

GEORGE PÓLYA

J. J. STOKER

E. G. STRAUS

KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY.

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

NATIONAL BUREAU OF STANDARDS,

INSTITUTE FOR NUMERICAL ANALYSIS

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors except Robinson, whose term expires with the completion of the present volume; they might also be sent to M.M. Schiffer, Stanford University, Stanford, California, who is succeeding Robinson. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1953 BY PACIFIC JOURNAL OF MATHEMATICS

Pacific Journal of Mathematics

Vol. 3, No. 2

April, 1953

William George Bade, <i>An operational calculus for operators with spectrum in a strip</i>	257
E. F. Beckenbach and Lloyd Kenneth Jackson, <i>Subfunctions of several variables</i>	291
David Blackwell, <i>Extension of a renewal theorem</i>	315
L. Carlitz, <i>Some theorems on the Schur derivative</i>	321
Paul Arnold Clement, <i>Generalized convexity and surfaces of negative curvature</i>	333
Merrill M. Flood, <i>On the Hitchcock distribution problem</i>	369
Watson Bryan Fulks, <i>On the unique determination of solutions of the heat equation</i>	387
John W. Green, <i>Length and area of a convex curve under affine transformation</i>	393
William Gustin, <i>An isoperimetric minimax</i>	403
Arthur Eugene Livingston, <i>Some Hausdorff means which exhibit the Gibbs' phenomenon</i>	407
Charles Loewner, <i>On generation of solutions of the biharmonic equation in the plane by conformal mappings</i>	417
Gábor Szegő, <i>Remark on the preceding paper of Charles Loewner</i>	437
Immanuel Marx and G. Piranian, <i>Lipschitz functions of continuous functions</i>	447
Ting-Kwan Pan, <i>The spherical curvature of a hypersurface in Euclidean space</i>	461
Ruth Lind Potter, <i>On self-adjoint differential equations of second order</i>	467
E. H. Rothe, <i>A note on the Banach spaces of Calkin and Morrey</i>	493
Eugene Schenkman, <i>A generalization of the central elements of a group</i>	501
A. Seidenberg, <i>A note on the dimension theory of rings</i>	505