

Pacific Journal of Mathematics

**THE SPHERICAL CURVATURE OF A HYPERSURFACE IN
EUCLIDEAN SPACE**

TING-KWAN PAN

THE SPHERICAL CURVATURE OF A HYPERSURFACE IN EUCLIDEAN SPACE

T. K. Pan

1. Introduction. Let V_n be a hypersurface immersed in a Euclidean space S_{n+1} . Let P be a point of V_n corresponding to the point P' of the hyperspherical representation G_n of V_n . Let V denote the extension of a region ϕ of V_n , and V' the extension of the corresponding hyperspherical region ϕ' of G_n . If the region around P tends to zero, the ratio V'/V tends to a limit Γ , which is called the *spherical curvature* of V_n at P [1, pp. 258-261]. It is found that $\Gamma = |\Omega/g|$, where $g = |g_{ij}|$ and $\Omega = |\Omega_{ij}|$ are respectively the determinants of the coefficients of the first and the second fundamental forms of V_n . In this note, some properties of the spherical curvature are studied, and new interpretations of the Gaussian curvature are derived.

The notation of Eisenhart [2] will be used for the most part.

2. Some properties. Let a real and analytic hypersurface V_n be defined by

$$y^\alpha = y^\alpha(x^1, \dots, x^n) \quad (\alpha = 1, \dots, n+1),$$

referred to a Cartesian coordinate system y^α in a Euclidean space S_{n+1} . Let a vector-field v in V_n be defined by

$$v^\alpha = p^i \partial y^\alpha / \partial x^i \quad (i = 1, \dots, n),$$

where the v^α are real and analytic functions of the x^i . Let C be a curve of V_n . The normal curvature vector of v with respect to C at P is defined as the normal component of the derived vector of the vector-field v along C at P [3]. Let κ denote a nonzero extreme value of the magnitudes of the normal curvature vectors of v with respect to all curves of V_n at P . Then κ , which is called a principal curvature of v at P , is defined by

$$(2.1) \quad |\Psi_{ij} - \kappa^2 g_{ij}| = 0,$$

Received May 13, 1952.

Pacific J. Math. 3 (1953), 461-466

where

$$\Psi_{ij} = \Omega_{ik} \Omega_{jl} p^k p^l / g_{kl} p^k p^l.$$

Since $\|\Psi_{ij}\|$ is of rank 1, there is one such extreme corresponding to a vector-field v . Its value is evidently equal to

$$(2.2) \quad \kappa = (\Psi_{ij} g^{ij})^{1/2} = (H_{ij} p^i p^j / g_{ij} p^i p^j)^{1/2},$$

where H_{ij} is the fundamental tensor of the hyperspherical representation G_n .

The extreme of the principal curvature of a vector-field v at P , as the field varies, is defined by

$$(2.3) \quad |H_{ij} - \kappa^2 g_{ij}| = 0.$$

There are n such extremes $\bar{\kappa}_i$ corresponding to the principal directions for the tensor H_{ij} . Their product is found to be

$$\prod_{i=1}^n \bar{\kappa}_i = |H/g|^{1/2} = |\Omega/g|,$$

since $H = |H_{ij}| = \Omega^2/g$, [1, p. 260]. The principal directions for the tensor H_{ij} and those determined by the tensor Ω_{ij} are identical, since the principal curvature of a principal vector-field can easily be shown equal to the normal curvature of the corresponding line of curvature. Hence we have:

THEOREM 2.1. *The spherical curvature of a V_n at P is equal to the product of the extreme principal curvatures of vector-fields in V_n at P , which is the same as the product of principal curvatures of V_n at P .*

Since S_{n+1} is Euclidean, the equations of Gauss are

$$(2.4) \quad R_{ijkl} = \Omega_{ik} \Omega_{jl} - \Omega_{il} \Omega_{jk}.$$

Multiplying (2.4) by g^{ik} and summing with respect to i and k , we obtain

$$(2.5) \quad H_{jl} = M \Omega_{jl} + R_{jl},$$

where M is the mean curvature of V_n , and where R_{jl} is the Ricci tensor. When V_n is a minimal hypersurface, we have $M = 0$, and the Ricci tensor is identical

with the fundamental tensor of G_n . If $M \neq 0$, we have

$$(2.6) \quad H_{ij} p^i p^j = R_{ij} p^i p^j$$

if and only if v is an asymptotic vector-field. If v is a unit asymptotic vector-field, we notice, from (2.2), (2.6), and the equality

$$R_{ij} \lambda_h^i \lambda_h^j = - \sum_{k=1}^n \gamma_{hk},$$

that the square of the principal curvature of v at P is numerically equal to the sum of the Riemannian curvatures determined by v and $n - 1$ other mutually orthogonal unit vectors orthogonal to v at P . Hence we have established the following result:

THEOREM 2.2. *The square of the principal curvature of an asymptotic vector-field at P in V_n is numerically equal to the mean curvature of V_n at P for the corresponding asymptotic direction.*

The extreme of the principal curvatures κ of asymptotic vector-fields at P in V_n is defined by

$$|R_{ij} - \kappa^2 g_{ij}| = 0.$$

There are n such extreme values corresponding to the principal directions for the Ricci tensor R_{ij} . Their product is evidently equal to $|\Omega/g|$, if V_n is minimal. Hence we have:

THEOREM 2.3. *The principal curvatures of asymptotic vector-fields at P in V_n attain their extreme values in the principal directions for the Ricci tensor.*

THEOREM 2.4. *The spherical curvature of a minimal V_n at P is the product of the principal curvatures of the n vector-fields at P corresponding to the principal directions for the Ricci tensor.*

3. The Gaussian curvature. When $n = 2$, Γ is called the spherical curvature of a surface S in an ordinary space. It coincides in absolute value with the Gaussian curvature K of S . The principal curvature of a vector-field v in V_n for $n = 2$ coincides in absolute value with the principal curvature of v in S , [3]. The extreme principal curvatures of vector-fields in V_n for $n = 2$ coincide in absolute value with the principal curvatures of S . The mean curvature of V_n for

$n = 2$ is identical with the Gaussian curvature of S . Hence Theorems 2.1 and 2.2 lead directly to the following new interpretations of the Gaussian curvature:

THEOREM 3.1. *The Gaussian curvature of S at P is the product of the extreme principal curvatures of vector fields of S at P , and is the negative of the square of the magnitude of the Gaussian representation of a unit arc along an asymptotic line from P in S .*

Let p^α and q^α be two distinct conjugate vector fields in S . Then we have

$$q^\beta = e^{\beta\mu} d_{\alpha\mu} p^\alpha \quad (\alpha, \beta, \mu = 1, 2),$$

where $d_{\alpha\mu}$ is the second fundamental tensor of S . The principal curvatures of the vector-fields p^α and q^α are respectively equal to

$$e\rho_p = (h_{\alpha\beta} p^\alpha p^\beta / g_{\alpha\beta} p^\alpha p^\beta)^{1/2},$$

$$e\rho_q = (hg_{\alpha\beta} p^\alpha p^\beta / gh_{\alpha\beta} p^\alpha p^\beta)^{1/2},$$

where $h_{\alpha\beta}$ is the third fundamental tensor of S . Hence their product is

$$(3.1) \quad (e\rho_p)(e\rho_q) = (h/g)^{1/2}.$$

The expression on the right side of (3.1) is equal to eK , where e is +1 or -1 according as K is positive or negative at the point under consideration. At an elliptic point, the principal curvatures of all vector-fields are of the same sign. At a hyperbolic point, the principal curvatures of two vector-fields are different in sign if they lie in different sections separated by the asymptotic lines of S . Consequently, the principal curvatures of two conjugate vector-fields have opposite signs, since conjugate directions are separated by the asymptotic directions of the surface. Hence at an elliptic point of S , the product of the principal curvatures of two conjugate vector-fields is positive; while at a hyperbolic point of S , it is negative. At a parabolic point the normal curvature of any vector-field with respect to any curve is zero. We may consider that every direction in S at a parabolic point is both an asymptotic direction and a principal direction of a vector-field which is to be considered. Hence at a parabolic point the principal curvature of any vector-field is zero; consequently, the product of the principal curvatures of two conjugate vector-fields is zero. Thus the following theorem is proved:

THEOREM 3.2. *The Gaussian curvature of S at P is the product of the principal curvatures of any two distinct conjugate vector-fields in S at P .*

The sum of the squares of the principal curvatures of the two conjugate vector-fields is found to be

$$(e\rho_p)^2 + (e\rho_q)^2 = M(\kappa_p + \kappa_q) - 2K,$$

where κ_p and κ_q are the normal curvatures of the curves of the two fields, and where M is the mean curvature of S . By Theorem 3.2 the above equation can be written as

$$(3.2) \quad (e\rho_p + e\rho_q)^2 = M(\kappa_p + \kappa_q).$$

Since the product of the normal radii at a point in conjugate directions is a maximum for characteristic lines, and a minimum for lines of curvature, and since the sum of normal radii in conjugate directions is constant, we obtain from (3.2) the following result:

THEOREM 3.3. *The sum of the principal curvatures of two conjugate vector-fields at P is the mean proportional between the mean curvature at P of S and the sum of the normal curvatures in the two conjugate directions at P . The square of the sum of the principal curvatures of two conjugate vector-fields at P is a maximum for the principal vector-fields of S , and a minimum for the characteristic vector-fields of S .*

Let m ($m > 2$) directions be such that the angle of two adjoining directions is $2\pi/m$. Let the principal curvatures of the vector-fields in such directions be denoted by $e\rho_1, e\rho_2, \dots, e\rho_m$. Then

$$\frac{1}{m} \sum_{i=1}^{m>2} (e\rho_i)^2 = \frac{1}{2} M^2 - K,$$

since

$$\frac{1}{m} \left(\sum_{i=1}^{m>2} \kappa_{p_i} \right) = \frac{1}{2} M,$$

where κ_{p_i} are the normal curvatures of the curves of the corresponding vector-fields.

THEOREM 3.4. *One m th of the sum of the squares of the principal curvatures of m (> 2) vector-fields at P , such that the angle of two adjoining vectors of these fields at P is $2\pi/m$, is constant and is the same for any m greater than two. The constant is half of the square of the mean curvature of S minus the Gaussian curvature of S at P .*

It is easy to prove that the principal direction of a vector-field in S is orthogonal to the curve of the field if and only if the vector-field is an asymptotic field. Let p^α be an asymptotic vector-field in S . Then its orthogonal trajectories are defined by

$$du^\beta = e^{\beta\mu} g_{\alpha\mu} p^\alpha.$$

The principal curvature of the asymptotic vector-field p^α is given by

$$(e\rho_p) = d_{\alpha\beta} p^\alpha e^{\beta\mu} g_{\gamma\mu} p^\gamma / [(g_{\alpha\beta} p^\alpha p^\beta) (g_{\alpha\beta} e^{\alpha\mu} g_{\gamma\mu} p^\gamma e^{\beta\lambda} g_{\sigma\lambda} p^\sigma)]^{1/2},$$

which after simplification becomes

$$(e\rho_p) = \epsilon^{\beta\mu} d_{\alpha\beta} g_{\gamma\mu} p^\alpha p^\gamma / g_{\alpha\beta} p^\alpha p^\beta = \tau_g,$$

where τ_g is the geodesic torsion of the curve of the asymptotic vector-field.

THEOREM 3.5. *The principal curvature of an asymptotic vector-field at P in S is equal to the geodesic torsion at P of the curve of the field, or simply the torsion at P of the corresponding asymptotic line.*

From Theorem 3.1 and Theorem 3.5 we immediately obtain the first part of the theorem of Enneper, that the square of the torsion of a real asymptotic line at a point is equal to the absolute value of the total curvature of the surface at the point. By the second part of the same theorem we notice that *the principal curvatures of the asymptotic vector-fields in S are different in sign.*

REFERENCES

1. T. Levi-Civita, *The absolute differential calculus*, Blackie & Son, London, 1947.
2. L. P. Eisenhart, *Riemannian geometry*, Princeton University Press, Princeton, 1949.
3. T. K. Pan, *Normal curvature of a vector field*, Amer. J. Math. 74 (1952), 955-966.

UNIVERSITY OF CALIFORNIA, BERKELEY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

R. M. ROBINSON

University of California
Berkeley 4, California

R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

E. HEWITT

University of Washington
Seattle 5, Washington

E. F. BECKENBACH

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN

HERBERT FEDERER

MARSHALL HALL

P. R. HALMOS

HEINZ HOPF

R. D. JAMES

BØRGE JESSEN

PAUL LÉVY

GEORGE PÓLYA

J. J. STOKER

E. G. STRAUS

KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY.

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

NATIONAL BUREAU OF STANDARDS,

INSTITUTE FOR NUMERICAL ANALYSIS

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors except Robinson, whose term expires with the completion of the present volume; they might also be sent to M.M. Schiffer, Stanford University, Stanford, California, who is succeeding Robinson. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1953 BY PACIFIC JOURNAL OF MATHEMATICS

Pacific Journal of Mathematics

Vol. 3, No. 2

April, 1953

William George Bade, <i>An operational calculus for operators with spectrum in a strip</i>	257
E. F. Beckenbach and Lloyd Kenneth Jackson, <i>Subfunctions of several variables</i>	291
David Blackwell, <i>Extension of a renewal theorem</i>	315
L. Carlitz, <i>Some theorems on the Schur derivative</i>	321
Paul Arnold Clement, <i>Generalized convexity and surfaces of negative curvature</i>	333
Merrill M. Flood, <i>On the Hitchcock distribution problem</i>	369
Watson Bryan Fulks, <i>On the unique determination of solutions of the heat equation</i>	387
John W. Green, <i>Length and area of a convex curve under affine transformation</i>	393
William Gustin, <i>An isoperimetric minimax</i>	403
Arthur Eugene Livingston, <i>Some Hausdorff means which exhibit the Gibbs' phenomenon</i>	407
Charles Loewner, <i>On generation of solutions of the biharmonic equation in the plane by conformal mappings</i>	417
Gábor Szegő, <i>Remark on the preceding paper of Charles Loewner</i>	437
Immanuel Marx and G. Piranian, <i>Lipschitz functions of continuous functions</i>	447
Ting-Kwan Pan, <i>The spherical curvature of a hypersurface in Euclidean space</i>	461
Ruth Lind Potter, <i>On self-adjoint differential equations of second order</i>	467
E. H. Rothe, <i>A note on the Banach spaces of Calkin and Morrey</i>	493
Eugene Schenkman, <i>A generalization of the central elements of a group</i>	501
A. Seidenberg, <i>A note on the dimension theory of rings</i>	505