

# Pacific Journal of Mathematics

**THE NEUMANN PROBLEM FOR THE HEAT EQUATION**

WATSON BRYAN FULKS

# THE NEUMANN PROBLEM FOR THE HEAT EQUATION

W. FULKS

**1. Introduction.** By the Neumann problem we mean the following boundary-value problem: to determine the solution  $u(x, t)$  of the equation

$$(1.1) \quad u_{xx}(x, t) - u_t(x, t) = 0$$

in the rectangle or semi-infinite strip  $R^{(b,c)}: \{b < x < c; a < t < T \leq \infty\}$ , given  $u(x, a)$  on  $b < x < c$  and  $u_x(b, t)$  and  $u_x(c, t)$  on  $a < t < T$ . There is a formula in terms of the Green's function (essentially given by Doetsch in [2, p. 361]) which gives the answer to this problem if the closed rectangle is in the interior of a larger region in which  $u(x, t)$  is a continuous solution of (1.1). This formula is as follows: let  $d = c - b$ , and let

$$F^{(b,c)}(x, t; y, s) = \frac{1}{2d} \left[ \vartheta_3 \left( \frac{x-y}{2d}, \frac{t-s}{d^2} \right) + \vartheta_3 \left( \frac{x+y-2b}{2d}, \frac{t-s}{d^2} \right) \right]$$

where  $\vartheta_3$  is the Jacobi Theta function; then

$$(1.2) \quad u(x, t) = \int_b^c F^{(b,c)}(x, t; y, a) u(y, a) dy - \int_a^t F^{(b,c)}(x, t; b, s) u_x(b, s) ds \\ + \int_a^t F^{(b,c)}(x, t; c, s) u_x(c, s) ds.$$

The purpose of this paper is to extend the use of formula (1.2) in the following manner: we will give conditions under which a solution of the heat equation can be written in the form (1.2) wherein  $u(a, y) dy$ , etc., are replaced by  $dA(y)$  or by  $a(y) dy$ , where  $A(y) \in BV$  (that is, of bounded variation) or  $a(y) \in L$ . And we will examine the senses in which these extensions of formula (1.2) solve the boundary-value problem; that is, the manner in which the solutions tend to the prescribed boundary data for approach to a boundary point. Furthermore, we will obtain criteria for the unique determination of the solutions of these generalized boundary-value problems.

---

Received June 30, 1952. The work on this paper was performed under sponsorship of the Office of Naval Research, Contract Nonr-386(00); (NR 044 004).

*Pacific J. Math.* 3 (1953), 567-583

We will normalize our rectangle to be  $R: \{0 < x < 1, 0 < t < T \leq \infty\}$ , and for this region we will delete the superscripts from the Green's function and denote it simply by  $F(x, t; y, s)$ . And we will denote by  $H$  the class of solutions of (1.1) for which both  $u_{xx}$  and  $u_t$  are continuous.

It will be convenient to display here the formula (see [2, p. 307])

$$(1.3) \quad \mathcal{D}_3(x/2, t) = (\pi t)^{-1/2} \sum_{n=-\infty}^{\infty} \exp \left[ \frac{-(x+n)^2}{4t} \right],$$

from which it is clear that  $F^{(b,c)}(x, t; y, s)$  is a uniformly continuous function of all six variables if  $d$  is bounded away from both zero and infinity, and if the point  $(x, t)$  is bounded away from the point  $(y, s)$ .

It also follows easily from (1.3) that

$$F_x(x, t; 0, s) = -G_y(x, t; 0, s) \quad \text{and} \quad F_x(x, t; 1, s) = -F_x(1-x, t; 0, s),$$

where

$$G(x, t; y, s) = \frac{1}{2} \left[ \mathcal{D}_3 \left( \frac{x-y}{2}, t-s \right) - \mathcal{D}_3 \left( \frac{x+y}{2}, t-s \right) \right]$$

is the Green's function for the corresponding Dirichlet problem. (See [3; 4; 5; 6; 7].)

**2. The Stieltjes integral representation.** Our first main theorem gives the solution to one of the generalized boundary-value problems.

**THEOREM 1.** For  $u(x, t)$  to be representable in  $R$  by

$$(2.1) \quad u(x, t) = \int_0^1 F(x, t; y, 0) dA(y) - \int_0^t F(x, t; 0, s) dB(s) \\ + \int_0^t F(x, t; 1, s) dC(s),$$

where  $A(y) \in BV$  ( $0 \leq y \leq 1$ ) and  $B(s), C(s) \in BV$  ( $0 \leq s \leq s_0$ ) for every  $s_0 < T$ , it is necessary and sufficient that

$$(1) \quad u(x, t) \in H \text{ in } R,$$

(2)  $\int_0^t |u_x(x, s)| ds < M_t$  uniformly for  $0 < x \leq x_0$  and  $x_1 \leq x < 1$  for some  $x_0, x_1$ , where  $M_t$  depends only on  $t$ ,

$$(3) \int_0^1 |u(y, t)| dy \leq M \text{ uniformly for } 0 < t \leq t_0 \text{ for some } t_0.$$

*Proof.* To prove the sufficiency, let  $(x, t) \in R$ . Then there exist  $a, b, c > 0$  such that  $u(x, t)$  is given by (1.2). But, by condition (3),

$$A_a(x) = \int_0^x u(y, a) dy \in BV[0 \leq x \leq 1]$$

uniformly in  $a$  for  $0 < a \leq t_0$ . Hence the uniformity holds for any sequence of values of  $a$  tending to zero, and thus by the well-known convergence theorems of Helly and Bray (see, for example, [9, p. 29-31]) there exists a subsequence  $\{a_n\}$  and a function  $A(x) \in BV(0 \leq x \leq 1)$ , to which  $A_{a_n}(x)$  converges substantially, such that

$$\lim_{n \rightarrow \infty} \int_b^c F^{(b,c)}(x, t; y, a_n) dA_{a_n}(y) = \int_b^c F^{(b,c)}(x, t; y, 0) dA(y).$$

Then (1.2) becomes

$$(2.2) \ u(x, t) = \int_b^c F^{(b,c)}(x, t; y, 0) dA(y) - \int_0^t F^{(b,c)}(x, t; b, s) u_x(b, s) ds + \int_0^t F^{(b,c)}(x, t; c, s) u_x(c, s) ds,$$

where the existence of the two latter integrals is guaranteed by condition (2).

Furthermore,

$$B_b(t) = \int_0^t u_x(b, s) ds \text{ and } C_c(t) = \int_0^t u_x(c, s) ds \in BV[0 \leq t \leq t_0]$$

for every  $t_0 < T$  uniformly for  $0 < b \leq x_0$  and  $x_1 \leq c < 1$ . Hence the uniformity holds for any sequence of values of  $b$  tending to zero and of  $c$  tending to one. Hence there exist subsequences  $\{b_n\}$  and  $\{c_n\}$  and functions

$$B(t), C(t) \in BV(0 \leq t \leq t_0)$$

such that

$$\lim_{n \rightarrow \infty} \int_0^t F^{(b_n, c_n)}(x, t; b_n, s) dB_{b_n}(s) = \int_0^t F(x, t; 0, s) dB(s)$$

and

$$\lim_{n \rightarrow \infty} \int_0^t F^{(b_n, c_n)}(x, t; c_n, s) dC_{c_n}(s) = \int_0^t F(x, t; 1, s) dC(s).$$

Hence  $u(x, t)$  has the representation asserted.

We will later show that  $A, B, C$  are independent of the particular sequences of  $a, b, c$  used here (see Theorem 3).

To prove the necessity of condition (1) we must differentiate under the integral sign. The only difficulty encountered in this is the disposition of the terms which arise from the variable upper limit. If, however, one forms a difference quotient it is easy to see that the contribution arising from the variability of the upper limit must always vanish, due to the strong convergence to zero of the kernel as  $s \rightarrow t - 0$ .

To establish (2) we write

$$\begin{aligned} u_x(x, t) &= \int_0^1 F_x(x, t; y, 0) dA(y) - \int_0^t F_x(x, t; 0, s) dB(s) \\ &\quad + \int_0^t F_x(x, t; 1, s) dC(s) \\ &= \int_0^1 F_x(x, t; y, 0) dA(y) + \int_0^t G_y(x, t; 0, s) dB(s) \\ &\quad + \int_0^t G_y(1-x, t; 0, s) dC(s) \\ &= U_1(x, t) + U_2(x, t) + U_3(x, t). \end{aligned}$$

Now

$$|U_2(x, t)| \leq \int_0^t G_y(x, t; 0, s) |dB(s)| = v_2(x, t)$$

and

$$|U_3(x, t)| \leq \int_0^t G_y(1-x, t; 0, s) |dC(s)| = v_3(x, t),$$

where  $v_2(x, t)$  and  $v_3(x, t)$  are nonnegative solutions of (1.1). Then, by [3, p. 22-23] and [7, p. 373],  $v_2(x, t)$  and  $v_3(x, t)$  must satisfy condition (2). Hence so must  $U_2(x, t)$  and  $U_3(x, t)$ .

To examine  $U_1(x, t)$  we need to note that, by (1.3), for  $0 < x, y < 1$ ,

$$F_x(x, t; y, 0) = -\frac{x-y}{4\pi^{1/2}t^{3/2}} \exp\left[-\frac{(x-y)^2}{4t}\right] - \frac{(x+y)^2}{4\pi^{1/2}t^{3/2}} \exp\left[-\frac{(x+y)^2}{4t}\right] \\ - \frac{(x+y-2)}{4\pi^{1/2}t^{3/2}} \exp\left[-\frac{(x+y-2)^2}{4t}\right] + \bar{u}_1(x, y, t),$$

where  $\bar{u}_1$  is bounded, say  $|\bar{u}_1| \leq B_1$ . Then

$$\int_0^t |U_1(x, s)| ds \leq \int_0^t \int_0^1 \sum_{n=1}^3 \frac{|a_n|}{4\pi^{1/2}s^{3/2}} \exp\left[-\frac{a_n^2}{4s}\right] |dA(y)| ds \\ + tB_1 V_A(1),$$

where  $a_1 = x - y$ ,  $a_2 = x + y$ ,  $a_3 = x + y - 2$ , and  $V_A(1)$  is the variation of  $A$ .  
Then

$$\int_0^t |U_1(x, s)| ds \leq \frac{1}{4\pi^{1/2}} \int_0^1 \int_0^t \sum_{n=1}^3 \frac{|a_n|}{s^{3/2}} \exp\left[-\frac{a_n^2}{4s}\right] ds |dA(y)| \\ + tB_1 V_A(1), \\ = \frac{1}{2\pi^{1/2}} \int_0^1 \sum_{n=1}^3 \int_{a_n^2/4t}^\infty e^{-s} s^{-1/2} ds |dA(y)| + tB_1 V_A(1), \\ \leq \frac{3}{2\pi^{1/2}} \int_0^1 \int_0^\infty e^{-s} s^{-1/2} ds |dA(y)| + tB_1 V_A(1), \\ = (3/2 + tB_1) V_A(1),$$

the change of order of integration being permissible by Fubini's theorem. Since  $U_1(x, t)$ ,  $U_2(x, t)$ , and  $U_3(x, t)$  separately satisfy condition (2), so must their sum,  $u_x(x, t)$ .

To verify condition (3) we write

$$u(x, t) = \int_0^1 - \int_0^t + \int_0^t = u_1(x, t) + u_2(x, t) + u_3(x, t),$$

and first consider

$$u_2(x, t) = - \int_0^t F(x, t; 0, s) dB(s).$$

But, by (1.3),

$$F(x, t; 0, s) = \pi^{-1/2}(t-s)^{-1/2} \exp\left[-\frac{x^2}{4(t-s)}\right] + \bar{u}_2(x, t, s),$$

where  $\bar{u}_2$  is bounded, say by  $B_2$ . Then

$$\begin{aligned} \int_0^1 |u_2(x, t)| dx &\leq \int_0^t \pi^{-1/2}(t-s)^{-1/2} \int_0^1 \exp\left[-\frac{x^2}{4(t-s)}\right] dx |dB(s)| \\ &\quad + B_2 V_B(t), \\ &\leq \int_0^t \pi^{-1/2}(t-s)^{-1/2} \int_0^\infty \exp\left[-\frac{x^2}{4(t-s)}\right] dx |dB(s)| + B_2 V_B(t), \\ &= (1 + B_2) V_B(t). \end{aligned}$$

Similarly,

$$\int_0^1 |u_3(x, t)| dx \leq (1 + B_3) V_C(t).$$

We turn now to  $u_1(x, t)$ :

$$\int_0^1 |u_1(x, t)| dx \leq \int_0^1 \int_0^1 F(x, t; y, 0) dx |dA(y)|.$$

But, again by (1.3),

$$\begin{aligned} F(x, t; y, 0) &= \frac{1}{2} (\pi t)^{-1/2} \left\{ \exp\left[-\frac{(x-y)^2}{4t}\right] + \exp\left[-\frac{(x+y)^2}{4t}\right] \right. \\ &\quad \left. + \exp\left[-\frac{(x+y-2)^2}{4t}\right] \right\} + \bar{u}_4(x, y, t), \end{aligned}$$

where  $\bar{u}_4$  is bounded by, say,  $B_4$ . Then

$$\begin{aligned} \int_0^1 |u_1(x, t)| dx &\leq \frac{1}{2} (\pi t)^{-1/2} \int_0^1 \int_0^1 \left\{ \exp\left[-\frac{(x-y)^2}{4t}\right] + \exp\left[-\frac{(x+y)^2}{4t}\right] \right. \\ &\quad \left. + \exp\left[-\frac{(x+y-2)^2}{4t}\right] \right\} dx |dA(y)| + B_4 V_A(1), \\ &\leq \frac{1}{2} (\pi t)^{-1/2} \int_0^1 \left\{ \int_{-1}^1 \exp\left[-\frac{(x-y)^2}{4t}\right] dx \right. \\ &\quad \left. + \int_0^1 \exp\left[-\frac{(x+y-2)^2}{4t}\right] dx \right\} |dA(y)| + B_4 V_A(1), \end{aligned}$$

$$\begin{aligned} &\leq \frac{1}{2} (\pi t)^{-1/2} \int_0^1 2 \int_{-\infty}^{\infty} \exp \left[ -\frac{x^2}{4t} \right] dx |dA(y)| + B_4 V_A(1), \\ &\leq V_A(1) (2 + B_4) \end{aligned}$$

Hence, for  $0 < t \leq t_0$ ,

$$\int_0^1 |u(x, t)| dx \leq V_A(1) (2 + B_4) + V_B(t_0) (1 + B_2) + V_C(t_0) (1 + B_3) = M$$

This completes the proof.

**3. The behavior at the boundary.** We are now prepared to examine in detail the behavior near the boundary of solutions of our generalized boundary value problem considered in section 2. The main result of the section is:

**THEOREM 2.** *If  $u(x, t)$  is representable in  $R$  by (2.1), then*

$$(1) \quad \lim_{t \rightarrow 0^+} u(x, t) = A'(x)$$

and

$$(2) \quad \lim_{x \rightarrow 0^+} u_x(x, t) = B'(t-0); \quad \lim_{x \rightarrow 1-0} u_x(x, t) = C'(t-0)$$

$$\left( \text{where } B'(t-0) = \lim_{h \rightarrow 0^+} \frac{B(t-0) - B(t-h)}{h}, \text{ and similarly for } C'(t-0) \right),$$

wherever the derivatives in question exist.

*Proof.* If  $u(x, t)$  is representable by (2.1), let

$$u(x, t) = \int_0^1 - \int_0^t + \int_0^t = u_1(x, t) + u_2(x, t) + u_3(x, t)$$

as before. Let  $I$  be any open interval whose closure is contained in  $\{0 < x < 1\}$ . Then for  $x \in I$ ,  $F(x, t; 0, s)$  and  $F(x, t; 1, s)$  both converge uniformly to zero as  $t \rightarrow 0^+$ , as can be seen from (1.3). Then clearly  $u_2(x, t), u_3(x, t) \rightarrow 0$  as  $t \rightarrow 0^+$ , for  $x \in I$ .

Also, for  $x \in I$ , by (1.3),

$$F(x, t; y, 0) = (4\pi t)^{-1/2} \exp \left[ -\frac{(x-y)^2}{4t} \right] + o(1)$$

uniformly as  $t \rightarrow 0^+$ . Hence



$$u_1(x, t) = \int_0^1 (4\pi t)^{-1/2} \exp \left[ -\frac{(x-y)^2}{4t} \right] dA(y) + o(1).$$

Then (see [3, p. 25-26 and 65-66] and [7, p. 393-394])

$$\lim_{t \rightarrow 0^+} u_1(x, t) = A'(x)$$

wherever this derivative exists. Since any  $x \in \{0 < x < 1\}$  can be caught in such an  $I$ , this establishes (1).

To verify conclusion (2) we write, as before,

$$\begin{aligned} u_x(x, t) &= \int_0^1 F_x(x, t; y, 0) dA(y) + \int_0^t G_y(x, t; 0, s) dB(s) \\ &\quad + \int_0^t G_y(1-x, t; 0, s) dC(s), \\ &= U_1(x, t) + U_2(x, t) + U_3(x, t). \end{aligned}$$

As  $x \rightarrow 0^+$ ,  $U_1(x, t)$  and  $U_3(x, t)$  vanish since the kernels converge uniformly to zero, and as  $x \rightarrow 1-0$ ,  $U_1(x, t)$  and  $U_2(x, t)$  vanish for the same reason. Then by [5],  $u_x(x, t)$  tends to  $B'(t-0)$  or  $C'(t-0)$  according as  $x$  tends to zero or one, whenever the derivatives exist.

We can now give criteria for the existence of boundary values of the function  $u(x, t)$  itself on the sides  $x=0$ , and  $x=1$ .

**COROLLARY 1.** *If  $u(x, t)$  is representable in  $R$  by (2.1), then  $u(0+, t)$  exists if  $B'(t-0)$  does.*

*Proof.* Let  $0 < x_0 < 1$ ; then

$$u(x, t) = \int_{x_0}^x u_x(y, t) dy + u(x_0, t) \quad (0 < x < 1),$$

and the integrand is bounded in  $0 < x \leq x_0$ . Hence the integral exists for  $x=0$  and defines  $u(0+, t)$ .

We might also note in passing that for such  $t$ , the  $x$  difference quotient at the boundary also tends to  $B'(t-0)$ ; for, by the mean value theorem,

$$\frac{u(h, t) - u(0+, t)}{h} = u_x(\bar{h}, t) \rightarrow B'(t-0)$$

as  $h \rightarrow 0$ .

From Theorem 1 we have:

COROLLARY 2. For  $u(0+, t)$  to exist it is sufficient that

$$\int_0^{t-0} (t-s)^{-1/2} |dB(s)|$$

converge.

*Proof.* Define

$$\begin{aligned} f(t) &= \int_0^1 F(0, t; y, 0) dA(y) + \int_0^t F(0, t, 1, s) dC(s) \\ &+ 2 \int_0^t \pi^{-1/2} (t-s)^{-1/2} \sum_{n=1}^{\infty} \exp\left[-\frac{n^2}{(t-s)}\right] dB(s) \\ &- \pi^{-1/2} \int_0^{t-0} (t-s)^{-1/2} dB(s), \end{aligned}$$

and consider

$$\begin{aligned} \limsup_{x \rightarrow 0+} |u(x, t) - f(t)| \\ \leq \pi^{-1/2} \limsup_{x \rightarrow 0+} \int_0^{t-0} (t-s)^{-1/2} \left\{ 1 - \exp\left[-\frac{x^2}{4(t-s)}\right] \right\} |dB(s)|. \end{aligned}$$

Now given  $\epsilon > 0$  there exists a  $\delta > 0$  such that

$$\begin{aligned} \pi^{-1/2} \int_{t-\delta}^{t-0} (t-s)^{-1/2} \exp\left[-\frac{x^2}{4(t-s)}\right] |dB(s)| \\ \leq \pi^{-1/2} \int_{t-\delta}^{t-0} (t-s)^{-1/2} |dB(s)| \leq \epsilon, \end{aligned}$$

so that

$$\begin{aligned} \limsup_{x \rightarrow 0+} |u(x, t) - f(t)| \\ \leq \pi^{-1/2} \limsup_{x \rightarrow 0+} \int_0^{t-\delta} (t-s)^{-1/2} \left\{ 1 - \exp\left[-\frac{x^2}{4(t-s)}\right] \right\} |dB(s)| + 2\epsilon = 2\epsilon. \end{aligned}$$

Let  $\epsilon \rightarrow 0$  to get

$$\lim_{x \rightarrow 0^+} u(x, t) = u(0^+, t) = f(t),$$

which completes the proof.

We may also note that if  $B(s)$  were monotone, then, since  $\exp[-x^2/4(t-s)]$  converges to unity in a monotone way, we could invoke, the monotone convergence theorem to obtain the convergence of the integral as a necessary and sufficient condition for the existence of  $u(0^+, t)$ .

Also with Theorem 2 at our disposal we can now prove:

**THEOREM 3.** *Let  $u(x, t)$  be representable by (2.1); then the functions  $A(x)$ ,  $B(t)$ ,  $C(t)$  are uniquely determined by  $u(x, t)$ , so that, at every point of continuity,*

$$A(x) = \lim_{a \rightarrow 0^+} \int_0^x u(y, a) dy$$

and

$$B(t) = \lim_{b \rightarrow 0^+} \int_0^t u_x(b, s) ds; \quad C(t) = \lim_{c \rightarrow 1^-} \int_0^t u_x(c, s) ds.$$

*Proof.* For suppose  $B_1(t)$  and  $B_2(t)$  arise from two distinct sequences. Then clearly if  $B_3(t) = B_1(t) - B_2(t)$ , we have

$$\int_0^t F(x, t; 0, s) dB_3(s) \equiv 0 \text{ in } R.$$

Hence, differentiating, we get

$$\int_0^t G_y(x, t; 0, s) dB_3(s) \equiv 0$$

We first show  $B_3(s)$  is continuous: suppose it has a jump  $\sigma$  at  $t_0$ ; then, for  $t > t_0$ ,

$$0 = \int_0^t G_y(x, t; 0, s) dB_4(s) + \sigma G_y(x, t; 0, t_0),$$

where  $B_4(s)$  is the boundary function remaining after the jump  $\sigma$  at  $t_0$  is removed. Then

$$0 = \int_0^t \frac{x}{2\pi^{1/2}(t-s)^{3/2}} \exp\left[-\frac{x^2}{4(t-s)}\right] dB_4(s) + \frac{x\sigma}{2\pi^{1/2}(t-t_0)^3} \exp\left[-\frac{x^2}{4(t-t_0)}\right] + o(1).$$

Choose  $\delta$  so small that

$$V_{B_4}(t_0 + \delta) - V_{B_4}(t_0 - \delta) < \frac{|\sigma|}{2},$$

set  $t - t_0 = x^2/4$ , and take  $x$  so small that  $x^2/4 < \delta$ . Then

$$0 = \int_{t_0 - \delta}^{t_0 + x^2/4} \frac{x}{2\pi^{1/2}(t_0 - s + x^2/4)^{3/2}} \exp\left[-\frac{x^2}{4(t_0 - s + x^2/4)}\right] dB_4(s) + \frac{4\sigma}{e\pi^{1/2}x^2} + o(1),$$

$$0 \geq \frac{4|\sigma|}{e\pi^{1/2}x^2} - \int_{t_0 - \delta}^{t_0 + x^2/4} \frac{x}{2\pi^{1/2}(t_0 - s + x^2/4)^{3/2}} \exp\left[-\frac{x^2}{4(t_0 - s + x^2/4)}\right] |dB_4(s)| + o(1).$$

The maximum of the integrand is at  $s = t_0$ , so that

$$0 \geq \frac{4|\sigma|}{e\pi^{1/2}x^2} - \frac{2|\sigma|}{e\pi^{1/2}x^2} + o(1) = \frac{2|\sigma|}{e\pi^{1/2}x^2} + o(1),$$

and we have a contradiction as  $x \rightarrow 0+$ .

Similarly the jumps of  $C(t)$  are determined.

Suppose  $A_1(x)$  and  $A_2(x)$  arise from two distinct sequences, and  $A_3(x) = A_1(x) - A_2(x)$ ; then, as before,

$$0 \equiv \int_0^1 F(x, t; y, 0) dA_3(y) \text{ in } R.$$

And suppose it has a jump of  $\sigma$  at  $x_0$ ; then, as before,

$$0 = \sigma(4\pi t)^{-1/2} + (4\pi t)^{-1/2} \int_0^1 \exp\left[-\frac{(y-x_0)^2}{4t}\right] dA_4(y) + o(1).$$

If  $\delta$  is so small that

$$V_{A_4}(x_0 + \delta) - V_{A_4}(x_0 - \delta) \leq |\sigma|/2,$$

then

$$0 = \sigma(4\pi t)^{-1/2} + (4\pi t)^{-1/2} \int_{x_0 - \delta}^{x_0 + \delta} \exp \left[ -\frac{(y - x_0)^2}{4t} \right] dA_4(y) + o(1),$$

$$0 \geq |\sigma| (4\pi t)^{-1/2} - |\sigma| (4\pi t)^{-1/2}/2 + o(1) = |\sigma| (4\pi t)^{-1/2}/2 + o(1),$$

and as  $t \rightarrow 0+$  we get a contradiction.

Then  $A_3, B_3, C_3$  are continuous functions of bounded variation, and by Theorem 2 their derivatives are zero almost everywhere. Each of them must then have an infinite derivative on a nondenumerable set. (See e.g. [8, p. 128].) This then implies that  $\lim u(x, t)$  and  $\lim u_x(x, t)$  must become infinite on a nondenumerable set, which is a contradiction, and the functions  $A_3, B_3, C_3$  are constants. Hence, since every sequence of  $a$ 's,  $b$ 's, or  $c$ 's contains a subsequence for which  $A_a(x)$ , etc., converges to a common limit, the limit must also be attained for continuous approach. Thus the last statement of the theorem is established.

**4. The Lebesgue integral representation.** We are now in a position to establish:

**THEOREM 4.** *For  $u(x, t)$  to be representable in  $R$  by*

$$(4.1) \quad u(x, t) = \int_0^1 F(x, t; y, 0) a(y) dy - \int_0^t F(x, t; 0, s) b(s) ds + \int_0^t F(x, t; 1, s) c(s) ds,$$

where  $a(y) \in L(0 \leq y \leq 1)$  and  $b(s), c(s) \in L(0 \leq s \leq s_0 < T \leq \infty)$  for every  $s_0, (0 < s_0 < T)$ , it is necessary and sufficient that

$$(1) \quad u(x, t) \in H \text{ in } R,$$

$$(2) \quad \lim_{y, y' \rightarrow 0+} \int_0^t |u_x(y, s) - u_x(y', s)| ds = 0$$

and

$$\lim_{y, y' \rightarrow 1-0} \int_0^t |u_x(y, s) - u_x(y', s)| ds = 0$$

for every  $t (0 < t < T)$ , and

$$(3) \quad \lim_{s, s' \rightarrow 0} \int_0^1 |u(y, s) - u(y, s')| dy = 0.$$

*Proof.* For the sufficiency, let the closed finite interval  $I \subset \{0 \leq s < T\}$  be prescribed, and let  $e$  be any measurable set in  $I$ . Given  $\epsilon > 0$ , there exists  $\delta = \delta(\epsilon, I)$  such that

$$\int_e |u_x(y, s) - u_x(y', s)| ds \leq \epsilon/2 \text{ for } y, y' < \delta.$$

Then

$$\begin{aligned} \int_e |u_x(y, s)| ds &\leq \int_e |u_x(y', s)| ds + \int_e |u_x(y, s) - u_x(y', s)| ds \\ &\leq \int_e |u_x(y', s)| ds + \epsilon/2. \end{aligned}$$

Now keep  $y'$  fixed and take  $m(e)$  so small that

$$\int_e |u_x(y', s)| ds \leq \epsilon/2,$$

so that, for  $0 < y < \delta$ ,

$$\int_e |u_x(y, s)| ds \leq \epsilon$$

if  $m(e)$  is sufficiently small. Hence  $B_b(s)$  are uniformly absolutely continuous; consequently, so is  $B(s)$ , and  $dB(s)$  can be replaced by  $b(s) ds$ , where  $B'(s) = b(s)$  almost everywhere. Similarly  $dC(s) = c(s) ds$  and  $dA(y) = a(y) dy$ .

The necessity of (1) follows by Theorem 1. To prove that of (2) we write

$$b(s) = b_1(s) - b_2(s),$$

where  $b_1(s)$  and  $b_2(s)$  are both nonnegative, say, for example,

$$b_1(s) = |b(s)| \text{ and } b_2(s) = |b(s)| - b(s).$$

Let

$$u^{(i)}(x, t) = - \int_0^t F(x, t; 0, s) b_i(s) ds \quad (i = 1, 2).$$

Then

$$u_x^{(i)}(x, t) = \int_0^t G_y(x, t; 0, s) b_i(s) ds \quad (i = 1, 2).$$

We know from Theorem 2 that

$$\lim_{x \rightarrow 0^+} u_x^{(i)}(x, t) = b_i(t) \quad (i = 1, 2),$$

almost everywhere, and, by Theorem 3,

$$\lim_{x \rightarrow 0^+} \int_0^t u_x^{(i)}(x, s) ds = \int_0^t b_i(s) ds \quad (i = 1, 2).$$

Since the  $u_x^{(i)}(x, t)$  are nonnegative (see [4, Remark 1, p.975]), we can say (see [4, p.977])

$$(4.2) \quad \lim_{x \rightarrow 0^+} \int_0^t |u^i(x, s) - b_i(s)| ds = 0 \quad (i = 1, 2).$$

Now consider

$$(4.3) \quad \int_0^t |u_x(x, s) - b(s)| ds \leq \int_0^t |u_x^{(1)}(x, s) - b_1(s)| ds \\ + \int_0^t |u_x^{(2)}(x, s) - b_2(s)| ds + \int_0^t \left| \int_0^1 F_x(x, s; y, 0) a(y) \right| dy ds \\ + \int_0^t \left| \int_0^s G_y(1-x, s; 0, \tau) c(\tau) d\tau \right| ds.$$

As  $x \rightarrow 0^+$ , the first and second integrals on the right vanish by (4.2), and the fourth since  $G_y(1-x, s; 0, \tau)$  tends to zero uniformly in  $s$  and  $\tau$  as  $x \rightarrow 0^+$ . To estimate the third we note

$$F_x(x, s; y, 0) = -\frac{1}{4\pi^{1/2}s^{3/2}} \left\{ (x-y) \exp \left[ -\frac{(x-y)^2}{4s} \right] \right. \\ \left. + (x+y) \exp \left[ -\frac{(x+y)^2}{4s} \right] \right\} + \bar{u}(x, y, s),$$

where  $\bar{u} = o(1)$  uniformly in  $y$  and  $s$  as  $x \rightarrow 0^+$ . Then

$$|F_x(x, s; y, 0)| \leq \frac{1}{4\pi^{1/2}s^{3/2}} \left\{ |x-y| \exp \left[ -\frac{(x-y)^2}{4t} \right] \right. \\ \left. + (x+y) \exp \left[ -\frac{(x+y)^2}{4t} \right] \right\} + |\bar{u}|.$$

But

$$\int_0^t s^{-3/2} \exp \left[ -\frac{a^2}{4s} \right] ds = \frac{2}{|a|} \int_{a^2/4t}^\infty e^{-v} v^{-1/2} dv \leq \frac{2\pi^{1/2}}{|a|}.$$

Hence

$$\int_0^t |F(x, s; y, 0)| ds \leq 1 + o(1) \leq 2$$

for  $x$  sufficiently small. Thus the third integral on the right side of (4.3) is dominated by

$$\int_0^1 |a(y)| \int_0^t |F_x(x, s; y, 0)| ds dy \leq 2 \int_0^1 |a(y)| dy.$$

Then by the dominated convergence theorem we can pass to the limit under the integral sign, by which we get zero as a limit, since  $F_x(x, s; y, 0)$  tends to zero. This proves

$$\lim_{x \rightarrow 0^+} \int_0^t |u_x(x, s) - b(s)| ds,$$

from which condition (2) follows immediately.

Condition (3) follows similarly, but more easily.

**5. Uniqueness.** We now turn to the question of the extent to which the boundary data uniquely determine the solution of the boundary-value problem. We get one result as an immediately corollary of our Theorem 4.

**COROLLARY 3.** *If  $u(x, t)$  is representable by (4.1) in  $R$ , and has zero boundary values almost everywhere for approach along the normal, then  $u(x, t) \equiv 0$  in  $R$ .*

*Proof.* By Theorem 2,  $a(y)$ ,  $b(s)$ , and  $c(s)$  vanish almost everywhere.

The situation in the case of the Stieltjes representation is not so simple (see [6]): We can have a function representable in  $R$  by (2.1) which has boundary values identically zero for approach along the normal, yet which is itself not identically zero; for example, for  $0 \leq t_0$ , let

$$u(x, t) = \begin{cases} 0 & (0 \leq t \leq t_0), \\ -F(x, t; 0, t_0) & (t_0 < t). \end{cases}$$

This is a nontrivial solution of the heat equation, representable by (2.1), for which



$$u(x, 0+) \equiv 0, \quad u_x(0+, t) \equiv 0, \quad u_x(1-0, t) \equiv 0.$$

However we can assert:

**THEOREM 5.** *Suppose  $u(x, t)$  is representable in  $R$  by (2.1), that*

$$u(x, 0+) \equiv 0 \quad (0 < x < 1),$$

and that  $B(s)$  and  $C(s)$  are monotone for  $0 \leq s < T \leq \infty$ . Let

$$\lim u_x(x, t) = 0 \quad \text{as } (x, t) \rightarrow (0, s)$$

along a parabolic arc of the form  $t - s = ax^2$ , ( $a > 0$ ) and

$$\lim u_x(x, t) = 0 \quad \text{as } (x, t) \rightarrow (1, s)$$

along a parabolic arc of the form  $t - s = b(x - 1)^2$  ( $b > 0$ ) for every  $s$  ( $0 \leq s < T$ ).

Then  $u(x, t) \equiv 0$  in  $R$

*Proof.* Let  $x$  be fixed,  $0 < x < 1$ . Then, by (2.1) and (1.3),

$$u(x, t) = \int_0^1 F(x, t; y, 0) dA(y) + o(1) \quad \text{as } t \rightarrow 0+.$$

Choose  $0 < \delta < (1/2) \min(x, 1 - x)$ , so that

$$\begin{aligned} u(x, t) &= \int_{x-\delta}^{x+\delta} (4\pi t)^{-1/2} \exp\left[-\frac{(y-x)^2}{4t}\right] dA(y) + o(1), \\ &= \int_{x-\delta}^{x+\delta} (4\pi t)^{-1/2} \exp\left[-\frac{(y-x)^2}{4t}\right] d[A(y) - A(x)] + o(1), \\ &= \int_{x-\delta}^{x+\delta} \frac{y-x}{4\pi^{1/2} t^{3/2}} \exp\left[-\frac{(y-x)^2}{4t}\right] [A(y) - A(x)] dy + o(1), \\ &= \int_{-\delta}^{\delta} \frac{z^2}{4\pi^{1/2} t^{3/2}} \exp\left[-\frac{z^2}{4t}\right] \frac{A(x+z) - A(x)}{z} dz + o(1). \end{aligned}$$

Then

$$\begin{aligned} u(x, t) &\geq \inf_{-\delta \leq z \leq \delta} \frac{A(x+z) - A(x)}{z} \int_{-\delta}^{\delta} \frac{z^2}{4\pi^{1/2} t^{3/2}} \exp\left[-\frac{z^2}{4t}\right] dz + o(1), \\ &= \inf_{-\delta \leq z \leq \delta} \frac{A(x+z) - A(x)}{z} \int_{-\delta/2}^{\delta/2} \frac{2}{\pi^{1/2}} \zeta^2 e^{-\zeta^2} d\zeta + o(1). \end{aligned}$$

Let  $t \rightarrow 0+$ :

$$u(x, 0+) = 0 \geq \inf_{-\delta \leq z \leq \delta} \frac{A(x+z) - A(x)}{z}$$

Let  $\delta \rightarrow 0$ :

$$0 \geq \underline{D}A(x).$$

Similarly,

$$0 \leq \bar{D}A(x)$$

for every  $x$  ( $0 < x < 1$ ). Now  $A(x)$  is continuous, for if it had a jump it would violate one or the other of these conditions. Then by [1, p. 580], it must be both nonincreasing and nondecreasing, and hence constant.

Furthermore,

$$u_x(x, t) = \int_0^t G_y(x, t; 0, s) dB(s) + o(1) \text{ as } (x, t) \rightarrow (0, s).$$

Then, by [6],  $B(s)$  is constant. Similarly one sees  $C(s)$  is constant. This completes the proof.

#### REFERENCES

1. C. Caratheodory, *Vorlesungen über reelle Funktionen*, Berlin, 1920.
2. G. Doetsch, *Theorie und Anwendung der Laplace-Transformation*, New York, 1943.
3. W. Fulks, *On integral representations and uniqueness of solutions of the heat equation*, University of Minnesota Thesis, June, 1949.
4. ———, *On the solutions of the heat equation*. Proc. Amer. Math. Soc. 2 (1951), 973-979.
5. ———, *On the boundary values of solutions of the heat equation*. Pacific J. Math. 2 (1952), 141-145.
6. ———, *On the unique determination of solutions of the heat equation*. Pacific J. Math. 3 (1953), 387-391.
7. P. Hartman and A. Wintner, *On the solutions of the equation of heat conduction*. Amer. J. Math. 72 (1950), 367-395.
8. S. Saks, *Theory of the Integral*, Warsaw-Lwow, 1937.
9. D. V. Widder, *The Laplace Transform*, Princeton, 1941.

THE UNIVERSITY OF MINNESOTA



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

R. M. ROBINSON

University of California  
Berkeley 4, California

R. P. DILWORTH

California Institute of Technology  
Pasadena 4, California

E. HEWITT

University of Washington  
Seattle 5, Washington

E. F. BECKENBACH

University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

H. BUSEMANN

HERBERT FEDERER

MARSHALL HALL

P. R. HALMOS

HEINZ HOPF

R. D. JAMES

BØRGE JESSEN

PAUL LÉVY

GEORGE PÓLYA

J. J. STOKER

E. G. STRAUS

KÔSAKU YOSIDA

## SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

\* \* \*

AMERICAN MATHEMATICAL SOCIETY

NATIONAL BUREAU OF STANDARDS,

INSTITUTE FOR NUMERICAL ANALYSIS

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors except Robinson, whose term expires with the completion of the present volume; they might also be sent to M. M. Schiffer, Stanford University, Stanford, California, who is succeeding Robinson. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1953 BY PACIFIC JOURNAL OF MATHEMATICS

# Pacific Journal of Mathematics

Vol. 3, No. 3

May, 1953

L. Carlitz, <i>Some theorems on generalized Dedekind sums</i> . . . . .	513
L. Carlitz, <i>The reciprocity theorem for Dedekind sums</i> . . . . .	523
Edward Richard Fadell, <i>Identifications in singular homology theory</i> . . . . .	529
Harley M. Flanders, <i>A method of general linear frames in Riemannian geometry. I</i> . . . . .	551
Watson Bryan Fulks, <i>The Neumann problem for the heat equation</i> . . . . .	567
Paul R. Garabedian, <i>Orthogonal harmonic polynomials</i> . . . . .	585
R. E. Greenwood and Andrew Mattei Gleason, <i>Distribution of round-off errors for running averages</i> . . . . .	605
Arthur Eugene Livingston, <i>The space <math>H^p</math>, <math>0 &lt; p &lt; 1</math>, is not normable</i> . . . . .	613
M. N. Mikhail, <i>On the order of the reciprocal set of a basic set of polynomials</i> . . . . .	617
Louis Joel Mordell, <i>On the linear independence of algebraic numbers</i> . . . . .	625
Leo Sario, <i>Alternating method on arbitrary Riemann surfaces</i> . . . . .	631
Harold Nathaniel Shapiro, <i>Iterates of arithmetic functions and a property of the sequence of primes</i> . . . . .	647
H. Shniad, <i>Convexity properties of integral means of analytic functions</i> . . . . .	657
Marlow C. Sholander, <i>Plane geometries from convex plates</i> . . . . .	667