

Pacific Journal of Mathematics

THE SPACE H^p , $0 < p < 1$, IS NOT NORMABLE

ARTHUR EUGENE LIVINGSTON

THE SPACE H^p , $0 < p < 1$, IS NOT NORMABLE

ARTHUR E. LIVINGSTON

1. **Introduction.** For $p > 0$, the space H^p is defined to be the class of functions $x(z)$ of the complex variable z , which are analytic in the interior of the unit circle, and satisfy

$$\sup_{0 \leq r < 1} \int_0^{2\pi} |x(re^{i\theta})|^p d\theta < \infty.$$

Set

$$A_p(r; x) = \left(\frac{1}{2\pi} \int_0^{2\pi} |x(re^{i\theta})|^p d\theta \right)^{1/p}$$

and

$$\|x\| = \sup_{0 \leq r < 1} A_p(r; x).$$

S. S. Walters has shown [2] that H^p , $0 < p < 1$, is a linear topological space under the topology: $U \subset H^p$ is open if $x_0 \in U$ implies the existence of a "sphere" $S: \|x - x_0\| < r$ such that $S \subset U$. He conjectured in [3] that H^p , $0 < p < 1$, does not have an equivalent normed topology, and it is shown here that this conjecture is correct. Since the conjugate space $(H^p)^*$ has sufficiently many members to distinguish elements of H^p , the space H^p , $0 < p < 1$, affords an interesting nontrivial example of a locally bounded linear topological space which is not locally convex.

2. *Proof.* For $x \in H^p$, $p > 0$, it is known [4, 160] that $A_p(r; x)$ is a non-decreasing function of r . Consequently, if $P(z)$ is a polynomial, then $P \in H^p$ and $\|P\| = A_p(1; P)$. This observation will be used below.

According to a theorem of Kolmogoroff [1], a linear topological space has an equivalent normed topology if and only if the space contains a bounded open convex set. It will be shown here that the "sphere" $K_1: \|x\| < 1$ of H^p ,

Received July 2, 1952. The work in this paper was supported by a grant from the Graduate School of the University of Oregon.

$0 < p < 1$, contains no convex neighborhood of the origin; this is clearly sufficient to show that H^p , $0 < p < 1$, contains no bounded open convex set, and hence is not normable.

To accomplish this, the contrary is assumed. Thus, it is assumed that K_1 contains a convex neighborhood V of the origin. Since V is open, V contains a "sphere" $K_\epsilon: \|x\| < \epsilon$. There will be exhibited $x_1, \dots, x_N \in K_\epsilon$, and $a_1 > 0, \dots, a_N > 0$, with $\sum a_k = 1$, such that $\sum a_k x_k \notin K_1$ and, *a fortiori*, $\sum a_k x_k \notin V$, in contradiction to the assumed convexity of V .

If $x(\theta)$ is a complex function of the real variable $\theta \in I: 0 \leq \theta \leq 2\pi$, define

$$A(x) = \left(\frac{1}{2\pi} \int_0^{2\pi} |x(\theta)|^p d\theta \right)^{1/p}.$$

Once and for all, k is any integer in the range $1, \dots, N$. Let I_k denote the interval

$$\frac{2\pi(k-1)}{N} < \theta < \frac{2k\pi}{N},$$

and let i_k denote the degenerate interval consisting of the point $(2\pi/N)(k-1/2)$. Define the continuous function $c_k(\theta)$ to be zero on $I - I_k$, to be equal to $\epsilon N^{1/p}$ on i_k , and to be linear on each of the two intervals in $I_k - i_k$. Let

$$a_k = B_N k^{-1/p}, \quad B_N = \left(\sum_1^N k^{-1/p} \right)^{-1},$$

so that $a_k > 0$ and $\sum a_k = 1$. It is easily verified that

$$A(c_k) = \epsilon(p+1)^{-1/p} < \epsilon$$

and

$$A(\sum a_k c_k) = \epsilon B_N (p+1)^{-1/p} \left(\sum_1^N k^{-1} \right)^{1/p}.$$

Since B_N is bounded away from zero below, N can be chosen such that

$$A(\sum a_k c_k) > 1.$$

Each $c_k(\theta)$ is absolutely continuous on I . Given $\alpha > 0$, it follows that

there is a trigonometrical polynomial

$$T_k(\theta) = \sum_{n=-m_k}^{m_k} a_{nk} e^{in\theta}$$

such that

$$|T_k(\theta) - c_k(\theta)| < \alpha$$

uniformly in θ . Setting

$$p_k(\theta) = e^{im_k\theta} T_k(\theta)$$

gives

$$|p_k(\theta) - e^{im_k\theta} c_k(\theta)| < \alpha$$

uniformly in θ . Set

$$C_k(\theta) = e^{im_k\theta} c_k(\theta).$$

It is clear that $A(C_k) = A(c_k)$ and $A(\sum a_k C_k) = A(\sum a_k c_k)$. Since $A(x)$ is a continuous function of x , it follows, if α is small enough, that $A(p_k) < \epsilon$ and $A(\sum a_k p_k) > 1$.

Let

$$P_k(z) = \sum_{n=-m_k}^{m_k} a_{nk} z^{n+m_k},$$

so that $P_k(e^{i\theta}) = p_k(\theta)$. As previously remarked,

$$\|P_k\| = A_p(1; P_k) = A(p_k)$$

and

$$\|\sum a_k P_k\| = A_p(1; \sum a_k P_k) = A(\sum a_k p_k).$$

Since $P_1, \dots, P_N \in K_\epsilon \subset V \subset K_1 \subset H^p$, $a_1 > 0, \dots, a_N > 0$, $\sum a_k = 1$, and $\sum a_k P_k \notin K_1$, we have obtained the required contradiction.

REFERENCES

1. A. Kolmogoroff, *Zur Normierbarkeit eines allgemeinen topologischen linearen Raumes*, *Studia Math.*, 5 (1934), 29-33.
2. S. S. Walters, *The space H^p with $0 < p < 1$* , *Proc. Amer. Math. Soc.* 1 (1950), 800-805.
3. _____, *Remarks on the space H^p* , *Pac. J. Math.*, 1 (1951), 455-471.
4. A. Zygmund, *Trigonometrical series*, Warsaw (1935).

UNIVERSITY OF OREGON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

R. M. ROBINSON

University of California
Berkeley 4, California

R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

E. HEWITT

University of Washington
Seattle 5, Washington

E. F. BECKENBACH

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN

HERBERT FEDERER

MARSHALL HALL

P. R. HALMOS

HEINZ HOPF

R. D. JAMES

BØRGE JESSEN

PAUL LÉVY

GEORGE PÓLYA

J. J. STOKER

E. G. STRAUS

KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA, BERKELEY

UNIVERSITY OF CALIFORNIA, DAVIS

UNIVERSITY OF CALIFORNIA, LOS ANGELES

UNIVERSITY OF CALIFORNIA, SANTA BARBARA

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD RESEARCH INSTITUTE

STANFORD UNIVERSITY

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

NATIONAL BUREAU OF STANDARDS,

INSTITUTE FOR NUMERICAL ANALYSIS

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors except Robinson, whose term expires with the completion of the present volume; they might also be sent to M. M. Schiffer, Stanford University, Stanford, California, who is succeeding Robinson. All other communications to the editors should be addressed to the managing editor, E. F. Beckenbach, at the address given above.

Authors are entitled to receive 100 free reprints of their published papers and may obtain additional copies at cost.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$8.00; single issues, \$2.50. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1953 BY PACIFIC JOURNAL OF MATHEMATICS

Pacific Journal of Mathematics

Vol. 3, No. 3

May, 1953

L. Carlitz, <i>Some theorems on generalized Dedekind sums</i>	513
L. Carlitz, <i>The reciprocity theorem for Dedekind sums</i>	523
Edward Richard Fadell, <i>Identifications in singular homology theory</i>	529
Harley M. Flanders, <i>A method of general linear frames in Riemannian geometry. I</i>	551
Watson Bryan Fulks, <i>The Neumann problem for the heat equation</i>	567
Paul R. Garabedian, <i>Orthogonal harmonic polynomials</i>	585
R. E. Greenwood and Andrew Mattei Gleason, <i>Distribution of round-off errors for running averages</i>	605
Arthur Eugene Livingston, <i>The space H^p, $0 < p < 1$, is not normable</i>	613
M. N. Mikhail, <i>On the order of the reciprocal set of a basic set of polynomials</i>	617
Louis Joel Mordell, <i>On the linear independence of algebraic numbers</i>	625
Leo Sario, <i>Alternating method on arbitrary Riemann surfaces</i>	631
Harold Nathaniel Shapiro, <i>Iterates of arithmetic functions and a property of the sequence of primes</i>	647
H. Shniad, <i>Convexity properties of integral means of analytic functions</i>	657
Marlow C. Sholander, <i>Plane geometries from convex plates</i>	667