

Pacific Journal of Mathematics

**MAPPING PROPERTIES OF CESÀRO SUMS OF ORDER TWO
OF THE GEOMETRIC SERIES**

S. E. RAUCH

MAPPING PROPERTIES OF CESÀRO SUMS OF ORDER TWO OF THE GEOMETRIC SERIES

S. E. RAUCH

1. Introduction. Previous investigations of the mappings

$$w = S_n^{(k)}(z)$$

of the unit circle $|z| \leq 1$, where

$$S_n^{(k)}(z) = \binom{n+k}{k} + \binom{n+k-1}{k}z + \cdots + \binom{k}{k}z^n$$

denotes the n th Cesàro sum of order k of the geometric series, have been made by Fejér, Schweitzer, Sidon, and Szegő. Knowledge of the properties of the sums $S_n^{(k)}(z)$ is valuable in the study of power series having coefficients monotonic of order $k+1$.

The present article provides additional asymptotic properties for

$$S_n^{(2)}(e^{i\phi}) = x_n(\phi) + iy_n(\phi).$$

The following results are established:

THEOREM 1. *For n sufficiently large, an α_n exists such that $y_n(\phi)$ is increasing for $0 < \phi < \alpha_n$ and decreasing for $\alpha_n < \phi < \pi$. Furthermore,*

$$\alpha_n = \alpha/n + O(n^{-2}), \text{ where } \pi < \alpha < 3\pi/2.$$

THEOREM 2. *For n sufficiently large, a β_n exists such that*

$$x_n'(\phi) \begin{cases} \leq 0, & 0 < \phi < \beta_n, & n \equiv 0 \pmod{3} \\ < 0, & 0 < \phi < \beta_n, & n \equiv 1 \pmod{3} \\ < 0, & 0 < \phi < \beta_n, & n \equiv 2 \pmod{3} \end{cases}$$

Received January 28, 1953.

Pacific J. Math. 4 (1954), 109-121

where

$$\beta_n = \frac{2\pi}{3} + \frac{\beta}{n} - O(n^{-3/2}),$$

and $\beta = 2\pi, 4\pi/3, 2\pi/3$ for $n \equiv 0, 1, 2 \pmod{3}$, respectively.

THEOREM 3. For n sufficiently large, the mapping of $|z| = 1$ by

$$w = S_n^{(2)}(e^{i\phi})$$

is convex for $0 < \phi < \gamma_n$, where γ_n is the maximum angle for which convexity holds, and $\gamma_n = \gamma/n + O(n^{-2})$ where $2\pi < \gamma < 3\pi$.

2. Proof of Theorem 1.

2.1. A closed expression for $y_n'(\phi)$ has been presented by Szegő [10]:

$$(2.1) \quad y_n'(\phi) =$$

$$\frac{1}{8 \sin^2 \phi/2} \left\{ -(n^2 + 3n + 3) - n \cdot \frac{\sin(n + 3/2)\phi}{\sin \phi/2} + 3 \frac{\sin^2(n + 1)\phi/2}{\sin^2 \phi/2} \right\}.$$

The inequality $y_n'(\phi) < 0$ is satisfied if

$$(2.2) \quad n^2 + 3n + 3 > -n \cdot \frac{\sin(n + 3/2)\phi}{\sin \phi/2} + 3 \frac{\sin^2(n + 1)\phi/2}{\sin^2 \phi/2},$$

or

$$(2.3) \quad n^2 + 3n + 3 > n \cdot \csc \phi/2 + 3 \cdot \csc^2 \phi/2.$$

Let δ be fixed, $\delta > 0$, and consider the restriction $\phi > \delta/n$. For n sufficiently large, $\sin(\delta/2n) > \delta/\pi n$, and the previous inequality is maintained if δ is chosen so that

$$n^2 + 3n + 3 > n^2\pi/\delta + 3\pi^2 n^2/\delta^2, \text{ or } \pi/\delta + 3\pi^2/\delta^2 < 1.$$

It is sufficient for the present problem to define $\delta = 3\pi$. Hence, if $\phi \geq 3\pi/n$, then $y_n'(\phi) < 0$ and $3\pi/n > \alpha_n$. Since

$$y'_n(\phi) = \sum_{m=1}^n a_m \cos m\phi,$$

where

$$a_m = m \binom{n+2-m}{2},$$

it at once follows that $y'_n(\phi) > 0$ for $0 \leq \phi \leq \pi/2n$.

2.2. In the next section it is shown that in the interval $\pi/2n < \phi < 3\pi/n$ there is exactly one $\phi = \alpha_n$ such that $y'_n(\phi) = 0$ if n is sufficiently large. More precisely, for the $\phi = \alpha_n$ the second derivative does not vanish and $\alpha_n \sim \alpha/n$, $\alpha > 0$, where $\pi < \alpha < 3\pi/2$. The magnitude of α is defined as the root of a transcendental equation.

It is possible to express (2.1) in the following form:

$$(2.4) \quad y'_n(\phi) = n^2/8 \cdot g_n(\phi) \cdot \csc^2 \phi/2,$$

where the function $g_n(\phi)$ is defined as

$$g_n(\phi) = -1 - \frac{1}{n} \cdot \frac{\sin(n+3/2)\phi}{\sin \phi/2} + \frac{3}{n^2} \cdot \frac{\sin^2(n+1)\phi/2}{\sin^2 \phi/2} - \left(\frac{3}{n} + \frac{3}{n^2} \right).$$

Let $\phi = c/n$, $\pi/2 < c < 3\pi$, and then $g_n(\phi)$ becomes a function of c , denoted by $G_n(c)$. In addition,

$$\lim_{n \rightarrow \infty} G_n(c) = -1/c^2 \cdot f(c),$$

where

$$f(c) = 2c \cdot \sin c + 6 \cdot \cos c + (c^2 - 6).$$

Furthermore, $G_n(c)$ converges uniformly to this limit for arbitrary values of c in the interval. It is sufficient to show that the function $f(c)$ has a unique simple zero in the interval $\pi/2 < c < 3\pi/2$ to assure that $g_n(c)$ has a simple zero in the same interval if n is sufficiently large.

An easy calculation yields

$$f'(c) = 8 \cdot \cos^2(c/2) \cdot (c/2 - \tan c/2).$$

Thus it is seen that $f(0) = 0$, $f'(c) < 0$ for $0 < c < \pi$; $f(\pi) = -(12 - \pi^2) < 0$, $f'(c) > 0$ for $\pi < c < 2\pi$; $f(2\pi) > 0$ and if $c > 2\pi$ then $f(c) > (c - 1)^2 - 13 > 0$. Since $f(3\pi/2) > 0$, there is a simple positive zero, $c = \alpha$, of the function $f(c)$, $\pi < \alpha < 3\pi/2$. In conclusion, $\alpha_n \sim \alpha/n$, $\pi < \alpha < 3\pi/2$, for n sufficiently large.

2.3. It is not difficult to find a more precise asymptotic expression for α_n . For this purpose let $\alpha_n = c/n$, where $c = \alpha + a/n$ and a is a bounded, real constant. Let $h_n(a)$ denote $g_n(c)$ when the latter is regarded as a function of a . Let $\phi = c/n$; a simplification yields

$$-n^2 h_n(a) = n^2(2c \cdot \sin c + 6 \cos c + c^2 - 6)/c^2 + n(3 + 3 \cos c - 6/c \cdot \sin c) \\ + (5/2 - 13c/6 \cdot \sin c - 5/2 \cdot \cos c) + O(1/n).$$

If

$$h(c) = 3c^2[1 + \cos c - 2/c \cdot \sin c]$$

and

$$k(c) = c^2[5/2 - 13c/6 \cdot \sin c - 5/2 \cdot \cos c],$$

then it is possible to rewrite the previous expression in the form

$$-n^2 c^2 h_n(a) = n^2 \cdot f(c) + n \cdot h(c) + k(c) + O(1/n).$$

Let the functions $f(c)$, $h(c)$, $k(c)$ be expanded by Taylor's formula for values of c near α . Then the previous equality becomes

$$-n^2 c^2 h_n(a) = n[a \cdot f'(\alpha) + h(\alpha)] + a^2/2 \cdot f''(\alpha) + a \cdot h'(\alpha) + k(\alpha) + O(1/n).$$

Thus one obtains

$$\lim_{n \rightarrow \infty} [-n \cdot c^2 h_n(a)] = a \cdot f'(\alpha) + h(\alpha) \text{ and } f'(\alpha) \neq 0.$$

Obviously the limit has a zero for the value $a = -h(\alpha)/f'(\alpha)$, or

$$a = -3\alpha^2/8 \cdot (1 + \cos \alpha - 2/\alpha \cdot \sin \alpha) \cdot \sec^2 \alpha/2 \cdot (\alpha/2 - \tan \alpha/2)^{-1},$$

and α is the simple zero of the function

$$f(c) = 2c \cdot \sin c + 6 \cdot \cos c + c^2 - 6$$

in the interval $\pi < \alpha < 3\pi/2$.

This shows that for n sufficiently large, $\gamma_n'(\phi) = 0$ for

$$\phi = \alpha_n = \alpha/n + (a + \epsilon_n)/n^2,$$

where $\epsilon_n \rightarrow 0$. Thus the assertion of Theorem 1 has been verified.

3. Proof of Theorem 2.

3.1. In the article by Szegő [10], a closed expression for $x_n'(\phi)$ is presented:

$$(3.1) \quad x_n'(\phi) = \frac{\csc \phi/2}{8 \sin^3 \phi/2} \left[-(2n + 3) - (n + 3/2) \frac{\cos (n + 3/2)\phi}{\cos \phi/2} + \frac{3}{2} \cdot \frac{\sin (n + 3/2)\phi}{\sin \phi/2} \right].$$

It immediately follows that $x_n'(\phi)$ is negative if

$$(3.2) \quad [3/(2n + 3)]^2 \cdot \csc^2 \phi/2 + \sec^2 \phi/2 < 4; \quad \cot \phi/2 > 0.$$

Let $0 < \phi \leq \pi/n$. As

$$x_n(\phi) = \sum_{m=1}^n b_m \cos m\phi,$$

where

$$b_m = \binom{n + 2 - m}{2},$$

then $x_n'(\phi) < 0$. Next consider the interval $\pi/n \leq \phi \leq 2\pi/3 - c/n$, where c is fixed, $c > 0$. Since

$$[3/(2n + 3)]^2 \cdot \csc^2 \phi/2 + [1 - \sin^2 \phi/2]^{-1},$$

as a function of $\sin^2 \phi/2$, is convex from below, it obtains its maximum at one or both end-points of the interval. Thus in order to prove the inequality (3.2) it is sufficient to consider only the end-point values of $\pi/n \leq \phi \leq 2\pi/3 - c/n$. It easily follows that (3.2) is satisfied by $\phi = \pi/n$. Now study $\phi = 2\pi/3 - c/n$. Since

$$\sin^{-2} \phi/2 = O(1), \quad \cos^2 \phi/2 = 1/4 \cdot (1 + \sqrt{3} \cdot c/n) + O(1/n^2),$$

the left side of (3.2) then can be written as

$$\begin{aligned} [3/(2n+3)]^2 \cdot O(1) + 4[1 + \sqrt{3} \cdot c/n + O(1/n^2)]^{-1} \\ = 4(1 - \sqrt{3} \cdot c/n) + O(1/n^2), \end{aligned}$$

which indeed is less than 4 provided n is sufficiently large. The minimum value of n is a function of c . Thus it now is established that $x_n''(\phi) < 0$ for $0 < \phi \leq 2\pi/3 - c/n$, if n is sufficiently large, $n \geq n_1(c)$, where c is an arbitrary positive fixed magnitude.

3.2. Next let $\phi = 2\pi/3$. By (3.1) it follows that

$$x_n'(2\pi/3) = -(2n+3)/6\sqrt{3} \cdot (1 - \cos 2\pi n/3) - 1/6 \cdot \sin 2\pi n/3.$$

Three possible cases for the n arise. For $n \equiv 0 \pmod{3}$, $x_n'(2\pi/3) = 0$; whereas for $n \equiv 1, 2 \pmod{3}$, $x_n'(2\pi/3) < 0$. Thus the behavior of $x_n'(\phi)$ in the neighborhood of $\phi = 2\pi/3$ must be examined more fully, $n \equiv 0 \pmod{3}$. Let

$$x_n'(\phi) = r(\phi) \cdot s(\phi),$$

where

$$r(\phi) = 1/8 \cdot \cos \phi/2 \cdot \csc^3 \phi/2$$

and

$$\begin{aligned} s(\phi) = -(2n+3) - (n+3/2) \cdot \cos(n+3/2)\phi \cdot \sec \phi/2 \\ + 3/2 \cdot \sin(n+3/2)\phi \cdot \csc \phi/2. \end{aligned}$$

As $s = 0$ for $\phi = 2\pi/3$, then

$$x_n''(2\pi/3) = r(2\pi/3) \cdot s'(2\pi/3).$$

Upon letting $N = n + 3/2$, we see that

$$s'(2\pi/3) = 0, \quad x_n''(2\pi/3) = 0.$$

An examination of the third derivative shows that

$$x_n'''(2\pi/3) = r(2\pi/3) \cdot s'''(2\pi/3).$$

As $r(2\pi/3) > 0$, $\text{sgn } x_n''(\phi) = \text{sgn } s''(\phi)$. Since

$$s''(\phi) = N^3 \sec \phi/2 \cdot \cos N\phi + O(N^2),$$

then $s''(2\pi/3) = -2N^3 + O(N^2)$, and for n sufficiently large $s''(2\pi/3) < 0$. It is now known that $x_n'(\phi) < 0$ for $0 < \phi < 2\pi/3$ if n is sufficiently large.

3.3. This section extends the investigation beyond $\phi = 2\pi/3$. For this purpose let $\phi = 2\pi/3 + c/N$, where again $N = n + 3/2$. The substitution of this value of ϕ into (3.1) yields

$$(3.3) \quad \frac{8 \sin^3 \phi/2}{\cos \phi/2} \cdot x_n'(\phi) = -2N \left[1 - \frac{1}{2} \frac{\cos(2\pi n/3 + c)}{\cos \phi/2} \right] - \frac{3}{2} \cdot \frac{(2\pi n/3 + c)}{\sin \phi/2}.$$

Any easy calculation shows that

$$\sin \phi/2 = \sqrt{3}/2 + c/4N + c^2 \cdot O(1/N^2),$$

$$\cos \phi/2 = 1/2 - \sqrt{3} \cdot c/4N + c^2 \cdot O(1/N^2).$$

The remainder of the section will study the separate cases of $n \pmod{3}$.

$n \equiv 0 \pmod{3}$. Let us rewrite (3.3) as follows:

$$(3.4) \quad \frac{8 \sin^3 \phi/2 \cdot x_n'(\phi)}{\cos \phi/2 \cdot 2(1 - \cos c)} \\ = -N + \frac{\sqrt{3}}{2} \left[\frac{c \cdot \cos c - \sin c}{1 - \cos c} \right] + \frac{c^2}{1 - \cos c} \cdot O\left(\frac{1}{N}\right).$$

Let

$$F(c) = [c \cdot \cos c - \sin c] \cdot [1 - \cos c]^{-1}.$$

Since

$$F'(c) = \sin c \cdot [\sin c - c] [1 - \cos c]^{-2},$$

it is easily seen that $F(c)$ is decreasing for $0 < c < \pi$ and increasing for $\pi < c < 2\pi$. It follows that $x_n'(\phi) \leq 0$ for $0 < \phi \leq 2\pi/3 + c/N$, where $\pi < c < 2\pi - \epsilon$, ϵ a fixed positive number. Now

$$c = 2\pi - \delta/\sqrt{N},$$

δ a fixed positive number for n sufficiently large. Then

$$F(c) = 4\pi/\delta^2 \cdot N + O(1/N),$$

so that, for the above value of ϕ , (3.4) becomes

$$\frac{4 \sin^3 \phi/2 \cdot x_n'(\phi)}{\cos \phi/2 \cdot (1 - \cos c)} = -N + 2\pi\sqrt{3} \cdot N/\delta^2 + O(1).$$

In addition,

$$(1 - \cos c)^{-1} = (1 - \cos \delta/\sqrt{N})^{-1} = O(N).$$

Thus

$$x_n'(\phi) < 0 \text{ if } 2\pi\sqrt{3}/\delta^2 < 1,$$

and

$$x_n'(\phi) > 0 \text{ if } 2\pi\sqrt{3}/\delta^2 > 1.$$

Thus

$$\delta = (2\pi)^{1/2} \cdot (3)^{1/4}$$

furnishes the critical value of ϕ . It has been shown that, for $n \equiv 0 \pmod{3}$,

$$x_n'(\phi) < 0 \text{ for } 0 < \phi < 2\pi/3$$

and

$$x_n'(\phi) \leq 0 \text{ for } 0 < \phi < 2\pi/3 + 2\pi/N - O(N^{-3/2}),$$

for n sufficiently large.

$n \equiv 1 \pmod{3}$. It is possible to rewrite (3.3) so that the right side becomes

$$-2N[1 - \cos(c + 2\pi/3)]$$

$$+ \sqrt{3}[c \cdot \cos(c + 2\pi/3) - \sin(c + 2\pi/3)] + c^2 \cdot O(1/N).$$

By reasoning as in the previous case, one finds $x_n'(\phi) < 0$ for $0 \leq c \leq 4\pi/3 - \epsilon$, $\epsilon > 0$, for n sufficiently large. Let

$$c = 4\pi/3 - \delta/\sqrt{N}.$$

Then the right side of (3.3) reduces to

$$-\delta^2 + 4\pi/\sqrt{3} + O(1/\sqrt{N}).$$

Therefore

$$x'_n(\phi) < 0 \text{ if } \delta > 2 \cdot \pi^{1/2} \cdot 3^{-1/4},$$

and

$$x'_n(\phi) > 0 \text{ if } \delta < 2 \cdot \pi^{1/2} \cdot 3^{-1/4},$$

for n sufficiently large. It follows that $x'_n(\phi) < 0$ for $0 < \phi < \beta_n$, where

$$\beta_n = 2\pi/3 + 4\pi/3N - O(N^{-3/2}),$$

for n sufficiently large.

$n \equiv 2 \pmod{3}$. In this case the right side of (3.3) becomes

$$\begin{aligned} & -2N[1 - \cos(c + 4\pi/3)] \\ & + \sqrt{3}[c \cdot \cos(c + 4\pi/3) - \sin(c + 4\pi/3)] + O(1/N). \end{aligned}$$

It follows that $x'_n(\phi) < 0$ for $0 \leq c \leq 2\pi/3 - \epsilon$, $\epsilon > 0$, for n sufficiently large. Let

$$c = 2\pi/3 - \delta/\sqrt{N}.$$

Then the right side of (3.3) is equivalent to

$$-\delta^2 + 2\pi/\sqrt{3} + O(N^{-1/2}).$$

Thus

$$x'_n(\phi) < 0 \text{ if } \delta > (2\pi)^{1/2} \cdot 3^{1/4}$$

and

$$x'_n(\phi) > 0 \text{ if } \delta < (2\pi)^{1/2} \cdot 3^{1/4},$$

for n sufficiently large. It has been shown that $x_n'(\phi) < 0$ for $0 < \phi < \beta_n$, where

$$\beta_n = 2\pi/3 + 2\pi/3N - O(N^{-3/2}),$$

for n sufficiently large.

If $n + 3/2$ is substituted for N , then the results expressed in Theorem 2 are proved.

4. Proof of Theorem 3.

4.1 The Curvature of an image is defined to be

$$1/\rho = [1 + \Re z \cdot f''(z)/f'(z)] \cdot [|z \cdot f'(z)|]^{-1}.$$

If the point $w = f(z)$ traverses a closed, single-valued curve in a preassigned positive direction, then the curve is called convex if

$$(4.1) \quad 1 + \Re [z \cdot f''(z)/f'(z)] > 0.$$

Let us examine the inequality (4.1) for the function

$$f(e^{i\phi}) = s_n^2(e^{i\phi}) = x_n(\phi) + i \cdot y_n(\phi)$$

if $z = e^{i\phi}$. By the employment of differentiation and elementary algebraic steps after substituting the derivatives in the left side of (4.1), one obtains

$$1 + \Re [z \cdot f''(z)/f'(z)] = [x_n' \cdot y_n'' - x_n'' \cdot y_n'] \cdot [x_n'^2 + y_n'^2]^{-1}.$$

Thus the condition for the mapping to be convex is satisfied if

$$(4.2) \quad x_n' \cdot y_n'' - x_n'' \cdot y_n' > 0.$$

4.2. The next section studies the previous condition of convexity for the function $w = s_n^2(z)$, $z = e^{i\phi}$, where $\phi = \gamma/n$, $\gamma > 0$, for n sufficiently large.

In the present case the expressions for $y_n'(\gamma/n)$ and $x_n'(\gamma/n)$, for which see (2.1) and (3.1), become

$$y_n'(\gamma/n) = n^4/\gamma^4 [-\gamma \sin \gamma - 3 \cos \gamma + 3 - \gamma^2/2 + O(1/n)],$$

$$x_n'(\gamma/n) = n^4/\gamma^4 [-2\gamma - \gamma \cos \gamma + 3 \sin \gamma + O(1/n)].$$

Substitution of the latter expressions into (4.2) yields directly

$$(2 \sin \gamma - \gamma \cos \gamma - \gamma) (-2\gamma - \gamma \cos \gamma + 3 \sin \gamma) - (-2 + \gamma \sin \gamma + 2 \cos \gamma) (-\gamma \sin \gamma - 3 \cos \gamma + 3 - \gamma^2/2) + O(1/n) > 0.$$

Further simplification of the previous inequality, which establishes the requirement for convexity of the image of $|z| = 1$, leads to the convenient form

$$(4.3) \quad \sin \gamma (\tan \gamma/2 - \gamma/2) (6 - \gamma^2/2 - 3\gamma \cot \gamma/2) + O(1/n) > 0.$$

The remainder of the section is devoted to determining the maximum value of $\phi = \gamma/n$ which satisfies (4.3). In particular it is shown that the maximum angle $\gamma_n = \gamma/n$ for which the mapping of $|z| = 1$ by $w = s_n^2(z)$ is convex, where $z = e^{i\phi}$, is determined by $2\pi < \gamma < 3\pi$, for n sufficiently large.

4.3. Consider the elementary function

$$v(\gamma) = \sin \gamma [\tan(\gamma/2) - \gamma/2].$$

Define γ_0 by the equality $\tan(\gamma_0/2) = \gamma_0/2$. Then it is easily shown that

$$(4.4) \quad v(\gamma) \begin{cases} > 0, & 0 < \gamma < 2\pi, \\ < 0, & 2\pi < \gamma < \gamma_0, \\ > 0, & \gamma_0 < \gamma < 3\pi, \end{cases}$$

Let us define

$$f(\gamma) = 6 - \gamma^2/2 - 3\gamma \cot(\gamma/2).$$

Then the image of $|z| = 1$ is convex if

$$(4.5) \quad f(\gamma) \begin{cases} > 0, & 0 < \gamma < 2\pi, \\ < 0, & 2\pi < \gamma < \gamma_0, \\ > 0, & \gamma_0 < \gamma < 3\pi, \end{cases}$$

for n sufficiently large.

Next it is shown that the first two inequalities for $f(\gamma)$ in (4.5) are satisfied, however, for $\gamma_0 < \gamma < 3\rho$, one finds that $f(\gamma) < 0$. Since

$$f'(\gamma) \cdot \sin^2(\gamma/2) = -\gamma \sin^2(\gamma/2) - 3/2 \cdot \sin \gamma + 3\gamma/2,$$

by a further differentiation with respect to γ one can obtain

$$\begin{aligned} d/d\gamma \{ f'(\gamma) \cdot \sin^2(\gamma/2) \} &= 1 - \cos \gamma - \gamma/2 \sin \gamma, \\ &= \sin \gamma \{ \tan(\gamma/2) - \gamma/2 \} = v(\gamma). \end{aligned}$$

Consider the interval $0 < \gamma < 2\pi$. By (4.4), $v(\gamma) > 0$. Also $f'(\gamma) \cdot \sin^2(\gamma/2) = 0$ if $\gamma = 0$. Thus $f'(\gamma) \sin^2(\gamma/2) > 0$ for $0 < \gamma < 2\pi$, and consequently $f'(\gamma) > 0$ for the same interval. Finally,

$$f(0) = \lim_{\gamma \rightarrow 0^+} f(\gamma) = 0,$$

which establishes the fact that $f(\gamma) > 0$ in the interval $0 < \gamma < 2\pi$.

In the interval $2\pi < \gamma < \gamma_0$, $v(\gamma) < 0$; therefore the function $f'(\gamma) \sin^2(\gamma/2)$ is decreasing. It follows that

$$f'(\gamma_0) \cdot \sin^2(\gamma_0/2) = \gamma_0 [3/2 - \sin^2(\gamma_0/2)] - 3/2 \cdot \sin \gamma_0,$$

and thus $f'(\gamma_0) \cdot \sin^2(\gamma_0/2) > 1/2(\gamma_0 - 3) > 0$. Consequently $f'(\gamma) \sin^2(\gamma/2)$ and also $f'(\gamma)$ are positive in the interval $2\pi < \gamma < \gamma_0$. Hence $f(\gamma)$ is increasing. As $f(\gamma)$ has no lower bound as γ approaches 2π from above, and

$$f(\gamma_0) = 6 - \gamma_0^2/2 - 3\gamma_0 \cot(\gamma_0/2) < -\gamma_0^2/2 < 0,$$

then it can be concluded that $f(\gamma) < 0$ for $2\pi < \gamma < \gamma_0$.

Finally consider the interval $\gamma_0 < \gamma < 3\pi$. Since $v(\gamma) > 0$, $f'(\gamma_0) \sin^2(\gamma_0/2) > 0$, and thus $f'(\gamma_0) > 0$, then $f'(\gamma) > 0$ holds for $\gamma_0 < \gamma < 3\pi$. Hence $f(\gamma)$ is increasing. But

$$f(3\pi) = 6 - 9\pi^2/2 < 0,$$

so that a γ exists such that $f(\gamma) < 0$ occurs in the interval.

It was shown in (4.5) that if the image of $|z| = 1$ was to be convex for $\gamma_0 < \gamma < 3\pi$, then $f(\gamma) > 0$. Thus the image is not convex for the complete interval, which completes the proof of Theorem 3.

REFERENCES

1. E. Egerváry, *Abbildungseigenschaften der arithmetischen Mittel der geometrischen Reihe*, Math. Z. 42 (1936-1937), 221-230.

2. L. Fejér, *Potenzreihen mit mehrfach monotoner Koeffizientenfolge und ihre Legendrepolynome*, Proc. Cambridge Philos. Soc. **31** (1935), 307-316.
3. ———, *Trigonometrische Reihen und Potenzreihen mit mehrfach monotoner Koeffizientenfolge*, Trans. Amer. Math. Soc. **39** (1936), 18-59.
4. ———, *Untersuchungen über Potenzreihen mit mehrfach monotoner Koeffizientenfolge*, Acta Sci. Math. Szeged **8** (1936), 90-116.
5. L. Fejér and G. Szegő, *Über die monotone Konvergenz von Potenzreihen mit mehrfach monotoner Koeffizientenfolge*, Prace Mat. Fiz. **44** (1935), 15-25.
6. ———, *Special conformal mappings*, Duke Math. J. **18** (1951), 535-547.
7. K. Knopp, *Mehrfach monotone Zahlenfolgen*, Math. Z. **22** (1925), 75-85.
8. M. Schweitzer, *The partial sums of second order of the geometric series*, Duke Math. J. **18** (1951), 527-533.
9. S. Sidon, *Über Potenzreihen mit monotoner Koeffizientenfolge*, Acta Sci. Math. Szeged **9** (1938-1940), 244-246.
10. G. Szegő, *Power series with multiply monotonic sequences of coefficients*, Duke Math. J. **8** (1941), 559-564.

UNIVERSITY OF CALIFORNIA
SANTA BARBARA COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

M. M. SCHIFFER*
Stanford University
Stanford, California

E. HEWITT
University of Washington
Seattle 5, Washington

R. P. DILWORTH
California Institute of Technology
Pasadena 4, California

E. F. BECKENBACH**
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN

HERBERT FEDERER

MARSHALL HALL

P. R. HALMOS

HEINZ HOPF

R. D. JAMES

BØRGE JESSEN

PAUL LÉVY

GEORGE PÓLYA

J. J. STOKER

E. G. STRAUS

KÔSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA, BERKELEY
UNIVERSITY OF CALIFORNIA, DAVIS
UNIVERSITY OF CALIFORNIA, LOS ANGELES
UNIVERSITY OF CALIFORNIA, SANTA BARBARA
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD RESEARCH INSTITUTE
STANFORD UNIVERSITY
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
HUGHES AIRCRAFT COMPANY
SHELL DEVELOPMENT COMPANY

* To be succeeded in 1955, by H.L. Royden, Stanford University, Stanford, California.

** To be succeeded in 1955, by E.G. Straus, University of California, Los Angeles 24, Calif.

Vari-Type Composition by
Elaine Barth

Printed in the United States of America by
Edwards Brothers, Inc., Ann Arbor, Michigan

UNIVERSITY OF CALIFORNIA PRESS * BERKELEY AND LOS ANGELES
COPYRIGHT 1954 BY PACIFIC JOURNAL OF MATHEMATICS

Hugh D. Brunk, <i>On the growth of functions having poles or zeros on the positive real axis</i>	1
J. Copping, <i>Application of a theorem of Pólya to the solution of an infinite matrix equation</i>	21
James Richard Jackson, <i>On the existence problem of linear programming</i>	29
Victor Klee, <i>Invariant extension of linear functionals</i>	37
Shu-Teh Chen Moy, <i>Characterizations of conditional expectation as a transformation on function spaces</i>	47
Hukukane Nikaidô, <i>On von Neumann's minimax theorem</i>	65
Gordon Marshall Petersen, <i>Methods of summation</i>	73
G. Power, <i>Some perturbed electrostatic fields</i>	79
Murray Harold Protter, <i>The two noncharacteristic problem with data partly on the parabolic line</i>	99
S. E. Rauch, <i>Mapping properties of Cesàro sums of order two of the geometric series</i>	109
Gerson B. Robison, <i>Invariant integrals over a class of Banach spaces</i>	123
Richard Steven Varga, <i>Eigenvalues of circulant matrices</i>	151