

# Pacific Journal of Mathematics

**ON THE DIMENSION THEORY OF RINGS. II**

A. SEIDENBERG

## ON THE DIMENSION THEORY OF RINGS (II)

A. SEIDENBERG

**1. Introduction.** As in [3], we shall say that an integral domain  $O$  is  $n$ -dimensional if in  $O$  there is a proper chain

$$(0) \subset P_1 \subset \dots \subset P_n \subset (1)$$

of prime ideals, but no such chain

$$(0) \subset P'_1 \subset \dots \subset P'_{n+1} \subset (1).$$

In Theorem 2 of [3] it was shown that if  $O$  is  $n$ -dimensional, then  $O[x]$  is at least  $(n+1)$ -dimensional and at most  $(2n+1)$ -dimensional: here, as throughout,  $x$  is an indeterminate. After preparatory constructions in Theorems 1 and 2 below, this theorem is completed in Theorem 3 by showing that for any integers  $m$  and  $n$  with  $n+1 \leq m \leq 2n+1$ , there exist  $n$ -dimensional rings  $O$  such that  $O[x]$  is  $m$ -dimensional. The other theorems mainly concern 1-dimensional rings. Such rings  $O$  can be divided into those for which  $O[x]$  is 2-dimensional and those for which this condition fails, the so-called  $F$ -rings. The paper [3] was concerned with the existence of  $F$ -rings and showed [3, Theorem 8] that the 1-dimensional ring  $O$  is not an  $F$ -ring if and only if every quotient ring of the integral closure of  $O$  is a valuation ring. Below, in Theorem 5, we show more generally that if  $O$  is 1-dimensional but not an  $F$ -ring, then  $O[x_1, \dots, x_n]$  is  $(n+1)$ -dimensional, where the  $x_i$  are indeterminates: this theorem depends on the essentially more general Theorem 4, which says that if  $O$  is an  $m$ -dimensional multiplication-ring, then  $O[x_1, \dots, x_n]$  is  $(m+n)$ -dimensional. In the case that the  $x_i$  are not indeterminates, one can still say (Theorem 10) that

$$\dim O[x_1, \dots, x_n] = 1 + \text{degree of transcendency of } O[x_1, \dots, x_n]/O,$$

provided that the intersection of the prime ideals ( $\neq (0)$ ) in  $O$  is  $(0)$ , where  $O$  is a 1-dimensional ring such that  $O[x]$  is 2-dimensional. For  $F$ -rings  $O$ , Theorem 6 shows that

$$n+2 \leq \dim O[x_1, \dots, x_n] \leq 2n+1,$$

---

Received April 10, 1953.

*Pacific J. Math.* 4 (1954), 603-614

where the  $x_i$  are indeterminates, while Theorem 7 constructs for any  $N$  and  $n$  with

$$n + 2 \leq N \leq 2n + 1$$

an  $F$ -ring  $O$  such that  $O[x_1, \dots, x_n]$  is  $N$ -dimensional. Similar results for rings of dimension greater than 1 would be interesting if one could get them.

**2. Simple extensions.** Let us call the integral domain  $O$  of type  $(n, m)$  if

$$\dim O = n \text{ and } \dim O[x] = m.$$

**THEOREM 1.** *Let  $O$  be integrally closed and of type  $(n, m)$ , let  $K$  be its quotient field, and let  $K'$  be a proper extension of  $K$  in which  $K$  is algebraically closed. Let  $\Sigma$  be any field having a discrete rank 1 valuation with  $K'$  as residue field. Let  $O^*$  be the set of elements whose residues are finite and in  $O$ . Then  $O^*$  is integrally closed and of type  $(n + 1, m + 2)$ .*

*Proof.* Let  $\alpha \in \Sigma$ , with  $\alpha$  integral over  $O^*$ ,

$$\alpha^s + a_1 \alpha^{s-1} + \dots + a_s = 0 \quad (a_i \in O^*),$$

an equation of integral dependence. Dividing this equation by  $\alpha^s$  and supposing  $1/\alpha$  to have residue 0, we get the contradiction  $1 = 0$ . So  $\alpha$  has finite residue, and

$$\bar{\alpha}^s + \bar{a}_1 \bar{\alpha}^{s-1} + \dots + \bar{a}_s = 0,$$

where the bars indicate residues. Since  $K$  is algebraically closed in  $K'$ , we have  $\bar{\alpha} \in K$ ; and  $\bar{\alpha} \in O$ , since  $O$  is integrally closed. Hence  $O^*$  is integrally closed.

Let  $P$  be the set of  $\alpha \in O^*$  having residue 0. Then  $P$  is a prime ideal. From the definitions one obtains

$$O^*/P \simeq 0,$$

whence  $O^*$  is at least  $(n+1)$ -dimensional. If  $P'$  is a prime ideal in  $O^*$ ,  $P' \neq (0)$ , then  $P' \supseteq P$ . In fact, let  $g \in P'$ ; since  $g$  is  $O^*$ , we have  $v(g) = s \geq 0$ , where  $v$  is the given valuation (and the group of integers is the valuation group). Then the  $(s+1)$ th power of any element in  $P$  is divisible by  $g$ , whence  $P \subseteq P'$ . From this it follows that  $O^*$  is at most  $(n+1)$ -dimensional.

The quotient ring  $O_p^*$  is integrally closed and has only one prime ideal

( $\neq (0)$ ). Moreover it is not a valuation ring. In fact, let  $\alpha \in \Sigma$  be an element having residue in  $K'$  but not in  $K$ . Since  $\alpha$  can clearly be written as a quotient of two elements of positive value, we have that  $\alpha$  is in the quotient field of  $O_P^*$ ; but neither  $\alpha$  nor  $1/\alpha$  has residue in  $K$ , so neither  $\alpha$  nor  $1/\alpha$  is in  $O_P^*$ . Thus  $O_P^*$  is not a valuation ring, and hence is an  $F$ -ring, by [3, Theorem 8]. It follows at once that  $O^*[x] \cdot P$  is not minimal in  $O^*[x]$ . Now

$$O^*[x]/O^*[x] \cdot P \simeq O^*/P[x] \simeq O[x],$$

so  $O^*[x]$  is at least  $(m + 2)$ -dimensional.

Finally, let  $(0) \subset P_1 \subset P_2 \subset \dots \subset P_s \subset (1)$  be a chain of prime ideals in  $O^*[x]$ . Let  $P_1$  be minimal; then  $P_1 \cap O^* = (0)$ , as otherwise

$$P_1 \cap O \supseteq P \text{ and } P_1 \supseteq O^*[x] \cdot P.$$

Similarly one concludes that if no chain of prime ideals  $P_1' \subset P_1''$  can be inserted between  $(0)$  and  $P_2$ , then

$$P_2 \cap O^* = P \text{ and } P_2 = O^*[x] \cdot P$$

(by [3, Theorem 1],  $P_2$  cannot contract in  $O$  to  $(0)$ ). From this it follows at once that  $O^*[x]$  is at most  $(m + 2)$ -dimensional, and the proof is complete.

REMARK. The above construction stems from an example of Krull showing that an integrally closed integral domain with only one proper prime ideal need not be a valuation ring; see [2, p. 670f].

THEOREM 2. *Let  $O, K, K', \Sigma, O^*$  be as in Theorem 1 except that we assume  $K = K'$ . Then  $O^*$  is integrally closed and of type  $(n + 1, m + 1)$ .*

*Proof.* The proof follows exactly the lines of the proof of Theorem 1, except that here  $O_P^*$  is a valuation ring, as one easily sees.

THEOREM 3. *For every  $n$  and  $m$  such that  $n + 1 \leq m \leq 2n + 1$  there exist integrally closed rings of type  $(n, m)$ .*

*Proof.* Any field is of type  $(0, 1)$ . Theorem 1 gives us an integrally closed ring of type  $(1, 3)$ , and Theorem 2 gives us one of type  $(1, 2)$ —the required valuations obviously exist. Suppose now by induction that for some  $n$  and each  $m, n + 1 \leq m \leq 2n + 1$ , we have an integrally closed ring of type  $(n, m)$ . If  $n + 3 \leq m \leq 2n + 3$ , then  $n + 1 \leq m - 2 \leq 2n + 1$ , and from an integrally closed ring of type  $(n, m - 2)$  we get, by Theorem 1, an integrally closed ring of type

$(n + 1, m)$ . If  $m = n + 2$ , we apply Theorem 2 similarly to get an integrally closed ring of type  $(n + 1, m)$ .

As for simple algebraic extensions  $O[\alpha]$  of an  $n$ -dimensional ring  $O$ , it is clear that  $\dim O[\alpha] \leq 2n$ . On the other hand, let  $O$  be an integrally closed ring of type  $(n, m)$  and let  $O^*$  be a ring constructed as in Theorem 1; also let  $\Sigma$  and  $P$  be as in Theorem 1. Let

$$\alpha \in \Sigma, \alpha \notin O_p^*, 1/\alpha \notin O_p^* .$$

Then

$$O^*[\alpha]/O^*[\alpha] \cdot P \simeq O^*/P[x] \simeq O[x],$$

by [3, Theorem 7], so  $O^*[\alpha]$  is at least  $(m + 1)$ -dimensional; it is also at most  $(m + 1)$ -dimensional, since  $O^*[x]$  is  $(m + 2)$ -dimensional. Hence

$$(n + 1) + 1 \leq \dim O^*[\alpha] \leq 2(n + 1) .$$

It is thus clear that for any  $n' > 0$  and  $m'$  with  $n' + 1 \leq m' \leq 2n'$ , there exists an  $n'$ -dimensional ring  $O^*$  such that for some  $\alpha$  in the quotient-field of  $O^*$  we have  $\dim O^*[\alpha] = m'$ . — Also

$$\dim O[\alpha] < \dim O$$

is possible. In fact, let  $O$  be a valuation ring of rank  $n$ ,  $(0) \subset p_1 \subset \dots \subset p_n \subset (1)$ , the chain of prime ideals in  $O$ . Let  $c \in p_{i+1}$ ,  $c \notin p_i$ ; then  $\dim O[1/c] = 1$ . In short,  $\dim O[\alpha]$  covers precisely the range from 0 to  $2n$  as  $O$  varies over the  $n$ -dimensional rings  $O$ .

**3. Multiple transcendental extensions.** We recall that a *multiplication-ring* may be defined as an integral domain  $O$  such that  $O_p$  is a valuation ring for each prime ideal  $p$  in  $O$  (see [2, p. 554]).

**THEOREM 4.** *If  $O$  is an  $m$ -dimensional multiplication-ring, then  $O[x_1, \dots, x_n]$  is  $(m + n)$ -dimensional, where the  $x_i$  are indeterminates.*

*Proof.* To facilitate the proof, we define the *dimension of a prime ideal  $P$*  in an extension  $O' = O[\alpha_1, \dots, \alpha_n]$  of a finite-dimensional ring  $O$  (relative to  $O$ ) as follows:

$$\dim P = \text{d.t. } (O'/P)/(O/P) + \dim O/p,$$

where  $p = P \cap O$  (and “d.t.” abbreviates “degree of transcendence”). The

following points (a), (b) do not assume  $O$  to be a multiplication-ring.

(a) Let  $\bar{O}'$ ,  $\bar{O}$ ,  $\bar{P}$ ,  $\bar{p}$  be the images of  $O'$ ,  $O$ ,  $P$ ,  $p$ , respectively, under a homomorphism with kernel contained in  $P$ . Then  $\dim \bar{P} = \dim P$ .

In fact,  $\bar{O}'/\bar{P} = O'/P$  and  $\bar{O}/\bar{p} = O/p$ ; also  $\bar{P} \cap \bar{O} = \bar{p}$ .

(b) Let  $M$  be a nonempty multiplicatively closed system in  $O$  not meeting  $p$ ,

$$O_M = \{ \alpha \mid \alpha = a/b, a \in O, b \in M \},$$

$$O'_M = \{ \alpha \mid \alpha = a/b, a \in O', b \in M \}.$$

Then

$$\dim P - \dim O/p = \dim O'_M \cdot P - \dim O_M/O_M \cdot p.$$

In fact, the rings  $O'/P$  and  $O'_M/O'_M \cdot P$  have the same quotient field, as do the rings  $O/p$  and  $O_M/O_M \cdot p$ . Note also that  $O'_M \cdot P \cap O_M = O_M \cdot p$ , whence the required equality follows.

Let  $P_1, P_2$  be two prime ideals in  $O'$ ,  $P_1 \subset P_2$ ,  $p_i = P_i \cap O$ ,  $i = 1, 2$ . We want to compare  $\dim P_1$  with  $\dim P_2$ . If  $p_1 = p_2$ , then, passing to a residue class ring, we may assume  $p_1 = p_2 = (0)$ . Taking  $M = O - (0)$ , we pass to the quotient-ring  $O'_M$ , which is a finite integral domain. Thus  $\dim P_1 > \dim P_2$  if  $p_1 = p_2$ . This conclusion holds also if  $p_1 \subset p_2$  provided  $O$  is a multiplication-ring.

(c) If  $P_1$  and  $P_2$  are prime ideals in  $O[x_1, \dots, x_n]$  and  $P_1 \subset P_2$ , then

$$\dim P_1 > \dim P_2;$$

also

$$\dim P_1 - \dim P_2 \geq \dim O/p_1 - \dim O/p_2,$$

provided that  $O$  is a multiplication-ring.

In fact, we may suppose  $p_1 \subset p_2$ , and have only to prove the second point. Also, by (b), we may pass to any quotient-ring  $O_M$ , where  $M$  does not meet  $p_2$ . Taking  $M = O - p_2$ , we may assume that  $O$  is a valuation-ring and that  $p_2$  is its ideal of non-units. Let  $z_1, \dots, z_r$  be elements of  $O'$  which are algebraically dependent mod  $P_1$  over  $O$ . Then they are also dependent mod  $P_2$ . In fact, let

$$f(z_1, \dots, z_r) \equiv 0(P_1),$$

where the coefficients of the polynomial  $f$  are in  $O$  but not all in  $p_1$ . Dividing by a coefficient of least value, we may suppose  $f$  to have a coefficient equal to unity. But then we have a relation mod  $P_2$ . This proves that

$$\text{d.t. } (O'/P_2)/(O/p_2) \leq \text{d.t. } (O'/P_1)/(O/p_1),$$

that is, (c) is proved.

The theorem now follows from (c) since  $\dim(0) = m + n$ .

**COROLLARY.** *If  $O$  is an  $m$ -dimensional multiplication-ring then*

$$\dim O[\alpha_1, \dots, \alpha_n] \leq m + r,$$

where

$$r = \text{d.t. } O[\alpha_1, \dots, \alpha_n]/O.$$

*Proof.* The foregoing proof shows that

$$\dim O[x_1, \dots, x_n] \leq \dim(0) = m + \text{d.t. } O[x_1, \dots, x_n]/O,$$

and in doing so makes no use of the fact that the  $x_i$  are indeterminates; this fact is used only to get that

$$\dim O[x_1, \dots, x_n] \geq m + n.$$

**THEOREM 5.** *If  $O$  is a 1-dimensional ring such that  $O[x]$  is 2-dimensional, then*

$$\dim O[\alpha_1, \dots, \alpha_n] \leq 1 + \text{d.t. } O[\alpha_1, \dots, \alpha_n]/O;$$

if the  $\alpha_i$  are indeterminates, then

$$\dim O[\alpha_1, \dots, \alpha_n] = 1 + n.$$

*Proof.* We may suppose  $O$  to be integrally closed. In that event,  $O$  is a multiplication-ring, by [3, Theorem 8]. The present theorem now follows immediately from the preceding corollary.

**THEOREM 6.** *If  $O$  is 1-dimensional, then  $O[x_1, \dots, x_n]$  is at most  $(2n+1)$ -dimensional, where the  $x_i$  are indeterminates.*

*Proof.* Let  $(0) \subset p_1 \subset p_2 \subset \dots \subset p_s \subset (1)$  be a chain of prime ideals in

$O[x_1, \dots, x_n]$ . Let  $K =$  quotient field of  $O$ . If  $p_s \cap O = (0)$ , then the above chain extends to a chain of  $s$  prime ideals in  $K[x_1, \dots, x_n]$ , so  $s \leq n$ . Suppose, then, that

$$p_i \cap O = (0), \quad p_{i+1} \cap O = p \neq (0),$$

whence also  $p_{i+k} \cap O = p$ , since  $O$  is 1-dimensional. Passing to  $K[x_1, \dots, x_n]$ , we see that  $i \leq n$ ; and passing to

$$O[x_1, \dots, x_n] / p_{i+1} = O/p [\bar{x}_1, \dots, \bar{x}_n],$$

we have  $s - (i + 1) \leq n$ , since  $O/p$  is a field. Hence  $s \leq 2n + 1$ .

**THEOREM 7.** *If  $O$  is an  $F$ -ring, then  $O[x_1, \dots, x_n]$  is at least  $(n + 2)$ -dimensional and at most  $(2n + 1)$ -dimensional. For any  $N$ ,  $n + 2 \leq N \leq 2n + 1$ , there is an  $F$ -ring  $O$  such that  $O[x_1, \dots, x_n]$  is  $N$ -dimensional, where the  $x_i$  are indeterminates.*

*Proof.* Let  $K$  be a field,  $x, y_1, \dots, y_m$  indeterminates. Let

$$K' = K(y_1, \dots, y_m), \quad \Sigma = K'(x),$$

and let  $v$  be the discrete rank 1 valuation of  $\Sigma$  obtained by placing

$$v(a_i x^i + a_{i+1} x^{i+1} + \dots + a_s x^s) = i,$$

where  $a_j \in K'$ ,  $a_i \neq 0$ . Let  $O^*$  be the set of elements whose residues are finite and in  $K$ . The ring  $O^*$  consists of the elements in  $K(x, y_1, \dots, y_m)$  which can be written in the form

$$\alpha(x, y_1, \dots, y_m) / \beta(x, y_1, \dots, y_m),$$

where

$$\alpha, \beta \in K[x, y_1, \dots, y_m], \quad \beta(0, y_1, \dots, y_m) \neq 0,$$

$$\alpha(0, y_1, \dots, y_m) / \beta(0, y_1, \dots, y_m) \in K.$$

By Theorem 1,  $O^*$  is an  $F$ -ring; and  $O^*$  contains only one proper prime ideal, namely the ideal  $P$  consisting of the elements  $\alpha/\beta$  with

$$\alpha(0, x_1, \dots, x_m) / \beta(0, x_1, \dots, x_n) = 0.$$



We shall prove that for  $m \leq n$ ,  $O^*[x_1, \dots, x_n]$  is  $(m+n+1)$ -dimensional. In  $O^*[x_1, \dots, x_n]$  let  $P_m$  be the ideal of polynomials which vanish for  $x_i = y_i$  ( $i = 1, \dots, m$ ). We claim this ideal is in

$$O^*[x_1, \dots, x_n] \cdot P = P'.$$

In fact, let

$$\sum a_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n} \in O^*[x_1, \dots, x_n]$$

be in  $P_m$ , and write

$$\sum a_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n} = \sum c_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n} + \sum d_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n},$$

where  $c_{i_1 \dots i_n} \in K$ ,  $d_{i_1 \dots i_n} \in P$ . This polynomial vanishes for  $x_i = y_i$ ,  $i = 1, \dots, m$ ; hence also for  $x_i = y_i$ ,  $i = 1, \dots, m$ ,  $x = 0$ . Hence

$$\sum c_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n}$$

vanishes for  $x_i = y_i$ ,  $i = 1, \dots, m$ , whence

$$\sum c_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n} = 0,$$

and

$$\sum a_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n} = \sum d_{i_1 \dots i_n} x_1^{i_1} \dots x_n^{i_n} \in O^*[x_1, \dots, x_n] \cdot P = P'.$$

Let  $P_j$  be the ideal of elements in  $O^*[x_1, \dots, x_n]$  which vanish for  $x_i = y_i$ ,  $i = 1, \dots, j$ . Then  $P_j$  is prime and  $(0) \subset P_1 \subset \dots \subset P_m \subset P'$ . Since any chain of  $n$  prime ideals in  $O^*/P[x_1, \dots, x_n]$  gives rise to such a chain in  $O^*[x_1, \dots, x_n]$  containing  $P'$ , we see that  $O^*[x_1, \dots, x_n]$  is at least  $(m+n+1)$ -dimensional. On the other hand,  $O^*[x_1, \dots, x_n]$  is of degree of transcendency  $m+n+1$  over  $K$ , and so  $O^*[x_1, \dots, x_n]$  is at most  $(m+n+1)$ -dimensional. This last point follows from the following lemma, the proof of which is exactly as in the well-known case that  $O$  is a valuation ring.

LEMMA. *Let  $O$  be an arbitrary integral domain containing a field  $K$ , and let  $O$  be of degree of transcendency  $r$  over  $K$ . Then  $O$  is at most  $r$ -dimensional.*

*Proof.* This follows at once if we can show that the degree of transcendency

of  $O/P$  over  $K$  is less than  $r$  for any proper prime ideal  $P$  in  $O$ . If  $\theta_1, \theta_2, \dots, \theta_s \in O$  map into (given) algebraically independent elements in  $O/P$ , and  $\theta \in P$ ,  $\theta \neq 0$ , then  $\theta, \theta_1, \dots, \theta_s$  are algebraically independent over  $K$ . Hence

$$\text{d.t. } O/K > \text{d.t. } (O/P)/K.$$

**4. Arbitrary finite extensions.** Let  $O$  be an arbitrary integral domain which is not a field. It is certainly possible, for appropriate  $O$ , that some simple ring extension  $O[\alpha]$  of  $O$  will be a field. In fact, let  $O$  be such that the intersection of all its prime ideals ( $\neq (0)$ ) is not the ideal  $(0)$ ; for example, any integral domain with a finite, positive number of prime ideals ( $\neq (0)$ ) will do. If  $c (\neq 0)$  is an element in all the prime ideals, then  $O[1/c]$  is a field; for if  $P$  is a prime ideal in  $O[1/c]$ ,  $P \neq (0)$ , then

$$P \cap O = p \neq (0)$$

and

$$1 = (1/c) \cdot c \in O[1/c] \cdot p \subseteq P.$$

We also have the converse.

**THEOREM 8.** *Given an integral domain  $O$ , there exists a field  $F$  which is a simple ring extension of  $O$  if and only if the intersection of all the prime ideals ( $\neq (0)$ ) in  $O$  is  $\neq (0)$ .*

*Proof.* Let  $F = O[\alpha]$ . Here  $\alpha$  must be algebraic over  $O$ , say

$$c_0 \alpha^m + c_1 \alpha^{m-1} + \dots + c_m = 0, \quad c_i \in O, \quad c_0 \neq 0.$$

Then  $c_0 \alpha$  is integral over  $O$ , as is the ring  $O_1 = O[c_0 \alpha]$ . Let  $F_1 = O_1[\alpha]$ ; then  $F_1$  is a field [1, p. 253]. Over every prime ideal in  $O$  there lies a prime ideal in  $O_1$ ; since  $O_1$  is algebraic over  $O$ , if the intersection of the prime ideals ( $\neq (0)$ ) in  $O_1$  is  $\neq (0)$ , then the like is true in  $O$ . Hence we may assume that  $O = O_1$ , that is, that  $\alpha$  is in the quotient field of  $O$ . By a similar reasoning we may suppose  $O$  is integrally closed. From the fact that  $1/\alpha \in O[\alpha]$ , one finds that  $1/\alpha$  is integral over  $O$ , hence in  $O$ . Thus  $\alpha = 1/b$ ,  $b \in O$ . The element  $b$  must be in every prime ideal  $p (\neq (0))$ ; in fact, if  $b \notin p$ , then  $O[1/b] \subseteq O_p$ , whence  $O[1/b] = O[\alpha]$  is not a field. This completes the proof. — This theorem has been previously proved in [4, p. 76].

A study of algebraic extensions of  $O$  must therefore separate the cases that

the intersection of the prime ideals ( $\neq (0)$ ) is  $= (0)$  or is  $\neq (0)$ .

**THEOREM 9.** *If  $O$  is a 1-dimensional ring such that  $O[x]$  is 2-dimensional, and the intersection of the prime ideals ( $\neq (0)$ ) in  $O$  is  $= (0)$ , then the like is true of any simple algebraic ring extension  $O[\alpha]$  of  $O$  (where it is assumed, of course, that  $O[\alpha]$  is an integral domain).*

*Proof.* By Theorem 5, we know that  $O[\alpha]$  is 0- or 1-dimensional, and the previous theorem excludes the first alternative. Also  $O[\alpha, x]$  is 2-dimensional, for otherwise  $O[y, x]$ ,  $y$  an indeterminate, would be of dimension more than 3, contradicting Theorem 5. Thus it remains to prove that the intersection of the prime ideals ( $\neq (0)$ ) in  $O[\alpha]$  is  $= (0)$ . Let

$$c_0 \alpha^n + c_1 \alpha^{n-1} + \dots + c_n = 0, \quad c_i \in O, \quad c_0 \neq 0,$$

and let  $S = \{p\}$  be the set of prime ideals ( $\neq (0)$ ) in  $O$  which do not contain  $c_0$ ;  $S$  is not empty. Then  $\bigcap p = (0)$ , for if  $d \in \bigcap p$ ,  $d \neq 0$ , then  $c_0 d$  is in every prime ideal ( $\neq (0)$ ) of  $O$ . Over every prime ideal  $p \in S$  there lies a prime ideal  $P$  in  $O[\alpha]$ . If  $T = \{P\}$  is the set of prime ideals in  $O[\alpha]$  contracting to prime ideals in  $S$ , then one concludes immediately that  $\bigcap P = (0)$ . A fortiori the intersection of all prime ideals ( $\neq (0)$ ) in  $O[\alpha]$  is  $= (0)$ . This completes the proof.

If  $O$  is an integral domain in which the intersection ( $\bigcap p$ ) of the prime ideals ( $\neq (0)$ ) is  $\neq (0)$ , then for every  $r$  it is possible to define a finite extension  $O[\alpha_1, \dots, \alpha_n]$  of  $O$  such that

$$\dim O[\alpha_1, \dots, \alpha_n] = r$$

and

$$\text{d.t. } O[\alpha_1, \dots, \alpha_n] / O = r;$$

namely, we adjoin to  $O$  an element  $1/c$ ,  $c \in \bigcap p$ , so that  $O[1/c]$  is the quotient field of  $O$ , and thereupon adjoin  $r$  indeterminates. The situation is different for a 1-dimensional ring which is not an  $F$ -ring and in which the intersection of the prime ideals ( $\neq (0)$ ) is  $= (0)$ .

**THEOREM 10.** *Let  $O$  be a 1-dimensional ring such that  $O[x]$  is 2-dimensional, and let the intersection of the prime ideals ( $\neq (0)$ ) in  $O$  be  $= (0)$ . Then for any integral domain  $O[\alpha_1, \dots, \alpha_n]$ ,*

$$\dim O[\alpha_1, \dots, \alpha_n] = 1 + \text{d.t. } O[\alpha_1, \dots, \alpha_n]/O.$$

*Proof.* Let

$$K = \text{quotient field of } O, \quad r = \text{d.t. } O[\alpha_1, \dots, \alpha_n]/O.$$

Then  $K[\alpha_1, \dots, \alpha_n]$  is  $r$ -dimensional and a chain  $(0) \subset P_1 \subset \dots \subset P_r \subset (1)$  of prime ideals in  $K[\alpha_1, \dots, \alpha_n]$  contracts to a chain

$$(0) \subset p_1 \subset \dots \subset p_r \subset (1), \text{ and } p_i \cap O = (0), \quad i = 1, \dots, r.$$

Moreover,  $p_r$  is not maximal, for if it were, then

$$O[\alpha_1, \dots, \alpha_n]/p_r = O[\bar{\alpha}_1, \dots, \bar{\alpha}_n]$$

would be a field; hence also  $K[\bar{\alpha}_1, \dots, \bar{\alpha}_n]$  would be a field, whence the  $\bar{\alpha}_i$  would be algebraic over  $K$ , therefore also over  $O$ . This contradicts the previous theorem. Hence

$$\dim O[\alpha_1, \dots, \alpha_n] \geq 1 + \text{d.t. } O[\alpha_1, \dots, \alpha_n]/O,$$

and we have already seen the reverse inequality.

Since the theory of 1-dimensional rings must separate the cases that the intersections of prime ideals ( $\neq (0)$ ) is  $= (0)$  or  $\neq (0)$ , it may be of interest to have an example of a 1-dimensional ring, not an  $F$ -ring, with infinitely many prime ideals ( $\neq (0)$ ) having intersection  $\neq (0)$ . We construct such a ring  $O$  as follows. Let  $K$  be a field containing all roots of unity,  $x$  an indeterminate,  $L$  the algebraic closure of  $K(x)$ ,  $S$  the integral closure in  $L$  of  $K[x]$ , and  $O$ , the quotient-ring of  $S$  with respect to the multiplicatively closed system of polynomials in  $K[x]$  which are not divisible by  $x$ . Infinitely many prime ideals in  $S$  lie over  $(x)$  in  $K[x]$ ; to see this, let  $n$  be any integer not divisible by the characteristic of  $K$ ,  $a_1, \dots, a_n$  the  $n$ th roots of unity,  $y = \sqrt[n]{1+x}$ . In  $K[x, y]$  there lie  $n$  prime ideals over  $(x)$ , namely  $(x, y - a_i)$ , since  $(0, a_i)$  is a point of  $y^n = 1 + x$ . Going up to  $S$ , we see that there exist at least  $n$  prime ideals over  $(x)$ . Every prime ideal in  $O$  which differs from  $(0)$  contains  $x$ , and there are infinitely many such ideals. We now verify immediately that  $O$  is a ring of the required type.

#### REFERENCES

1. I. S. Cohen and A. Seidenberg, *Prime ideals and integral dependence*, Bull. Amer. Math. Soc. 52 (1946), 252-261.

2. W. Krull, *Beiträge zur Arithmetik kommutativer Integritätsbereiche*, I. *Multiplikationsringe, ausgezeichnete Idealsysteme, Kroneckersche Funktionalringe*, Math. Z. 41(1936), 545-577. II. *v-Ideale und vollständige ganz abgeschlossene Integritätsbereiche*, Math. Z. 41 (1936), 665-679.

3. A. Seidenberg, *A note on the dimension theory of rings*, Pacific J. Math. 3 (1953), 505-512.

4. E. Artin and J. T. Tate, *A note on finite ring extensions*, J. Math. Soc. Japan, 3 (1951), 74-77.

UNIVERSITY OF CALIFORNIA, BERKELEY

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

M.M. SCHIFFER\*  
Stanford University  
Stanford, California

R. P. DILWORTH  
California Institute of Technology  
Pasadena 4, California

E. HEWITT  
University of Washington  
Seattle 5, Washington

E.F. BECKENBACH\*\*  
University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

H. BUSEMANN	P. R. HALMOS	BØRGE JESSEN	J. J. STOKER
HERBERT FEDERER	HEINZ HOPF	PAUL LÉVY	E. G. STRAUS
MARSHALL HALL	R. D. JAMES	GEORGE PÓLYA	KÔSAKU YOSIDA

## SPONSORS

UNIVERSITY OF BRITISH COLUMBIA	UNIVERSITY OF SOUTHERN CALIFORNIA
CALIFORNIA INSTITUTE OF TECHNOLOGY	STANFORD RESEARCH INSTITUTE
UNIVERSITY OF CALIFORNIA, BERKELEY	STANFORD UNIVERSITY
UNIVERSITY OF CALIFORNIA, DAVIS	WASHINGTON STATE COLLEGE
UNIVERSITY OF CALIFORNIA, LOS ANGELES	UNIVERSITY OF WASHINGTON
UNIVERSITY OF CALIFORNIA, SANTA BARBARA	* * *
UNIVERSITY OF NEVADA	AMERICAN MATHEMATICAL SOCIETY
OREGON STATE COLLEGE	HUGHES AIRCRAFT COMPANY
UNIVERSITY OF OREGON	

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. Manuscripts intended for the outgoing editors should be sent to their successors. All other communications to the editors should be addressed to the managing editor, E.G. Straus, at the University of California Los Angeles 24, California.

50 reprints of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50; back numbers (Volumes 1, 2, 3) are available at \$2.50 per copy. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

\* To be succeeded in 1955, by H.L. Royden, Stanford University, Stanford, California.

\*\* To be succeeded in 1955, by E.G. Straus, University of California, Los Angeles 24, Calif.

UNIVERSITY OF CALIFORNIA PRESS · BERKELEY AND LOS ANGELES

COPYRIGHT 1954 BY PACIFIC JOURNAL OF MATHEMATICS

Paul Civin, <i>Orthonormal cyclic groups</i> .....	481
Kenneth Lloyd Cooke, <i>The rate of increase of real continuous solutions of algebraic differential-difference equations of the first order</i> .....	483
Philip J. Davis, <i>Linear functional equations and interpolation series</i> .....	503
F. Herzog and G. Piranian, <i>Sets of radial continuity of analytic functions</i> ...	533
P. C. Rosenbloom, <i>Comments on the preceding paper by Herzog and Piranian</i> .....	539
Donald G. Higman, <i>Remarks on splitting extensions</i> .....	545
Margaret Jackson, <i>Transformations of series of the type <math>{}_3\Psi_3</math></i> .....	557
Herman Rubin and Patrick Colonel Suppes, <i>Transformations of systems of relativistic particle mechanics</i> .....	563
A. Seidenberg, <i>On the dimension theory of rings. II</i> .....	603
Bertram Yood, <i>Difference algebras of linear transformations on a Banach space</i> .....	615