

Pacific Journal of Mathematics

CANTOR-TYPE UNIQUENESS OF MULTIPLE TRIGONOMETRIC INTEGRALS

VICTOR LENARD SHAPIRO

CANTOR-TYPE UNIQUENESS OF MULTIPLE TRIGONOMETRIC INTEGRALS

VICTOR L. SHAPIRO

1. Introduction. It is the purpose of this paper to obtain results in Cantor-type uniqueness for multiple trigonometric integrals similar to those obtained previously for multiple trigonometric series ([5, 11, 12]). As might be expected, the results in the integral case are a bit more difficult to obtain.

Vectorial notation is used for the most part throughout this paper. Thus u designates the point in n -dimensional euclidean space, E_n , with coordinates (u_1, \dots, u_n) , the scalar product $(u, x) = u_1 x_1 + \dots + u_n x_n$, with $|u| = (u, u)^{1/2}$ and $u + \alpha x$ is the point $(u_1 + \alpha x_1, \dots, u_n + \alpha x_n)$.

Previously the author [13], using equisummability between trigonometric integrals and trigonometric series, has obtained in the special case of double trigonometric integrals the following result:

Let $c(u)$, in L_2 on any bounded domain, be $O(|u|^{-\epsilon})$, $\epsilon > 0$. Suppose the double trigonometric integral $\int_{E_2} e^{i(x, u)} c(u) du$ is circularly summable $(C, 1)$ to $f(x)$. Furthermore suppose $f(x)$ is in $\text{Lip } \alpha$, $\alpha > 0$, on every bounded domain (α depending on the domain). Then the double trigonometric integral

$$\frac{1}{4\pi^2} \int_{E_2} e^{-i(x, u)} f(x) dx$$

is spherically summable $(C, 1)$ to $c(u)$ for almost every u .

Specializing $f(x)$ to be the zero function (which is what is meant by Cantor-type uniqueness, [15, p. 274]) and using a more direct attack on the problem, we are able in this paper both to weaken the hypotheses of the above theorem as well as to extend the results to n -dimensional integrals.

2. Definitions and notation. The open n -dimensional sphere with center x and radius r will be designated by $D_n(x, r)$, and the surface of the sphere by $C_n(x, r)$.

Received November 26, 1953. Presented to the American Mathematical Society November 27, 1953. This investigation was supported in part by a grant from the Rutgers University Research Fund.

Pacific J. Math. 5 (1955), 607-622

Following Bochner [1], we shall say that the multiple trigonometric integral $\int_{E_n} c(u) e^{i(x,u)} du$ is spherically convergent at the point x to the finite value $L(x)$ if the spherical partial integrals of rank R converge to $L(x)$, that is if

$$(1) \quad I_R(x) = \int_{D_n(0,R)} e^{i(x,u)} c(u) du \longrightarrow L(x) \quad (\text{as } R \longrightarrow \infty.)$$

The integral

$$(2) \quad \sigma_R^{(\alpha)}(x) = 2\alpha R^{-2\alpha} \int_0^R I_r(x) (R^2 - r^2)^{\alpha-1} r dr, \quad \alpha > 0,$$

is called the (C, α) -mean of rank R of the multiple trigonometric integral $\int_{E_n} c(u) e^{i(x,u)} du$, and this integral is said to be spherically summable (C, α) to $L(x)$ if $\sigma_R^{(\alpha)}(x) \longrightarrow L(x)$ as $R \longrightarrow \infty$.

Given $F(x)$ integrable on $D_n(x_0, r)$, we designate the mean value of F in this sphere by $A(F; x_0; r)$. Given $F(x)$ integrable on $C_n(x_0; r)$, we designate the mean value of F on this surface by $L(F; x_0; r)$. Thus, designating the volume of the unit n -dimensional sphere, $2\pi^{n/2}/n\Gamma(n/2)$, by Ω_n and the $(n-1)$ -dimensional volume of its surface, $2\pi^{n/2}/\Gamma(n/2)$, by ω_n , we have

$$(3) \quad \begin{aligned} A(F; x_0; r) &= (\Omega_n r^n)^{-1} \int_{D_n(x_0, r)} F(x) dx \\ L(F; x_0; r) &= \omega_n^{-1} \int_{C_n(0,1)} F(x_0 + rx) dS_{n-1}(x) \end{aligned}$$

where dS_{n-1} is the $(n-1)$ dimensional volume element of $C_n(0, 1)$.

We set

$$\nabla_1(F; x_0; r) = L(F; x_0; r) - F(x_0) \quad \text{and} \quad \nabla_2(F; x_0; r) = A(F; x_0; r) - F(x_0)$$

and say that $F(x)$ has a generalized Laplacian of the first or second kind at the point x_0 equal to α_1 or α_2 , respectively, if

$$\lim_{r \rightarrow 0} 2n \nabla_1(F; x_0; r)/r^2 = \alpha_1$$

or

$$\lim_{r \rightarrow 0} 2(n+2) \nabla_2(F; x_0; r)/r^2 = \alpha_2.$$

The generalized Laplacian of the first and second kind of F at x_0 will be designated by $\Delta_1 F(x_0)$ and $\Delta_2 F(x_0)$, respectively. It is known, [6, p. 261], that if $F(x)$ is in class $C^{(2)}$ on $D_n(x_0, r_0)$, then $\Delta F(x_0) = \Delta_1 F(x_0) = \Delta_2 F(x_0)$ where $\Delta F(x)$ is the ordinary Laplacian of F at x .

The closure of the set W is designated by \bar{W} ; and its characteristic function by $\chi_W(x)$. The set Z is said to be a closed set of vanishing capacity if for every r , $Z\bar{D}_n(0, r)$ is a closed set of capacity zero. It is known, [4], that if Z is a set of vanishing capacity then $D_n(x_0, r) - ZD_n(x_0, r)$ is a domain.

The trigonometric integral $\int_{E_n} e^{i(x,u)} c(u) du$ is said to be of type (U) on a domain G if

$$\int_{E_n - D_n(0,1)} e^{i(x,u)} c(u) |u|^{-2} du$$

converges spherically on G to a function $F(x)$ which is continuous on G .

Throughout this paper E_n stands for n -dimensional euclidean space where $n \geq 2$, and $\mu = (n - 2)/2$.

The function $J_i(r)$ is the Bessel function of the first kind of order i .

3. Statement of main results. We shall prove the following two theorems concerning Cantor-type uniqueness for multiple trigonometric integrals.

THEOREM 1. *Given the multiple trigonometric integral $\int_{E_n} e^{i(x,u)} c(u) du$ where $c(u)$ is a complex-valued function which is integrable on every bounded domain. Let Z be a closed set of vanishing capacity. Suppose that*

- (i) *The integral is spherically summable $(C, 1)$ to zero almost everywhere.*
- (ii) *The $(C, 1)$ spherical mean of rank R , $\sigma_R^{(1)}(x)$, is such that $\overline{\lim}_{R \rightarrow \infty} |\sigma_R^{(1)}(x)| < \infty$ in $E_n - Z$.*
- (iii) *$c(u)(|u|^2 + 1)^{-1}$ is in L_1 on E_n .*

Then $c(u)$ vanishes almost everywhere.

THEOREM 2. *Given the multiple trigonometric integral $\int_{E_n} e^{i(x,u)} c(u) du$ where $c(u)$ is a complex-valued function which is integrable on every bounded domain. Let Z be a closed set of vanishing capacity. Suppose that*

- (i) and (ii) *The same as (i) and (ii) of Theorem 1.*
- (iii) *The integral is of type (U) on E_n .*
- (iv) *$c(u)(|u|^2 + 1)^{-1}$ is in L_2 on E_n .*

Then $c(u)$ vanishes almost everywhere.

For the special case of the plane, we prove the following theorem.

THEOREM 3. *Given the double trigonometric integral $\int_{E_n} e^{i(x,u)} c(u) du$ where $c(u)$ is a complex-valued function which is integrable on every bounded domain. Let Z be a closed set of vanishing capacity and W be a closed denumerable set such that $WZ = 0$. Suppose that*

- (i) *The integral is spherically summable $(C, 1)$ to zero in $E_2 - Z$.*
- (ii) *The integral is of type (U) on $E_2 - W$.*
- (iii) *$c(u) = o(|u|)$ as $|u| \rightarrow \infty$*
- (iv) *$c(u)(|u|^2 + 1)^{-1}$ is in L_2 on E_n .*

Then $c(u)$ vanishes almost everywhere.

4. Fundamental lemmas. Before proving the main theorems of this paper, it is first necessary to establish a connection between the $(C, 1)$ spherical summability of the integral $\int_{E_n} e^{i(x,u)} c(u) du$ and the generalized Laplacians of the "anti-Laplacian" of this integral. In short, we shall now establish some Riemann-type, [15, p. 270], results for the multiple trigonometric integrals.

We need prove the following lemma only for the plane, since the conclusion is hypothesized for Theorems 1 and 2.

LEMMA 1. *Let $c(u)$ be a complex-valued function which is integrable on every bounded domain in the plane, vanishes in $D_2(0, r_0)$, $r_0 > 0$, and is $o(|u|)$. Suppose that $\sigma_R^{(1)}(x_0) = o(R)$ where $\sigma_R^{(1)}(x)$ is the $(C, 1)$ spherical mean of rank R of $\int_{E_2} e^{i(x,u)} c(u) du$. Then $\int_{E_2} e^{i(x,u)} c(u) |u|^{-2} du$ is spherically convergent.*

Without loss of generality, we assume x_0 to be the origin. Then with $I_R(x)$ given by (1) and $\sigma_R^{(1)}(x)$ by (2), we have

$$\begin{aligned} \int_{D_2(0,R)} c(u) |u|^{-2} du &= 2 \int_{D_2(0,R)} c(u) du \int_0^R \chi_{D_2(0,r)}(u) r^{-3} dr + R^{-2} I_R(0) \\ &= 2 \int_0^R r^{-3} dr \int_{D_2(0,R)} c(u) \chi_{D_2(0,r)}(u) du + R^{-2} I_R(0) \\ &= 2 \int_0^R r^{-3} I_r(0) dr + R^{-2} I_R(0) \\ &= 4 \int_0^R \sigma_r^{(1)}(0) r^{-3} dr + \sigma_R^{(1)}(0) R^{-2} + R^{-2} I_R(0). \end{aligned}$$

Since by assumption $\sigma_R^{(1)}(0) = o(R)$, to prove the lemma it only remains to show that $I_R(0) = o(R^2)$. But

$$I_R(0) = [(R + 1)^2 \sigma_{R+1}^{(1)}(0) - R^2 \sigma_R^{(1)}(0)](2R + 1)^{-1} - (2R + 1)^{-1} \int_{D_2(0, R+1) - D_2(0, R)} c(u) [R+1)^2 - |u|^2] du = o(R^2),$$

and the proof is complete.

LEMMA 2. Let $c(u)$ be a complex-valued function which is integrable on every bounded domain in E_n and which vanishes in $D_n(0, r_0)$, $r_0 > 0$. Suppose that

(i) $\overline{\lim}_{R \rightarrow \infty} |\sigma_R^{(1)}(x_0)| = d$ where $\sigma_R^{(1)}(x_0)$ is the $(C, 1)$ spherical mean of rank R of $\int_{E_n} e^{i(x,u)} c(u) du$ and d is finite-valued.

(ii) $-\int_{E_n} e^{i(x,u)} c(u) |u|^{-2} du$

is spherically uniformly convergent in $D_n(x_0, t_0)$, $t_0 > 0$, to $F(x)$.

Then $\overline{\lim}_{t \rightarrow 0} |2n \nabla_1(F; x_0; t)/t^2| \leq Kd$ where K is a constant independent of x_0 and d .

Observing that for fixed u

$$L(e^{i(x,u)}; x_0; t) = 2^\mu \Gamma(\mu + 1) e^{i(u, x_0)} (|u|t)^{-\mu} J_\mu(|u|t)$$

(see [1, p. 177]), we have by assumption (ii) for t sufficiently small that

$$L(F; x_0; t)$$

$$= -2^\mu \Gamma(\mu + 1) \lim_{R \rightarrow \infty} \int_{D_n(0, R)} e^{i(x_0, u)} c(u) |u|^{-2} J_\mu(|u|t) (|u|t)^{-\mu} du$$

and consequently that

$$(4) \quad (2n) \nabla_1(F; x_0; t)/t^2 = \lim_{R \rightarrow \infty} \int_{D_n(0, R)} e^{i(x_0, u)} c(u) \eta(|u|t) du$$

where

$$\eta(r) = 4(\mu + 1) [1 - 2^\mu \Gamma(\mu + 1) J_\mu(r) r^{-\mu}] / r^2 \text{ for } r > 0, \eta(0) = 1,$$

and $\eta(r)$ is in $C^{(\infty)}$.

Making the following observations:

(a) By the second mean-value theorem applied to the real and imaginary parts of $I_R(x_0)$ given by (1) and hypothesis (ii) we have $I_R(x_0) = o(R^2)$,

(b) For fixed t , $\eta(Rt) = O(R^{-2})$ and $\eta'(Rt) = O(R^{-5/2})$ where $\eta'(r) = d\eta(r)/dr$, we obtain from (4) that

$$(5) \quad 2n\nabla_1(F; x_0; t)/t^2 = 2^{-1} \int_0^\infty r^2 \sigma_r^{(1)}(x_0) t^3 \alpha(rt) dr$$

where $\alpha(r) = dr^{-1}\eta'(r)/dr$.

From the fact that $\alpha(z)$ is an entire function of the form $\sum_{i=0}^\infty b_i z^{2i+1}$, we have that there exists a constant K_1 such that

$$(6) \quad |\alpha(r)| \leq K_1 r \text{ for } r \leq 1$$

From the fact that $J_\mu(r) = O(r^{-1/2})$ as $r \rightarrow \infty$, and

$$dr^{-\mu} J_\mu(r)/dr = -r^{-\mu} J_{\mu+1}(r),$$

we obtain that there exists a constant K_2 such that

$$(7) \quad |\alpha(r)| \leq K_2 [r^{-(\mu+7/2)} + r^{-5}] \quad \text{for } r \geq 1$$

From (5), (6), and (7), the conclusion of the lemma follows readily. For given an $\epsilon > 0$, choose R_0 so large that $|\sigma_R^{(1)}(x_0)| \leq d + \epsilon$ for $R > R_0$. Then for $t < R_0^{-1}$, it is easily seen that

$$(8) \quad |2n\nabla_1(F; x_0, t)/t^2| \leq K(d + \epsilon) + O(t^4)$$

where K is a constant independent of x_0 , d , and ϵ . Taking the limit superior of the left side of (8) as $t \rightarrow 0$ and then the limit of the right side as $\epsilon \rightarrow 0$, we have the proof of the lemma.

LEMMA 3. *Let the hypotheses be the same as in Lemma 2 except that $\lim_{R \rightarrow \infty} \sigma_R^{(1)}(x_0) = d$. Then $\Delta_1 F(x_0) = d$.*

For if $d = 0$, the lemma follows immediately from Lemma 2. If $d \neq 0$, choose $c_1(u)$ integrable on E_n , vanishing for u in $[E_n - D_n(0, 2)] + D_n(0, 1)$, and such that $\int_{E_n} c_1(u) e^{i(x_0, u)} = d$. Set $F_1(x) = -\int_{E_n} c_1(u) |u|^{-2} e^{i(x, u)} du$. Then

$$0 = \Delta_1 F(x_0) - \Delta_1 F_1(x_0) = \Delta_1 F(x_0) - \Delta F_1(x_0) = \Delta_1 F(x_0) - d.$$

LEMMA 4. Let $c(u)$ be a complex-valued function which is integrable on every bounded domain in E_n and which vanishes in $D_n(0, r_0)$, $r_0 > 0$. Suppose that

(i) $\sigma_R^{(1)}(x)$, the $(C, 1)$ spherical mean of rank R of $\int_{E_n} e^{i(x,u)} c(u) du$, is such that $\lim_{R \rightarrow \infty} |\sigma_R^{(1)}(x_0)| = d$.

(ii) $c(u) |u|^{-2}$ is in L_2 on E_n .

(iii) $-\int_{E_n} e^{i(x_0, u)} |u|^{-2} c(u) du$

is spherically convergent to $F(x_0)$. Set

$$F(x) = -\text{l.i.m.}_{R \rightarrow \infty} \int_{D_n(0, R)} e^{i(x, u)} c(u) |u|^{-2} du.$$

Then

$$\overline{\lim}_{t \rightarrow 0} |2(n+1) \nabla_2(F; x_0; t)/t^2| \leq Kd$$

where K is a constant independent of x_0 and d .

Setting

$$T_R(x) = -\int_{D_n(0, R)} e^{i(x, u)} c(u) |u|^{-2} du,$$

observing that $A(F; x_0; t) = \lim_{R \rightarrow \infty} A(T_R; x_0; t)$ and that for fixed u ,

$$\begin{aligned} A(e^{i(x, u)}; x_0; t) &= \omega_n / \Omega_n t^n \int_0^t r^{n-1} L(e^{i(x, u)}; x_0; r) dr \\ &= 2^{\mu+1} \Gamma(\mu+2) (|u|t)^{-(\mu+1)} J_{\mu+1}(|u|t) e^{i(x_0, u)}, \end{aligned}$$

we obtain

$$(9) \quad A(F; x_0; t)$$

$$= -\lim_{R \rightarrow \infty} 2^{\mu+1} \Gamma(\mu+2) \int_{D_n(0, R)} e^{i(x_0, u)} c(u) |u|^{-2} (|u|t)^{-(\mu+1)} J_{\mu+1}(|u|t) du$$

and consequently

$$2(n + 2)\nabla_2(F; x_0; t)/t^2 = \lim_{R \rightarrow \infty} \int_{D_n(0, R)} e^{i(x_0, u)} \gamma(|u|t) c(u) du$$

where

$$\gamma(r) = 4(\mu + 2)[1 - 2^{(\mu+1)}\Gamma(\mu + 2)r^{-(\mu+1)}J_{\mu+1}(r)]/r^2 \text{ for } r > 0, \gamma(0) = 1,$$

and $\gamma(r)$ is in $C^{(\infty)}$.

Since $\gamma(r)$ has the same form as $\eta(r)$ in Lemma 2 with μ replaced by $\mu + 1$, we can proceed as in that lemma and obtain

$$2(n + 2)\nabla_2(F; x_0; t)/t^2 = 2^{-1} \int_0^\infty t^3 r^2 \sigma_r^{(1)}(x_0) \beta(rt) dr$$

where $\beta(r) = dr^{-1} \gamma'(r)/dr$. Then we can proceed in a similar manner to obtain that for $\epsilon > 0$

$$\overline{\lim}_{t \rightarrow \infty} |2(n + 2)\nabla_2(F; x_0; t)/t^2| \leq K(d + \epsilon)$$

where K is a constant independent of x_0, d , and ϵ . Since ϵ is arbitrary the conclusion of the lemma follows.

LEMMA 5. *Let the hypotheses be the same as in Lemma 4 except that $\lim_{R \rightarrow \infty} \sigma_R^{(1)}(x_0) = d$. Then $\Delta_2 F(x_0) = d$.*

In the same manner that we obtained Lemma 3 from Lemma 2, we obtain Lemma 5 from Lemma 4.

LEMMA 6. *Let $F(x)$ be real-valued and continuous on $\overline{D}_n(x_0, r_0)$, $r_0 > 0$. Suppose that*

- (i) $\Delta_2 F(x) = 0$ almost everywhere in $D_n(x_0, r_0)$
- (ii) $\overline{\lim}_{r \rightarrow \infty} |2(n + 2)\nabla_2(F; x; r)/r^2| < \infty$ for all x in $D_n(x_0, r_0)$.

Then $F(x)$ is harmonic in $D_n(x_0, r_0)$.

Following the pattern of proof in [9], we give a proof for $n \geq 3$.

To prove the lemma, it is sufficient to show that $F(x)$ is subharmonic in $D_n(x_0, r_0)$.

Set

$$f(x) = 2(n + 2) \left[\overline{\lim}_{r \rightarrow 0} \nabla_2(F; x; r)/r^2 + \underline{\lim}_{r \rightarrow 0} \nabla_2(F; x; r)/r^2 \right] / 2$$

for x in $D_n(x_0, r_0)$. Then $f(x) = 0$ almost everywhere in $D_n(x_0; r_0)$.

By the theorem of Vitali-Caratheodory [10, p. 75], there exists a sequence of nondecreasing upper semicontinuous functions $\{g_m(x)\}$ such that $g_m(x) \leq f(x)$ for all x in $D_n(x_0, r_0)$, $g_m(x) \rightarrow f(x)$ almost everywhere in $D_n(x_0, r_0)$, $g_m(x)$ is integrable on $D_n(x_0, r_0)$, and such that

$$\lim_{m \rightarrow \infty} \int_{D_n(x_0, r)} g_m(x) dx = \int_{D_n(x_0, r)} f(x) dx \quad \text{for } r \leq r_0.$$

Set

$$\Delta^{-1} g_m(x) = -[\omega_n(n - 2)]^{-1} \int_{D_n(x_0, r_0)} g_m(u) |u - x|^{2-n} du.$$

Then $\Delta^{-1} g_m(x)$ is superharmonic, since $g_m(u) \leq 0$ for almost all u in $D_n(x_0, r_0)$. Furthermore, we observe that for fixed u

$$\begin{aligned} A(|x - u|^{2-n}; x_0; r) &= |x_0 - u|^{2n} \text{ if } |x_0 - u| > r \\ &= nr^{-n} 2^{-1} [r^2 + |x_0 - u|^2 (2 - n)n^{-1}] \text{ if } |x_0 - u| \leq r. \end{aligned}$$

Consequently, for x_1 in $D_n(x_0, r)$ with r sufficiently small,

$$\begin{aligned} (11) \quad \nabla_2(\Delta^{-1} g_m; x_1, r) &= [\omega_n(n - 2)]^{-1} \int_{D_n(x_1, r)} g_m(u) \{ |u - x_1|^{2-n} \\ &\quad - nr^{-n} 2^{-1} [r^2 + |x_1 - u|^2 (2 - n)n^{-1}] \} du. \end{aligned}$$

Suppose $g_m(x_1)$ is finite. Then by the upper semi-continuity of $g_m(u)$ at x_1 , for $\epsilon > 0$ and r sufficiently small, we have from (11) that

$$\nabla_2(\Delta^{-1} g_m; x_1; r) \leq [g_m(x_1) + \epsilon] [\omega_n(n - 2)]^{-1} [\omega_n(n - 2)] r^2 / 2(n + 2).$$

Consequently, we conclude that

$$(12) \quad \overline{\lim}_{r \rightarrow 0} 2(n + 2) \nabla_2(\Delta^{-1} g_m; x_1; r) / r^2 \leq g_m(x_1).$$

Similarly, in case $g_m(x_1) = -\infty$, conclusion (12) remains valid.

From the fact that $\Delta^{-1}g_m(x)$ is superharmonic, we have that $F - \Delta^{-1}g_m$ is upper semi-continuous in $D_n(x_0, r_0)$. From (12) we conclude that

$$\overline{\lim}_{r \rightarrow 0} 2(n+2)\nabla_2(F - \Delta^{-1}g_m; x; r)/r^2 \geq 0 \quad \text{for } x \text{ in } D_n(x_0, r_0).$$

Therefore by [8, p. 14], $\{F - \Delta^{-1}g_m\}$ is a nondecreasing sequence of subharmonic functions in $D_n(x_0, r_0)$. But $\lim_{m \rightarrow \infty} \Delta^{-1}g_m(x) = 0$ almost everywhere. Therefore $F(x)$ is almost everywhere equal to a subharmonic function, $G(x)$, by [8, p. 22]. But $A(F; x; r) = A(G; x; r) \rightarrow G(x)$ for all x in $D_n(x_0, r_0)$. However from the continuity of F we have $A(F; x; r) \rightarrow F(x)$, and the proof of the lemma is complete for $n \geq 3$. For $n = 2$ a similar proof can be given with the Newtonian potential replaced by the logarithmic potential.

For the case of the generalized Laplacian of the first kind, we have a similar lemma with a similar proof, see [9].

LEMMA 7. *Let $F(x)$ be real-valued and continuous on $\overline{D}_n(x_0, r_0)$, $r_0 > 0$. Suppose that*

- (i) $\Delta_1 F(x) = 0$ almost everywhere in $D_n(x_0, r_0)$.
- (ii) $\overline{\lim}_{r \rightarrow 0} |2n\nabla_1(F; x; r)/r^2| < \infty$ for all x in $D_n(x_0, r_0)$.

Then $F(x)$ is harmonic in $D_n(x_0, r_0)$.

We now prove some lemmas concerning the spherical summability (C, n) of Fourier transforms.

LEMMA 8. *Let $G(x)$ be a function in L_1 on E_n which vanishes in $D_n(0, r_0)$, $r_0 > 0$. Suppose that $F(x) = \int_{E_n} e^{i(x,u)} G(u) du$ is in $C^{(2)}$ on E_n . Then for u in $D_n(o, r_0/2) - 0$*

$$(13) \quad \int_{E_n} [e^{-i(x,u)} F(x) - (-e^{-i(x,u)} |u|^{-2} \Delta F(x))] dx$$

is spherically summable (C, n) to zero.

For, by Green's second identity, we have

$$(14) \quad \begin{aligned} I_R(u) &= \int_{D_n(0,R)} [e^{-i(x,u)} F(x) - (-e^{-i(x,u)} |u|^{-2} \Delta F(x))] dx \\ &= |u|^{-2} \left\{ R^{n-1} \int_{C_n(0,1)} F(Rx) i(x,u) e^{-iR(x,u)} dS_{n-1}(x) \right\} \end{aligned}$$

$$+ R^{n-1} \int_{C_n(0,1)} \frac{\partial F(Rx)/\partial R}{e^{-iR(x,u)}} dS_{n-1}(x) \Big\} \equiv |u|^{-2} (A_R + B_R)$$

where $dS_{n-1}(x)$ is the $(n - 1)$ dimensional volume element on the unit sphere $C_n(0, 1)$.

We shall now show that both A_R and B_R are (C, n) summable to zero. For, by Fubini's theorem, we have

$$(15) \quad (MR^2)^{-1} \int_0^\infty r \phi_n\left(\frac{r}{R}\right) A_r dr \\ = (MR^2)^{-1} i \int_{E_n - D_n(0, r_0)} G(y) \left[\int_{E_n} \phi_n\left(\frac{|x|}{R}\right) e^{-i(x, u-y)}(x, u) dx \right] dy$$

where $M = (2\pi)^{n/2}/2^{n-1}(n - 1)!$ and

$$\phi_1(r) = \begin{cases} 1 & \text{for } 0 \leq r \leq 1 \\ 0 & \text{for } r > 1 \end{cases} \quad \phi_n(r) = \begin{cases} (1 - r^2)^{n-1} & \text{for } 0 \leq r \leq 1 \\ 0 & \text{for } r > 1 \end{cases} .$$

Since for fixed $u \neq 0$, (x, u) is a homogeneous polynomial which is also a harmonic function in x , we have by [2, p. 806] and [14, p. 373] that the right side of (15) is equal to

$$(16) \quad -R^{-1} \int_{E_n - D_n(0, r_0)} G(y) \frac{(y - u, u)}{|y - u|^{n+1}} \frac{J_{n+\mu+1}(R|y - u|) dy}{(R|y - u|)^{n-\mu-2}} .$$

Clearly (16) tends to zero as $R \rightarrow \infty$; so A_R is (C, n) summable to zero for u in $D_n(0, r_0/2) - 0$.

We also observe after integrating by parts that

$$(17) \quad (MR^2)^{-1} \int_0^R r \phi_n\left(\frac{r}{R}\right) B_r dr \\ = (MR^2)^{-1} \int_0^R dr \int_{C_n(0,1)} F(rx) \frac{\partial}{\partial r} \left[r^n \phi_n\left(\frac{r}{R}\right) e^{-ir(x,u)} \right] dS_{n-1}(x) .$$

From the above discussion concerning A_R and from [1, Theorem 1], to show that B_R is (C, n) summable to zero for u in $D_n(0, r_0/2) - 0$, it is sufficient to show that

$$(18) \quad (MR^2)^{-1} \int_{E_n} F(x) \phi_{n-1} \left(\frac{|x|}{R} \right) \frac{|x|^2}{R^2} e^{-i(x,u)} dx \rightarrow 0 \text{ as } R \rightarrow \infty.$$

But by [2, p. 806] and [14, p. 373] the expression in (18) is equal to

$$(19) \quad R^{-1} \int_{E_n - D_n(0, r_0)} \frac{G(y)}{|y-u|^{n-1}} \left[K_1 \frac{J_{n+\mu}(R|y-u|)}{(R|y-u|)^{n-\mu-1}} - K_2 \frac{J_{n+\mu-1}(R|y-u|)}{(R|y-u|)^{n-\mu-2}} \right] dy.$$

where K_1 and K_2 are two constants depending on n .

Clearly (19) tends to zero as $R \rightarrow \infty$ for u in $D_n(0, r_0/2) - 0$; so B_R is (C, n) summable to zero and the lemma is proved.

LEMMA 9. Let $G(x)$ be a function in L_2 on E_n which vanishes in $D_n(0, r_0)$, $r_0 > 0$. Suppose that $\int_{E_n} e^{i(x,u)} G(x) dx$ is spherically convergent to a function $F(x)$ which is in $C^{(2)}$ on E_n . Then for u in $D_n(0, r_0/2) - 0$

$$\int_{E_n} [e^{-i(x,u)} F(x) - (-e^{-i(x,u)} |u|^{-2}) \Delta F(x)] dx$$

is spherically summable (C, n) to zero.

For (14) also holds in this case, and as in Lemma 8, we have to show that both A_R and B_R are (C, n) summable to zero.

Since both $F(x)$ and $\phi_n(|x|/R)(x, u)$ are in L_2 on E_n , Parseval's formula gives us both (15) and (16). We therefore conclude as before that A_R is (C, n) summable to zero for u in $D_n(0, r_0/2) - 0$.

To show that B_R is summable (C, n) to zero, we obtain (17) as in Lemma 8. Then from the fact that A_R is (C, n) summable to zero and from [3, Theorem 55], it is sufficient once again to show that (18) holds.

But by Parseval's formula, we obtain that the expression in (18) is equal to (19). Observing that for u in $D_n(0, r_0/2) - 0$ and for y in $E_n - D_n(0, r_0)$ there exists a constant \bar{K}_n such that

$$|J_{n+\mu-1}(R|y-u|)| \leq \bar{K}_n (R|y-u|)^{-1/2} \quad \text{for } R > 1$$

and that for such u , $|y-u|^{1/2-n}$ is in L_2 on $E_n - D_n(0, r_0)$, we conclude that

(18) holds and consequently that B_R is (C, n) summable to zero, which proves the lemma.

5. Proof of Theorem 1. To prove Theorem 1, it is sufficient to show that for any $r_0 > 0$, $c(u) = 0$ almost everywhere in $D_n(0, r_0/2)$. Set

$$F_1(x) = - \int_{D_n(0, r_0)} \frac{[e^{i(x, u)} - 1 - i(x, u)]}{|u|^2} c(u) du$$

Then, $F_1(x)$ is in $C^{(\infty)}$ on E_n and

$$\Delta F_1(x) = \int_{D_n(0, r_0)} e^{i(x, u)} c(u) du.$$

Set

$$F_2(x) = - \int_{E_n - D_n(0, r_0)} e^{i(x, u)} \frac{c(u)}{|u|^2} du$$

which is by (iii) continuous in E_n . Then by Lemma 2 and (ii),

$$\overline{\lim}_{r \rightarrow 0} |2n \nabla_1(F_2; x; r)/r^2| < \infty$$

in $E_n - Z$ and by Lemma 3 and (i), $\Delta_1 F_2(x) = -\Delta F_1(x)$ almost everywhere.

Set $F(x) = F_1(x) + F_2(x)$. Take any x_0 in E_n and consider $D_n(x_0, r_1)$, $r_1 > 0$. From the definition of a closed set of vanishing capacity, we see that there is a closed bounded set of capacity zero Z_1 such that

$$\overline{\lim}_{r \rightarrow 0} |2n \nabla_1(F; x; r)/r^2| \leq |\Delta F_1(x)| + \overline{\lim}_{r \rightarrow 0} |2n \nabla_1(F_2; x; r)/r^2| < \infty$$

for x in the domain $G = D_n(x_0, r_1) - Z_1 D_n(x, r_1)$. Furthermore almost everywhere in G , $\Delta_1 F(x) = \Delta F_1(x) + \Delta_1 F_2(x) = 0$. Consequently it follows from Lemma 7, that $F(x)$ is harmonic in the domain $G = D_n(x_0, r_1) - Z_1 D_n(x_0, r_1)$. But $F(x)$ is continuous in $D_n(x_0, r_1)$. Therefore by [7, p. 335], $F(x)$ is harmonic in $D_n(x_0, r_1)$ and since x_0 is arbitrary, $F(x)$ is harmonic in E_n .

From the fact that $F(x)$ is harmonic in E_n , we now have that $F_2(x) = F(x) - F_1(x)$ is in $C^{(\infty)}$ on E_n and that $\Delta F_2(x) = -\Delta F_1(x)$ for all x . Also by [1, Theorem 1] we obtain that

$$(2\pi)^{-n} \int_{E_n} e^{-i(x,u)} F_2(x) dx$$

is spherically summable (C, n) to zero for u in $D_n(0, r_0/2) - 0$. Therefore by Lemma 8 for such u ,

$$\int_{E_n} e^{-i(x,u)} [-\Delta F_2(x)] dx$$

is spherically summable (C, n) to zero. But for almost all such u , we have that

$$(2\pi)^{-n} \int_{E_n} e^{-i(x,u)} \Delta F_1(x) dx$$

is spherically summable (C, n) to $c(u)$. Since $\Delta F_1(x) = -\Delta F_2(x)$, we conclude that for almost all u in $D_n(0, r_0/2)$, $c(u) = 0$, which proves the theorem.

6. Proof of Theorem 2. The proof is quite similar to that of Theorem 1. Once again it is sufficient to prove that for any $r_0 > 0$, $c(u) = 0$ almost everywhere in $D_n(0, r_0/2)$.

Set

$$F_1(x) = - \int_{D_n(0, r_0)} [e^{i(x,u)} - 1 - i(x, u)] \frac{c(u)}{|u|^2} du,$$

and

$$F_2(x) = - \lim_{R \rightarrow \infty} \int_{D_n(0, r_R) - D_n(0, r_0)} e^{i(x,u)} \frac{c(u)}{|u|^2} du.$$

By (iii), $F_2(x)$ is continuous. Then in a manner exactly analogous to the proof of Theorem 1 except that Lemmas 4, 5, and 6 are used instead of 2, 3, and 7, we obtain that $F_2(x)$ is in $C^{(\infty)}$ and that $\Delta F_2(x) = -\Delta F_1(x)$. By Lemma 9 and [3, Theorem 55], we obtain that $\int_{E_n} e^{-i(x,u)} [-\Delta F_2(x)] dx$ is spherically summable (C, n) to zero for u in $D_n(0, r_0/2) - 0$. But by [1, Theorem 1] for almost all such u , we have that

$$(2\pi)^{-n} \int_{E_n} e^{-i(x,u)} \Delta F_1(x)$$

is spherically summable (C, n) to $c(u)$. Since $-\Delta F_2(x) = \Delta F_1(x)$, we conclude that $c(u) = 0$ almost everywhere in $D_n(0, r_0/2)$ and the theorem is proved.

7. Proof of Theorem 3. Let $F_1(x)$ be as in Theorem 2 with n replaced by 2, and let

$$F_2(x) = - \lim_{R \rightarrow \infty} \int_{D_2(0,R) - D_2(0,r_0)} \frac{c(u)}{|u|^2} e^{i(x,u)}$$

where $r_0 > 0$. This limit exists for x in Z by (ii) and for x not in Z by (i), (iii), and Lemma 1. Furthermore by (ii) $F_2(x)$ is assumed continuous in $E_2 - W$. It is clear from the proof of Theorem 2 that to prove this theorem we need only show that $F_2(x)$ is continuous in E_2 or what is the same thing that $F(x) = F_1(x) + F_2(x)$ is continuous in E_2 .

By (ii) $F(x)$ is continuous in $E_2 - W$, and by Lemmas 5, $\Delta_2 F(x) = 0$ in $E_2 - Z$. Let $D_2(x_0, r_1)$ be any disc which has a null intersection with W . Then as in the proof to Theorem 1, $F(x)$ is harmonic in this disc and consequently in $E_2 - W$. We also observe that now $\Delta_2 F(x) = 0$ in the whole plane and furthermore that $F(x)$ is in L_2 on any bounded domain.

Let W_1 be the set of discontinuity points of $F(x)$ and let x_0 be an isolated point of W_1 . Then there is a closed disc $\bar{D}_2(x_0, r_2)$ whose intersection with W_1 is x_0 . Then by the above discussion we have that $F(x)$ is in L_2 on $D_2(x_0, r_2)$, harmonic in $D_2(x_0, r_2) - x_0$, and satisfies the further condition that $\Delta_2 F(x_0) = 0$. Consequently by [12, Lemma 4], $F(x)$ is then harmonic in the whole disc and, a fortiori, continuous at x_0 .

Therefore W_1 , has no isolated points and \bar{W}_1 is a perfect set. But $\bar{W}_1 \subset W$ is at most denumerable, and by [10, p. 55], \bar{W}_1 is then the empty set. Thus $F(x)$ is continuous in the whole plane, and, as mentioned above, the proof of this theorem is reduced to that of Theorem 2.

8. Appendix. In closing we point out that the assumption W and Z have a null intersection in Theorem 3 is a necessary one. For consider the double trigonometric integral $\int_{E_2} c(u) e^{i(x,u)} du$ with $c(u) = 1$. (iii) and (iv) of Theorem 3 are clearly satisfied. Observing that the spherical mean of rank R , $\sigma_R^{(1)}(x)$, with $x \neq 0$ is given by

$$\sigma_R^{(1)}(x) = 4\pi J_2(|x|R) |x|^{-2} = O(R^{-1/2}),$$

we see that (i) is satisfied with Z equal to the origin. Furthermore, we observe that for $x \neq 0$

$$\lim_{R \rightarrow \infty} \int_{D_2(0,R) - D_2(0,1)} |u|^{-2} e^{i(x,u)} du = 2\pi \int_{|x|}^{\infty} J_0(r) r^{-1} dr.$$

Consequently (ii) is satisfied with W consisting of the origin. But W and Z do not have a null intersection, and the conclusion of Theorem 3 does not hold.

REFERENCES

1. S. Bochner, *Summation of multiple Fourier series by spherical means*, Trans. Amer. Math. Soc. **40** (1936), 175-207.
2. ———, *Theta relations with spherical harmonics*, Proc. Nat. Acad. Sci., **37** (1951), 804-808.
3. S. Bochner and K. Chandrasekharan, *Fourier transforms*, Princeton, 1949.
4. M. Brelot, *Sur la structure des ensembles de capacité nulle*, C. R. Acad. Sci. Paris **192** (1931), 206-208.
5. M. T. Cheng, *Uniqueness of multiple trigonometric series*, Annals of Math. **52** (1950), 403-416.
6. R. Courant and D. Hilbert, *Methoden der mathematischen Physik*, vol. 2 Berlin, 1937.
7. O. D. Kellogg, *Foundation of potential theory*, Berlin, 1929.
8. T. Rado, *Subharmonic functions*, Ergebnisse der Mathematik, vol. 5, no. 1, Berlin, 1937.
9. W. Rudin, *Integral representations of continuous functions*, Trans. Amer. Math. Soc. **68** (1950), 278-286.
10. S. Saks, *Theory of the integral*, 2d. ed., Warsaw, 1937.
11. V. L. Shapiro, *An extension of results in the uniqueness theory of double trigonometric series*, Duke Math. J. **20** (1953), 359-366.
12. ———, *A note on the uniqueness of double trigonometric series*, Proc. Amer. Math. Soc. **4** (1953), 692-695.
13. ———, *Summability and uniqueness of double trigonometric integrals*, Trans. Amer. Math. Soc., **77** (1954), 322-339.
14. G. N. Watson, *A treatise on the theory of Bessel functions*, Cambridge, 1944.
15. A. Zygmund, *Trigonometrical series*, Warsaw, 1935.

RUTGERS UNIVERSITY, NEW BRUNSWICK, NEW JERSEY AND
THE INSTITUTE FOR ADVANCED STUDY, PRINCETON, NEW JERSEY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H.L. ROYDEN
Stanford University
Stanford, California

E. HEWITT
University of Washington
Seattle 5, Washington

R. P. DILWORTH
California Institute of Technology
Pasadena 4, California

* Alfred Horn
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

H. BUSEMANN	P.R. HALMOS	R.D. JAMES	GEORGE PÓLYA
HERBERT FEDERER	HEINZ HOPF	BØRGE JESSEN	J.J. STØKER
MARSHALL HALL	ALFRED HORN	PAUL LEVÝ	KOSAKU YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA, BERKELEY
UNIVERSITY OF CALIFORNIA, DAVIS
UNIVERSITY OF CALIFORNIA, LOS ANGELES
UNIVERSITY OF CALIFORNIA, SANTA BARBARA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
HUGHES AIRCRAFT COMPANY
SHELL DEVELOPMENT COMPANY

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. Manuscripts intended for the outgoing editors should be sent to their successors. All other communications to the editors should be addressed to the managing editor, Alfred Horn, at the University of California Los Angeles 24, California.

50 reprints of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50; back numbers (Volumes 1, 2, 3) are available at \$2.50 per copy. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to the publishers, University of California Press, Berkeley 4, California.

Printed at Ann Arbor, Michigan. Entered as second class matter at the Post Office, Berkeley, California.

* During the absence of E.C. Straus.

UNIVERSITY OF CALIFORNIA PRESS • BERKELEY AND LOS ANGELES

COPYRIGHT 1955 BY PACIFIC JOURNAL OF MATHEMATICS

Richard Horace Battin, <i>Note on the "Evaluation of an integral occurring in servomechanism theory"</i>	481
Frank Herbert Brownell, III, <i>An extension of Weyl's asymptotic law for eigenvalues</i>	483
Wilbur Eugene Deskins, <i>On the homomorphisms of an algebra onto Frobenius algebras</i>	501
James Michael Gardner Fell, <i>The measure ring for a cube of arbitrary dimension</i>	513
Harley M. Flanders, <i>The norm function of an algebraic field extension. II</i>	519
Dieter Gaier, <i>On the change of index for summable series</i>	529
Marshall Hall and Lowell J. Paige, <i>Complete mappings of finite groups</i>	541
Moses Richardson, <i>Relativization and extension of solutions of irreflexive relations</i>	551
Peter Scherk, <i>An inequality for sets of integers</i>	585
W. R. Scott, <i>On infinite groups</i>	589
A. Seidenberg, <i>On homogeneous linear differential equations with arbitrary constant coefficients</i>	599
Victor Lenard Shapiro, <i>Cantor-type uniqueness of multiple trigonometric integrals</i>	607
Leonard Tornheim, <i>Minimal basis and inessential discriminant divisors for a cubic field</i>	623
Helmut Wielandt, <i>On eigenvalues of sums of normal matrices</i>	633