

Pacific Journal of Mathematics

NOTE ON THE LERCH ZETA FUNCTION

F. OBERHETTINGER

NOTE ON THE LERCH ZETA FUNCTION

F. OBERHETTINGER

1. Introduction. The functional equation for the Riemann zeta function and Hurwitz's series were derived by a formal application of Poisson's summation formula in some previous papers [9], [10], [12, p. 17]. The results are correct although obtained by equating two infinite series whose regions of convergence exclude each other. It was shown later that this difficulty can be overcome if a generalized form of Poisson's formula is used [6]. It is the purpose of this note to show that the application of Poisson's summation formula in its ordinary form to the more general case of Lerch's zeta function [1], [2], [3], [5, pp. 27-30], [8] does not present the difficulties with respect to convergence which arise in its special cases. Thus a very simple and direct method for developing the theory of this function can be given. It may be noted that Hurwitz's series can be obtained immediately upon specializing one of the parameters. (cf. also [1]).

2. Lerch's zeta function $\varPhi(z, s, v)$ is usually defined by the power series

$$(1a) \quad \varPhi(z, s, v) = \sum_{m=0}^{\infty} z^m (v+m)^{-s} \quad |z| < 1, \quad v \neq 0, -1, -2, \dots$$

Equivalent to this is the integral representation [5, pp. 27-30]

$$(1b) \quad \begin{aligned} \varPhi(z, s, v) = & \frac{1}{2} v^{-s} + \int_0^{\infty} (v+t)^{-s} z^t dt \\ & + 2 \int_0^{\infty} \sin \left(s \tan^{-1} \frac{t}{v} - t \log z \right) (v^2 + t^2)^{-s/2} (e^{2\pi t} - 1)^{-1} dt \\ & |z| < 1, \quad \Re v > 0, \end{aligned}$$

which can be written as

$$(1c) \quad \begin{aligned} \varPhi(z, s, v) = & \frac{1}{2} v^{-s} + z^{-v} \Gamma(1-s) \left(\log \frac{1}{z} \right)^{s-1} - z^{-v} v^{1-s} \sum_{n=0}^{\infty} \frac{\left(-v \log \frac{1}{z} \right)^n}{n! (n+1-s)} \\ & + 2 \int_0^{\infty} \sin \left(s \tan^{-1} \frac{t}{v} - t \log z \right) (v^2 + t^2)^{-s/2} (e^{2\pi t} - 1)^{-1} dt \\ & z \neq 1, \quad \Re v > 0 \end{aligned}$$

upon replacing the first integral on the right of (1b) by the second and

Received February 11, 1954.

third expression on the right of (1c). Hence it follows that the point $z=1$ is a singular point of ϕ such that

$$(2) \quad \lim_{z \rightarrow 1} \left[\sum_0^{\infty} z^m (v+m)^{-s} - z^{-v} \Gamma(1-s) \left(\log \frac{1}{z} \right)^{s-1} \right] \\ = \frac{1}{2} v^{-s} + \frac{v^{1-s}}{s-1} + 2 \int_0^{\infty} \sin \left(s \tan^{-1} \frac{t}{v} \right) (v^2+t^2)^{-s/2} (e^{2\pi t}-1)^{-1} dt = \zeta(s, v)$$

by Hermite's formula [5 p. 26]. This relation was first proved by Hardy [7] in a different manner. For $\Re s > 1$, (2) is obviously a consequence of Abel's theorem. We apply now Poisson's summation formula [4, p. 33], [11, p. 60]

$$(3) \quad \sum_{n=-\infty}^{\infty} e^{i2\pi nv} f(2\pi n) = \frac{1}{2\pi} \sum_{m=-\infty}^{\infty} G(v+m) \quad 0 < v \leq 1$$

where

$$(4) \quad G(t) = \int_{-\infty}^{\infty} f(y) e^{iyt} dy$$

to the function

$$(5) \quad f(x) = (\alpha + ix)^{-s}, \quad \Re s > 1, \quad \Re \alpha > 0, \quad x \text{ real}, \quad -\frac{1}{2}\pi < \arg(\alpha + ix) < \frac{1}{2}\pi.$$

According to (4)

$$(6) \quad G(t) = \int_{-\infty}^{\infty} (\alpha + iy)^{-s} e^{iyt} dy = 2\pi e^{-\alpha t} \frac{1}{2\pi i} \int_{\alpha-i\infty}^{\alpha+i\infty} p^{-s} e^{pt} dp.$$

The last integral multiplied by $(2\pi i)^{-1}$ is the inverse Laplace transform of p^{-s} and is equal to $\frac{t^{s-1}}{\Gamma(s)}$. Thus for $\Re s > 1$

$$(7) \quad G(t) = \frac{2\pi}{\Gamma(s)} t^{s-1} e^{-\alpha t}, \quad t > 0, \\ = 0, \quad t < 0.$$

From (3) to (7) we obtain

$$(8) \quad \sum_{n=-\infty}^{\infty} (\alpha + i2\pi n)^{-s} e^{i2\pi nv} = \frac{1}{\Gamma(s)} \sum_{m=0}^{\infty} (v+m)^{s-1} e^{-(v+m)\alpha}$$

or, with s replaced by $1-s$

$$(9) \quad \sum_{m=0}^{\infty} (v+m)^{-s} e^{-m\alpha} = \Gamma(1-s) e^{v\alpha} \sum_{n=-\infty}^{\infty} e^{i2\pi nv} (\alpha + i2\pi n)^{s-1} \\ \Re s < 0, \quad 0 < v \leq 1, \quad 0 \leq \Im \alpha < 2\pi.$$

The last restriction means no loss of generality since both sides of (9) are periodic in α with the period $2\pi i$. We write (9)

$$(10) \quad \begin{aligned} & \sum_{m=0}^{\infty} (v+m)^{-s} e^{-m\alpha} - \Gamma(1-s)e^{v\alpha}\alpha^{s-1} \\ & = \Gamma(1-s)e^{v\alpha} \left[\sum_{n=1}^{\infty} \frac{e^{i2\pi nv}}{(\alpha + i2\pi n)^{1-s}} + \sum_{n=1}^{\infty} \frac{e^{-i2\pi nv}}{(\alpha - i2\pi n)^{1-s}} \right] \\ & \quad \Re s < 0, \quad 0 < v \leq 1, \quad 0 \leq \Im \alpha < 2\pi, \end{aligned}$$

where according to the condition in (5) for $n=1, 2, 3, \dots$

$$0 < \arg\left(\frac{\alpha}{2\pi} + in\right) < \frac{1}{2}\pi, \quad -\frac{1}{2}\pi < \arg\left(\frac{\alpha}{2\pi} - in\right) < 0.$$

Hence,

$$\arg\left(\frac{\alpha}{2\pi} \pm in\right) = \pm \frac{1}{2}\pi + \arg\left(n \pm \frac{\alpha}{2\pi i}\right)$$

and consequently

$$(\alpha \pm i2\pi n)^{s-1} = (2\pi)^{s-1} e^{\pm i\pi(s-1)/2} \left(n \pm \frac{\alpha}{2\pi i} \right)^{s-1}.$$

Using (1a) we obtain finally from (10)

$$(11) \quad \begin{aligned} & \varphi(e^{-\alpha}, s, v) - \Gamma(1-s)e^{v\alpha}\alpha^{s-1} \\ & = -i(2\pi)^{s-1} \Gamma(1-s)e^{v\alpha} \left[\sum_{n=1}^{\infty} \frac{e^{i(2\pi nv + \frac{1}{2}\pi s)}}{\left(n + \frac{\alpha}{2\pi i}\right)^{1-s}} - \sum_{n=1}^{\infty} \frac{e^{-i(2\pi nv + \frac{1}{2}\pi s)}}{\left(n - \frac{\alpha}{2\pi i}\right)^{1-s}} \right] \\ & \quad \Re s < 0, \quad 0 < v \leq 1, \quad 0 \leq \Im \alpha < 2\pi, \end{aligned}$$

which may be written as

$$(12) \quad \begin{aligned} & \varphi(e^{-\alpha}, s, v) - \Gamma(1-s)e^{v\alpha}\alpha^{s-1} \\ & = -i(2\pi)^{s-1} \Gamma(1-s)e^{v\alpha} \left[e^{i(2\pi v + \frac{1}{2}\pi s)} \varphi\left(e^{i2\pi v}, 1-s, 1 + \frac{\alpha}{2\pi i}\right) \right. \\ & \quad \left. - e^{-i(2\pi v + \frac{1}{2}\pi s)} \varphi\left(e^{-i2\pi v}, 1-s, 1 - \frac{\alpha}{2\pi i}\right) \right]. \end{aligned}$$

This is Lerch's transformation formula and (11) corresponds to Hurwitz's series. Now it follows that the right hand side of (11) is also regular for $\alpha=0$. If in (11) we let α tend to zero, the left side becomes $\zeta(s, v)$ by (2) and the right hand side becomes Hurwitz's series. Thus

$$(13) \quad \zeta(s, v) = 2(2\pi)^{s-1} \Gamma(1-s) \sum_{n=1}^{\infty} n^{s-1} \sin \left(2\pi nv + \frac{1}{2}\pi s \right)$$

$$0 < v \leq 1, \quad \Re s < 0.$$

REFERENCES

1. T. M. Apostol, *Remark on the Hurwitz zeta function*, Proc. Amer. Math. Soc., **2** (1951), 690-693.
2. ———, *On the Lerch zeta function*, Pacific J. Math., **1** (1951), 161-167.
3. E. W. Barnes, *On certain functions defined by Taylor series of finite radius of convergence*, Proc. London Math. Soc., **4** (1906), 284-316.
4. S. Bochner, *Vorlesungen über Fouriersche Integrale*, Leipzig, 1932.
5. A. Erdelyi, *Higher transcendental functions*, vol. 1, New-York, 1953.
6. A. P. Guinand, *On Poisson's summation formula*, Ann. Math., **42** (1941), 591-603.
- 6a. ———, *Some formulae for the Riemann zeta function*, J. London Math. Soc., **22** (1947), 14-18.
7. G. H. Hardy, *Method for determining the behavior of certain classes of power series near a singular point on the circle of convergence*, Proc. London Math. Soc., **3** (1905), 381-389.
8. M. Lerch, *Note sur la fonction $K(w, x, s) = \sum_{k=0}^{\infty} e^{2\pi i k x} (w+k)^{-s}$* , Acta Math., **11** (1887), 19-24.
9. L. J. Mordell, *Some applications of Fourier series in the analytic theory of numbers*, Proc. Cambridge Phil. Soc., **24** (1925), 585-596.
10. ———, *Poisson's summation formula and the Riemann zeta function*, J. London Math. Soc., **4** (1929), 285-291.
11. E. C. Titchmarsh, *Introduction to the theory of Fourier Integrals*, Oxford, 1943.
12. ———, *The theory of Riemann's zeta function*, Oxford, 1951.

NATIONAL BUREAU OF STANDARDS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. L. ROYDEN

Stanford University
Stanford, California

R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

E. HEWITT

University of Washington
Seattle 5, Washington

E. G. STRAUS

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

C. E. BURGESS

H. BUSEMANN

H. FEDERER

M. HALL

P. R. HALMOS

V. GANAPATHY IYER

R. D. JAMES

M. S. KNEBELMAN

I. NIVEN

T. G. OSTROM

M. M. SCHIFFER

J. J. STOKER

G. SZEKERES

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF UTAH

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

CALIFORNIA RESEARCH CORPORATION

HUGHES AIRCRAFT COMPANY

Printed in Japan by Kokusai Bunken Insatsusha
(International Academic Printing Co., Ltd.), Tokyo, Japan

Pacific Journal of Mathematics

Vol. 6, No. 1

November, 1956

David Blackwell, <i>An analog of the minimax theorem for vector payoffs</i>	1
L. W. Cohen, <i>A non-archimedean measure in the space of real sequences</i>	9
George Bernard Dantzig, <i>Constructive proof of the Min-Max theorem</i>	25
Jim Douglas, <i>On the numerical integration of quasilinear parabolic differential equations</i>	35
James Michael Gardner Fell, <i>A note on abstract measure</i>	43
Isidore Isaac Hirschman, Jr., <i>A note on orthogonal systems</i>	47
Frank Harary, <i>On the number of dissimilar line-subgraphs of a given graph</i>	57
Newton Seymour Hawley, <i>Complex bundles with Abelian group</i>	65
Alan Jerome Hoffman, Morris Newman, Ernst Gabor Straus and Olga Taussky, <i>On the number of absolute points of a correlation</i>	83
Ernst Gabor Straus and Olga Taussky, <i>Remark on the preceding paper. Algebraic equations satisfied by roots of natural numbers</i>	97
Ralph D. James, <i>Summable trigonometric series</i>	99
Gerald R. Mac Lane, <i>Limits of rational functions</i>	111
F. Oberhettinger, <i>Note on the Lerch zeta function</i>	117
Gerald C. Preston, <i>On locally compact totally disconnected Abelian groups and their character groups</i>	121
Vikramaditya Singh and W. J. Thron, <i>On the number of singular points, located on the unit circle, of certain functions represented by C-fractions</i>	135
Sherman K. Stein, <i>The symmetry function in a convex body</i>	145
Edwin Weiss, <i>Boundedness in topological rings</i>	149
Albert Leon Whiteman, <i>A sum connected with the series for the partition function</i>	159
Alfred B. Willcox, <i>Some structure theorems for a class of Banach algebras</i>	177
Joseph Lawrence Zemmer, <i>Some remarks on p-rings and their Boolean geometry</i>	193