

Pacific Journal of Mathematics

**SOME INTEGRAL FORMULAS FOR CLOSED
HYPERSURFACES IN RIEMANNIAN SPACE**

CHUAN CHIH HSIUNG

SOME INTEGRAL FORMULAS FOR CLOSED HYPERSURFACES IN RIEMANNIAN SPACE

CHUAN-CHIH HSIUNG

Introduction. Let V^n be a hypersurface twice differentially imbedded in a Riemannian space R^{n+1} of $n+1$ ($n \geq 2$) dimensions, and $\kappa_1, \dots, \kappa_n$ the n principal curvatures at a point P of the hypersurface V^n . It is known that the i th mean curvature M_i of the hypersurface V^n at the point P is defined by

$$(0.1) \quad C_{n,i} M_i = \sum \kappa_1 \kappa_2 \cdots \kappa_i \quad (i=1, \dots, n),$$

where the expression on the right side is the i th elementary symmetric function of $\kappa_1, \dots, \kappa_n$, and $C_{n,i}$ denotes the number of combinations of n different things taken i at a time. Let dA be the area element of the hypersurface V^n at the point P , and p the scalar product of the unit normal vector of the hypersurface V^n at the point P and the position vector of the point P with respect to any orthogonal frame in the space R^{n+1} .

The purpose of this paper is to prove the following four theorems concerning closed hypersurfaces by first showing that:

a) If V^n is an orientable hypersurface, with a closed boundary V^{n-1} of dimension $n-1$ ($n \geq 2$), which is twice differentially imbedded in an $(n+1)$ -dimensional Riemannian space R^{n+1} , then the integral $\int_{V^n} (1 + M_1 p) dA$ can be expressed as an integral over the boundary V^{n-1} .

b) If in addition V^n is of class C^3 and the space R^{n+1} is of constant Riemannian curvature, then the integral $\int_{V^n} (M_{n-1} + M_n p) dA$ can also be expressed as an integral over V^{n-1} .

These results have been obtained in a previous paper [2] by the author for an orientable hypersurface V^n twice differentially imbedded in a Euclidean space E^{n+1} of $n+1$ ($n \geq 2$) dimensions.

THEOREM 1. *Let V^n be a closed orientable hypersurface twice differentially imbedded in a Riemannian space R^{n+1} of $n+1$ ($n \geq 2$) dimensions, then*

$$(I) \quad A + \int_{V^n} M_1 p dA = 0 .$$

Received June 28, 1954 and in revised forms January 3, 1955 and May 10, 1955.

THEOREM 2. *Let V^n be a closed orientable hypersurface of class C^3 imbedded in an $(n+1)$ -dimensional ($n \geq 2$) Riemannian space R^{n+1} of constant Riemannian curvature K , then*

$$(II) \quad \int_{V^n} M_{n-1} dA + \int_{V^n} M_n p dA = 0 .$$

THEOREM 3. *Let V^n be a hypersurface satisfying the conditions of Theorem 2. Suppose that the principal curvatures $\kappa_1, \dots, \kappa_n$ at each point of the hypersurface V^n are positive and that in the space R^{n+1} there exists a point O for which either $p \leq -1/M_1$ or $p \geq -1/M_1$ at all points of the hypersurface V^n . Then every point of the hypersurface V^n is umbilic.*

THEOREM 4. *Let V^n be a hypersurface satisfying the conditions of Theorem 2. Suppose that the principal curvatures $\kappa_1, \dots, \kappa_n$ at each point of the hypersurface V^n are positive and M_{n-1} is constant, and that in the space R^{n+1} there exists a point O for which the function p is of the same sign at all points of the hypersurface V^n . Then every point of the hypersurface V^n is umbilic.*

1. Preliminaries. In a Riemannian space R^{n+1} of dimension $n+1$ ($n \geq 2$) with a positive definite fundamental form we consider a fixed orthogonal frame $Oe_1 \dots e_{n+1}$, where e_1, \dots, e_{n+1} form an ordered set of $n+1$ mutually orthogonal contravariant unit vectors at a point O in R^{n+1} . With respect to this orthogonal frame let y^α ($\alpha=1, \dots, n+1$) be¹ the coordinates of a point in R^{n+1} and $a_{\alpha\beta} dy^\alpha dy^\beta$ the fundamental form for R^{n+1} , where $a_{\alpha\beta} = a_{\beta\alpha}$ and the matrix $\|a_{\alpha\beta}\|$ is positive definite so that the determinant $a = |a_{\alpha\beta}| > 0$.

Let A_{i1} ($i=1, \dots, n$) be n vectors at a point in the space R^{n+1} whose contravariant components with respect to the frame $Oe_1 \dots e_{n+1}$ are A_{i1}^α ($\alpha=1, \dots, n+1$). First we define the vector product of the n vectors A_{i1} ($i=1, \dots, n$) to be a vector in R^{n+1} , denoted by $A_{11} \times \dots \times A_{n1}$, whose contravariant components are given by

$$(1.1) \quad A_{11} \times \dots \times A_{n1} = (-1)^n \begin{vmatrix} e_1 & e_2 & \dots & e_{n+1} \\ a_{\alpha 1} A_{11}^\alpha & a_{\alpha 2} A_{11}^\alpha & \dots & a_{\alpha, n+1} A_{11}^\alpha \\ a_{\alpha 1} A_{21}^\alpha & a_{\alpha 2} A_{21}^\alpha & \dots & a_{\alpha, n+1} A_{21}^\alpha \\ \dots & \dots & \dots & \dots \\ a_{\alpha 1} A_{n1}^\alpha & a_{\alpha 2} A_{n1}^\alpha & \dots & a_{\alpha, n+1} A_{n1}^\alpha \end{vmatrix} .$$

From the definition of the scalar product of any two vectors A_{i1} and A_{j1} ,

¹ Throughout this paper Greek indices take the values 1 to $n+1$, and Latin indices the values 1 to n unless stated otherwise. We use the convention that repeated indices imply summation.

namely, $A_{i_1} \cdot A_{j_1} = a_{\alpha\beta} A_{i_1}^\alpha A_{j_1}^\beta$, it follows immediately that $A_{i_1} \times \cdots \times A_{n_1}$ is orthogonal to A_{i_1} ($i=1, \dots, n$).

Now we consider a hypersurface V^n twice differentially imbedded in the space E^{n+1} . With respect to the orthogonal frame $Oe_1 \cdots e_{n+1}$ the hypersurface V^n can be given by the parametric equations

$$(1.2) \quad y^\alpha = f^\alpha(x^1, \dots, x^n) \quad (\alpha=1, \dots, n+1),$$

or the vector equation

$$(1.3) \quad Y = F(x^1, \dots, x^n),$$

where y^α and f^α are respectively the contravariant components of the two vectors Y and F , the parameters x^1, \dots, x^n take values in a simply connected domain D of the n -dimensional real number space, and $f^\alpha(x^1, \dots, x^n)$ is of rank n at all points of D . Let the first fundamental form of the hypersurface V^n at a point P be

$$(1.4) \quad ds^2 = g_{ij} dx^i dx^j,$$

where the matrix $\|g_{ij}\|$ is positive definite so that the determinant $g = |g_{ij}| > 0$, and

$$(1.5) \quad g_{ij} = a_{\alpha\beta} y_i^\alpha y_j^\beta,$$

$$(1.6) \quad y_{,i}^\alpha = \partial y^\alpha / \partial x^i.$$

Let $A_{\beta i}^\alpha$ be a mixed tensor of the second order in the y 's, and a covariant vector in the x 's, as indicated by the Greek and Latin indices. Then following Tucker [3], the generalized covariant derivative of $A_{\beta i}^\alpha$ with respect to the x 's is defined as

$$(1.7) \quad A_{\beta i,j}^\alpha = \frac{\partial A_{\beta i}^\alpha}{\partial x^j} + \left\{ \begin{matrix} \alpha \\ \beta \gamma \end{matrix} \right\} A_{\beta i}^\gamma y_{,j}^\delta - \left\{ \begin{matrix} \gamma \\ \beta \delta \end{matrix} \right\} A_{\gamma i}^\alpha y_{,j}^\delta - \left\{ \begin{matrix} k \\ ij \end{matrix} \right\} A_{\beta k}^\alpha,$$

where the Christoffel symbols $\left\{ \begin{matrix} \alpha \\ \beta \gamma \end{matrix} \right\}$ with Greek indices are formed with respect to the $a_{\alpha\beta}$ and the y 's, and those $\left\{ \begin{matrix} k \\ ij \end{matrix} \right\}$ with Latin indices with respect to the g_{ij} and the x 's. It should be noted that the definition of generalized covariant differentiation can be applied to any tensor in the x 's and y 's and that the generalized covariant differentiation of sums and products obeys the ordinary rules. If a tensor is one with respect to the x 's only, so that only Latin indices appear, its generalized covariant derivative is the same as its covariant derivative with respect to the x 's. Moreover, in generalized covariant differentiation the fundamental tensors $a_{\alpha\beta}$ and g_{ij} can be treated as constants. Since y^α is an invariant for transformation of the x 's, its generalized covariant derivative

is the same as its covariant derivative with respect to the x 's; so that

$$(1.8) \quad y_{;i}^\alpha = y_{,i}^\alpha = \partial y^\alpha / \partial x^i .$$

By (1.7) the generalized covariant derivative of $y_{;i}^\alpha$ is

$$(1.9) \quad y_{;i;j}^\alpha = \frac{\partial^2 y^\alpha}{\partial x^i \partial x^j} - \left\{ \begin{matrix} h \\ ij \end{matrix} \right\} y_{,h}^\alpha + \left\{ \begin{matrix} \alpha \\ \beta \gamma \end{matrix} \right\} y_{,i}^\beta y_{,j}^\gamma ,$$

which is symmetric in the indices i and j .

Let N be the unit normal vector at a point P of the hypersurface V^n , then

$$(1.10) \quad a_{\alpha\beta} N^\alpha N^\beta = 1 ,$$

$$(1.11) \quad a_{\alpha\beta} N_\alpha y_{,i}^\beta = 0 \quad (i=1, \dots, n).$$

We can easily obtain (see, for instance, [4, Chap. VIII]):

$$(1.12) \quad y_{;i;j}^\alpha = \Omega_{i,j} N^\alpha ,$$

$$(1.13) \quad \Omega_{i,j} = y_{,i;j}^\alpha a_{\alpha\beta} N^\beta ,$$

$$(1.14) \quad N_{;i}^\alpha = -\Omega_{i,j} g^{jk} y_{,k}^\alpha ,$$

where $\Omega_{i,j} = \Omega_{j,i}$ are the coefficients of the second fundamental form of the hypersurface V^n at the point P , and g^{ij} denotes the cofactor of g_{ij} in g divided by g so that

$$(1.15) \quad g^{ij} g_{jk} = \delta_k^i ,$$

δ_k^i being the Kronecker delta. Moreover, we have

$$(1.16) \quad R_{iij;k} = (\Omega_{i,j} \Omega_{i,k} - \Omega_{i,k} \Omega_{i,j}) + \bar{R}_{\beta\gamma\delta\varepsilon} y_{,i}^\beta y_{,i}^\gamma y_{,j}^\delta y_{,k}^\varepsilon ,$$

$$(1.17) \quad \Omega_{i,j,k} - \Omega_{i,k,j} = -\bar{R}_{\beta\gamma\delta\varepsilon} N^\beta y_{,i}^\gamma y_{,j}^\delta y_{,k}^\varepsilon ,$$

where $R_{iij;k}$ and $\bar{R}_{\beta\gamma\delta\varepsilon}$ are Riemann symbols formed with the tensors g_{ij} and $a_{\alpha\beta}$ respectively. In particular, if the space R^{n+1} is of constant Riemannian curvature K , it follows from the definition of Riemannian curvatures of the space R^{n+1} that

$$(1.18) \quad \bar{R}_{\beta\gamma\delta\varepsilon} = K(a_{\beta\delta} a_{\gamma\varepsilon} - a_{\beta\varepsilon} a_{\gamma\delta}) ,$$

and therefore (1.16), (1.17) reduce to

$$(1.19) \quad R_{iij;k} = (\Omega_{i,j} \Omega_{i,k} - \Omega_{i,k} \Omega_{i,j}) + K(g_{ij} g_{ik} - g_{ik} g_{ij}) ,$$

$$(1.20) \quad \Omega_{i,j,k} - \Omega_{i,k,j} = 0 .$$

Taking the generalized covariant derivative of each side of (1.14) and

making use of (1.12), (1.19), (1.20) we thus obtain

$$(1.21) \quad N_{;ji}^\alpha - N_{;ij}^\alpha = N^\alpha g^{lk} (R_{jilk1} - \bar{R}_{\beta\gamma\delta\epsilon} y_j^\beta y_i^\gamma y_l^\delta y_1^\epsilon) .$$

The n principal curvatures $\kappa_1, \dots, \kappa_n$ of the hypersurface V^n at the point P are the roots of the determinant equation

$$(1.22) \quad |\Omega_{ij} - \kappa g_{ij}| = 0 .$$

From (0.1) and (1.22) it follows immediately that

$$(1.23) \quad M_n = \Omega/g , \quad nM_1 = \Omega_{ij} g^{ij} , \quad nM_{n-1} = g_{ij} \Omega^{ij} /g ,$$

where $\Omega = |\Omega_{ij}|$ and Ω^{ij} is the cofactor of Ω_{ij} in Ω .

Consider the two matrices

$$(1.24) \quad \phi = \|\phi_\gamma^i\| , \quad \psi = \|\psi_i^\gamma\| ;$$

where

$$(1.25) \quad \phi_\gamma^i = a_{\beta\gamma} y_{,i}^\beta , \quad \psi_i^\gamma = y_{,i}^\gamma \quad (i=1, \dots, n ; \gamma=1, \dots, n+1) ,$$

the superscript of the element ϕ_γ^i or ψ_i^γ indicating the row to which the element belongs and the subscript indicating the column. Solving (1.11) for N^α , we obtain

$$(1.26) \quad N^\alpha = (-1)^{n-\alpha+1} c A^\alpha \quad (\alpha=1, \dots, n+1) ,$$

where c is a constant and A^α the determinant of n th order obtained by deleting the α th column from the matrix ϕ . Substitution of (1.26) in (1.10) gives

$$(1.27) \quad c^2 = \frac{1}{aA} ,$$

where

$$(1.28) \quad A = \begin{vmatrix} A^1 & -A^2 & \dots & (-1)^n A^{n+1} \\ y_{,1}^1 & y_{,1}^2 & \dots & y_{,1}^{n+1} \\ \dots & \dots & \dots & \dots \\ y_{,n}^1 & y_{,n}^2 & \dots & y_{,n}^{n+1} \end{vmatrix} ,$$

which is equal to the sum of the products of the corresponding determinants of n th order of the two matrices (1.24). By an elementary theorem on determinants (see, for instance, [1, p. 102]), from (1.5) it follows immediately that

$$(1.29) \quad A = |\phi_\gamma^i \psi_j^\gamma| = g .$$

Now we choose the direction of the unit normal vector N in such

a way that the two frames $PY_{,1}\cdots Y_{,n}N$ and $Oe_1\cdots e_{n+1}$ have the same orientation. Then from (1.10), (1.26), (1.27), (1.29) we obtain

$$(1.30) \quad \sqrt{g\bar{a}}N = Y_{,1} \times \cdots \times Y_{,n} ,$$

$$(1.31) \quad |Y_{,1}, \cdots, Y_{,n}, N| = \sqrt{g/\bar{a}} .$$

The area element of the hypersurface V^n at the point P is given by

$$(1.32) \quad dA = \sqrt{g} dx^1 \cdots dx^n .$$

Let A_{i_1} ($i=1, \cdots, n$) be n vectors at a point in the space R^{n+1} , whose contravariant components with respect to the frame $Oe_1\cdots e_{n+1}$ are differentiable functions of x^1, \cdots, x^n , then by (1.1) and the differentiation of determinants

$$(1.33) \quad (A_{i_1} \times \cdots \times A_{i_n})_{;i} = \sum_j (A_{i_1} \times \cdots \times A_{j-1} \times A_{j;i} \times A_{j+1} \times \cdots \times A_{i_n}) .$$

2. Proof of the formula (I). First we observe that the vector $Y_{,1} \times \cdots \times Y_{,i-1} \times N \times Y_{,i+1} \times \cdots \times Y_{,n}$ is orthogonal to the normal vector N and can therefore be written in the form

$$(2.1) \quad Y_{,1} \times \cdots \times Y_{,i-1} \times N \times Y_{,i+1} \times \cdots \times Y_{,n} = c^{ij} Y_{,j} \quad (i=1, \cdots, n).$$

Taking the scalar products of both sides of (2.1) with the vector $Y_{,k}$ and making use of (1.2), (1.5), (1.31), we obtain

$$(2.2) \quad c^{ij} g_{jk} = -\sqrt{g\bar{a}} \delta_k^i \quad (i, k=1, \cdots, n).$$

Solving (2.2) for c^{ij} for each fixed i and substituting the results in (2.1), we are led to

$$(2.3) \quad Y_{,1} \times \cdots \times Y_{,i-1} \times N \times Y_{,i+1} \times \cdots \times Y_{,n} = -\sqrt{g\bar{a}} g^{ij} Y_{,j} \quad (i=1, \cdots, n).$$

Making use of the relation $Y_{;ij} = Y_{;ji}$ and (1.14), (1.23), (1.30), (1.33), it is easily seen that

$$(2.4) \quad \begin{aligned} & \sum_i (Y_{,1} \times \cdots \times Y_{,i-1} \times N \times Y_{,i+1} \times \cdots \times Y_{,n})_{;i} \\ &= \sum_i Y_{,1} \times \cdots \times Y_{,i-1} \times N_{;i} \times Y_{,i+1} \times \cdots \times Y_{,n} \\ &= -n\sqrt{g\bar{a}} M_1 N . \end{aligned}$$

Thus, from (2.3) and (2.4),

$$(2.5) \quad n\sqrt{g} M_1 N = (\sqrt{g} g^{ij} Y_{,i};_j) .$$

Taking the scalar products of both sides of (2.5) with the vector Y , we obtain in consequence of (1.5) and (1.15)

$$(2.6) \quad nM_1 p \sqrt{g} = (\sqrt{g} g^{ij} \eta_i)_{,j} - n \sqrt{g} ,$$

where we have put

$$(2.7) \quad p = Y \cdot N , \quad \eta_i = Y \cdot Y_{,i} \quad (i=1, \dots, n).$$

Now let us consider a hypersurface V^n , with a closed boundary V^{n-1} of dimension $n-1$ ($n \geq 2$), twice differentiably imbedded in a Riemannian space R^{n+1} of dimension $n+1$. Integrating (2.6) with respect to x^1, \dots, x^n over this hypersurface V^n and applying the general theorem of Stokes to the first term on the right side of (2.6), we obtain

$$(2.8) \quad A + \int_{V^n} M_1 p dA = \frac{1}{n} \int_{V^{n-1}} \sum_j (-1)^{j-1} \sqrt{g} g^{ij} \eta_i dx^1 \dots dx^{j-1} dx^{j+1} \dots dx^n .$$

In particular, when the hypersurface V^n is closed and orientable, the integral on the right side of (2.8) vanishes and hence we obtain the formula (I).

3. Proof of the formula (II). For the same reason as in the preceding section, the vector $N_{,1} \times \dots \times N_{,i-1} \times N \times N_{,i+1} \times \dots \times N_{,n}$ is orthogonal to the normal vector N and can therefore be written in the form

$$(3.1) \quad N_{,1} \times \dots \times N_{,i-1} \times N \times N_{,i+1} \times \dots \times N_{,n} = c^{ij} Y_{,j} \quad (i=1, \dots, n).$$

Taking the scalar products of both sides of (3.1) with the vector $Y_{,k}$ and making use of (1.1), (1.14), (1.31), we obtain

$$c^{ij} g_{jk} = (-1)^{n+i} a |N, Y_{,1}, \dots, Y_{,n}|$$

$$= (-1)^{n+i+k} \sqrt{ga} \begin{vmatrix} \Omega_{11} & \Omega_{12} & \dots & \Omega_{1n} \\ \dots & \dots & \dots & \dots \\ \Omega_{i-1,1} & \Omega_{i-1,2} & \dots & \Omega_{i-1,n} \\ \dots & \dots & \dots & \dots \\ \Omega_{i+1,1} & \Omega_{i+1,2} & \dots & \Omega_{i+1,n} \\ \dots & \dots & \dots & \dots \\ \Omega_{n1} & \Omega_{n2} & \dots & \Omega_{nn} \\ g_{k1} & g_{k2} & \dots & g_{kn} \end{vmatrix} \begin{vmatrix} g^{11} & \dots & g^{1,k-1} & g^{1,k+1} & \dots & g^{1n} & g^{1k} \\ g^{21} & \dots & g^{2,k-1} & g^{2,k+1} & \dots & g^{2n} & g^{2k} \\ \dots & \dots & \dots & \dots & \dots & \dots & \dots \\ g^{n1} & \dots & g^{n,k-1} & g^{n,k+1} & \dots & g^{nn} & g^{nk} \end{vmatrix} ,$$

and therefore

$$(3.2) \quad c^{ij}g_{jk} = (-1)^n \sqrt{a/g} g_{kj} \Omega^{ij} \quad (i, k=1, \dots, n).$$

Solving (3.2) for c^{ij} for each fixed i and substituting the results in (3.1), we find

$$(3.3) \quad N_{;i} \times \dots \times N_{;i-1} \times N \times N_{;i+1} \times \dots \times N_{;n} = (-1)^n \sqrt{a/g} \Omega^{ij} Y_{,j}.$$

Making use of (1.1), (1.14), (1.21), (1.23), (1.30), (1.33), it is easily seen that

$$\begin{aligned} & \sum_i (N_{;1} \times \dots \times N_{;i-1} \times N \times N_{;i+1} \times \dots \times N_{;n})_{;i} \\ &= \sum_i (N_{;1} \times \dots \times N_{;i-1} \times N_{;i} \times N_{;i+1} \times \dots \times N_{;n}) \\ &= n \begin{vmatrix} e_1 & a_{\alpha 1} y_{,1}^\alpha & \dots & a_{\alpha 1} y_{,n}^\alpha \\ e_2 & a_{\alpha 2} y_{,1}^\alpha & \dots & a_{\alpha 2} y_{,n}^\alpha \\ \dots & \dots & \dots & \dots \\ e_{n+1} & a_{\alpha, n+1} y_{,1}^\alpha & \dots & a_{\alpha, n+1} y_{,n}^\alpha \end{vmatrix} \cdot \begin{vmatrix} 1 & 0 & 0 & \dots & 0 \\ 0 & \Omega_{1j} g^{j1} & \Omega_{2j} g^{j1} & \dots & \Omega_{nj} g^{j1} \\ 0 & \Omega_{1j} g^{j2} & \Omega_{2j} g^{j2} & \dots & \Omega_{nj} g^{j2} \\ \dots & \dots & \dots & \dots & \dots \\ 0 & \Omega_{1j} g^{jn} & \Omega_{2j} g^{jn} & \dots & \Omega_{nj} g^{jn} \end{vmatrix} \\ &= (-1)^n n \sqrt{ga} M_n N. \end{aligned}$$

Thus, from the above equation and (3.3),

$$(3.4) \quad n \sqrt{g} M_n N = (\Omega^{ij} Y_{,j} / \sqrt{g})_{;i}.$$

Taking the scalar products of both sides of (3.4) with the vector Y , we obtain in consequence of (1.23) and (2.7)

$$(3.5) \quad n M_n p \sqrt{g} = (\Omega^{ij} \eta_j / \sqrt{g})_{;i} - n M_{n-1} \sqrt{g}.$$

As in the preceding section, let us consider a hypersurface V^n , with a closed boundary V^{n-1} of dimension $n-1$ ($n \geq 2$), differentiably of class C^3 imbedded in an $(n+1)$ -dimensional Riemannian space R^{n+1} of constant Riemannian curvature K . Integrating (3.5) with respect to x^1, \dots, x^n over this hypersurface V^n and applying Stokes' theorem to the first term on the right side of (3.5), we then obtain

$$(3.6) \quad \int_{V^n} M_{n-1} dA + \int_{V^n} M_n p dA = \frac{1}{n} \int_{V^{n-1}} \sum_j (-1)^{j-1} \frac{\Omega^{ij} \eta_j}{\sqrt{g}} dx^1 \dots dx^{j-1} dx^{j+1} \dots dx^n.$$

In particular, when the hypersurface V^n is closed and orientable, the integral on the right side of (3.6) vanishes and hence the formula (II).

4. Proofs of Theorems 3 and 4. For $M_1 > 0$, the assumptions $p \leq -1/M_1$ and $p \geq -1/M_1$ are respectively equivalent to $1 + M_1 p \leq 0$ and $1 + M_1 p \geq 0$. From formula (I) it follows that each of the above two assumptions implies that $p = -1/M_1$. Substituting this in (II) we obtain

$$(4.1) \quad \int_{V^n} \frac{1}{M_1} (M_1 M_{n-1} - M_n) dA = 0,$$

which holds when and only when $M_1 M_{n-1} - M_n = 0$, since

$$(4.2) \quad \begin{aligned} M_1 M_{n-1} - M_n &= \frac{1}{n^2} \left(\sum_i \kappa_i \sum \kappa_{i_1} \kappa_{i_2} \cdots \kappa_{i_{n-2}} - n^2 \kappa_1 \kappa_2 \cdots \kappa_n \right) \\ &= \frac{1}{n^2} \sum [\kappa_{i_1} \kappa_{i_2} \cdots \kappa_{i_{n-2}} (\kappa_{i_{n-1}} - \kappa_{i_n})^2] \geq 0, \end{aligned}$$

where i_1, i_2, \dots, i_n are distinct and run from 1 to n . From (4.1), (4.2) it follows that $\kappa_1 = \kappa_2 = \dots = \kappa_n$ at each point of the hypersurface V^n and therefore that the quantity defined by

$$(\Omega_{ij} q^i q^j) / (g_{ij} q^i q^j)$$

at each point of the hypersurface V^n for an arbitrary direction q in the hypersurface V^n with contravariant components q^i is independent of the direction q . Hence $\Omega_{ij} = c g_{ij}$ for all i and j at each point of the hypersurface V^n , where c is a scalar invariant, so that every point of the hypersurface V^n is umbilic.

If M_{n-1} is constant, multiplying the formula (I) by M_{n-1} and subtracting the formula (II) by the resulting equation we obtain

$$(4.3) \quad \int_{V^n} p (M_1 M_{n-1} - M_n) dA = 0.$$

From this and the assumption that p is of the same sign at all points of the hypersurface V^n , Theorem 4 follows by exactly the same argument as above.

REFERENCES

1. L. P. Eisenhart, *Riemannian geometry*, Princeton, 1926.
2. C. C. Hsiung, *Some integral formulas for closed hypersurfaces*, Math. Scand., **2** (1954), 286-294.
3. A. W. Tucker, *On generalised covariant differentiation*, Ann. of Math., **32** (1931), 451-460.
4. C. E. Weatherburn, *An introduction to Riemannian geometry and the tensor calculus*, Cambridge, 1938.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. L. ROYDEN

Stanford University
Stanford, California

E. HEWITT

University of Washington
Seattle 5, Washington

R. P. DILWORTH

California Institute of Technology
Pasadena 4, California

A. HORN*

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

C. E. BURGESS

H. BUSEMANN

H. FEDERER

M. HALL

P. R. HALMOS

V. GANAPATHY IYER

R. D. JAMES

M. S. KNEBELMAN

I. NIVEN

T. G. OSTROM

M. M. SCHIFFER

J. J. STOKER

G. SZEKERES

F. WOLF

K. YOSIDA

SPONSORS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
CALIFORNIA RESEARCH CORPORATION
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
HUGHES AIRCRAFT COMPANY

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. Manuscripts intended for the outgoing editors should be sent to their successors. All other communications to the editors should be addressed to the managing editor, Alfred Horn at the University of California, Los Angeles 24, California.

50 reprints of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, c/o University of California Press, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 10, 1-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

* During the absence of E. G. Straus.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION
COPYRIGHT 1956 BY PACIFIC JOURNAL OF MATHEMATICS

Pacific Journal of Mathematics

Vol. 6, No. 2

December, 1956

Louis Auslander, <i>Remark on the use of forms in variational calculations</i>	209
Hubert Spence Butts, Jr. and Henry B. Mann, <i>Corresponding residue systems in algebraic number fields</i>	211
L. Carlitz and John Herbert Hodges, <i>Distribution of matrices in a finite field</i>	225
Paul Civin and Bertram Yood, <i>Invariant functionals</i>	231
David James Dickinson, Henry Pollak and G. H. Wannier, <i>On a class of polynomials orthogonal over a denumerable set</i>	239
Bernard Friedman and Luna Mishoe, <i>Eigenfunction expansions associated with a non-self-adjoint differential equation</i>	249
Luna Mishoe and G. C. Ford, <i>On the uniform convergence of a certain eigenfunction series</i>	271
John W. Green, <i>Mean values of harmonic functions on homothetic curves</i>	279
Charles John August Halberg, Jr. and Angus E. Taylor, <i>On the spectra of linked operators</i>	283
Chuan Chih Hsiung, <i>Some integral formulas for closed hypersurfaces in Riemannian space</i>	291
Norman D. Lane, <i>Differentiable points of arcs in conformal n-space</i>	301
Louis F. McAuley, <i>A relation between perfect separability, completeness, and normality in semi-metric spaces</i>	315
G. Power and D. L. Scott-Hutton, <i>The slow shearing motion of a liquid past a semi-infinite plane</i>	327
A. C. Schaeffer, <i>Entire functions</i>	351
Edward Silverman, <i>An intrinsic inequality for Lebesgue area</i>	363
Choy-Tak Taam, <i>Asymptotic relations between systems of differential equations</i>	373
Ti Yen, <i>Quotient algebra of a finite AW^*-algebra</i>	389