

Pacific Journal of Mathematics

**MONOTONE COMPLETENESS OF NORMED SEMI-ORDERED
LINEAR SPACES**

SADAYUKI YAMAMURO

MONOTONE COMPLETENESS OF NORMED SEMI-ORDERED LINEAR SPACES

SADAYUKI YAMAMURO

Introduction. Let R be a continuous semi-ordered linear space, namely, a semi-ordered linear space where, for any sequence $x_\nu \geq 0$ ($\nu=1, 2, \dots$), $\bigcap_{\nu=1}^{\infty} x_\nu$ exists.¹ R is said to be a normed semi-ordered linear space, if a norm $\|x\|$ ($x \in R$) is defined and satisfies the condition:

$$|x| \leq |y| \quad \text{implies} \quad \|x\| \leq \|y\|$$

in addition to the usual conditions.

A norm $\|x\|$ ($x \in R$) on a normed semi-ordered linear space is said to be *monotone complete*, if, when $0 \leq x_\nu \uparrow_{\nu=1}^{\infty}$ and $\sup_{\nu \geq 1} \|x_\nu\| < +\infty$, there exists $\bigcup_{\nu=1}^{\infty} x_\nu$.

A norm on R is said to be *continuous*, if $x_\nu \downarrow_{\nu=1}^{\infty} 0$ implies $\lim_{\nu \rightarrow \infty} \|x_\nu\| = 0$ and *semi-continuous*, if $0 \leq x_\nu \uparrow_{\nu=1}^{\infty} x$ implies $\sup_{\nu \geq 1} \|x_\nu\| = \|x\|$. It is clear that continuity implies semi-continuity.

Kantorovitch [4] has proved that, if a norm on R is monotone complete and continuous, then it is complete, namely, R is a Banach lattice. Nakano [5; Theorem 31.7] has proved that, if a norm on R is monotone complete and semi-continuous, then the norm is complete, and, recently, Amemiya [1] has proved that, if a norm on R is monotone complete, it is complete.² In this connection, see also [2].

In this paper, we will consider several problems concerning monotone completeness and completeness of normed semi-ordered linear spaces and Nakano spaces.

1. Monotone completeness of normed semi-ordered linear spaces.
In this section, we will consider two problems.

As usual, let (c_0) be the set of all null-sequences of real numbers. This is a normed semi-ordered linear space by the usual ordering and

Received December 12, 1956. In revised form April 22, 1957.

¹ Namely, a conditionally σ -complete vector lattice. In this paper we use the terminology and notation of [5].

² In this paper, Amemiya also proved the following lemma: Let R be a monotone complete normed semi-ordered linear space. Then there exists a number $\gamma > 0$ such that $0 \leq x_\nu \uparrow_{\nu=1}^{\infty} x$ implies $\gamma \|x\| \leq \sup_{\nu \geq 1} \|x_\nu\|$.

the norm: $\|x\| = \sup_{\nu=1} |\xi_\nu|$ for $x = (\xi_\nu) \in (e_0)$. The fact that this norm is complete is well known. But, it is not monotone complete, because, for the sequence of elements:

$$e_1 = (1, 0, 0, \dots), \quad e_2 = (1, 1, 0, \dots), \quad e_3 = (1, 1, 1, 0, \dots), \quad \dots,$$

we have $0 \leq e_\nu \uparrow_{\nu=1}^\infty$ and $\sup_{\nu \geq 1} \|e_\nu\| \leq 1$, but $\bigcup_{\nu=1}^\infty e_\nu$ does not exist in the space (e_0) .

Among function spaces, we can also find an example of this type. Let $L_{\xi/t}^f$ be the set of all measurable functions $x(t)$ ($0 \leq t \leq 1$) such that

$$\int_0^1 |\xi x(t)|^{1/t} dt < +\infty \quad \text{for all} \quad \xi > 0.$$

Then $L_{\xi/t}^f$ is a Banach lattice by the norm:

$$\|x\| = \inf_{m(\xi x) \leq 1} \frac{1}{|\xi|} \quad \text{where} \quad m(x) = \int_0^1 |x(t)|^{1/t} dt,$$

but this norm is not monotone complete.

In § 1.1, we will state a necessary and sufficient condition in order that a complete norm be monotone complete.

It is well known that every (norm) closed subset of a Banach lattice is also complete. But, we have a monotone complete semi-ordered linear space which contains a closed, but not monotone complete subspace. Namely, let $L_{1/t}$ be the set of all measurable functions $x(t)$ ($0 \leq t \leq 1$) such that

$$\int_0^1 |\xi x(t)|^{1/t} dt < +\infty \quad \text{for some} \quad \xi > 0.$$

This is a monotone complete normed semi-ordered linear space and $L_{\xi/t}^f$ is a (norm) closed subspace of $L_{1/t}$.

In § 1.2, we will state a necessary and sufficient condition in order that every closed subspace of a monotone complete semi-ordered linear space be monotone complete.

1.1. Let R be a continuous semi-ordered linear space. A sequence x_ν ($\nu=1, 2, \dots$) is said to be *bounded*, if there exists an element $x \in R$ such that $x_\nu \leq x$ ($\nu=1, 2, \dots$). If $0 \leq x_\nu \uparrow_{\nu=1}^\infty$ and this sequence is not bounded, then we write $0 \leq x_\nu \uparrow_{\nu=1}^\infty + \infty$.

DEFINITION. R is said to be *K-bounded* (bounded in the sense of Kantorovitch), if $0 \leq x_\nu \uparrow_{\nu=1}^\infty + \infty$ implies we can find a sequence of real

numbers ξ_ν ($\nu=1, 2, \dots$) such that $\xi_\nu \downarrow_{\nu=1}^\infty 0$ and the sequence $\xi_\nu x_\nu$ ($\nu=1, 2, \dots$) is not bounded.

DEFINITION. R is said to be K^2 -bounded, if $0 \leqq x_{\mu,\nu} \uparrow_{\mu=1}^\infty +\infty$ for every ν implies we can find a sequence of indices μ_ν ($\nu=1, 2, \dots$) such that the sequence $x_{\mu_\nu,\nu}$ ($\nu=1, 2, \dots$) is not bounded.

These concepts were introduced by Kantorovitch [4]. It is easily seen that K^2 -boundedness implies K -boundedness. If R is reflexive in the sense of [5] § 24, then it is easily seen that R is K -bounded. Therefore, for any R , its conjugate space is always K -bounded.

The K -boundedness can be expressed in other ways, namely, the following three conditions are mutually equivalent:

- (1) R is K -bounded;
- (2) if $0 \leqq x_\nu \uparrow_{\nu=1}^\infty$ and $\sum_{\nu=1}^\infty \xi_\nu x_\nu$ is order-convergent for all sequences (ξ_ν) with $\sum_{\nu=1}^\infty |\xi_\nu| < +\infty$, then the sequence x_ν ($\nu=1, 2, \dots$) is bounded;
- (3) if $x_\nu \geqq 0$ and $\sum_{\nu=1}^\infty \xi_\nu x_\nu$ is order-convergent for all sequences (ξ_ν) with $\xi_\nu \downarrow_{\nu=1}^\infty 0$, then $\sum_{\nu=1}^\infty x_\nu$ is order-convergent.

For example, we will prove that (1) implies (2). Let $0 \leqq x_\nu \uparrow_{\nu=1}^\infty +\infty$. Then there exists a sequence of real numbers $\xi_\nu \downarrow_{\nu=1}^\infty 0$ such that $\xi_\nu x_\nu$ ($\nu=1, 2, \dots$) are not bounded. Since

$$x_\nu = \sum_{\mu=2}^\nu (x_\mu - x_{\mu-1}) + x_1 \uparrow_{\nu=1}^\infty +\infty,$$

and

$$\xi_\nu x_\nu \leqq \sum_{\mu=2}^\nu \xi_\mu (x_\mu - x_{\mu-1}) + \xi_1 x_1,$$

the sequence:

$$\sum_{\mu=1}^\nu (\xi_\mu - \xi_{\mu+1}) x_\mu \quad (\nu=1, 2, \dots)$$

is not bounded and

$$\sum_{\mu=1}^\infty |\xi_\mu - \xi_{\mu+1}| < +\infty.$$

This is inconsistent with the hypothesis of (2).

THEOREM 1.1. *Let R be a normed semi-ordered linear space. Then the following three conditions are mutually equivalent;*

- (1) *The norm on R is monotone complete;*
- (2) *the norm is complete and R is K -bounded;*
- (3) *the norm is complete and R is K^2 -bounded.*

Proof. We have only to prove that (2) implies (1).

Let $0 \leq x_\nu \uparrow_{\nu=1}^\infty$ and $\sup_{\nu \geq 1} \|x_\nu\| < +\infty$. Then, for any sequence of numbers $\xi_\nu > 0$ ($\nu=1, 2, \dots$) such that $\sum_{\nu=1}^\infty \xi_\nu < +\infty$, we have $\sum_{\nu=1}^\infty \xi_\nu \|x_\nu\| < +\infty$. Since the norm is complete by assumption, $\sum_{\nu=1}^\infty \xi_\nu x_\nu$ is convergent in norm, and so, in order convergence. Therefore, x_ν ($\nu=1, 2, \dots$) is bounded, because R is K -bounded.

1.2. Let R be a continuous semi-ordered linear space. For any element $p \geq 0$ and for all $x \geq 0$, the projector $[p]$ is defined as

$$[p]x = \bigcup_{\nu=1}^\infty (x \cap \nu p).$$

$[p] \geq [q]$ means $[p]x \geq [q]x$ for any $x \geq 0$.

Let R be a normed semi-ordered linear space. A norm $\|x\|$ on R is continuous if and only if $x \geq 0$ and $[p_\nu] \downarrow_{\nu=1}^\infty 0$ implies $\lim_{\nu \rightarrow \infty} \|[p_\nu]x\| = 0$ ([Nakano] Theorem 30.8). We will call a subset A of R monotone complete, if $0 \leq x_\nu \uparrow_{\nu=1}^\infty$ and $\sup_{\nu \geq 1} \|x_\nu\| < +\infty$ for $x_\nu \in A$ implies $\bigcup_{\nu=1}^\infty x_\nu \in A$.

If a norm on R is monotone complete and continuous, then every (norm) closed subset is monotone complete in the sense described above. Here, we will prove the converse. A subset A is said to be semi-normal, if $x \in A$, $|y| \leq |x|$ implies $y \in A$.

THEOREM 1.2. *Let R be a normed semi-ordered linear space and suppose every (norm) closed, semi-normal subset of R is monotone complete. Then the norm is continuous.*

Proof. Let us assume that there exist $[p_\nu]$ ($\nu=1, 2, \dots$) and $x_0 \in R$ such that $[p_\nu] \downarrow_{\nu=1}^\infty 0$ and $\lim_{\nu \rightarrow \infty} \|[p_\nu]x_0\| \geq \varepsilon$ for some $\varepsilon > 0$. Then the least closed set A containing all $x \in R$ such that $\lim_{\nu \rightarrow \infty} \|[p_\nu]x\| = 0$ is semi-normal and $(1 - [p_\nu])x_0 \in A$. On the other hand,

$$0 \leq (1 - [p_\nu])x_0 \uparrow_{\nu=1}^\infty x_0 \quad \text{and} \quad \|(1 - [p_\nu])x_0\| \leq \|x_0\|.$$

Therefore, since A is monotone complete, $x_0 \in A$. This is inconsistent with the definition of A .

2. Monotone completeness of Nakano spaces. It will be necessary to state here the definition and several properties of Nakano spaces.

A semi-ordered linear space is said to be *universally continuous*, if for any system of positive elements x_λ ($\lambda \in \Lambda$) there exists $\bigcap_{\lambda \in \Lambda} x_\lambda$. A *Nakano space* is a universally continuous semi-ordered linear space where a functional $m(x)$ ($x \in R$) is defined and satisfies the following conditions:

- (1) $0 \leq m(x) \leq +\infty$ ($x \in R$);
- (2) for any $x \in R$ we can find a number $\xi > 0$ such that $m(\xi x) < +\infty$;
- (3) if $m(\xi x) = 0$ for every $\xi > 0$, then $x = 0$;
- (4) $|x| \leq |y|$ implies $m(x) \leq m(y)$;
- (5) $m\left(\frac{\xi + \eta}{2}x\right) \leq \frac{1}{2}\{m(\xi x) + m(\eta x)\}$ for numbers $\xi, \eta > 0$ and for every

element $x \in R$;

- (6) $|x| \wedge |y| = 0$ implies $m(x+y) = m(x) + m(y)$;
- (7) $0 \leq x_\lambda \uparrow_{\lambda \in \Lambda} x$ implies $m(x) = \sup_{\lambda \in \Lambda} m(x_\lambda)$.

This functional $m(x)$ is called a *modular* on the Nakano space R . In the Nakano space R , we can define two kinds of norms:

the first norm:
$$\|x\| = \inf_{\xi > 0} \frac{1 + m(\xi x)}{\xi};$$

the second norm:
$$\|x\| = \inf_{m(\xi x) \leq 1} \frac{1}{|\xi|}.$$

It is easily seen that $\|x\| \leq \|x\| \leq 2\|x\|$. The modular is said to be complete or monotone complete, if these norms are complete or monotone complete. Namely, a modular m on R is said to be monotone complete, if, when $0 \leq x_\nu \uparrow_{\nu=1}^\infty$ and $\sup_{\nu \geq 1} m(x_\nu) < +\infty$, then there exists $\bigcup_{\nu=1}^\infty x_\nu$.

A modular m is said to be *simple*, if $m(x) = 0$ implies $x = 0$. If m is simple, we can define in R a convergence by this modular. Namely, a sequence x_ν ($\nu = 1, 2, \dots$) is said to be *modular-convergent* to $x \in R$, if $\lim_{\nu \rightarrow \infty} m(x_\nu - x) = 0$. If a sequence x_ν ($\nu = 1, 2, \dots$) is convergent to $x \in R$ by the norms defined above, then it is modular-convergent to the same limit. But the converse is not always true. In order that the modular-convergence be equivalent to the norm convergence, it is necessary and sufficient that the modular is *uniformly simple*: $\inf_{0 \neq x \in R} m\left(\xi \frac{x}{\|x\|}\right) > 0$ for any $\xi > 0$ ([5] Theorem 48.1)

The norms defined above are not always continuous. If the modular

is *finite*, namely, $m(x) < +\infty$ for every $x \in R$, then the norms are continuous ([5] Theorem 44.4)

A modular m is said to be *uniformly finite*, if $\sup_{0+x \in R} m\left(\xi \frac{x}{\|x\|}\right) < +\infty$ for every $\xi > 0$. It is clear that uniform finiteness is stronger than finiteness.³

2.1. In this section, we will consider the relations between monotone completeness and completeness of Nakano spaces. In the sequel, let R be a Nakano space and $m(x)$ ($x \in R$) be its modular.

The following lemma is a generalization of the essential part of Kalugyna's results [3].

LEMMA 2.1. *If m is monotone complete, simple, and its norms are continuous, then m is uniformly simple.*

Proof. If m is not uniformly simple, we can find a sequence $x_\nu \geq 0$ ($\nu = 1, 2, \dots$) such that $\lim_{\nu \rightarrow \infty} m(x_\nu) = 0$ and $\|x_\nu\| \geq \varepsilon > 0$ for all ν . Hence, we can select a subsequence x_{ν_μ} ($\mu = 1, 2, \dots$) such that

$$m(x_{\nu_\mu}) \leq 1/2^\mu .$$

Then, for the elements:

$$y_{\mu,\lambda} = x_{\nu_\mu} \cup x_{\nu_{\mu+1}} \cup \dots \cup x_{\nu_{\mu+\lambda}} ,$$

we have

$$\begin{aligned} m(y_{\mu,\lambda}) &\leq m(x_{\nu_\mu}) + m(x_{\nu_{\mu+1}}) + \dots + m(x_{\nu_{\mu+\lambda}}) \\ &\leq \frac{1}{2^\mu} + \dots + \frac{1}{2^{\mu+\lambda}} . \end{aligned}$$

Namely, we have $y_{\mu,\lambda} \uparrow_{\lambda=1}^\infty$ and $\sup_{\lambda \geq 1} m(y_{\mu,\lambda}) < +\infty$. Since m is monotone complete, there exist y_μ ($\mu = 1, 2, \dots$) such that $y_\mu = \bigcup_{\lambda=1}^\infty y_{\mu,\lambda}$ and $m(y_\mu) \leq 1/2^{\mu+1}$.

It is clear that $y_\mu \downarrow_{\mu=1}^\infty$. On the other hand, for any $x \geq 0$ such that $x \leq y_\mu$ ($\mu = 1, 2, \dots$), we have

$$m(y_\mu - x) \leq m(y_\mu) , \quad \text{thus,} \quad \lim_{\mu \rightarrow \infty} m(y_\mu - x) = 0 .$$

³ More details of the theory of Nakano spaces are given in [5]. As examples of Nakano spaces, we cite two representative types. The first is an Orlicz space. The second is the space $L_{p(t)}(p(t) \geq 1)$, namely, the set of measurable functions $x(t)$ ($0 \leq t \leq 1$) such that $\int_0^1 |\xi x(t)|^{p(t)} dt$ is finite for some $\xi > 0$. Here $p(t)$ is a measurable function on $0 \leq t \leq 1$.

Therefore,

$$\begin{aligned}
 m\left(\frac{1}{2}x\right) &= m\left(\frac{1}{2}(y_\mu - x) + \frac{1}{2}y_\mu\right) \\
 &\leq \frac{1}{2}\left\{m(y_\mu - x) + m(y_\mu)\right\} \rightarrow 0 \quad (\mu \rightarrow \infty),
 \end{aligned}$$

that is to say, $m\left(\frac{1}{2}x\right) = 0$. Since m is simple, $x = 0$. This means that $y_\mu \downarrow_{\mu=1}^\infty 0$. As the norm is continuous, we have $\lim_{\mu \rightarrow \infty} \|y_\mu\| = 0$, which contradicts the assumption, because

$$\|y_\mu\| \geq \|x_{\nu_\mu}\| \geq \varepsilon.$$

Therefore, m is uniformly simple.

The next two lemmas constitute the converse of the above.

LEMMA 2.2. *If m is uniformly simple, then its norms are continuous.*

Proof. Let $x_\nu \downarrow_{\nu=1}^\infty 0$. Then there exists a number $\xi > 0$ such that $m(\xi x_\nu) < +\infty$ for all ν . For the elements $y_\nu = \xi x_1 - \xi x_\nu$, since $y_\nu \geq 0$ and $\xi x_\nu \geq 0$, we have

$$m(y_\nu + \xi x_\nu) \geq m(y_\nu) + m(\xi x_\nu),$$

so,

$$m(\xi x_\nu) \leq m(y_\nu + \xi x_\nu) - m(y_\nu) = m(\xi x_1) - m(y_\nu).$$

On the other hand, we have $m(\xi x_1) = \sup_{\nu \geq 1} m(y_\nu)$, because $0 \leq y_\nu \uparrow_{\nu=1}^\infty \xi x_1$. Therefore, $\lim_{\nu \rightarrow \infty} m(\xi x_\nu) = 0$, and hence it follows that $\lim_{\nu \rightarrow \infty} \|x_\nu\| = 0$, because m is uniformly simple.

LEMMA 2.3. *If m is uniformly simple and its norms are complete, then m is monotone complete.*

Proof. Let $0 \leq x_\nu \uparrow_{\nu=1}^\infty$ and $\sup_{\nu \geq 1} m(x_\nu) < +\infty$. Then

$$m(x_\nu - x_\mu) \leq m(x_\nu) - m(x_\mu) \quad (\nu \geq \mu),$$

and hence, we have

$$\lim_{\nu, \mu \rightarrow \infty} m(x_\nu - x_\mu) = 0.$$

Since m is uniformly simple, we have

$$\lim_{\nu, \mu \rightarrow \infty} \|x_\nu - x_\mu\| = 0,$$

so that there exists an element $x \in R$ such that $\lim_{\nu \rightarrow \infty} \|x_\nu - x\| = 0$. For this x , it is easily seen that $x = \bigcup_{\nu=1}^{\infty} x_\nu$, which shows that m is monotone complete.

From these lemmas, we obtain the following theorem:

THEOREM 2.1. *A modular on a Nakano space is monotone complete, simple, and its norms are continuous, if and only if it is uniformly simple and complete.*

Next, we will consider the case when m is finite.

DEFINITION. A modular $m(x)$ ($x \in R$) is said to be totally finite, if $0 \leq x_\nu \uparrow_{\nu=1}^{\infty}$ and $\sup_{\nu \geq 1} m(x_\nu) < +\infty$ implies $\sup_{\nu \geq 1} m(\xi x_\nu) < +\infty$ for every $\xi > 0$.

LEMMA 2.4. *If m is monotone complete and finite, then it is totally finite.*

Proof. $0 \leq x_\nu \uparrow_{\nu=1}^{\infty}$ and $\sup_{\nu \geq 1} m(x_\nu) < +\infty$. Then, since m is monotone complete, there exists $x \in R$ such that $x = \bigcup_{\nu=1}^{\infty} x_\nu$. Therefore $\xi x = \bigcup_{\nu=1}^{\infty} \xi x_\nu$ for every $\xi > 0$. Hence it follows that $m(\xi x) = \sup_{\nu \geq 1} m(\xi x_\nu) < +\infty$, because m is finite.

LEMMA 2.5. *If m is totally finite and complete, then it is monotone complete.*

Proof. Let $0 \leq x_\nu \uparrow_{\nu=1}^{\infty}$ and $\sup_{\nu \geq 1} m(x_\nu) < +\infty$. Then, by the assumption, we have $\sup_{\nu \geq 1} m(\xi x_\nu) < +\infty$ for every $\xi > 0$. Since

$$m(\xi x_\nu - \xi x_\mu) \leq m(\xi x_\nu) - m(\xi x_\mu) \quad (\nu \geq \mu),$$

we have $\lim_{\nu, \mu \rightarrow \infty} m(\xi x_\nu - \xi x_\mu) = 0$ for every $\xi > 0$, therefore we have

$$\lim_{\nu, \mu \rightarrow \infty} \|x_\nu - x_\mu\| = 0.$$

Hence, there exists an element $x \in R$ such that $\lim_{\nu \rightarrow \infty} \|x_\nu - x\| = 0$. Therefore, we have $x = \bigcup_{\nu=1}^{\infty} x_\nu$, which shows that m is monotone complete.

Thus we obtain the following theorem:

THEOREM 2.2. *A modular on a Nakano space is monotone complete and finite if and only if it is totally finite and complete.*

REMARK. It is easily seen that uniform finiteness implies total finiteness and the latter implies finiteness. The converses are not always true. In fact, $L_{1/t}^f$ is a finite Nakano space by the following modular:

$$m(x) = \int_0^1 |x(t)|^{1/t} dt \quad \text{for } x(t) \in L_{1/t}^f .$$

But, this is not totally finite, because, if it were totally finite, then, by Theorem 2.2, it would be monotone complete, which is impossible. Next, let $f_\nu(\xi)$ ($\nu=1, 2, \dots$) be a sequence of convex functions such that

$$f_\nu(\xi) = \begin{cases} \xi & \text{if } 0 \leq \xi \leq 1 ; \\ \nu(\xi-1)+1 & \text{if } \xi > 1 . \end{cases}$$

Then, the space⁴ $U(f_1, f_2, \dots)$ with the modular

$$m(x) = \sum_{\nu=1}^{\infty} f_\nu(|\xi_\nu|) \quad \text{for } x = (\xi_\nu)$$

is totally finite, but not uniformly finite. To see this, we need only take the elements:

$$e_1 = (1, 0, 0, \dots), \quad e_2 = (0, 1, 0, \dots), \quad e_3 = (0, 0, 1, 0, \dots), \quad \dots .$$

It is easily proved that $\|e_\nu\| = 1$ and $m(2e_\nu) = \nu + 1 \rightarrow +\infty$. But, this sequence space is uniformly simple by Theorem 2.1. The relations between uniform simplicity and uniform finiteness were considered by my colleagues. If a modular on a Nakano space is uniformly finite and simple, then, by considering the monotone completion and applying Theorem 2.1, we can prove that it is uniformly simple. On the other hand, T. Shimogaki has proved in an unpublished paper that, if a modular is uniformly simple and the space has no atomic elements, then it is uniformly finite.

2.2. In this section, we will consider relations between monotone completeness and finiteness.

An element x is said to be *finite*, if $m(\xi x) < +\infty$ for every $\xi > 0$. The set of all finite elements is called a *finite manifold* of R and denoted by F . F is a (norm) closed subspace of R and the norms are continuous in F ([5] Theorem 44.5.). If the norms are continuous in R and m is monotone complete, then F is *universally monotone complete*, that is, if $0 \leq x_\lambda \uparrow_{\lambda \in \Lambda}$ and $\sup_{\lambda \in \Lambda} m(x_\lambda) < +\infty$ then there exists $\bigcup_{\lambda \in \Lambda} x_\lambda$.

m is said to be *almost finite*, if F is complete in R (that is, if $|x \cap y| = 0$ for all $y \in F$, then $x = 0$).

⁴ For the definition of this sequence space, see [6].

THEOREM 2.3. *If m is almost finite and monotone complete, then m is finite if and only if F is, as a space, universally monotone complete.*

Proof. We need only prove the sufficiency. For any $x \in R$, since m is almost finite, there exists a system of projectors $[p_\lambda] \uparrow_{\lambda \in \Lambda} [x]$ such that $[p_\lambda]x \in F$, and there exists a number $\xi > 0$ such that $m(\xi x) < +\infty$. Therefore we have

$$\bigcup_{\lambda \in \Lambda} \xi [p_\lambda]x = \xi x \in F,$$

since m is monotone complete. Hence it follows that m is finite.

THEOREM 2.4. *If m is almost finite, monotone complete and separable in its norm topology, then m is finite.*

Proof. It is well known that if m is almost finite and norms are continuous, then m is finite. Therefore, we need only prove that if m is monotone complete and separable, then its norms are continuous.

For this purpose, let us suppose that there exists an element $x \geq 0$ and a sequence of projectors $[p_\nu] \downarrow_{\nu=1}^{\infty} 0$ such that

$$\inf_{\nu \geq 1} \|[p_\nu]x\| > \varepsilon \quad \text{for some} \quad \varepsilon > 0.$$

Then, by Amemiya's lemma, we can find a number $\xi > 0$ such that

$$\lim_{\nu \rightarrow \infty} \|[p_\mu]a - [p_\nu]a\| \geq \xi \|[p_\mu]a\| > \xi \varepsilon > 0,$$

and here, we can select μ_ν ($\nu = 1, 2, \dots$) such that

$$\|[p_{\mu_\nu}]x - [p_{\mu_{\nu+1}}]x\| > \xi \varepsilon.$$

Putting $p_\nu = [p_{\mu_\nu}]x - [p_{\mu_{\nu+1}}]x$ ($\nu = 1, 2, \dots$), we see easily that

$$p_\nu \geq 0, \quad p_\nu \cap p_\lambda = 0 (\nu \neq \lambda) \quad \text{and} \quad \|p_\nu\| > \xi \varepsilon,$$

and, for any subsequence p_{ν_λ} ($\lambda = 1, 2, \dots$), we have

$$\sum_{\lambda=1}^{\infty} p_{\nu_\lambda} \leq x.$$

Moreover, the set of all such sequences is not denumerable and

$$\left\| \sum_{\lambda=1}^{\infty} p_{\nu_\lambda} - \sum_{\rho=1}^{\infty} p_{\nu_\rho} \right\| > \xi \varepsilon$$

for different sequences $\{p_{\nu_\lambda}\}$ and $\{p_{\nu_\rho}\}$. This contradicts the separability. Therefore, norms are continuous and the proof is established.

REMARK. In order that m be finite, it is necessary and sufficient that its norms be continuous and all atomic elements belong to F .

REFERENCES

1. I. Amemiya, *A Generalization of Riesz-Fischer's Theorem*, Jour. Math. Soc. Japan, **5** (1953), 353-354.
2. I. Halperin and W. A. J. Luxemburg, *The Riesz-Fischer completeness theorem for function spaces and vector lattices*, Trans. Royal Soc. Canada, **50** (1956), 33-39.
3. E. P. Kalugyna, *Class L_ϕ as a convex set of functions* (Russian), Doklady A. N. URSS, **98** (1954), 13-16.
4. L. V. Kantorovitch, *Lineare halbgeordnete Räume*, Recueil Math., **44** (1937), 121-165.
5. H. Nakano, *Modulated Semi-ordered Linear Spaces*, 1950.
6. S. Yamamuro, *Modulated Sequence Spaces*, Jour. Fac. Science Hokkaido Univ., **13** (1954), 1-12.

HOKKAIDŌ UNIVERSITY AND
THE INSTITUTE FOR ADVANCED STUDY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. L. ROYDEN
Stanford University
Stanford, California

R. A. BEAUMONT
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

E. G. STRAUS
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
C. E. BURGESS
M. HALL
E. HEWITT

A. HORN
V. GANAPATHY IYER
R. D. JAMES
M. S. KNEBELMAN

L. NACHBIN
I. NIVEN
T. G. OSTROM
M. M. SCHIFFER

G. SZEKERES
F. WOLF
K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
THE RAMO-WOOLDRIDGE CORPORATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. G. Straus at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 10, 1-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 7, No. 4

April, 1957

Robert Geroge Buschman, <i>A substitution theorem for the Laplace transformation and its generalization to transformations with symmetric kernel</i>	1529
S. D. Conte, <i>Numerical solution of vibration problems in two space variables</i>	1535
Paul Dedecker, <i>A property of differential forms in the calculus of variations</i>	1545
H. Delange and Heini Halberstam, <i>A note on additive functions</i>	1551
Jerald L. Ericksen, <i>Characteristic direction for equations of motion of non-Newtonian fluids</i>	1557
Avner Friedman, <i>On two theorems of Phragmén-Lindelöf for linear elliptic and parabolic differential equations of the second order</i>	1563
Ronald Kay Gettoor, <i>Additive functionals of a Markov process</i>	1577
U. C. Guha, <i>(γ, k)-summability of series</i>	1593
Alvin Hausner, <i>The tauberian theorem for group algebras of vector-valued functions</i>	1603
Lester J. Heider, <i>T-sets and abstract (L)-spaces</i>	1611
Melvin Henriksen, <i>Some remarks on a paper of Aronszajn and Panitchpakdi</i>	1619
H. M. Lieberstein, <i>On the generalized radiation problem of A. Weinstein</i>	1623
Robert Osserman, <i>On the inequality $\Delta u \geq f(u)$</i>	1641
Calvin R. Putnam, <i>On semi-normal operators</i>	1649
Binyamin Schwarz, <i>Bounds for the principal frequency of the non-homogeneous membrane and for the generalized Dirichlet integral</i>	1653
Edward Silverman, <i>Morrey's representation theorem for surfaces in metric spaces</i>	1677
V. N. Singh, <i>Certain generalized hypergeometric identities of the Rogers-Ramanujan type. II</i>	1691
R. J. Smith, <i>A determinant in continuous rings</i>	1701
Drury William Wall, <i>Sub-quasigroups of finite quasigroups</i>	1711
Sadayuki Yamamuro, <i>Monotone completeness of normed semi-ordered linear spaces</i>	1715
C. T. Rajagopal, <i>Simplified proofs of "Some Tauberian theorems" of Jakimovski: Addendum and corrigendum</i>	1727
N. Aronszajn and Prom Panitchpakdi, <i>Correction to: "Extension of uniformly continuous transformations in hyperconvex metric spaces"</i>	1729
Alfred Huber, <i>Correction to: "The reflection principle for polyharmonic functions"</i>	1731