

# Pacific Journal of Mathematics

**REMARKS ON THE MAXIMUM PRINCIPLE FOR PARABOLIC  
EQUATIONS AND ITS APPLICATIONS**

AVNER FRIEDMAN

# REMARKS ON THE MAXIMUM PRINCIPLE FOR PARABOLIC EQUATIONS AND ITS APPLICATIONS

AVNER FRIEDMAN

**Introduction.** In [3] Nirenberg has proved maximum principles, both weak and strong, for parabolic equations. In § 1 of this paper we give a generalization of his strong maximum principle (Theorem 1). Hopf [2] and Olainik [4] have proved that if  $Lu \geq 0$  and  $L$  is a linear elliptic operator of the second order, if the coefficient of  $u$  in  $L$  is non-positive, and if  $u$  ( $\neq \text{const.}$ ) assumes its positive maximum at a point  $P'$  (which necessarily belongs to the boundary) then  $\partial u / \partial \nu < 0$ , where  $\nu$  is the inwardly directed normal. In § 2 we extend this result to parabolic operators (Theorem 2). A further discussion of the assumptions made in Theorem 2 is given in § 3. Application of Theorem 2 to the Neumann problem is given in § 4. In § 5 we apply the weak maximum principle to prove a uniqueness theorem for certain nonlinear parabolic equations with nonlinear boundary conditions, and thus extend the special case considered by Ficken [1]. An even more special case arises in the theory of diffusion (for references, see [1]).

## 1. Consider the operator

$$(1) \quad Lu \equiv \sum_{i,j=1}^n a_{ij}(x, t) \frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_{i=1}^n a_i(x, t) \frac{\partial u}{\partial x_i} + a(x, t)u - \frac{\partial u}{\partial t}$$

with  $a(x, t) \leq 0$ . Here,  $(x, t) = (x_1, \dots, x_n, t)$  varies in the closure  $\bar{D}$  of a given  $(n+1)$ -dimensional domain  $D$ . Assume that  $L$  is parabolic in  $\bar{D}$ , that is, for every real vector  $\lambda \neq 0$  and for every  $(x, t) \in \bar{D}$  we have

$$\sum a_{ij}(x, t) \lambda_i \lambda_j > 0.$$

All the coefficients of  $L$  are assumed to be continuous in  $\bar{D}$  and  $u$  is assumed to be continuous in  $\bar{D}$  and to have a continuous  $t$ -derivative and continuous second  $x$ -derivatives in  $D$ . From [3; Th. 5] it follows that, under the above assumptions, if  $Lu \geq 0$  and if  $u$  assumes its positive maximum at an interior point  $P^0$ , then  $u \equiv \text{const.}$  in  $S(P^0)$ . Here,  $S(P^0)$  denotes the set of all points  $Q$  in  $D$  which can be connected to  $P^0$  by a simple continuous curve in  $D$  along which the coordinate  $t$  is non-decreasing from  $Q$  to  $P^0$ . In the following theorem we consider the case

---

Received October 29, 1957. Prepared under ONR Contract Nonr-908 (09), NR 041 037 with Indiana University.

in which  $P^0$  is a boundary point of  $D$ . We may assume that  $P^0$  is the origin. Let  $t=\varphi(x)$  be the equation of the boundary of  $D$  near  $P^0$ . Assume that  $t=0$  is the tangent hyperplane to the boundary of  $D$  at  $P^0$ . Therefore  $\partial\varphi/\partial x_i|_{P^0}=0$ . Let  $D$  be on the side  $t<\varphi(x)$ .

**THEOREM 1.** *If  $Lu \geq 0$  in  $D$ , if  $u$  assumes its positive maximum  $M$  at  $P^0$ , if*

$$(2) \quad \lim_{P \rightarrow P^0} \frac{\partial u(P)}{\partial x_i} = 0, \lambda \equiv \lim_{P \rightarrow P^0} \sum a_{ij}(P) \frac{\partial^2 u(P)}{\partial x_i \partial x_j} \leq 0 \quad P \in D$$

and if

$$(3) \quad 1 + \sum a_{ij} \frac{\partial^2 \varphi}{\partial x_i \partial x_j} \Big|_{P^0} > 0 \quad \varphi \in C''$$

then  $u \equiv M$  in  $S(P^0)$ .

**REMARK 1.** Without making any use of (3) one can deduce the following :

Put  $\mu \equiv \limsup_{P \rightarrow P^0} \frac{\partial u(P)}{\partial t}$  ( $P \in D$ ), then  $\mu \geq 0$  since  $\mu < 0$  will contradict  $u(P^0) \geq u(P)$ . Letting  $P \rightarrow P^0$  in  $Lu(P) \geq 0$  and using (2), we obtain  $\lambda + a(P^0)M - \mu \geq 0$ , from which it follows that  $\lambda \geq 0$ . Since, by (2),  $\lambda \leq 0$ , we conclude that  $\lambda = 0$ . Hence  $a(P^0)M - \mu \geq 0$ , from which it follows that  $\mu \leq 0$  and, therefore, (since  $\mu \geq 0$ )  $\mu = 0$ . We also get  $a(P^0) = 0$ .

**REMARK 2.** The assumptions (2) and (3) can be verified if we assume that  $\varphi(x) = o(|x|^2)$  and that  $u$  belongs to  $C''$  in the closure of the domain  $V \cap \{t < 0\}$ , where  $V$  is some neighborhood of  $P^0$ . Indeed, by making an appropriate orthogonal transformation we can assume that  $a_{ij}(P^0) = \delta_{ij}$ . By the mean value theorem we have

$$u(x, t) - u(0, 0) = \sum x_i \frac{\partial}{\partial x_i} u(\tilde{x}, \tilde{t}) + t \frac{\partial}{\partial t} u(\tilde{x}, \tilde{t}).$$

Taking  $(x, t) \in \overline{D} \cap V \cap \{t < 0\}$  such that  $|t| = o(|x|)$  and noting that  $u(x, t) \leq u(0, 0)$ , one can show that  $\partial u(P^0)/\partial x_i = 0$ . Noting that  $\varphi(x) = o(|x|^2)$  and expanding  $[u(x, t) - u(0, 0)]$  in terms of the first and second derivatives of  $u$ , one can show that  $\partial^2 u(P^0)/\partial x_i^2 \leq 0$ , and (2) is thereby proved. The proof of (3) is immediate.

**PROOF OF THEOREM 1.** For simplicity we shall prove the theorem only in case  $n=1$ ; the proof of the general case is analogous.  $Lu$  takes the form

$$(4) \quad Lu \equiv A \frac{\partial^2 u}{\partial x^2} + a \frac{\partial u}{\partial x} + cu - \frac{\partial u}{\partial t} \quad c \leq 0, A > 0.$$

From the strong maximum principle [3; Th. 5] it follows that all we need to prove is that  $u(P) \equiv M$  if  $P \in V' \cap S(P^0)$  where  $V'$  is some neighborhood of  $P^0$ .

There are two possibilities: Either there exists a sequence  $\{P^k\}$  such that  $P^k \in S(P^0)$ ,  $P^k \rightarrow P^0$ ,  $u(P^k) = M$ , or there exists a neighborhood  $V = \{x^2 + t^2 < R^2\}$  of  $P^0$  such that  $u(P) < M$  for all  $P \in V \cap S(P^0)$ ,  $P \neq P^0$ . In the first case we can use [3; Th. 5] to conclude that  $u(P) \equiv M$  if  $P \in V' \cap S(P^0)$  where  $V'$  is some neighborhood of  $P^0$  (since  $u(P) = M$  for all  $P \in S(P^k)$ ).

It remains therefore to consider the case in which  $u(P) < M$  for all  $P \in V \cap S(P^0)$ ,  $P \neq P^0$ . We shall prove that this case cannot occur by deriving a contradiction. Writing

$$\varphi(x) = Kx^2 + o(x^2),$$

we define a domain  $D_\delta$  ( $\delta > 0$ ) as the intersection of  $S(P^0)$  with the set of points  $(x, t)$  in  $V$  for which

$$t < \tilde{\varphi}(x) = (K - \delta)x^2.$$

If  $K < 0$  then, because of (3), we can choose  $\delta$  sufficiently small such that

$$(5) \quad 1 + A \frac{\partial^2}{\partial x^2} \tilde{\varphi}(x)|_{x=0} > 0.$$

If  $K \geq 0$ , we can obviously take  $\delta$  such that  $K - \delta < 0$  and such that (5) holds.

We now can take  $R$  sufficiently small such that  $\tilde{\varphi}(x) < \min(0, \varphi(x))$  for all  $(x, t)$  in  $D_\delta$ ,  $x \neq 0$ . Consequently,  $u(x, t) < M$  if  $t = \tilde{\varphi}(x)$ ,  $x \neq 0$ . The function  $h(x, t) = -t + \tilde{\varphi}(x)$  vanishes on  $t = \tilde{\varphi}(x)$  and is positive in  $D_\delta$ . Therefore, if  $\varepsilon > 0$  is sufficiently small, then  $v = u + \varepsilon h$  is smaller than  $M$  at all points on the boundary of  $D_\delta$  with the exception of  $P^0$ , where  $v(P^0) = M$ . Noting that  $\tilde{\varphi}'(0) = 0$  and using (5), we conclude that

$$Lh = 1 + A\tilde{\varphi}''(x) + a\tilde{\varphi}'(x) + ch > 0$$

if  $R$  has been chosen sufficiently small. Hence,  $Lv = Lu + \varepsilon Lh > 0$ . It follows that  $v$  cannot assume its positive maximum at interior points of  $D_\delta$  and, therefore, it assumes its maximum  $M$  at  $P^0$ . We thus obtain  $\partial v / \partial t \geq 0$  at  $P^0$  and, consequently,

$$\frac{\partial u}{\partial t} = \frac{\partial v}{\partial t} - \varepsilon \frac{\partial h}{\partial t} \geq \varepsilon > 0$$

(Here

$$\frac{\partial g}{\partial t} = \liminf_{t \rightarrow 0} \frac{g(0, 0) - g(0, t)}{-t}.$$

On the other hand, letting in (4)  $P \rightarrow P^0$  in an appropriate way and using (2) and the inequality  $Lu(P) \geq 0$ , we get

$$0 \leq \lim A(P) \frac{\partial^2 u(P)}{\partial x^2} + \lim a(P) \frac{\partial u(P)}{\partial x} + C(P^0)M - \limsup \frac{\partial u(P)}{\partial t} \leq \\ - \limsup \frac{\partial u(P)}{\partial t} .$$

We have thus obtained

$$\limsup_{P \rightarrow P^0} \partial u(P)/\partial t \leq 0 < \varepsilon \leq \partial u/\partial t .$$

This is however a contradiction (since

$$\frac{\partial u}{\partial t} = \lim_{t_k \rightarrow 0} \frac{\partial u(0, t_k)}{\partial t} \leq \limsup_{P \rightarrow P^0} \frac{\partial u(P)}{\partial t}$$

for an appropriate sequence  $\{t_k\}$ ), and the proof is completed.

REMARK (a) Consider the following example:  $n=1$ ,  $P^0=(0, 0)$  and  $D$  defined by

$$x^2 + t^2 < R, \quad t < \gamma_1 x, \quad t < \gamma_2 x \quad \gamma_1 > 0 > \gamma_2 .$$

The function  $u(x, t) = (t - \gamma_1 x)(\gamma_2 x - t)$  satisfies the following properties:  $u < 0$  in  $D$ ,  $u=0$  at  $P^0$ , and

$$Lu \equiv A \frac{\partial^2 u}{\partial x^2} + a \frac{\partial u}{\partial x} - \frac{\partial u}{\partial t} = -2A\gamma_1\gamma_2 + 0(|x| + |t|) \geq 0 ,$$

provided  $R$  is sufficiently small. Consequently, (3) and the second assumption in (2) are not satisfied and also the assertion of Theorem 1 is false.

REMARK (b). Consider now the case in which the tangent hyperplane at  $P^0$  is not of the form  $t = \text{const.}$ . We shall prove that in this case Theorem 1 is false. Take  $n=1$  and consider first the case in which  $D$  is defined by

$$x > 0, \quad x^2 + t^2 < R^2 .$$

If  $Lu \equiv \partial^2 u / \partial x^2 - \partial u / \partial t$ , then the function  $u(x, t) = -x$  takes its maximum in  $\bar{D}$  at  $P^0 = (0, 0)$ ,  $Lu = 0$ , but  $u \neq 0$  in  $S(P^0)$ .

Consider next the case in which  $\bar{D}$  is defined by

$$x > \alpha t, \quad x^2 + t^2 < R^2 .$$

and take  $Lu = \partial^2 u / \partial x^2 - \alpha \partial u / \partial x - \partial u / \partial t$ .

The transformation  $t'=t, x'=x-\alpha t$  carries the present case into the previous one.

Note that if the tangent hyperplane  $H$  at  $P^0$  is not the plane  $t=0$  and the axes are rotated so as to give  $H$  the equation  $t'=0$  (in new  $x', t'$  coordinate), then  $Lu$  loses the form (1), for  $u_{x't'}$  and  $u_{t't'}$  will appear in it.

REMARK (c). If in Theorem 1 the domain  $D$  is on the side  $t > \varphi(x)$ , then the theorem is false. Indeed, as a counter-example take  $u = -t$ , and  $D$  bounded from below by  $t=0$ .

## 2. Consider the linear operator

$$(6) \quad \begin{aligned} L'u = & \sum_{i,j=1}^n a_{ij}(x, t) \frac{\partial^2 u}{\partial x_i \partial x_j} + \sum_{i,j=1}^m b_{ij}(x, t) \frac{\partial^2 u}{\partial t_i \partial t_j} + \sum_{i=1}^n a_i(x, t) \frac{\partial u}{\partial x_i} \\ & + \sum_{i=1}^m b_i(x, t) \frac{\partial u}{\partial t_i} + a(x, t)u \end{aligned} \quad a(x, t) \leq 0,$$

where  $x = (x_1, \dots, x_n)$  and  $t = (t_1, \dots, t_m)$  vary in the closure of a given  $(n+m)$ -dimensional domain  $D$ . We assume that  $L'$  is elliptic in the variables  $x$  and parabolic in the variables  $t$ , that is, for every real vector  $\lambda \neq 0$ ,

$$(7) \quad \sum a_{ij} \lambda_i \lambda_j > 0, \quad \sum b_{ij} \lambda_i \lambda_j \geq 0.$$

All the coefficients appearing in (6) are assumed to be continuous in  $\bar{D}$  and  $u$  is assumed to be continuous in  $\bar{D}$  and to have a continuous  $t$ -derivative and continuous second  $x$ -derivatives in  $D$ . Under these assumptions, Nirenberg [3 ; Th. 2] has proved a weak maximum principle from which it follows that, if  $L'u \geq 0$  in  $D$  then  $u$  must assume its positive maximum on the boundary.

Let  $P^0 = (x^0, t^0)$  be a point on the boundary of  $D$  such that  $u(P^0) = M > 0$  is the maximum of  $u$  in  $\bar{D}$ . Assume that there exists a neighborhood  $V: |x - x^0|^2 + |t - t^0|^2 < R_0^2$  of  $P^0$  such that  $u(x, t) < M$  in  $V \cap D$ . We then can prove the following theorem.

**THEOREM 2.** *If there exists a sphere  $S: |x - x'|^2 + |t - t'|^2 < R^2$  passing through  $P^0$  and contained in  $\bar{D}$ , and if  $x^0 \neq x'$  then, under the assumptions made above (in particular,  $L'u \geq 0, u(x, t) < M$  in  $V \cap D$ ), every non-tangential derivative  $\partial u / \partial \tau$  at  $(x^0, t^0)$ , understood as the limit inferior of  $\Delta u / \Delta \tau$  along a non-tangential direction  $\tau$ , is negative.*

By a non-tangential direction we mean a direction from  $P^0$  into the interior of the sphere  $S$ .

REMARK (a). If  $a(x, t) \equiv 0$  then the assumption  $M > 0$  is superfluous.

REMARK (b). In § 3 we shall show that the assumption  $x^0 \neq x'$  is essential. We shall also discuss the case in which  $u(x, t)$  is not smaller than  $M$  at all the points of  $V \cap D$ .

*Proof.* For simplicity we give the proof in the case  $m = n = 1$ , so that

$$(8) \quad L'u \equiv A \frac{\partial^2 u}{\partial x^2} + B \frac{\partial^2 u}{\partial t^2} + a \frac{\partial u}{\partial x} + b \frac{\partial u}{\partial t} + cu \quad A > 0, B \geq 0, c \leq 0;$$

the proof of the general case is quite similar. Without loss of generality we can take  $(x', t') = (0, 0)$  and  $x^0 > 0$ . Furthermore, we may assume that, with the exception of  $P^0$ ,  $S$  lies in  $V \cap D$ , so that  $u(x, t) < M$  in  $S - P^0$ . Denote by  $C$  the intersection of  $S$  with the plane  $x > \delta$ , where  $0 < \delta < x^0$ . The function

$$h(x, t) = \exp(-\alpha(x^2 + t^2)) - \exp(-\alpha R^2)$$

satisfies the following properties:  $h = 0$  on the boundary of  $S$ ,  $h \geq 0$  in  $C$ ; if  $\alpha$  is large enough, then

$$L'h = \exp(-\alpha(x^2 + t^2))[4\alpha^2(Ax^2 + Bt^2) - 2\alpha(A + B + ax + bt) + c] - c \exp(-\alpha R^2) > 0.$$

(Here we used  $x > \delta > 0$ ,  $c \leq 0$ .)

If  $\varepsilon$  is sufficiently small, then the function  $v = u + \varepsilon h$  is smaller than  $M$  at all points of the boundary of  $C$  with the exception of  $P^0$ , where  $v(P^0) = M$ . Since  $L'v = L'u + \varepsilon L'h > 0$ ,  $v$  cannot assume its positive maximum in  $\bar{C}$  at the interior of  $C$  (since, otherwise, at such interior points  $L'v$  would be non-positive). Hence,  $v$  assumes its maximum at  $P^0$  and, consequently,  $\partial v / \partial \tau = \liminf (\Delta v / \Delta \tau) \leq 0$ . Since along the normal  $\nu$  (i. e., along the radius through  $P^0$ )  $\partial h / \partial \nu > 0$  and since along the tangential direction  $\sigma$   $\partial h / \partial \sigma = 0$ , it follows that  $\partial h / \partial \tau > 0$ . Using the definition of  $v$ , we conclude that  $\partial u / \partial \tau = \partial v / \partial \tau - \varepsilon \partial h / \partial \tau < 0$ , and the proof is completed.

*Added in proof.* Theorem 2 was recently and independently proved also by R. Viborni, *On properties of solutions of some boundary value problems for equations of parabolic type*, Doklady Akad. Nauk SSSR, 117 (1957), 563-565.

3. From now on we shall consider only parabolic operators of the form (1). Suppose the assumption  $u < M$  in  $V \cap D$ , made in Theorem 2, is replaced by  $u \leq M$ . If there exists a sequence of points  $\{P^k\}$  such

that  $P^k \rightarrow P^0$ ,  $P^k \in D$ ,  $P^k = (x^k, t^k)$  and  $t^k \geq t^0$ ,  $u(P^k) = M$ , then, by [3; Th. 5],  $u \equiv M$  in  $S(P^k)$ . Hence, if the boundary of  $D$  near  $P^0$  is sufficiently smooth,  $u \equiv M$  in some set  $V' \cap D$  where  $V'$  is some neighborhood of  $P^0$ . Consequently  $\partial u / \partial \tau = 0$  for every  $\tau$ .

If  $u(P) \leq M$  for all  $P \in V \cap D$ , if  $u(P)$  is not strictly smaller than  $M$  for all  $P \in V \cap D$ ,  $P \neq P^0$ , and if the previous situation does not arise, then one and only one of the following cases must occur:

(i)  $u < M$  at all points  $(x, t)$  in  $V \cap D$  with  $t \geq t^0$ . Using [3; Th. 5] one can easily conclude that there exists a neighborhood  $V'$  of  $P$  such that  $u < M$  in  $V' \cap D$ , and Theorem 2 remains true.

(ii)  $u < M$  at all points  $(x, t)$  in  $V \cap D$  with  $t > t_0$  and  $u \equiv M$  at all points  $(x, t)$  in  $V \cap D$  with  $t \leq t_0$ . We then consider only those directions  $\tau$  along which  $u < M$ . We claim that *Theorem 2 is not true for the present situation*. To prove this, consider the following simple counter-example:

$$P^0 = (0, 0), M = 0, Lu = \frac{\partial^2 u}{\partial x^2} - \frac{\partial u}{\partial t}, u(x, t) = \begin{cases} -t^2 & \text{if } t > 0 \\ 0 & \text{if } t \leq 0. \end{cases}$$

$u$  satisfies  $Lu \geq 0$  and assumes its maximum 0 for  $t \leq 0$ . But, the derivative  $\partial u / \partial \tau$  at  $P^0 = (0, 0)$ , along any direction  $\tau$ , is zero.

As another counter-example (with  $Lu = 0$ ) one can take a fundamental solution of the heat equation.

Note that the preceding counter-examples are valid without any assumptions on the behavior of the boundary of  $D$  near  $P^0$ .

We shall now consider the case  $x^1 = x^0$  which was excluded by the assumptions of Theorem 2. We shall assume that at  $P^0 = (0, 0)$  there passes a tangent hyperplane  $t = 0$ . If  $D$  is above this hyperplane, then the preceding counter-examples show that Theorem 2 is not true. It remains to consider the case in which  $D$  is "essentially" below  $t = 0$ , that is, if we denote by  $t = \varphi(x)$  the equation of the boundary of  $D$  near  $P^0$ , then  $D$  is on the side  $t < \varphi(x)$ . In this case, however, Theorem 1 tells us that in general we cannot assume both  $u(P^0) = \max u(P) > 0$  ( $P \in \overline{D}$ ) and  $u < u(P^0)$  in  $V \cap D$ .

The example in § 1 Remark (a) can also serve as a counter-example to Theorem 2 in case  $P^0$  is a vertex-point. Indeed, along the  $t$ -direction

$$\frac{\partial u}{\partial t} \Big|_{P^0} = \frac{\partial}{\partial t} [(t - \gamma_1 x)(\gamma_2 x - t)] \Big|_{x=0, t=0} = 0.$$

By a small modification of this counter-example one can get a counter-example to the analogue of Theorem 2 for elliptic operators [2] [4] in case  $P^0$  is a vertex. Indeed, define  $D$  by

$$x^2 + y^2 < R^2, y < \gamma_1 x, y > \gamma_2 x \quad \gamma_1 > 0 > \gamma_2,$$



and take  $Lu = \partial^2 u / \partial x^2 + A \partial^2 u / \partial y^2$ , where  $A > |\gamma_1 \gamma_2|$ . The function  $u(x, y) = (y - \gamma_1 x)(y - \gamma_2 x)$  satisfies:  $u < 0$  in  $D$ ,  $u = 0$  at the origin,  $Lu = 2\gamma_1 \gamma_2 + 2A > 0$ . But along any direction  $\tau$  within  $D$ ,  $\partial u / \partial \tau|_{x=0, y=0} = 0$ .

4. Let  $D$  be a domain bounded by the two hyperplanes  $t=0$ ,  $t=T > 0$  and a surface  $B$  between them. Assume that the intersection  $\{t=T\} \cap \bar{D}$  is the closure of an open set on  $t=T$ , and denote by  $A$  the boundary of  $D$  on  $t=0$ . The Neumann problem for the parabolic equation  $Lu=0$  consists in finding a solution to the equation  $Lu=0$  which satisfies the following initial and boundary conditions:

$$u=f \text{ on } A, \quad \frac{\partial u}{\partial \nu} = g \text{ on } B$$

( $f, g$  are given functions).

From Theorem 2 and from the strong maximum principle [3; Th. 5] we conclude: *If for every point  $P^0 = (x^0, t^0)$  of  $B$  (i) there exists a sphere with center  $(x', t')$ ,  $x' \neq x^0$ , passing through  $P^0$  and contained in  $\bar{D}$ , and (ii)  $\bar{S}(P^0)$  contains interior points of  $A$ , then the Neumann problem has at most one solution.* Clearly, this uniqueness property holds also for the more general problem where  $\partial u / \partial \nu$  is replaced by  $\partial u / \partial \tau$  and  $\tau$  is a non-tangential direction which varies on  $B$ .

As another application to Theorem 2, one can deduce the positivity of  $\partial G / \partial \nu$ , where  $G$  is the Green's function of  $Lu=0$ .

5. Let  $D$  be a domain bounded by  $t=0$ ,  $t=T$  ( $0 < T \leq \infty$ ) and surfaces  $\Gamma_k$ ,  $0 \leq k \leq m$ ,  $\Gamma_0$  being the outer boundary. Suppose further that the intersection of each  $\Gamma_k$  with  $t=t_0$  ( $0 \leq t_0 < T$ ) is a simple closed curve  $\gamma_k(t_0)$  which belongs to  $C^{(3)}$  and does not reduce to a single point. Write  $u_{x_i} = \partial u / \partial x_i$ ,  $u_t = \partial u / \partial t$ . We shall consider the following problem  $P$ :

$$(9) \quad \sum_{i,j=1}^n a_{ij}(x, t) u_{x_i x_j} - u_t = c(x, t, u, \nabla u)$$

(where  $\nabla u$  denotes the vector  $\partial u / \partial x_i$ ),

$$(10) \quad \frac{\partial u}{\partial \tau} = \sum_{i=1}^n \alpha_i(x, t) u_{x_i} + \alpha(x, t) u_t = \varphi(x, t, u) \quad (x, t) \in \Gamma = \sum_{k=0}^m \Gamma_k$$

$$(11) \quad u(x, 0) = \psi(x) \text{ on } A \quad A = \bar{D} \cap \{t=0\}$$

We make the following assumptions:

(a)  $a_{ij}(x, t)$  is continuous in  $\bar{D}$ ;  $c(x, t, u, \nabla u)$  and its first derivatives with respect to  $u$ ,  $\nabla u$  are continuous for  $(x, t) \in \bar{D}$  and for all values of  $u$ ,  $\nabla u$ .

- (b)  $\varphi$  and  $\partial\varphi/\partial u$  are continuous for all  $(x, t) \in \Gamma$  and for all  $u$ .  
 (c)  $\alpha_i(x, t), \alpha(x, t)$  are continuous for  $(x, t) \in \Gamma$ ;  $\psi(x)$  is continuous in  $A$ .  
 (d) (9) is parabolic in  $\bar{D}$ , that is, there exists a positive constant  $\delta$  such that

$$(12) \quad \sum a_{ij}(x, t) \xi_i \xi_j \geq \delta \sum \xi_i^2$$

holds for all real  $\xi$  and for all  $(x, t) \in \bar{D}$ .

- (e) On each surface  $\Gamma_k$  ( $k=0, 1, \dots, m$ ) either all the directions  $\tau=(\alpha_i, \alpha)$  are exterior or all are interior, and in the exterior case  $\alpha \geq 0$  and the directions  $(\alpha_i, 0)$  are exterior while in the interior case  $\alpha \leq 0$  and the directions  $(\alpha_i, 0)$  are interior.

Denote by  $\Sigma$  the class of functions  $u(x, t)$  defined and continuous in  $\bar{D}$  and satisfying the following conditions:

- ( $\alpha$ )  $\partial u/\partial t, \partial u/\partial x_i, \partial^2 u/\partial x_i \partial x_j$  are continuous in  $D$ ;  
 ( $\beta$ ) For every  $R > 0$ ,  $\partial u/\partial x_i$  is bounded in  $D \cap \{|x|^2 + t^2 < R^2\}$ .

**THEOREM 3.** *Under the assumptions (a)–(e) the problem  $P$  cannot have two different solutions in the class  $\Sigma$ .*

We shall need the following lemma.

**LEMMA.** *There exists a function  $\zeta(x)$  defined in  $A$  and having the following properties: (i)  $\zeta$  has continuous first derivatives in  $A$  and continuous second derivatives in the interior of  $A$ ; (ii)  $\partial\zeta/\partial\nu = -1$  and  $\partial\zeta/\partial\mu = 0$  on  $\gamma_0(0), \dots, \gamma_m(0)$ , where  $\partial/\partial\nu$  and  $\partial/\partial\mu$  denote the derivatives with respect to the interior normal and to any tangential direction, respectively.*

**PROOF OF THE LEMMA.** It will be enough to construct a function  $\chi_0(x)$  which is  $C''$  in  $A$ , which vanishes in a neighborhood of  $\gamma_i(0)$  ( $i=1, \dots, m$ ) and for which  $\partial\chi_0/\partial\nu = -1$ ,  $\partial\chi_0/\partial\mu = 0$  along  $\gamma_0(0)$ ; constructing  $\chi_1(x)$  in a similar manner, we can then take  $\zeta(x) = \sum \chi_i(x)$ . Since  $\gamma_0(0)$  belongs to  $C^{(3)}$ , the normals issuing from  $\gamma_0(0)$  and inwardly directed cover in a one-to-one manner a small inner neighborhood of  $\gamma_0(0)$ , call it  $A_0$ . To each point  $x$  in  $A_0$  there corresponds a unique point  $x^0$  on the boundary of  $\gamma_0(0)$ , such that  $x$  lies on the normal through  $x^0$ . Denote by  $\sigma(x)$  the distance  $|x - x^0|$ . It is elementary to show that  $\sigma(x)$  has continuous second derivatives in  $A_0$ . Denote by  $A_1$  the domain  $0 \leq \sigma \leq \varepsilon_0$ , where  $\varepsilon_0 > 0$  is small enough so that  $A_1 \subset A_0$ . The function

$$\chi_0(x) = \begin{cases} \frac{1}{3\varepsilon_0^2} (\varepsilon_0 - \sigma(x))^3 & \text{if } x \in \bar{A}_1 \\ 0 & \text{if } x \in A - A_1 \end{cases}$$

belongs to  $C''$  in  $A$  and satisfies:  $\partial\chi_0/\partial\nu = \partial\chi_0/\partial\sigma = -1$  and  $\partial\chi_0/\partial\mu = 0$  on  $\gamma_0(0)$ , and  $\chi_0$  vanishes near  $\gamma_k(0)$ , ( $1 \leq k \leq m$ ); the proof is completed.

PROOF OF THEOREM 3. We first consider the case  $n > 1$ . We may suppose that the vectors  $(\alpha_i, \alpha)$  are exterior directions on  $\Gamma_0, \dots, \Gamma_q$  and that  $(\alpha_i, \alpha)$  are interior directions on  $\Gamma_{q+1}, \dots, \Gamma_m$ . Suppose now that  $u$  and  $v$  are two solutions in  $\Sigma$  of the problem  $P$ , and define  $w = v - u$ . Writing

$$\begin{aligned} C(x, t, u, v) &= \int_0^1 \frac{\partial}{\partial u} c(x, t, u + \lambda w, \nabla u + \lambda \nabla w) d\lambda \\ C_i(x, t, u, v) &= \int_0^1 \frac{\partial}{\partial u_{x_i}} c(x, t, u + \lambda w, \nabla u + \lambda \nabla w) d\lambda \\ \Phi(x, t, u, v) &= \int_0^1 \frac{\partial}{\partial u} \varphi(x, t, u + \lambda w) d\lambda \end{aligned}$$

and using (9), (10) and (11), we obtain for  $w$  the following system :

$$(13) \quad \sum a_{ij} w_{x_i x_j} - w_t = Cw + \sum C_i w_{x_i}$$

$$(14) \quad \frac{\partial w}{\partial \tau} = \sum \alpha_i w_{x_i} + \alpha w_t = \Phi w$$

$$(15) \quad w(x, 0) = 0.$$

Substituting  $w(x, t) = z(x, t) \exp(Kt + M\zeta(x))$ , where  $\zeta(x)$  is the function constructed in the lemma and  $K, M$  are constant to be determined later, we get for  $z$  the following system :

$$\begin{aligned} (13') \quad \sum a_{ij} z_{x_i x_j} - z_t &= -M \sum a_{ij} \zeta_{x_i x_j} z - M^2 \sum a_{ij} \zeta_{x_i} \zeta_{x_j} z \\ &\quad - 2M \sum a_{ij} \zeta_{x_i} z_{x_j} + Kz + Cz + M \sum C_i \zeta_{x_i} z + \sum C_i z_{x_i} \end{aligned}$$

$$(14) \quad \frac{\partial z}{\partial \tau} = \sum \alpha_i z_{x_i} + \alpha z_t = -M \sum \alpha_i \zeta_{x_i} z - \alpha Kz + \Phi z$$

$$(15') \quad z(x, 0) = 0.$$

If  $0 \leq k \leq q$ ,  $\alpha \geq 0$  and  $\sum \alpha_i(x, 0) \zeta_{x_i}(x) > 0$  on  $\gamma_k(0)$ , since the angle between the vectors  $(\alpha_i)$  and  $\text{grad } \zeta$  is  $< \pi/2$ . By continuity we get  $\sum \alpha_i(x, t) \zeta_{x_i}(x) \geq \eta > 0$  on  $\gamma_k(t)$ , provided  $0 \leq t \leq T'$  and  $T'$  is sufficiently small. Hence, we can choose  $M$  sufficiently large such that

$$(16) \quad -M \sum \alpha_i \zeta_{x_i} - \alpha K + \Phi < 0$$

holds on  $\gamma_k(t)$ , provided  $K \geq 0$  and  $0 \leq t \leq T'$ .

If  $q+1 \leq k \leq m$ ,  $\alpha \leq 0$  and  $\sum \alpha_i(x, 0) \zeta_{x_i}(x) < 0$ , since the angle between  $(\alpha_i)$  and  $-\text{grad } \zeta$  is  $< \pi/2$ . Again, if  $K \geq 0$  and  $M$  is sufficiently large, then

$$(17) \quad -M \sum \alpha_i \zeta_{x_i} - \alpha K + \Phi > 0$$

on  $\gamma_k(t)$ ,  $0 \leq t \leq T'$ .

Having fixed  $M$ , we now choose  $K$  sufficiently large so that the coefficient of  $z$  on the right side of (13') becomes positive in the domain  $D_{T'} = D \cup \{0 < t < T'\}$ . We claim that  $z \equiv 0$  in  $D_{T'}$ . Indeed, if this is not the case then, using (15') and the weak maximum principle [3; Th. 2] we conclude that  $z$  assumes either its positive maximum or its negative minimum on the boundary  $\sum_{k=0}^m \gamma_k(t)$ ,  $0 \leq t \leq T'$ , of  $D_{T'}$ . It will be enough to consider the case in which  $z$  assumes its positive maximum at a point  $P^0$  on  $\gamma_k(t)$ . If  $0 \leq k \leq q$ , then  $\partial z / \partial \tau \geq 0$  since  $\tau$  is outwardly directed. On the other hand, using (14') and (16) we get  $\partial z / \partial \tau < 0$ , which is a contradiction. If  $q+1 \leq k \leq m$ , then  $\partial z / \partial \tau \leq 0$  since  $\tau$  is inwardly directed. On the other hand, using (14') and (17) we get  $\partial z / \partial \tau > 0$  which is a contradiction. We have thus proved that  $z \equiv w \equiv 0$  in  $D_{T'}$ . We can now apply a classical procedure of continuation and thus complete the proof of the theorem for the case  $n > 1$ .

In the case  $n=1$ ,  $\Gamma = \Gamma_0$  is composed of two curves  $\Gamma_{01}$  and  $\Gamma_{02}$ . Suppose  $\Gamma_{0k}$  intersects  $t=0$  at  $a_k$ ,  $a_1 < a_2$ . The function

$$\zeta(x) = \frac{(x-a_1)(x-a_2)}{a_2-a_1}$$

can be used in the preceding proof. Note that it is not necessary to make any assumptions on the smoothness of the curves  $\Gamma_{0k}$ .

## REFERENCES

1. F. A. Ficken, *Uniqueness theorems for certain parabolic equations*, J. Rational Mechanics and Analysis, **1** (1952), 573-578.
2. E. Hopf, *A remark on linear elliptic differential equations of second order*, Proc. Amer. Math. Soc., **3** (1952), 791-793.
3. L. Nirenberg, *A strong maximum principle for parabolic equations*, Comm. Pure and Appl. Math., **6** (1953), 167-177.
4. O. A. Olainik, *On properties of some boundary problems for equations of elliptic type*, Mat. Sb. (N.S.), **30** (72) (1952), 695-702.

INDIANA UNIVERSITY



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. L. ROYDEN

Stanford University  
Stanford, California

R. A. BEAUMONT

University of Washington  
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California  
Los Angeles 7, California

E. G. STRAUS

University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

E. F. BECKENBACH

C. E. BURGESS

M. HALL

E. HEWITT

A. HORN

V. GANAPATHY IYER

R. D. JAMES

M. S. KNEBELMAN

L. NACHBIN

I. NIVEN

T. G. OSTROM

M. M. SCHIFFER

G. SZEKERES

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA

OREGON STATE COLLEGE

UNIVERSITY OF OREGON

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF UTAH

WASHINGTON STATE COLLEGE

UNIVERSITY OF WASHINGTON

\* \* \*

AMERICAN MATHEMATICAL SOCIETY

CALIFORNIA RESEARCH CORPORATION

HUGHES AIRCRAFT COMPANY

THE RAMO-WOOLDRIDGE CORPORATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any of the editors. All other communications to the editors should be addressed to the managing editor, E. G. Straus at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 10, 1-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

# Pacific Journal of Mathematics

Vol. 8, No. 2

April, 1958

John Herbert Barrett, <i>Second order complex differential equations with a real independent variable</i> .....	187
Avner Friedman, <i>Remarks on the maximum principle for parabolic equations and its applications</i> .....	201
Richard Robinson Goldberg, <i>An inversion of the Stieltjes transform</i> .....	213
Olavi Hellman, <i>On the periodicity of the solution of a certain nonlinear integral equation</i> .....	219
Gilbert Helmsberg, <i>A theorem on equidistribution on compact groups</i> .....	227
Lloyd Kenneth Jackson, <i>Subfunctions and the Dirichlet problem</i> .....	243
Naoki Kimura, <i>The structure of idempotent semigroups. I</i> .....	257
Stephen Kulik, <i>A method of approximating the complex roots of equations</i> .....	277
Ancel Clyde Mewborn, <i>A note on a paper of L. Guttman</i> .....	283
Zeev Nehari, <i>On the principal frequency of a membrane</i> .....	285
G. Pólya and I. J. Schoenberg, <i>Remarks on de la Vallée Poussin means and convex conformal maps of the circle</i> .....	295
B. M. Stewart, <i>Asymmetry of a plane convex set with respect to its centroid</i> .....	335
Hans F. Weinberger, <i>Lower bounds for higher eigenvalues by finite difference methods</i> .....	339
Edwin Weiss and Neal Zierler, <i>Locally compact division rings</i> .....	369
Bertram Yood, <i>Homomorphisms on normed algebras</i> .....	373