

Pacific Journal of Mathematics

ON THE NUMBER OF BI-COLORED GRAPHS

FRANK HARARY

ON THE NUMBER OF BI-COLORED GRAPHS

FRANK HARARY

1. Introduction. This is an extension of papers [2, 3, 4] whose notation and terminology will be used. The main result is a formulation of the generating function or counting polynomial of bicolored graphs, obtained by the enumeration methods of Pólya [6]. A modification of the method yields the number of balanced signed graphs, solving a problem proposed in [5]. In the process of enumerating bicolored graphs, we consider two binary operations on permutation groups called “cartesian product” and “exponentiation” which are abstractly but not permutationally equivalent to the direct product and Pólya’s “Gruppenkranz” [6], respectively.

A *graph* consists of a finite set of *points* together with a prescribed subset of the collection of all *lines*, i.e., unordered pairs of distinct points. Two points are *adjacent* if there is a line joining them. A graph is *k-chromatic*¹ if each of the points can be assigned one of *k* given colors so that any two adjacent points have different colors. A graph is *k-colored* if it is *k-chromatic* and its points are colored so that all *k* colors are used. More precisely, a *k-colored* graph is a pair (G, f) where G is a graph and f is a function from the set of points of G onto the set of numbers $1, 2, \dots, k$ such that if a and b are adjacent points, then $f(a) \neq f(b)$. Two graphs are *isomorphic* if there exists a one-to-one adjacency preserving transformation between their sets of points. Two *k-colored* graphs are *chromatically isomorphic* if there is a color preserving isomorphism between them. Thus (G_1, f_1) is chromatically isomorphic with (G_2, f_2) if there is an isomorphism $\theta: G_1 \rightarrow G_2$ and a permutation $\omega: \{1, \dots, k\} \rightarrow \{1, \dots, k\}$ such that $\omega(f_1(a)) = f_2(\theta(a))$ for every point a in G_1 . Let $g_{p,q}^{(k)}$ be the number of chromatically nonisomorphic *k-colored* graphs with p points and q lines, and let the corresponding generating function be

$$(1) \quad g_p^{(k)}(x) = \sum_{q=0}^{p(p-1)/2} g_{p,q}^{(k)} x^q.$$

We first derive the number of bicolored graphs, $k = 2$, and then discuss the formula for $k = 3$. The problem remains open for $k > 2$.

Received November 4, 1957, and in revised forms March 28, 1958, and June 16, 1958. This work was supported by a grant from the National Science Foundation. The author is deeply grateful to the referee for making several insightful comments of clarification.

¹ This definition is different from that of Dirac [1]. According to Dirac, a graph has *chromatic number* k if it is k -chromatic but not $(k-1)$ -chromatic as defined here.

In precisely the form in which we require it, Pólya's enumeration theorem is reviewed briefly in §2 of [2]. Therefore, we shall not repeat here the definitions leading up to it, but shall only restate the theorem itself.

PÓLYA'S THEOREM. *The configuration counting series $F(x)$ is obtained by substituting the figure counting series $\varphi(x)$ into the cycle index $Z(\Gamma)$ of the configuration group Γ . Symbolically,*

$$(2) \quad F(x) = Z(\Gamma, \varphi(x)) .$$

This theorem reduces the problem of finding the configuration counting series to the determination of the figure counting series and the cycle index of the configuration group.

2. Bicolored graphs; the cartesian product of permutation groups.

Let K_n be the complete graph of n points, in which any two points are adjacent. Let K_{mn} be the bicolored graph whose $m+n$ points are $a_1, a_2, \dots, a_m, b_1, b_2, \dots, b_n$ and whose mn lines are all those of the form $a_i b_j$.

Clearly if a graph is k -colored then its point set is partitioned into k disjoint non-empty subsets such that no two points in the same subset are adjacent. Hence a bicolored graph with p points is a "line-subgraph" (as in [3]) or a spanning subgraph² of a graph K_{mn} for which $m+n=p$. Let $g_{mn,p}$ be the number of chromatically nonisomorphic spanning subgraphs of K_{mn} having q lines, and let

$$(3) \quad g_{mn}(x) = \sum_{q=0}^{mn} g_{mn,q} x^q .$$

Then

$$(4) \quad g_p^{(2)}(x) = \sum_{1 \leq m \leq n} g_{mn}(x) ,$$

where the sum is taken over all m and n such that $m+n=p$. Therefore in order to obtain a formula for the counting polynomial (4), it is sufficient to find that for (3). In this section, we find $g_{mn}(x)$ for the case $m \neq n$ using the "cartesian product" of two permutation groups. In the next section we see that this combinatorial technique is not valid for $m=n$ and formulate $g_{nn}(x)$ in terms of the "exponentiation" of the appropriate two permutation groups.

By Theorem 1 of [3], the counting polynomial $g_{mn}(x)$ for the

² A *spanning subgraph* of a graph G is one whose set of points coincides with that of G .

number of spanning subgraphs of K_{mn} is obtained by substituting $1 + x$ into the cycle index of the line-group³ of K_{mn} :

$$(5) \quad g_{mn}(x) = Z(\Gamma_1(K_{mn}), 1 + x).$$

We note that this equation can also be obtained from the main result, equation (5), of [4]. For the subgraphs of K_{mn} correspond to the different supergraphs of the union $K_m \cup K_n$ of two complete graphs on disjoint point sets. The derivation of $Z(\Gamma_1(K_{mn}))$ for the case $m \neq n$ is parallel to and algebraically simpler than that of $Z(\Gamma_1(K_p))$, which appears in §3 of [2]. Throughout the rest of this section we assume $m \neq n$.

The line group of K_{mn} may be described as an appropriately formulated product of the symmetric groups S_m and S_n . This product can be generally defined for any two permutation groups in the following way. Let A and B be any two permutation groups with object sets X and Y , degrees d and e and orders m and n respectively. The *cartesian product* $A \times B$ of these two permutation groups has degree de and order mn . Its object set is the cartesian product of X and Y and each of its permutations (α, β) is the cartesian product of permutations α and β from A and B defined by $(\alpha, \beta)(x, y) = (\alpha x, \beta y)$. As an abstract group, the cartesian product is isomorphic to the direct product AB , but they are not permutationally equivalent. For the degree of the direct product is $d + e$ since the group AB has $X \cup Y$ as its object set.

There is a precise method for finding the cycle index of a cartesian product in terms of the cycle indices of the two permutation groups. We first illustrate the method by finding $Z(\Gamma_1(K_{23}))$. The line group of K_{23} is the cartesian product of S_2 and S_3 which is a permutation group of degree 6 and order 12, written $\Gamma_1(K_{23}) = S_2 \times S_3$. Let a_1, a_2 be the indeterminates occurring in $Z(S_2)$ and b_1, b_2, b_3 be those in $Z(S_3)$ so that

$$Z(S_2) = \frac{1}{2} (a_1^2 + a_2) \text{ and } Z(S_3) = \frac{1}{6} (b_1^3 + 3b_1b_2 + 2b_3).$$

Then we write⁴

$$\begin{aligned} Z(\Gamma_1(K_{23})) &= Z(S_2 \times S_3) = Z(S_2) \times Z(S_3) \\ &= \frac{1}{12} (a_1^2 \times b_1^3 + 3a_1^2 \times b_1b_2 + 2a_1^2 \times b_3 + a_2 \times b_1^3 + 3a_2 \times b_1b_2 + 2a_2 \times b_3), \end{aligned}$$

³ The *line group* $\Gamma_1(G)$ of a graph G is the collection of all permutations on the set of lines of G consistent with the automorphism group $\Gamma(G)$ of G ; see [3].

⁴ By the following formulas we mean that the cartesian product of two permutation groups can be extended to the cartesian product of their cycle indices in the indicated manner.

and give each of these six terms in Table 1, in which c_1 to c_6 denote the indeterminates in $Z(\Gamma_1(K_{23}))$.

Table 1

Term of $Z(S_2 \times S_3)$	$a_1^2 \times b_1^3$	$a_1^2 \times b_1 b_2$	$a_1^2 \times b_3$	$a_2 \times b_1^3$	$a_2 \times b_1 b_2$	$a_2 \times b_3$
Term of $Z(\Gamma_1(K_{23}))$	c_1^6	$c_1^2 c_2^2$	c_3^2	c_2^3	c_2^2	c_3

We illustrate Table 1 for the term $a_2 \times b_1 b_2$. Let the 2-cycle $(p_1 p_2)$ stand for a_2 and the 1-cycle and 2-cycle $(q_1)(q_2 q_3)$ for $b_1 b_2$. The admissible lines of K_{23} are only those of the form $p_i q_j$. The pair $p_1 q_1$ is transformed into $p_2 q_1$, and then back again to give the cycle of length 2 in the corresponding permutation of $\Gamma_1(K_{23})$ of the form $(p_1 q_1 p_2 q_1)$. Similarly the transpositions $(p_1 q_2 p_2 q_3)$ and $(p_1 q_3 p_2 q_2)$ are factors of this element of $\Gamma_1(K_{23})$. Altogether there are three transpositions, so the corresponding term of $Z(\Gamma_1(K_{23}))$ is c_2^3 .

In general, we have

$$(6) \quad a_1^i a_2^j \dots a_m^m \times b_1^i b_2^j \dots b_n^n = \prod_{\alpha, \beta} (a_{\alpha}^i \times b_{\beta}^j)$$

and

$$(7) \quad a_{\alpha}^i \times b_{\beta}^j = c_{m(\alpha, \beta)}^{i a_j^j \beta^{d(\alpha, \beta)}}$$

where $d(\alpha, \beta)$ and $m(\alpha, \beta)$ are the greatest common divisor and least common multiple.

The cycle index of S_p is

$$(8) \quad Z(S_p) = \frac{1}{p!} \sum_{(j)} \frac{p!}{1^{j_1} j_1! \dots p^{j_p} j_p!} f_1^{j_1} \dots f_p^{j_p}$$

where the sum is taken over all partitions $(j) = (j_1, j_2, \dots, j_p)$ of p such that

$$1j_1 + 2j_2 + \dots + pj_p = p.$$

The last four equations together with

$$(9) \quad Z(\Gamma_1(K_{m,n})) = Z(S_m) \times Z(S_n)$$

provide a formula for $g_{mn}(x)$ when $m \neq n$.

We use Table 1 to illustrate equation (9) by finding $g_{23}(x)$. Here

$$Z(\Gamma_1(K_{23})) = \frac{1}{12} (c_1^6 + 3c_1^2 c_2^2 + 2c_3^2 + 4c_2^3 + 2c_3),$$

so that

$$g_{23}(x) = 1 + x + 3x^2 + 3x^3 + 3x^4 + x^5 + x^6.$$

The bicolored graphs with two points of one color and three points of the other color which correspond to the coefficients in the preceding counting polynomial are shown in Figure 1.




Fig. 1.

3. Bicolored graphs; exponentiation of permutation groups. We now turn to the enumeration of bicolored graphs for the case $m = n$. As in the preceding section, we again have equation (5) holding for this special case:

$$g_{nn}(x) = Z(\Gamma_1(K_{nn}), 1 + x).$$

However, it is not true that $\Gamma_1(K_{nn}) = S_n \times S_n$ since $S_n \times S_n$ is a proper subgroup of $\Gamma_1(K_{nn})$. The remaining $(n!)^2$ permutations in $\Gamma_1(K_{nn})$ are obtained on interchanging the two colors in accordance with the definition of chromatic isomorphism.




Fig. 2.

For example, all the (chromatically nonisomorphic) bicolored graphs for $m = n = 2$ are shown in Figure 2, so that $g_{22}(x) = 1 + x + 2x^2 + x^3 + x^4$. However, the formulas in the preceding section give

$$Z(S_2 \times S_2, 1 + x) = 1 + x + 3x^2 + x^3 + x^4,$$

since the permutations in $S_2 \times S_2$ distinguish between the two bicolored graphs in Figure 3, in which the color assigned to each point is indicated by one of the integers 1 or 2.

With the appropriate definition of group exponentiation, we will express $\Gamma_1(K_{nn})$ as S_n raised to the power S_2 . We first review the definition of the "composition" of two permutation groups (the "Gruppenkranz" of Pólya [6]). Let **A** and **B** be any two permutation groups as in the preceding section. Then using the notation of Pólya [6] the

composition $A[B]$ of A with B has object set $X \times Y$ (as for the cartesian product). However, it is more convenient to regard the object set here as a d by e matrix $M = (x_{ij})$. Then the elements of $A[B]$ are the permutations of the entries of M constructed as follows. First permute the rows of M in accordance with an element of A . Then permute the column indices in each row separately using one element



Fig. 3.

of B for each row, repetitions permitted. Hence the degree of $A[B]$ is de and the order is mn^d .

The *exponentiation* B^A of A with B is that permutation group whose object set is Y^X , the collection of all functions from X into Y , and whose elements are constructed as follows. It is assumed that the objects x_1, x_2, \dots, x_d in X are indexed. First permute the objects in X in accordance with an element α of A . Then for each object x_i in X , permute the e objects of Y into which it can be mapped, using a permutation β_i from B . More precisely each selection of $\alpha \in A$ and $\beta_1, \beta_2, \dots, \beta_d \in B$ (not necessarily distinct) determines a permutation of Y^X which takes the function f into the function f^* defined by:

$$f^*(x_i) = \beta_i f(\alpha x_i) \text{ for all } x_i \in X; \quad i = 1, 2, \dots, d.$$

It can easily be shown that distinct selections of $\alpha, \beta_1, \dots, \beta_d$ lead to distinct permutations of Y^X and that these permutations form a group.

The degree of B^A is e^d and the order is mn^d . It follows at once from their constructions that the group B^A and $A[B]$ are isomorphic as abstract groups. But they are not equivalent as permutation groups since they have different degrees.

With this definition of exponentiation, it follows at once that the line group of K_{mn} is given by

$$(10) \quad \Gamma_1(K_{mn}) = S_n^{S_m^2}.$$

Before calculating the cycle index of $S_n^{S_m^2}$, we illustrate for $n = 2$ and 3 . Since $S_2^2 = S_2[S_2] = D_4$, the dihedral group of degree 4 and order 8, its cycle index in terms of the indeterminates c_1, c_2, c_3, c_4 is given by

$$Z(D_4) = \frac{1}{8}(c_1^4 + 3c_2^2 + 2c_1^2c_2 + 2c_4).$$

The correct polynomial $g_{22}(x)$ which verifies Figure 2 follows at once from this cycle index.

For $n = 3$, let the object set of $\Gamma_1(K_{33})$ be denoted:

$$X = \{11', 12', 13', 21', 22', 23', 31', 32', 33'\} .$$

Then $\Gamma_1(K_{33}) = \mathbf{S}_3^{\mathbf{S}_2}$ contains the $(3!)^2$ permutations in $\mathbf{S}_3 \times \mathbf{S}_3$ and also the $(3!)^2$ permutations obtained from these on multiplying each of them by the following reflection ρ which interchanges primed and unprimed digits in the objects in X :

$$\rho = (11')(22')(33')(12' 21')(23' 32')(31' 13') .$$

Symbolically, we write

$$\Gamma_1(K_{33}) = \mathbf{S}_3^{\mathbf{S}_2} = (\mathbf{S}_3 \times \mathbf{S}_3) \cup \rho(\mathbf{S}_3 \times \mathbf{S}_3) .$$

Then

$$Z(\mathbf{S}_3 \times \mathbf{S}_3) = \frac{1}{(3!)^2} (c_1^9 + 6c_1^3c_2^3 + 8c_3^3 + 9c_1c_2^4 + 12c_3c_6) ,$$

and a straightforward calculation gives (using not quite proper notation since cycle index is defined for groups rather than cosets):

$$Z(\rho(\mathbf{S}_3 \times \mathbf{S}_3)) = \frac{1}{(3!)^2} (6c_1^3c_2^3 + 18c_1c_4^2 + 12c_3c_6) .$$

Combining these, we have

$$Z(\mathbf{S}_3^{\mathbf{S}_2}) = \frac{1}{2 \cdot (3!)^2} (c_1^9 + 12c_1^3c_2^3 + 8c_3^3 + 9c_1c_2^4 + 18c_1c_4^2 + 24c_3c_6) ,$$

from which one readily calculates using (10) and (5),

$$g_{33}(x) = 1 + x + 2x^2 + 4x^3 + 5x^4 + 5x^5 + 4x^6 + 2x^7 + x^8 + x^9 .$$

We now proceed to obtain a closed formula for $Z(\mathbf{S}_n^{\mathbf{S}_2})$, thereby completing the explicit solution of the enumeration of bicolored graphs. The process of finding this cycle index is also analogous to the calculation of $Z(\Gamma_1(K_p))$ which appears in §3 of [2]. Clearly, the automorphism group of K_{nn} is $\mathbf{S}_2[\mathbf{S}_n]$. For the complement K'_{nn} consists of two disjoint copies of K_n . By a result in Pólya [6], the cycle index of the composition of two permutation groups is the composition of their cycle indices. For example,

$$Z(\mathbf{S}_2[\mathbf{S}_3]) = \frac{1}{2} \left[\left(\frac{1}{6} (a_1^3 + 3a_1a_2 + 2a_3) \right)^2 + \frac{1}{6} (a_2^3 + 3a_2a_1 + 2a_6) \right]$$

But we require here the cycle index of the line group of K_{nn} . There is a one-to-one correspondence between the terms of the cycle indices

$Z(S_n^{S_2})$ and $Z(S_2[S_n])$ with the same integral coefficients. Analogous to the terms in the above illustration of $Z(S_2[S_3])$, let us write

$$(11) \quad Z(S_2[S_n]) = \frac{1}{2} [(Z(S_n))^2 + Z(S_n(2))].$$

Thus $Z(S_n(2))$ is obtained from $Z(S_n)$ on replacing each indeterminate f_k by f_{2k} .

The term of $Z(S_n^{S_2})$ corresponding to the first term of the right hand member of equation (11) is $Z(S_n \times S_n)$. For bicolored graphs with $m \neq n$, this is the result of the preceding section. The term of $Z(S_n^{S_2})$ corresponding to the term $Z(S_n(2))$ of (11) is derived as follows. Let the general term of $Z(S_n(2))$ be given by

$$(12) \quad f_{2^1}^{j_1} f_{4^2}^{j_2} \cdots f_{2^n}^{j_n}.$$

This term (12) occurs in the cycle index of the point group of K_{nn} . We require the corresponding term in the cycle index of the line group obtained by calculating the induced permutation on pairs of points from two disjoint sets. Let the letters c_i be the indeterminates in the cycle index $Z(S_n^{S_2})$. There are two contributions to $Z(S_n^{S_2})$ arising from (12): those from each of the n factors $f_{2k}^{j_k}$ separately, and those from pairs of factors $f_{2r}^{j_r} f_{2s}^{j_s}$, $r \neq s$.

The contribution to the cycle index due to each factor in (12) is

$$f_{2^1}^{j_1} \rightarrow c_1^{j_1} c_2^{\binom{j_1}{2}}, \quad f_{4^2}^{j_2} \rightarrow c_4^{j_2} c_2^{\binom{j_2}{2}}, \quad f_{6^3}^{j_3} \rightarrow (c_3 c_6)^{j_3} c_6^{\binom{j_3}{3}}, \quad \dots$$

It is convenient to express the contribution of $f_{2k}^{j_k}$ separately for k even and k odd:

$$f_{2k}^{j_k} \rightarrow \begin{cases} (c_{2k}^{k/2})^{j_k} c_{2k}^{k \binom{j_k}{2}}, & k \text{ even} \\ (c_k c_{2k}^{(k-1)/2})^{j_k} c_{2k}^{k \binom{j_k}{2}}, & k \text{ odd.} \end{cases}$$

Similarly, the contribution from pairs is given by

$$f_{2r}^{j_r} f_{2s}^{j_s} \rightarrow c_{2m(r,s)}^{j_r j_s d(r,s)}, \quad r < s,$$

where $m(r, s)$ and $d(r, s)$ are the least common multiple and greatest common divisor respectively.

Collecting these observations, we find

$$Z(S_n^{S_2}) = \frac{1}{2} [Z(S_n \times S_n) + Z'], \quad \text{where}$$

$$Z' = \frac{1}{n!} \sum_{(j)} \frac{n!}{\prod_k k^j j_k!} \prod_{k \text{ even}} c_k^{j_k/2+k} \binom{j_k}{2} \prod_{k \text{ odd}} (c_k c_{2k}^{(k-1)/2})^{j_k} c_{2k}^{j_k/2} \prod_{r < s} c_{2m(r,s)}^{j_r j_s A(r,s)}.$$

This formula for $Z(S_n^2)$ together with equations (10) and (5) give the number of bicolored graphs for $m = n$. For $n = 3$, this expression for Z' specializes to that for $Z(\rho(S_3 \times S_3))$ in the above example.

The only other known cycle index of the exponentiation of two permutation groups also involves complex combinatorial calculations and is worked out in Slepian [8]. Consider the counting polynomial $b_n(x) = \sum b_{nm} x^m$, where b_{nm} is the number of symmetry types of boolean functions of n variables having m nonzero terms when written in disjunctive normal form. Pólya [7] showed that

$$b_n(x) = Z(Q_n, 1 + x),$$

where Q_n is the automorphism group of the n -cube. It is easily seen that $Q_n = S_n^2$ and in fact Pólya [7, footnote 7] comments that Q_n and $S_n[S_2]$ are isomorphic as abstract groups. Slepian [8] completed the enumeration problem for $b_n(x)$ by providing a calculus for an explicit formulation of $Z(S_n^2)$, although using different terminology and notation.

It would be interesting to solve the general problem of obtaining an expression for $Z(B^A)$ in terms of $Z(A)$ and $Z(B)$. This would be analogous to equations (6) and (7) which give $Z(A \times B)$ in terms of $Z(A)$ and $Z(B)$.

To summarize, the counting polynomial $g_{mnt}(x)$ for bicolored graphs is given by

$$(14) \quad g_{mnt}(x) = \begin{cases} Z(S_m \times S_n, 1 + x) & \text{when } m \neq n \\ Z(S_n^2, 1 + x) & \text{when } m = n. \end{cases}$$

4. k-colored graphs. We illustrate the general problem for $k = 3$. Here we have, analogous to equations (3), (4) and (5), and with similar notation:

$$(3') \quad g_{mnt}(x) = \sum_{q=0}^{mnt} g_{mnt,q} x^q,$$

$$(4') \quad g_p^{(3)}(x) = \sum_{1 \leq m \leq n \leq t} g_{mnt}(x),$$

$$(5') \quad g_{mnt}(x) = Z(\Gamma_1(K_{mnt}), 1 + x).$$

Thus K_{mnt} is the complete tricolored graph with $m + n + t$ points $p_1, p_2, \dots, p_m, q_1, q_2, \dots, q_n, r_1, r_2, \dots, r_t$ and all $mn + nt + tm$ lines of the

form $p_i q_j, q_j r_k,$ and $r_k p_i.$ Similarly $g_{mnt,q}$ is the number of spanning subgraphs of K_{mnt} having q lines, etc. We distinguish between three cases: (a) m, n, t distinct, (b) $m = n \neq t,$ and (c) $m = n = t.$ These are illustrated in Figures 4 (a), (b), and (c).




Fig. 4.

Only in case (a) have we obtained an algorithm for $Z(\Gamma_1(K_{mnt}))$ in closed form. The result analogous to (9), derived in same manner, is as follows. Let a_1 to a_m, b_1 to $b_n,$ and c_1 to c_t be the indeterminates in $Z(S_m), Z(S_n),$ and $Z(S_t)$ respectively. Let the indeterminates in $(\Gamma_1(K_{mnt}))$ be $d_1, d_2, \dots.$ Let $A, B,$ and C denote arbitrary terms of $Z(S_m), Z(S_n),$ and $Z(S_t)$ respectively. Then in this notation the left-hand member of equation (6) is $A \times B$ and the term of $Z(\Gamma_1(K_{mnt}))$ obtained from $A, B,$ and C is,

$$(15) \quad (A \times B)(B \times C)(C \times A),$$

where each of the three factors in the expression (15) is a product of the indeterminates d_i using equations (6) and (7). For example,

$$Z(\Gamma_1(K_{123})) = \frac{1}{12} (d_1^{11} + 3d_2^3 d_1^5 + 2d_3^3 d_1^2 + d_2^4 d_1^3 + 3d_2^5 d_1 + 2d_3 d_3 d_2)$$

is the cycle index of the line group of the tricolored graph K_{123} shown in Figure 4 (a).

Referring to Figure 4(b), one can find

$$Z(\Gamma_1(K_{122})) = \frac{1}{8} (d_1^8 + 2d_2^2 + 4d_1^2 d_2^3 + d_2^4).$$

The group $\Gamma_1(K_{122})$ appears to be irreducible by any of the operations of direct product, cartesian product, composition, or exponentiation. However, it is abstractly isomorphic to D_4 and can be obtained from two copies of D_4 defined on disjoint object sets $\{1, 2, 3, 4\}$ and $\{5, 6, 7, 8\}$ by the following operation.

Let the set X be the union of the disjoint sets X_1 and $X_2.$ Let A_1 and A_2 be permutation groups defined on X_1 and X_2 respectively, such that h is an abstract isomorphism of A_1 onto $A_2.$ Then the permutation group $A_1 \oplus_h A_2$ can be defined as follows: The function f from X onto X belongs to $A_1 \oplus_h A_2$ if and only if there exist $\alpha_1 \in A_1$ and $\alpha_2 \in A_2$

with $\alpha_2 = h\alpha_1$ such that $f(x) = \alpha_1(x)$ if $x \in X_1$ and $f(x) = \alpha_2(x)$ if $x \in X_2$. Clearly $A = A_1 \oplus_h A_2$ is abstractly isomorphic to A_1 .

Now let $D_{4,1}$ be the dihedral group of degree 4 generated by the permutations (1234) and (12)(34), let $D_{4,2}$ be generated by (5678) and (57)(6)(8), and let h be the isomorphism between them which preserves respective generators. Then

$$\Gamma_1(K_{122}) = D_{4,1} \oplus_h D_{4,2} .$$

Finally, it is easy to see that $\Gamma_1(K_{222})$ is abstractly isomorphic to $S_3[S_2]$ and that

$$\begin{aligned} Z(\Gamma_1(K_{222})) &= \frac{1}{3! 2^3} [(c_1^{12} + 3c_1^4 c_2^4 + 4c_2^6) + 3 \cdot 2(2c_1^2 c_2^5 + 2c_3^3) + 2 \cdot 2^2(c_3^4 + c_6^2)] \\ &= \frac{1}{48} (c_1^{12} + 3c_1^4 c_2^4 + 4c_2^6 + 8c_3^4 + 8c_6^2 + 12c_1^2 c_2^5 + 12c_3^3) . \end{aligned}$$

It is clear that the line group of K_{nnn} is abstractly isomorphic to the automorphism group of K_{nnn} . Its complement K'_{nnn} consists of three disjoint copies of K_n , so that the group of K_{nnn} is $S_3[S_n]$. But an explicit expression for $Z(\Gamma_1(K_{nnn}))$ does not appear to be obvious. (For the particular case $n = 3$, it can be shown that $\Gamma_1(K_{333})$ is permutationally equivalent to S_3^3 .) It does not appear that the operations considered here will suffice to enumerate even the tricolored graphs.

5. Connected k -colored graphs. Let

$$g(x, y) = \sum_p g_p^{(k)}(x)y^p$$

be the generating function for *all* (connected or not) k -colored graphs, and let $c(x, y)$ be that for the connected ones only. Then to find the number of connected k -colored graphs, we substitute into equation (33) of [2] to get

$$(16) \quad 1 + g(x, y) = \exp \left(\sum_{n=1}^{\infty} \frac{1}{n} c(x^n, y^n) \right)$$

or equivalently,

$$(16') \quad \sum_{n=1}^{\infty} \frac{1}{n} c(x^n, y^n) = \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n} g^n(x, y) .$$

From either of these last two equations, the number of connected k -colored graphs of p points can be computed recursively in terms of the total number of k -colored graphs.

6. Balanced signed graphs. *Signed graphs* are obtained by assigning either a positive or a negative sign to each line of a graphs. It was indicated in [5] how one could enumerate all signed graphs by a variation in one of the formulas of [2]. The *sign of a cycle* is the product of the signs of its lines, and a signed graph is *balanced* if all its cycles are positive. The problem of enumerating balanced signed graphs was proposed in [5]. The result is derivable by an appropriate modification of the generating function for bicolored graphs.

It was shown in [5] that a signed graph is balanced if and only if its set of points can be partitioned into two disjoint subsets such that each positive line and no negative line joins two points in the same subset. In view of this characterization, called the "structure theorem for balance", on deleting all the positive lines of a balanced signed graph one obtains a bicolored graph. Let G_1 and G_2 be arbitrary graphs with m and n points respectively, $m \leq n$, and let $p = m + n$. Let $\Gamma(G_1)$ and $\Gamma(G_2)$ be the groups of G_1 and G_2 respectively. Let $b_q(G_1, G_2)$ be the number of nonisomorphic balanced signed graphs with q negative lines, whose positive lines generate the (disjoint) graphs G_1 and G_2 in accordance with the structure theorem for balance. Let

$$b(G_1, G_2, x) = \sum_{q=0}^{mn} b_q(G_1, G_2) x^q$$

be the desired configuration counting series. Then the figure counting series is $1 + x$. For the figures are the mn pairs of points (c, d) where $c \in G_1$ and $d \in G_2$. The content of a figure (c, d) is 0 if c and d are not joined by a negative line and is 1 if they are.

Analogously to the situation for bicolored graphs there are two possibilities. If G_1 and G_2 are not isomorphic, then the configuration group is $\Gamma(G_1) \times \Gamma(G_2)$. But if they are isomorphic, the configuration group is $\Gamma(G_1)^{S_2}$. Hence an application of Pólya's Theorem yields

$$(17) \quad b(G_1, G_2, x) = \begin{cases} Z(\Gamma(G_1) \times \Gamma(G_2), 1 + x) & \text{when } G_1 \not\cong G_2 \\ Z(\Gamma(G_1)^{S_2}, 1 + x) & \text{when } G_1 \cong G_2 \end{cases}$$

It is clear for the special case where G_1 and G_2 are the totally disconnected graphs of m and n points that $b(G_1, G_2, x) = g_{mn}(x)$, $\Gamma(G_1) = S_m$, and $\Gamma(G_2) = S_n$. Thus the formula (17) is a generalization of that for bicolored graphs.

REFERENCES

1. G. A. Dirac, *The structure of k -chromatic graphs*, Fund. Math. **40** (1953), 42-55.
2. F. Harary, *The number of linear, directed rooted and connected graphs*. Trans. Amer. Math. Soc. **78** (1955), 445-463.

3. ———, *On the number of dissimilar line-subgraphs of a given graph*, Pacific J. Math., **6** (1956), 57-64.
4. ———, *The number of dissimilar supergraphs of a linear graph*, Pacific J. Math, **7** (1957), 903-911.
5. ———, *On the notion of balance of a signed graph*, Michigan Math. J. **2** (1953-54), 143-146.
6. G. Pólya, *Kombinatorische Anzahlbestimmungen für Gruppen, Graphen, und chemische Verbindungen*. Acta Math. **68** (1937), 145-254.
7. ———, *Sur les types des propositions composees*, J. Symbolic Logic **5** (1940), 98-103.
8. D. Slepian, *On the number of symmetry types of boolean functions of n variables*, Canadian J. Math. **5** (1953), 185-193.

UNIVERSITY OF MICHIGAN AND THE INSTITUTE FOR ADVANCED STUDY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG
Stanford University
Stanford, California

R. A. BEAUMONT
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

E. G. STRAUS
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
C. E. BURGESS
M. HALL
E. HEWITT

A. HORN
V. GANAPATHY IYER
R. D. JAMES
M. S. KNEBELMAN

L. NACHBIN
I. NIVEN
T. G. OSTROM
H. L. ROYDEN

M. M. SCHIFFER
G. SZEKERES
F. WOLF
K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
THE RAMO-WOOLDRIDGE CORPORATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, E. G. Straus at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 2120 Oxford Street, Berkeley 4, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 8, No. 4

June, 1958

Richard Arens, <i>The maximal ideals of certain functions algebras</i>	641
Glen Earl Baxter, <i>An operator identity</i>	649
Robert James Blattner, <i>Automorphic group representations</i>	665
Steve Jerome Bryant, <i>Isomorphism order for Abelian groups</i>	679
Charles W. Curtis, <i>Modules whose annihilators are direct summands</i>	685
Wilbur Eugene Deskins, <i>On the radical of a group algebra</i>	693
Jacob Feldman, <i>Equivalence and perpendicularity of Gaussian processes</i>	699
Marion K. Fort, Jr. and G. A. Hedlund, <i>Minimal coverings of pairs by triples</i>	709
I. S. Gál, <i>On the theory of (m, n)-compact topological spaces</i>	721
David Gale and Oliver Gross, <i>A note on polynomial and separable games</i>	735
Frank Harary, <i>On the number of bi-colored graphs</i>	743
Bruno Harris, <i>Centralizers in Jordan algebras</i>	757
Martin Jurchescu, <i>Modulus of a boundary component</i>	791
Hewitt Kenyon and A. P. Morse, <i>Runs</i>	811
Burnett C. Meyer and H. D. Sprinkle, <i>Two nonseparable complete metric spaces defined on $[0, 1]$</i>	825
M. S. Robertson, <i>Cesàro partial sums of harmonic series expansions</i>	829
John L. Selfridge and Ernst Gabor Straus, <i>On the determination of numbers by their sums of a fixed order</i>	847
Annette Sinclair, <i>A general solution for a class of approximation problems</i>	857
George Szekeres and Amnon Jakimovski, <i>(C, ∞) and (H, ∞) methods of summation</i>	867
Hale Trotter, <i>Approximation of semi-groups of operators</i>	887
L. E. Ward, <i>A fixed point theorem for multi-valued functions</i>	921
Roy Edwin Wild, <i>On the number of lattice points in $x^t + y^t = n^{t/2}$</i>	929