

Pacific Journal of Mathematics

ERROR BOUNDS FOR AN APPROXIMATE SOLUTION TO THE VOLTERRA INTEGRAL EQUATION

JOHN HILZMAN

ERROR BOUNDS FOR AN APPROXIMATE SOLUTION TO THE VOLTERRA INTEGRAL EQUATION

JOHN HILZMAN

In 1945 Michal [2] obtained several results which he asserted were useful for approximating the solution to the Volterra integral equation. These results were concerned with certain equations in Fréchet differentials having as their unique solutions the resolvent kernel and the exact solution to the Volterra integral equation of the second kind. Michal treated the resolvent kernel $S[K|x, t]$ and the solution $y[K|x]$ as functions¹ of the given kernel $K(x, t)$, the setting being the Banach spaces

$$T = \{G(x, t) \mid G(x, t) \text{ is real and continuous on } a \leq t \leq x \leq b\}$$

and

$$I = \{g(x) \mid g(x) \text{ is real and continuous on } a \leq x \leq b\}$$

with the norms

$$(1) \quad \begin{aligned} \|G(x, t)\| &= \max |G(x, t)| \quad (a \leq t \leq x \leq b), \\ \|g(x)\| &\equiv \max |g(x)| \quad (a \leq x \leq b), \end{aligned}$$

respectively. In another work [3, pp. 16-17] Michal showed that the solution $y[K|x]$ can be expressed by a Taylor-type expansion in Fréchet differentials of $y[K|x]$ about an arbitrary $K_0(x, t)$ from T . In this paper we shall use Michal's results to obtain approximations to the solution of the Volterra integral equation with error bounds.

I wish to thank Professor A. T. Lonseth for suggesting this course of investigation and the Referee for recommendations which have improved this paper.

Consider the integral equation

$$(2) \quad y(x) + \int_a^x K(x, t)y(t)dt = f(x)$$

where $K(x, t)$ is in T and $f(x)$ is in I . It is known that the exact solution to (2) is given by

$$(3) \quad y(x) = f(x) + \int_a^x S(x, t)f(t)dt$$

Received January 15, 1959, and in revised form March 30, 1959. This work was partially supported by the Office of Ordnance Research, U.S. Army, under contract with Oregon State College.

¹ The symbols $S[K|x, t]$ and $y[K|x]$ were used to indicate the functional dependence of $S(x, t)$ and $y(x)$ on $K(x, t)$.

where the resolvent kernel $S(x, t)$ is in T . Let $K_0(x, t)$ from T be another kernel such that $S_0(x, t)$, the resolvent of $K_0(x, t)$, is known and that $\|h(x, t)\| = \|K(x, t) - K_0(x, t)\|$ is small in the sense of (1). Then by (3) the solution to (2) with kernel $K_0(x, t)$ is

$$(4) \quad y_0(x) = f(x) + \int_a^x S_0(x, t)f(t)dt .$$

Now treat $y(x)$ as a function of the kernel $K(x, t)$. The first Fréchet differential $dy(x)$ of $y(x)$ with increment $h(x, t)$ (applied to $K(x, t)$) is

$$dy(x) = - \int_a^x \left[h(x, t) + \int_t^x S(x, z)h(z, t)dz \right] y(t)dt$$

[2, p. 253]. In particular, the Fréchet differential of $y(x)$ evaluated at $K_0(x, t)$ with increment $h(x, t) = K(x, t) - K_0(x, t)$ will be

$$(5) \quad dy_0(x) = - \int_a^x \left[h(x, t) + \int_t^x S_0(x, z)h(z, t)dz \right] y_0(t)dt .$$

Furthermore, by Theorem 2 of [2] the differential system

$$\begin{cases} dy_0(x) = - \int_a^x \left[h(x, t) + \int_t^x S_0(x, z)h(z, t)dz \right] y_0(t)dt \\ y_0(x) = f(x) \quad (K_0(x, t) = 0) \end{cases}$$

has a unique solution which is given by (4). Thus a first order approximation to the solution $y(x)$ of (2) will be

$$y_0(x) + dy_0(x) .$$

The exact solution to (2) is given by the Taylor expansion [3; 1. p. 112]

$$(6) \quad y(x) = y_0(x) + \sum_{j=1}^{\infty} (j!)^{-1} d^j y_0(x)$$

where, in terms of composition powers²,

$$(7) \quad d^j y_0(x) = (-1)^j j! [h + S_0 h]^j * y_0 .$$

Thus knowledge of the higher order differentials will allow closer approximations to $y(x)$.

We now take up the problem of establishing error bounds for any order of approximation to $y(x)$ from (6). If A_j ($j = 1, 2, \dots, n$) is in T and g is in I , and

² $VW = \int_t^x V(x, z)W(z, t)dz$, $W^2 = \int_t^x W(x, z)W(z, t)dz$, $W^n = \int_t^x W(x, z)W^{n-1}(z, t)dz$, and $W^n * g = \int_a^x W^n(x, t)g(t)dt$

$$A = A_1 A_2 \cdots A_n = \int_t^x \int_t^{z_1} \cdots \int_t^{z_{n-2}} A_1(x, z_1) A_2(z_1, z_2) \cdots A_n(z_{n-1}, t) dz_{n-1} \cdots dz_1,$$

it is seen that

$$(8) \quad \|A\| \leq \frac{|b - a|^{n-1}}{(n - 1)!} \prod_{j=1}^n \|A_j\|$$

and

$$(9) \quad \|A * g\| \leq \frac{\|g\| |b - a|^n}{n!} \prod_{j=1}^n \|A_j\|.$$

Let $P_{n-i,i}[h(S_0h)]$ denote the sum of terms obtained from the composition $h^{n-i}(S_0h)^i$ by a permutation on the n places occupied by

$$\underbrace{hh \cdots h}_{n-i} \underbrace{h(S_0h)(S_0h) \cdots (S_0h)}_i = h^{n-i}(S_0h)^i.$$

For example, by setting

$$P_{2,1}[h(S_0h)] = h^2(S_0h) + h(S_0h)h + (S_0h)h^2$$

and

$$P_{1,2}[h(S_0h)] = h(S_0h)^2 + (S_0h)h(S_0h) + (S_0h)^2h$$

we can write with brevity

$$[h + S_0h]^3 = h^3 + P_{2,1}[h(S_0h)] + P_{1,2}[h(S_0h)] + (S_0h)^3.$$

Now let

$$c = \|h(x, t)\|, m = \|y_0(x)\|, B = \|S_0(x, t)\|, \text{ and } u = |b - a|.$$

Then from (7), (8), (9), and the mechanics of composition we obtain

$$\begin{aligned} & \| (n!)^{-1} d^n y_0(x) \| = \| (-1)^n [h + S_0h]^n * y_0 \| \\ & = \| h^n * y_0 + P_{n-1,1}[h(S_0h)] * y_0 + \cdots + P_{1,n-1}[h(S_0h)] * y_0 + (S_0h)^n * y_0 \| \\ & \leq \| h^n * y_0 \| + \| P_{n-1,1}[h(S_0h)] * y_0 \| + \cdots + \| (S_0h)^n * y_0 \| \\ (10) & \leq \frac{mc^n u^n}{n!} + \binom{n}{1} \frac{mc^n u^{n+1} B}{(n+1)!} + \cdots + \binom{n}{n} \frac{mc^n u^{2n} B^n}{(2n)!} \\ & \leq mc^n u^n \sum_{j=0}^n \binom{n}{j} \frac{(uB)^j}{(n+j)!} \\ & \leq \frac{m[cu(1 + uB)]^n}{n!}. \end{aligned}$$

Thus transposing the desired n th order approximation to $y(x)$ from the right side of (6) to the left side and applying (10) we get

$$\begin{aligned}
 (11) \quad \left\| y(x) - y_0(x) - \sum_{j=1}^n \frac{1}{j!} d^j y_0(x) \right\| &= \left\| \sum_{j=n+1}^{\infty} \frac{1}{j!} d^j y_0(x) \right\| \\
 &\leq \sum_{j=n+1}^{\infty} m(j!)^{-1\theta^j} \\
 &\leq m \left[e^\theta - \sum_{j=0}^n (j!)^{-1\theta^j} \right]
 \end{aligned}$$

where $\theta = cu[1 + uB]$, For small values of θ we readily discern the asymptotic relation

$$(12) \quad \left\| y(x) - y_0(x) - \sum_{j=1}^{n-1} \frac{1}{j!} d^j y_0(x) \right\| = 0(\theta^n).$$

A simple numerical example will be given next.

Consider the Volterra equation

$$(13) \quad y(x) + \frac{1}{3} \int_0^x xt[3 + x^3 - t^3]y(t)dt = x \exp [1/3x^3]$$

where $K(x, t) = 1/3 xt[3 + x^3 - t^3]$ is in T , $f(x) = x \exp [1/3x^3]$ is in I and $a = 0, b = 1$. Take $K_0(x, t) = xt \exp [1/3(x^3 - t^3)]$. The resolvent kernel for $K_0(x, t)$ is $S_0(x, t) = -xt$. By (4) the solution to (13) with kernel $K_0(x, t)$ is

$$(14) \quad y_0(x) = x \exp [1/3x^3] + \int_0^x -xt^2 \exp [1/3t^3]dt = x.$$

By virtue of (5), the Fréchet differential of $y(x)$ evaluated at $K_0(x, t)$ with increment

$$h(x, t) = K(x, t) - K_0(x, t) = \frac{1}{3} xt[3 + x^3 - t^3] - xt \exp [1/3(x^3 - t^3)]$$

is

$$\begin{aligned}
 (15) \quad dy_0(x) &= - \int_0^x \left\{ \frac{1}{3} xt(3 + x^3 - t^3 - 3) \exp [1/3(x^3 - t^3)] \right. \\
 &\quad \left. + \int_t^x -xz \left(\frac{1}{3} zt(3 + z^3 - t^3 - 3) \exp [1/3(z^3 - t^3)] \right) dz \right\} t dt \\
 &= \frac{x^{10}}{162}.
 \end{aligned}$$

Thus a first order approximation to $y(x)$ will be

$$(16) \quad y(x) \approx x + \frac{x^{10}}{162} .$$

It is easily established that

$$\|h(x, t)\| < 0.04, \|S_0(x, t)\| = 1, \|y_0(x)\| = 1 .$$

Hence, with $\theta = 0.08$, it follows from (11) that

$$(17) \quad \|y(x) - y_0(x) - dy_0(x)\| < 0.0033 .$$

REFERENCES

1. E. Hille and R. Phillips, *Functional analysis and semi-groups*, Amer. Math. Soc. Coll. Publ. **31** (1957).
2. A. D. Michal, *Differential equations in Frechet differentials occurring in integral equations*, Proc. Nat. Acad. Sci., **31** (1945), 252-258.
3. ———, *On a non-linear total differential equation in normed linear spaces* Acta Mathematica, **80** (1948), 1-21.

HARPUR COLLEGE

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG

Stanford University
Stanford, California

F. H. BROWNELL

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

L. J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

G. M. CHERRY

D. DERRY

E. HEWITT

A. HORN

L. NACHBIN

M. OHTSUKA

H. L. ROYDEN

M. M. SCHIFFER

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Printed in Japan by Kokusai Bunken Insatsusha
(International Academic Printing Co., Ltd.), Tokyo, Japan

Richard Arens, <i>Extensions of Banach algebras</i>	1
Fred Guenther Brauer, <i>Spectral theory for linear systems of differential equations</i>	17
Herbert Busemann and Ernst Gabor Straus, <i>Area and normality</i>	35
J. H. Case and Richard Eliot Chamberlin, <i>Characterizations of tree-like continua</i>	73
Ralph Boyett Crouch, <i>Characteristic subgroups of monomial groups</i>	85
Richard J. Driscoll, <i>Existence theorems for certain classes of two-point boundary problems by variational methods</i>	91
A. M. Duguid, <i>A class of hyper-FC-groups</i>	117
Adriano Mario Garsia, <i>The calculation of conformal parameters for some imbedded Riemann surfaces</i>	121
Irving Leonard Glicksberg, <i>Homomorphisms of certain algebras of measures</i>	167
Branko Grünbaum, <i>Some applications of expansion constants</i>	193
John Hilzman, <i>Error bounds for an approximate solution to the Volterra integral equation</i>	203
Charles Ray Hobby, <i>The Frattini subgroup of a p-group</i>	209
Milton Lees, <i>von Neumann difference approximation to hyperbolic equations</i>	213
Azriel Lévy, <i>Axiom schemata of strong infinity in axiomatic set theory</i>	223
Benjamin Muckenhoupt, <i>On certain singular integrals</i>	239
Kotaro Oikawa, <i>On the stability of boundary components</i>	263
J. Marshall Osborn, <i>Loops with the weak inverse property</i>	295
Paulo Ribenboim, <i>Un théorème de réalisation de groupes réticulés</i>	305
Daniel Saltz, <i>An inversion theorem for Laplace-Stieltjes transforms</i>	309
Berthold Schweizer and Abe Sklar, <i>Statistical metric spaces</i>	313
Morris Weisfeld, <i>On derivations in division rings</i>	335
Bertram Yood, <i>Faithful $*$-representations of normed algebras</i>	345