

Pacific Journal of Mathematics

FAITHFUL *-REPRESENTATIONS OF NORMED ALGEBRAS

BERTRAM YOOD

FAITHFUL*-REPRESENTATIONS OF NORMED ALGEBRAS

BERTRAM YOOD

1. Introduction. Let B be a complex Banach algebra with an involution $x \rightarrow x^*$ in which, for some $k > 0$, $\|xx^*\| \geq k\|x\|\|x^*\|$ for all x in B . Kaplansky [8, p. 403] explicitly made note of the conjecture that all such B are symmetric. An equivalent formulation is the conjecture that all such B are B^* -algebras in an equivalent norm. In 1947 an affirmative answer had already been provided by Arens [1] for the commutative case. We consider in § 2 the general (non-commutative) case. It is shown that the answer is affirmative if k exceeds the sole real root of the equation $4t^3 - 2t^2 + t - 1 = 0$. This root lies between .676 and .677. In any case these algebras are characterized spectrally as those Banach algebras with involution for which self-adjoint elements have real spectrum and there exists $c > 0$ such that $\rho(h) \geq c\|h\|$, h self-adjoint (where $\rho(h)$ is the spectral radius of h).

A basic question concerning a given complex Banach algebra B with an involution is whether or not it has a faithful*-representation as operators on a Hilbert space. In § 3 we give a necessary and sufficient condition entirely in terms of algebraic and linear space notions in B . This is that $\rho(h) = 0$ implies $h = 0$ for h self-adjoint and that $R \cap (-R) = (0)$. Here R is the set of all self-adjoint elements linearly accessible [11, p. 448] from the set of all finite sums of elements of the form x^*x . This is related to a previous criterion of Kelley and Vaught [10] which however involves topological notions (in particular, the assumption that the involution is continuous).

If B is semi-simple with minimal one-sided ideals a simpler discussion of *-representations (§ 5) is possible even if B is incomplete. For example if B is primitive then B has a faithful*-representation if and only if $xx^* = 0$ implies $x^*x = 0$. The incomplete case has features not present in the Banach algebra case. In the former case, unlike the latter, a^* -representation may be discontinuous. A class of examples is provided in § 5.

2. Arens*-algebras. Let B be a complex normed algebra with an involution $x \rightarrow x^*$. An *involution* is a conjugate linear anti-automorphism of period two. Elements for which $x = x^*$ are called *self-adjoint* (s. a.) and the set of s. a. elements is denoted by H . Let \mathfrak{H} be a Hilbert space and

Received May 4, 1959. This research was supported by the National Science Foundation, research grant NSF G 5865, and by the United States Air Force through the Office of Scientific Research of the Air Research and Development Command under Contract No. SAR/AF - 49(636) - 153.

$\mathfrak{C}(\mathfrak{F})$ be the algebra of all bounded linear operators on \mathfrak{F} . By a $*$ -representation of B we mean a homomorphism $x \rightarrow T_x$ of B into some $\mathfrak{C}(\mathfrak{F})$ where T_{x^*} is the adjoint of T_x . A $*$ -representation which is one-to-one is called *faithful*.

We shall be mainly, but not exclusively, interested in the case where B is complete (a Banach algebra). In § 2 we shall assume throughout that B is a Banach algebra with an involution $x \rightarrow x^*$.

As in [5, p. 8] we set $x \circ y = x + y - xy$ and say that x is quasi-regular with quasi-inverse y if $x \circ y = y \circ x = 0$. The quasi-inverse of x is unique, if it exists, and is denoted by x' . As, for example, in [16, p. 617] we define the *spectrum* of x , $sp(x)$, to be the set consisting of all complex numbers $\lambda \neq 0$ such that $\lambda^{-1}x$ is not quasi-regular, plus $\lambda = 0$ provided there does not exist a subalgebra of B with an identity element and containing x as an invertible element. (The treatment of zero as a spectral value plays no role below.) The *spectral radius* $\rho(x)$ if x is defined to be $\sup |\lambda|$ for $\lambda \in sp(x)$.

We say that B is an *Arens*-algebra* [1] if there exists $k > 0$ such that $\|x^*x\| \geq k \|x\| \|x^*\|$, $x \in B$. As usual, we say that B is a *B*-algebra* if $\|x^*x\| = \|x\|^2$, $x \in B$.

2.1. LEMMA. *Let B an Arens*-algebra with $\|xx^*\| \geq k \|x\| \|x^*\|$, $x \in B$. Then for each s. a. element h , $\rho(h) \geq k \|h\|$ and $sp(h)$ is real.*

That the spectrum of a s. a. element h is real is shown in [1, p. 273]. By use of the inequality $\|h^{2^n}\| \geq k \|h^{2^{n-1}}\|^2$ as in [16, p. 626] it follows that $\rho(h) \geq k \|h\|$. We shall show (Theorem 2.4) that the spectral conditions of Lemma 2.1 imply that B is an Arens*-algebra.

2.2. LEMMA. *Suppose that for each s. a. element h , $\rho(h) \geq c \|h\|$ and $sp(h)$ is real, where $c > 0$. Let h be s. a., $sp(h) \subset [-a, b]$ where $a \geq 0$, $b \geq 0$ and let $r > 0$. Then*

- (1) $\|(-t^{-1}h)'\| < r$ if $t > (1 - cr)b/cr$ and $t > (1 + cr)a/cr$,
- (2) $\|(t^{-1}h)'\| < r$ if $t > (1 - cr)a/cr$ and $t > (1 + cr)b/cr$.

Note that (2) follows from (1) as applied to the element $-h$. By [18, Theorem 3.4] the involution is continuous on B . Therefore h generates a closed $*$ -subalgebra B_0 . Let \mathfrak{M} be the space of regular maximal ideals of B_0 . For $t > a$ set $u = (-t^{-1}h)'$. By [8, Theorem 4.2], $u \in B_0$. It is readily seen that u is s. a. Since $-t^{-1}h + u + t^{-1}hu = 0$ we have, for each $M \in \mathfrak{M}$, $u(M) = h(M)/(t + h(M))$. By [8, p. 402] the spectrum of h is the same whether computed in B or in B_0 so that $-a \leq h(M) \leq b$. Since $\lambda/(t + \lambda)$ is an increasing function of λ we see that $-a/(t - a) \leq u(M) \leq b/(t + b)$. Now $\rho(u) = \sup |u(M)|$, $M \in \mathfrak{M}$. Therefore, since u is s.a.,

$$(2.1) \quad c \|u\| \leq \rho(u) \leq \max [a/(t - a), b/(t + b)] .$$

From formula (2.1), $\|u\| < r$ if $a/(t - a) < cr$ and $b/(t + b) < cr$. This yields (1).

Note that, under the given hypotheses, $c \leq 1$.

2.3. LEMMA. *Let x and y be quasi-regular. Then $x + y$ is quasi-regular if and only if $x'y'$ is quasi-regular.*

The formulas $x' \circ (x + y) \circ y' = x'y'$ and $x + y = x \circ (x'y') \circ y$ yield the desired result. Let $r > 0$. If $\|x'\| < r$ and $\|y'\| < r^{-1}$ it follows from Lemma 2.3 and [12, p. 66] that $(x + y)'$ exists.

Consider the situation of Lemma 2.2 and let h_k be s. a., $k = 1, 2$ where $N = \max(\rho(h_1), \rho(h_2))$. By Lemma 2.2, $\|(t^{-1}h_k)'\| < 1$ and $\|(-t^{-1}h_k)'\| < 1$ if $t > (1 + c)N/c$. Then, by Lemma 2.3,

$$(2.2) \quad sp(h_1 + h_2) \subset [-(1 + c)N/c, (1 + c)N/c].$$

Suppose next that $sp(h_k) \subset [0, \infty)$, $k = 1, 2$. Then $\|(t^{-1}h_k)'\| < 1$ if $t > (1 + c)N/c$ and $\|(-t^{-1}h_k)'\| < 1$ if $t > (1 - c)N/c$. Then by Lemma 2.3,

$$(2.3) \quad sp(h_1 + h_2) \subset [-(1 - c)N/c, (1 + c)N/c].$$

2.4. THEOREM. *Suppose that for each s. a. element h , $\rho(h) \geq c \|h\|$ and $sp(h)$ is real, where $c > 0$. Then B is an Arens*-algebra with $\|xx^*\| \geq k \|x\| \|x^*\|$, $x \in B$, where k can be chosen to be $c^5/(1 + c)(1 + 2c^2)$.*

Let $x = u + iv$ where u and v are s. a. Then $x^*x = u^2 + v^2 + i(uv - vu)$, $xx^* = u^2 + v^2 + i(vu - uv)$ and $xx^* + x^*x = 2u^2 + 2v^2$. We next compare $\rho(u^2) = [\rho(u)]^2$ and $\rho(v^2)$ with $\rho(xx^*)$. For this purpose we may suppose that $\rho(u) \geq \rho(v)$ for otherwise we can replace x by $ix = -v + iu$. If $\lambda \neq 0$ then $\lambda \in sp(xx^*)$ if and only if $\lambda \in sp(x^*x)$. Thus $\rho(xx^*) = \rho(x^*x)$. By (2.2), $sp(xx^* + x^*x) \subset [-(1 + c)\rho(xx^*)/c, (1 + c)\rho(xx^*)/c]$. Now $2u^2 = xx^* + x^*x - 2v^2$. Let $r > 0, t > 0$. By Lemma 2.2,

$$(2.4) \quad \|[t^{-1}(xx^* + x^*x)]'\| < r, t > (1 + cr)(1 + c)\rho(xx^*)/c^2r.$$

Since $sp(-2v^2) \subset (-\infty, 0]$ and $\rho(2v^2) \leq \rho(2u^2)$, by Lemma 2.2 we have, for $t > 0$,

$$(2.5) \quad \|[t^{-1}(-2v^2)]'\| < r^{-1}, t > (r - c)\rho(2u^2)/c.$$

we select $c < r < 2c$. For such r , Lemma 2.3 and formulas (2.4) and (2.5) show that $[t^{-1}(2u^2)]'$ exists if $t > \max\{(1 + cr)(1 + c)\rho(xx^*)/c^2r, (r - c)\rho(2u^2)/c\}$. Now $(r - c)/c < 1$ and $sp(2u^2) \subset [0, \infty)$. Therefore, letting $r \rightarrow 2c$, we have

$$(2.6) \quad \rho(2u^2) \leq (1 + 2c^2)(1 + c)\rho(xx^*)/(2c^3).$$

On the other hand $\|x\| \leq \|u\| + \|v\| \leq [\rho(u) + \rho(v)]/c \leq 2\rho(u)/c$ and $\|x^*\| \leq 2\rho(u)/c$. Therefore, by (2.6),

$$(2.7) \quad \|x\| \|x^*\| \leq 4\rho(u^2)/c^2 \leq (1+2c)(1+c)\rho(xx^*)/c^5.$$

But $\rho(xx^*) \leq \|xx^*\|$. This together with (2.7) completes the proof.

2.5. COROLLARY. *Under the hypotheses of Theorem 2.4, the norm of the involution as an operator on B does not exceed $(1+c)(1+2c^2)/c^5$.*

In (2.7) we may replace $\|x\| \|x^*\|$ by $\|x^*\|^2$ and $\rho(xx^*)$ by $\|x\| \|x^*\|$. This gives $\|x^*\| \leq (1+c)(1+2c^2)\|x\|/c^5$.

We denote by $P(N)$ the set of $x \in B$ such that $sp(x^*x) \subset [0, \infty)(sp(x^*x) \subset (-\infty, 0])$.

2.6. LEMMA. *For an Arens*-algebra B the following are equivalent.*

- (a) B is a B^* -algebra in an equivalent norm.
- (b) $N = (0)$.
- (c) $P = B$.

Suppose that $N = (0)$. Let $y \in B$. Since the involution on B is continuous, the element y^*y generates a closed*-subalgebra B_0 . Let \mathfrak{M} be the space of regular maximal ideals of B_0 . By [1, p. 279] the commutative algebra B_0 is *-isomorphic to $C(\mathfrak{M})$. Also $sp(y^*y)$ is real. Hence there exist $u, v \in B_0$ such that $u(M) = \sup(y^*y(M), 0)$ and $v(M) = -\inf(y^*y(M), 0)$, $M \in \mathfrak{M}$. Then u and v are s. a., $y^*y = u - v$ and $uv = 0$. As in [14, p. 281], $(yv)^*(yv) = -v^3$ so that $yv = 0$ by hypothesis. Then $v = 0$ and $sp(y^*y) \subset [0, \infty)$.

A theorem of Gelfand and Neumark [13] asserts that if B is semi-simple, has a continuous involution, is symmetric ($B = P$) and has an identity then there exists a faithful*-representation $x \rightarrow T_x$ of B . This theorem is also valid when B has no identity [4, Theorem 2.16]. In our situation, B is semi-simple [18, Lemma 3.5] and the involution is continuous. Thus a faithful*-representation exists. This representation is bi-continuous by [18, Corollary 4.4].

That (a) implies (b) follows from the well-known fact that any B^* -algebra is symmetric [14, p. 207 and p. 281].

The equation $4t^3 - 2t^2 + t - 1 = 0$ has exactly one real root a . This root a lies between .676 and .677.

2.7. THEOREM. *Suppose that for each s. a. element h , $\rho(h) \geq c\|h\|$ and $sp(h)$ is real, where $c > 0$. Then there is an equivalent norm for B in which B is a B^* -algebra if $c > a$.*

Suppose that $sp(x^*x) \subset (-\infty, 0]$. By Lemma 2.6 it is sufficient to show that $x = 0$. Suppose that $x \neq 0$. By Theorem 2.4 it is clear that $x^*x \neq 0$ and $\rho(x^*x) \neq 0$. Set $x = u + iv$ where u and v are s. a. As in the proof of Theorem 2.4, $xx^* + x^*x = 2u^2 + 2v^2$ and we may assume that $\rho(u) \geq \rho(v)$. Since $sp(u^2) \subset [0, \infty)$, $sp(v^2) \subset [0, \infty)$ formula 2.3 shows that $sp(2u^2 + 2v^2) \subset [-(1-c)\rho(2u^2)/c, (1+c)\rho(2u^2)/c]$. Let $r > 0, t > 0$. From Lemma 2.2,

$\|[-t^{-1}(2u^2 + 2v^2)]'\| < r$ if $t > (1 - cr)(1 + c)\rho(2u^2)/(c^2r)$ and $t > (1 + cr)(1 - c)\rho(2u^2)/(c^2r)$.

We write $x^*x = 2u^2 + 2v^2 + (-xx^*)$. By Lemma 2.2, $\|[-t^{-1}(-xx^*)]'\| < r^{-1}$ if $t > 0$ and $t > (r - c)\rho(x^*x)/c$ since $sp(-xx^*) \subset [0, \rho(x^*x)]$. By Lemma 2.3, $(-t^{-1}x^*x)'$ exists if $t > \max\{(1 + cr)(1 - c)\rho(2u^2)/c^2r, (1 - cr)(1 + c)\rho(2u^2)/c^2r, (r - c)\rho(x^*x)/c\}$. Since $sp(x^*x) \subset (-\infty, 0]$, $\rho(x^*x)$ cannot exceed this maximum. Now select $r, 1 \leq r < 2c$ which is possible since $c > a$. Then $(r - c)/c < 1$ and $(1 + cr)(1 - c) \geq (1 - cr)(1 + c)$. Therefore $\rho(x^*x) \leq (1 + cr)(1 - c)\rho(2u^2)/c^2r$. Letting $r \rightarrow 2c$ we obtain

$$(2.8) \quad \rho(x^*x) \leq (1 + 2c^2)(1 - c)\rho(2u^2)/2c^3 .$$

Next we express $-2u^2 = 2v^2 + (-xx^* - x^*x)$. By formula (2.3), $sp(-xx^* - x^*x) \subset [-(1 - c)\rho(x^*x)/c, (1 + c)\rho(x^*x)/c]$. Recall that $\rho(2v^2) \leq \rho(2u^2)$. Repeating the above reasoning we see that for $r > 0, t > 0$, $(-t^{-1}(-2u^2))'$ exists for $t > \max\{1 - cr, (1 + c)\rho(x^*x)/c^2r, (1 + cr)(1 - c)\rho(x^*x)/c^2r, (r - c)\rho(2u^2)/c\}$. But $sp(-2u^2) \subset (-\infty, 0]$. Then by the argument above we obtain

$$(2.9) \quad \rho(2u^2) \leq (1 + 2c^2)(1 - c)\rho(x^*x)/2c^3 .$$

From formulas (2.8) and (2.9) we see that $(1 + 2c^2)(1 - c) \geq 2c^3$ or $4c^3 - 2c^2 + c - 1 \leq 0$. This gives $c \leq a$ which is impossible by hypothesis.

Thus if $c > a$ we have $N = (0)$. We subsequently show (Corollary 2.11) that, in any case, N and P are closed in an Arens*-algebra B .

Following Rickart [16, p. 625] we say that B is an A^* -algebra if there exists in B an auxiliary normed-algebra norm $|x|$ (B need not be complete in this norm) such that, for some $c > 0, |x^*x| \geq c|x|^2$. He raises the question of whether every A^* -algebra has a faithful*-representation.

2.8. COROLLARY. *An A^* -algebra B where $|x^*x| \geq c|x|^2, x \in B$, in the auxiliary norm has a faithful*-representation if $c > a$.*

Observe that $|x^*||x| \geq c|x|^2$ so that $|x^*| \leq c^{-1}|x|, x \in B$. Thus the involution on B is continuous in the topology provided by the norm $|x|$. Let B_0 be the completion of B in the norm $|x|$. We extend the function $|x|$ from B to B_0 by continuity. Likewise the involution $x \rightarrow x^*$ can be extended by continuity to provide a continuous involution $y \rightarrow y^*$ on B_0 . We then have $|y^*y| \geq c|y|^2, y \in B_0$. As in [16, p. 626] we obtain $\rho(h) \geq c|h|$ for h s. a. in B_0 where $\rho(h)$ is the spectral radius computed for h as an element of the Banach algebra $B_0, \rho(h) = \lim |h^n|^{1/n}$. Also $|y^*y| \geq c^2|y^*||y|, y \in B_0$, so that B_0 is an Arens*-algebra. Hence, by Lemma 2.1, the spectrum of each s. a. element of B_0 is real. By Theorem 2.7, B_0 is a B^* -algebra in an equivalent norm. Therefore B has the desired faithful*-representation.

We have no information on the truth or falsity of Theorem 2.7 for $c \leq a$.

To prove Theorem 2.7 without restriction on the size of c one can assume without loss of generality that B has an identity. For suppose that B has no identity, $\|x^*x\| \geq k\|x^*\| \|x\|$, $x \in B$. Adjoin an identity e to B to form the algebra B_1 with the norm defined in B_1 by the rule

$$\|\lambda e + x\| = \sup_{\substack{\|y\|=1 \\ y \in B}} \|\lambda y + xy\|.$$

Then B_1 is a Banach algebra with the involution $(\lambda e + x)^* = \bar{\lambda}e + x^*[1, p. 275]$. By changing in minor ways arguments in [14, p. 207] we see that B_1 is an Arens*-algebra. There is a constant K such that $\|x^*\| \leq K\|x\|$, $x \in B$. Choose $0 < r < 1$. Given $\lambda e + x \in B_1$ there exists $y \in B$, $\|y\|=1$, such that

$$\begin{aligned} r^2 \|\lambda e + x\|^2 &< \|\lambda y + xy\|^2 \leq K \|(\lambda y + xy)^*\| \|\lambda y + xy\| \\ &\leq Kk^{-1} \|y^*(\lambda e + x)^*(\lambda e + x)y\| \\ &\leq K^2k^{-1} \|(\lambda e + x)^*(\lambda e + x)\|. \end{aligned}$$

Then

$$\|(\lambda e + x)^*(\lambda e + x)\| \geq kK^{-2} \|\lambda e + x\|^2 \geq (kK^{-2})^2 \|\lambda e + x\| \|(\lambda e + x)^*\|.$$

We use this fact later.

Some results on spectral theory in Arens*-algebras were obtained by Newburgh [15]. In a B^* -algebra $\rho(x)$ is a continuous function on the set H of s. a. elements since $\rho(h) = \|h\|$, $h \in H$. This property holds for all Arens*-algebras.

2.9. THEOREM. *In any Arens*-algebra, $\rho(x)$ is a continuous function on H .*

We assume that $\rho(h) \geq c\|h\|$ and $sp(h)$ is real, $h \in H$. We shall use the following principle [12, p. 67]. If y' exists and $\|z\| < (1 + \|y'\|)^{-1}$ then $(y + z)'$ exists.

Let $h \in H$, $h \neq 0$. Select $t > \rho(h)$ and set $u = (t^{-1}h)'$. We proceed as in the proof of Lemma 2.2. Let B_0 be the closed*-subalgebra generated by h and let \mathfrak{M} be its space of regular maximal ideals. Then $u \in B_0$. Since $t^{-1}h \circ u = 0$ we obtain, for each $M \in \mathfrak{M}$, $u(M) = h(M)/(h(M) - t)$. Since $\lambda/(\lambda - t)$ is a decreasing function of λ , $\sup |u(M)|$ can be majorized by $\rho(h)/(t - \rho(h))$. Then $(1 + \|u\|)^{-1} \geq (1 + c^{-1}\rho(u))^{-1} \geq c(t - \rho(h))/(ct + (1 - c)\rho(h)) = a(t)$, say.

Therefore $t^{-1}h + t^{-1}h_1$ is quasi-regular if $\|t^{-1}h_1\| < a(t)$ or if

$$(2.10) \quad ct^2 - c[\rho(h) + \|h_1\|]t - (1 - c)\rho(h) \|h_1\| > 0.$$

We apply this to $h_1 \in H$, $\|h_1\| < \rho(h)$. The larger zero d of the left hand side of (2.10) is given by

$$(2.11) \quad 2d = \rho(h) + \|h_1\| + [(\rho(h) - \|h_1\|)^2 + 4c^{-1}\rho(h)\|h_1\|]^{1/2}.$$

The radical term of (2.11) is majorized by $\rho(h) - \|h_1\| + 2(c^{-1}\rho(h)\|h_1\|)^{1/2}$. Hence $d \leq \rho(h) + (c^{-1}\rho(h)\|h_1\|)^{1/2}$. Thus $t \notin sp(h + h_1)$ if $t > \rho(h) + (c^{-1}\rho(h)\|h_1\|)^{1/2}$. Likewise $t \notin sp(-h - h_1)$ under the same condition. This shows that

$$(2.12) \quad \rho(h + h_1) \leq \rho(h) + (c^{-1}\rho(h)\|h_1\|)^{1/2}.$$

provided $h_1 \in H$ and $\|h_1\| < \rho(h)$.

Note that $\rho(h + h_1) \geq c\|h + h_1\| \geq c(\|h\| - \|h_1\|) \geq c(\rho(h) - \|h_1\|)$. Therefore if $\|h_1\| < c(\rho(h) - \|h_1\|)$ or equivalently if $\|h_1\| < c\rho(h)/(1 + c)$ we have $\|h_1\| < \rho(h + h_1)$. We may then apply the above analysis to the pair of s. a. elements $(h + h_1), -h_1$, to obtain (if $\|h_1\| < c\rho(h)/(1 + c)$)

$$(2.13) \quad \rho(h) \leq \rho(h_1 + h_2) + (c^{-1}\rho(h + h_1)\|h_1\|)^{1/2}.$$

From (2.12), $\rho(h + h_1) \leq [c^{-1/2} + 1]\rho(h)$. Inserting this estimate in the radical term of (2.13) we obtain

$$(2.14) \quad \rho(h) \leq \rho(h + h_1) + (c^{-1} + c^{-3/2})^{1/2}(\rho(h)\|h_1\|)^{1/2}$$

Combining (2.12) and (2.14) we obtain

$$|\rho(h + h_1) - \rho(h)| \leq [(c^{-1} + c^{-3/2})\rho(h)\|h_1\|]^{1/2}$$

provided $\|h_1\| < c\rho(h)/(1 + c)$.

This show that $\rho(x)$ is continuous on H at $x = h$. Clearly we have continuity on H at $x = 0$.

For x s.a. in an Arens*-algebra let $[a(x), b(x)]$ be the smallest closed interval containing $sp(x)$.

2.10. COROLLARY. *For an Arens*-algebra B , $a(x)$ and $b(x)$ are continuous functions of x on H .*

As remarks above indicate, there is no loss of generality in supposing that B has an identity e . Let h be s.a. Choose $\lambda > 0$ such that $sp(\lambda e + h) \subset [1, \infty)$. Let $h_n \rightarrow h$, where each h_n is s.a., and choose $0 < \varepsilon < 1$. We have $\rho(\lambda e + h) = b(\lambda e + h) = \lambda + b(h)$. By the "spectral continuity theorem" (see e. g. [15, Theorem 1]) for all n sufficiently large $sp(\lambda e + h_n) \subset (1 - \varepsilon, b(\lambda e + h) + \varepsilon)$. Also for all n sufficiently large $|\rho(\lambda e + h_n) - \rho(\lambda e + h)| < \varepsilon$ by Theorem 2.9. Since, for such n , $sp(\lambda e + h_n) \subset (0, \infty)$, then $\lambda + b(h_n) = \rho(\lambda e + h_n) \rightarrow \lambda + b(h)$. Therefore $b(h_n) \rightarrow b(h)$. A similar argument shows that $a(h_n) \rightarrow a(h)$.

2.11. COROLLARY. *For an Arens*-algebra B , N and P are closed sets.*

This follows directly from the continuity of the involution on B and Corollary 2.10. Likewise the set H^+ of all s.a. elements whose spectrum is non-negative is closed.

3. Faithful*-representations. Let B be a Banach algebra with an involution $x \rightarrow x^*$. Our aim here is to give necessary and sufficient conditions for B to possess a faithful*-representation. Our criterion (Theorem 3.4) is in terms of algebraic and linear space properties of B . A criterion of Kelley and Vaught [10] is largely topological in nature. To discuss this we first prove a simple lemma. We adopt the following notation. Let R_0 be the collection of all finite sums of elements of B of the form x^*x . Let $R = \{x \in H \mid \text{there exists } y \in R_0 \text{ such that } ty + (1-t)x \in R_0, 0 < t \leq 1\}$. In the notation of Klee [11, p. 448], $R = \text{lin } R_0$ (computed in the real linear space H , the union in H of R_0 and the points of H linearly accessible from R_0). Let P be the closure in B of R_0 . If B has an identity e and the involution is continuous then H is closed, e is an interior point of R_0 [10] and $R = P$ [11, p. 448]. If B has no identity or if the involution is not assumed continuous we see no relation, in general, between R and P other than $R \subset P$.

3.1. LEMMA. *Suppose that B has a continuous involution $x \rightarrow x^*$ and an identity e . Then there is an equivalent Banach algebra norm $\|x\|_1$ where $\|x^*\|_1 = \|x\|_1$, $x \in B$, and $\|e\|_1 = 1$.*

We first introduce an equivalent norm $\|x\|_0$ in which $\|x^*\|_0 = \|x\|_0$, $x \in B$, by setting $\|x\|_0 = \max(\|x\|, \|x^*\|)$. Let $L_x(R_x)$ be the operator on B defined by left (right) multiplication by x ; $L_x(y) = xy$ and $R_x(y) = yx$. Let $\|L_x\|$ be the norm of L_x as an operator on B where the norm $\|y\|_0$ is used for B . $\|R_x\|$ is defined in the same way. We set $\|x\|_1 = \max(\|L_x\|, \|R_x\|)$. Then $\|x + y\|_1 \leq \|x\|_1 + \|y\|_1$ and $\|xy\|_1 \leq \|x\|_1 \|y\|_1$. Clearly $\|x\|_1 \leq \|x\|_0$. Moreover $\|L_x\| \geq \|x\|_0 / \|e\|_0$ and the norms $\|x\|_0$ and $\|x\|_1$ are equivalent. Trivially $\|e\|_1 = 1$. Also

$$\|L_{x^*}\| = \sup_{\|y\|_0=1} \|x^*y\|_0 = \sup_{\|y\|_0=1} \|y^*x\|_0 = \|R_x\|.$$

Then $\|x^*\|_1 = \max(\|L_{x^*}\|, \|R_{x^*}\|) = \max(\|L_x\|, \|R_x\|) = \|x\|_1$.

In view of Lemma 3.1 the result [10, p. 51] of Kelley and Vaught in question may be expressed as follows.

3.2. THEOREM. *Let B be a Banach algebra with an identity and an involution $x \rightarrow x^*$. Then B has a faithful*-representation if and only if $*$ is continuous and $P \cap (-P) = (0)$.*

As it stands this criterion breaks down if B has no identity. For let $B = C([0, 1])$ with the usual involution $x \rightarrow x^*$ and norm. Let B_0 be the algebra obtained from B by keeping the norm and involution but defining all products to be zero. Then $*$ is still continuous and $P \cap (-P) = (0)$. But B_0 has no faithful*-representation, for otherwise B_0 would be semi-simple [16, p. 626].

As in [4] we call the involution $x \rightarrow x^*$ in B *regular* if, for h s.a., $\rho(h) = 0$ implies $h = 0$. By [4, Lemma 2.15]. $*$ is regular if and only if every

maximal commutative $*$ -subalgebra of B is semi-simple. Also every maximal commutative $*$ -subalgebra of B is closed [4, Lemma 2.13].

By a *positive linear functional* f on B we mean a linear functional such that $f(x^*x) \geq 0, x \in B$. The functional f is not assumed to be continuous. If B has an identity then [13, p. 115], $f(h)$ is real for h s.a. and $f(x^*) = \overline{f(x)}$. Trivial examples show this to be false, in general. However, from the positivity of $f, f(x^*y)$ and $f(y^*x)$ are complex conjugates which is the fact really needed for the introduction of the inner product in Theorem 3.4.

3.3 LEMMA. *Let the involution on B be regular. Then*

(1) *a positive linear f satisfies the inequalities*

$$(3.1) \quad f(y^*hy) \leq f(y^*y) \|h\|, y \in B, h \in H,$$

$$(3.2) \quad f(y^*x^*xy) \leq f(y^*y) \|x^*x\|, x, y \in B,$$

(2) *if B has an identity e , any $h \in H, \|e - h\| \leq 1$ has a s.a. square root and, moreover, any positive linear functional is continuous on H .*

Suppose first that B has an identity $e, \|e - h\| \leq 1, h$ s.a. In the course of the proof of [4, Theorem 2.16] it was shown that h has a s.a. square root. Next do not assume that B has an identity. Let B_1 be the Banach algebra obtained by adjoining an identity e to B . Consider the power series $(1 - t)^{1/2} = 1 - t/2 - t^2/8 \dots$. Let $h \in B, h$ s.a. and $\|h\| \leq 1$. Then the expansion $-h/2 - h^2/8 - \dots$ converges to an element $z \in B$. Let B_0 be a maximal abelian $*$ -subalgebra of B containing h . As noted above, B_0 is a semi-simple Banach algebra. The involution is continuous on B_0 ([16, Corollary 6.3]). Therefore z is s.a. Also $(e + z)^2 = e - h$. Let $y \in B$ and set $k = y + zy$. Then $k^*k = (y^* + y^*z)(y + zy) = y^*(e + z)^2y = y^*y - y^*hy$. For any positive linear functional f on $B, f(k^*k) \geq 0$ which yields (3.1). Formula (3.2) is a special case.

Suppose that B has an identity e . If we set $y = e$ in (3.1) we obtain $|f(h)| \leq f(e) \|h\|$ which shows that f is continuous on H .

3.4. THEOREM. *B has a faithful $*$ -representation if and only if f^* is regular and $R \cap (-R) = (0)$.*

Suppose that B has a faithful $*$ -representation $x \rightarrow T_x$ as operators on a Hilbert space \mathfrak{H} . Let h be s.a. and $\rho(h) = 0$. Then $\rho(T_h) = 0$. As T_h is a s.a. operator on a Hilbert space, $T_h = 0$ and therefore $h = 0$. Thus the involution is regular. Let $x \in R \cap (-R)$ and let f be a positive linear functional on B . Then clearly $f(y) \geq 0, y \in R_0$. From the definition of R there exists $y \in R_0$ such that $tf(y) + (1 - t)f(x) \geq 0, 0 < t \leq 1$. It follows that $f(x) \geq 0$ and hence $f(x) = 0$. Let $\xi \in \mathfrak{H}$ and set $f(x) = (T_x\xi, \xi)$. Then $(T_x\xi, \xi) = 0$ for all $\xi \in \mathfrak{H}$. Since T_x is a s.a. operator we see that $T_x = 0$ and $x = 0$.

Suppose now that $*$ is regular and $R \cap (-R) = (0)$. We show first that the regularity of the involution makes available a general representation procedure of Gelfand and Neumark [13].

Let f be a positive linear functional on B . Let $I_f = \{x | f(x^*x) = 0\}$. I_f is a left ideal of B . Let π be the natural homomorphism of B onto B/I_f . Since $f(x^*y) = f(y^*x)$, $\mathfrak{H}'_f = B/I_f$ is a pre-Hilbert space if we define $(\pi(x), \pi(y)) = f(y^*x)$. As in [13, p. 120] we associate with $y \in B$ an operator A'_y on \mathfrak{H}'_f defined by $A'_y[\pi(x)] = \pi(yx)$. Formula (3.2) yields

$$(3.3) \quad \|A'_y[\pi(x)]\|^2 = f(x^*y^*yx) \leq \|y^*y\| \|\pi(x)\|^2.$$

Thus A'_y is a bounded operator with norm not exceeding $\|y^*y\|^{1/2}$. It may then be extended to T'_y , a bounded operator on the completion \mathfrak{H}_f of \mathfrak{H}'_f . The mapping $x \rightarrow T'_x$ is a $*$ -representation of B with kernel $\{y \in B | yx \in I_f, \text{ for all } x \in B\} = K$. Note that $K^* = K$.

Now take the direct sum \mathfrak{H} of the Hilbert spaces \mathfrak{H}_f as f ranges over all positive linear functionals on B ([13, p. 95]). Since $\|T'_y\| \leq \|y^*y\|^{1/2}$ by (3.3) and this estimate is independent of f , the direct sum ([13, p. 113]) $x \rightarrow T_x$ of the representations $x \rightarrow T'_x$ yields a $*$ -representation of B as bounded operators on \mathfrak{H} with kernel $\{y \in B | yx \in \cap I_f, \text{ for all } x \in B\}$. If B has an identity, the kernel is the reducing ideal of B ([13, p. 130]), namely $\cap I_f$.

Suppose first that B has an identity e . The set R_0 has the property that $x, y \in R_0, \lambda, \mu \geq 0$ imply $\lambda x + \mu y \in R_0$. By Lemma 3.3, $R_0 \supset \{x \in H | \|e - x\| \leq 1\}$. Thus e is an interior point of R_0 . By the theory of convex sets in normed linear spaces, R is the closure in H of R_0 and R is a closed cone in H ([11, p. 448]).

Let f be a positive linear functional on B . By Lemma 3.2, f is continuous on H . Also $f(w) \geq 0, w \in R$. Let H' be the conjugate space of H and $G = \{g \in H' | g(w) \geq 0, w \in R\}$. It is easy to see ([10, p. 48]) that G , the dual cone of R , is the set of linear functionals on H which are the restrictions to H of positive linear functional on B . There is no loss generality in assuming that $\|e\| = 1$. Let $x \in H$. By [10, Lemma 1.3], $\text{dist}(-x, R) = \sup \{g(x) | g \in G, g(e) \leq 1\}$.

We show that $R \cap (-R) = H \cap (\cap I_f)$. Let $y \in H, y \in \cap I_f$. For any fixed $f, T'_y = 0$ and $(T'_y\xi, \xi) = 0, \xi \in \mathfrak{H}_f$. Then $(\pi(yx), \pi(x)) = 0$ for all $x \in B$ in the notation used above. Therefore $f(x^*yx) = 0, x \in B$. Setting $x = e$ we see that $f(y) = 0$. Then by the distance formula, $-y \in R$. Likewise $y \in R$. Suppose conversely that $y \in R \cap (-R)$. It is easy to see that for each $z \in B, z^*R_0z \subset R_0$. Therefore $z^*Rz \subset R$. Hence $z^*yz \in R \cap (-R), z \in B$. From the distance formula, $\sup \{f(z^*yz) | f \text{ positive}, f(e) \leq 1\} = 0 = \sup \{f(-z^*yz) | f \text{ positive}, f(e) \leq 1\}$. Hence $f(z^*yz) = 0$ for each positive linear functional. Then $(T'_y\pi(z), \pi(z)) = 0$ for all z whence $T'_y = 0$. Therefore $T_y = 0$ and $y \in H \cap (\cap I_f)$.

This proves the theorem in case B has an identity. Suppose that B has no identity. Let B_1 be the algebra obtained by adjoining an identity e to B . We extend the involution to B_1 by setting $(\lambda e + x)^* = \overline{\lambda e + x}^*$. The involution on B_1 is regular [4, Lemma 2.14]. Let R'_0 and R' be the sets R_0 and R respectively computed for the algebra B_1 . By the above it is sufficient to show that $R \cap (-R) = (0)$ implies $R' \cap (-R') = (0)$. Suppose that $R \cap (-R) = (0)$.

Let $x, y \in B$. Then $y^*(\lambda e + x)^*(\lambda e + x)y = (\lambda y + xy)^*(\lambda y + xy)$. This shows that $y^*R'_0y \subset R_0$ which implies $y^*R'y \subset R$. Note also that B is semi-simple [18, Lemma 3.5] which implies that $zB = (0)$, or $Bz = (0)$, $z \in B$, can hold only for $z = 0$.

Suppose that $\lambda e + x \in R' \cap (-R')$ where $x \in B$ and λ is a scalar. We derive a contradiction from $\lambda \neq 0$. For every $y \in B$, $y^*(\lambda e + x)y \in R \cap (-R)$. Setting $u = -x/\lambda$ we have $y^*(e - u)y = 0$ or $y^*y = y^*uy$ for all $y \in B$. Then

$$(3.4) \quad h^2 = huh, \text{ h s.a.}$$

Let h_1 and h_2 be s.a. Then $(h_1 + h_2)^2 = (h_1 + h_2)u(h_1 + h_2)$. From (3.4) we obtain

$$(3.5) \quad h_1h_2 + h_2h_1 = h_1uh_2 + h_2uh_1.$$

Also $(h_1 - ih_2)(h_1 + ih_2) = (h_1 - ih_2)u(h_1 + ih_2)$ From (3.4) we get

$$(3.6) \quad h_2h_1 - h_1h_2 = h_2uh_1 - h_1uh_2$$

From (3.5) and (3.6) we see that $h_1h_2 = h_1uh_2$. Consequently for h_k s.a., $k = 1, 2, 3, 4$, we see that $(h_1 + ih_2)(h_3 + ih_4) = (h_1 + ih_2)u(h_3 + ih_4)$. In other words

$$(3.7) \quad zw = zuw, z, w \in B.$$

From (3.7) $(z - zu)w = 0$ for all $w \in B$ so that $z = zu$ for each z . Hence u is a right identity for B . Likewise from $z(w - uw) = 0$ for all $z \in B$ we see that u is an identity for B . But this is impossible since we are considering the case where B has no identity.

We now have $x \in R' \cap (-R')$. Then $y^*xy = 0$ for all $y \in B$. Therefore $h_xh = 0$, h s.a. Also for h_k s.a., $k = 1, 2$, $(h_1 + h_2)x(h_1 + h_2) = 0$ so that $h_2xh_1 + h_1xh_2 = 0$. Also $(h_1 - ih_2)x(h_1 + ih_2) = 0$ so that $h_1xh_2 - h_2xh_1 = 0$. Therefore $h_1xh_2 = 0$. It follows that $zxw = 0$ for all $z, w \in B$. This implies that $x = 0$ and completes the proof.

4. Preliminary ring theory. Let R be a semi-simple ring with minimal one-sided ideals. For a subset A of R let $\mathfrak{L}(A) = \{x \in R \mid xA = (0)\}$ and $\mathfrak{R}(A) = \{x \in R \mid Ax = (0)\}$. Consider a two-sided I of R . If $x \in R(I)$, $y \in R$, $z \in I$ then $zy \in I$, $z(yx) = 0$ so that $\mathfrak{R}(I)$ is a two-sided ideal of R . Therefore $\mathfrak{R}(I)I$ is an ideal. But $[\mathfrak{R}(I)I]^2 = (0)$. Thus, by semi-simplicity, $\mathfrak{R}(I)I = (0)$

and $\mathfrak{R}(I) \subset \mathfrak{L}(I)$. Likewise we have $\mathfrak{L}(I) \subset \mathfrak{R}(I)$ and thus $\mathfrak{R}(I) = \mathfrak{L}(I)$. Let S be the socle [5, p. 64] of R . This is the algebraic sum of the minimal left (right) ideals of R . S is a two-sided ideal. Therefore $\mathfrak{L}(S) = \mathfrak{R}(S)$. This set we denote by S^\perp . Note that $S \cap S^\perp = (0)$.

We call an idempotent e of R a *minimal idempotent* if eR is a minimal right ideal.

4.1. LEMMA. (a) *Let I be a left (right) ideal of R , $I \neq (0)$. Then I contains no minimal left (right) ideal of R if and only if $I \subset S^\perp$.*

(b) *R/S^\perp is semi-simple. If S_0 is the socle of R/S^\perp then $S_0^\perp = (0)$.*

Let $I \neq (0)$ be a left ideal of R . Suppose that $I \subset S^\perp$. Then I cannot contain a minimal left ideal J of R for any such J would be contained in $S \cap S^\perp$. Next suppose that $I \not\subset S^\perp$. We must show that I contains a minimal left ideal of R . There exists a minimal idempotent e such that $eI \neq (0)$. Choose $u \in I$ such that $eu \neq 0$. By semi-simplicity and the minimality of eR , $eR = euR$. Thus there exists $z \in R$ such that $eu z = e$. Since $(eu z)^2 = e$, we have $j \neq 0$ where $j = zu$. Note that $j^2 = j$. As $u \in I$ we have $Rj \subset I$. To see that Rj is the desired minimal ideal it is sufficient to see that jRj is a division ring [5, p. 65].

Note that $jz = zu z = ze \neq 0$. Then $Rze = Re$ so that there exists $v \in R$ where $vze = e$. Then $vj = vzeu = eu$ and $vz = e$.

We assert that $jx_1j = jx_2j$ if and only if $eux_1ze = eux_2ze$. For if $jx_1j = jx_2j$, multiply on the left by v and on the right by z and use the relations $vj = eu$ and $jz = ze$. If $eux_1ze = eux_2ze$ multiply on the left by z and on the right by u and use $zeu = j$.

Therefore the mapping $\tau: \tau(jxj) = euxze$ is a well-defined one-to-one mapping of jRj into eRe . The mapping is onto. For let $ewe \in eRe$. Then $ewe = eu z w v z e = \tau(jz w v j)$. τ is clearly additive. But also $\tau[(jxj)(jyj)] = \tau(jxjyj) = euxjyze = (euxze)(euyze) = \tau(jxj)\tau(jyj)$. Therefore τ is a ring isomorphism of jRj onto eRe . Since eRe is a division ring so is jRj .

Let J be the radical of R/S^\perp and π be the natural homomorphism of R onto R/S^\perp . Suppose that $J \neq 0$. Then $\pi^{-1}(J) \supset S^\perp$ and $\pi^{-1}(J) \neq S^\perp$. By (a), $\pi^{-1}(J)$ contains a minimal idempotent e of R . We then have $\pi(e) \in J$, $\pi(e) \neq 0$. This is impossible since the radical of a ring contains no non-zero idempotents.

Let S_0 be the socle of R/S^\perp and e be a minimal idempotent of R . Clearly $\pi(e) \neq 0$ and π is one-to-one on eRe . Then $\pi(e)\pi(R)\pi(e)$ is a division ring so that, since R/S^\perp is semi-simple, $\pi(e) \in S_0$. Let $\pi(x) \in S_0^\perp$. Then $\pi(ex) = 0$ so that $ex \in S^\perp \cap S = (0)$. Hence $x \in S^\perp$ and $\pi(x) = 0$.

The following result is due to Rickart [17, Lemma 2.1.]:

4.2. LEMMA. *Let A be any ring. Let $x \rightarrow x^*$ be a mapping of A onto A such that $x^{**} = x$, $(xy)^* = y^*x^*$ and $xx^* = 0$ implies $x = 0$. Then any*

minimal right (left) ideal I of A can be written in the form $I=eA(I=Ae)$ where $e^2 = e \neq 0, e^* = e$.

We improve this result by relaxing the conditions on $x \rightarrow x^*$ but at the expense of assuming the ring to be semi-simple.

4.3. LEMMA. *Let R be semi-simple with minimal one-sided ideals. Let $x \rightarrow x^*$ be a mapping of R onto R satisfying $x^{**} = x$ and $(xy)^* = y^*x^*$. Then the following statements are equivalent.*

- (1) *Every minimal right ideal is generated by a s.a. idempotent.*
- (2) *Every minimal left ideal is generated by a s.a. idempotent.*
- (3) *$jj^* \neq 0$ for each minimal idempotent j of R .*
- (4) *$xx^* = 0$ implies $x \in S^\perp$*

We say that the idempotent e is s.a. if $e^* = e$. Note that $x \rightarrow x^*$ is one-to-one and $0^* = 0$. As a preliminary we show that j^* is a minimal idempotent if j is a minimal idempotent. The ideal $I = jR$ is a minimal right ideal. Then $I^* = Rj^*$ is a left ideal $\neq (0)$. Suppose $I^* \supset K \neq (0), I^* \neq K$ where K is a left ideal of R . By semi-simplicity there exists $x \in K$ such that $x^2 \neq 0$. Then $I^* \supset Rx \neq (0), I^* \neq Rx$. This implies that $I \supset x^*R \neq (0), I \neq x^*R$. This is impossible. Therefore I^* is a minimal left ideal and j^* is a minimal idempotent. It is clear from this argument that (1) and (2) imply each other.

Assume (1). Let j be a minimal idempotent, $I = Rj$ a minimal left ideal. We can write $I = Re$ where e is a s.a. idempotent. Then for some $v \in R, vj = e$. But $e = ee^* = vjj^*v$. Therefore $jj^* \neq 0$. Thus (1) implies (3).

Assume (3). Suppose that $xx^* = 0, x \neq 0$. Let $I = Rx$. Then $I \neq (0)$. Suppose that I contains a minimal left ideal Rj of R where j is a minimal idempotent. We can write $j = yx, y \in R$. Then $0 \neq jj^* = yxx^*y^* = 0$. This shows that I contains no minimal left ideal of R . By Lemma 4.1, $I \subset S^\perp$. Then for any minimal idempotent $e, 0 = e(ex)$ and $x \in S^\perp$. Thus (3) implies (4).

Assume (4). If j is a minimal idempotent and $jj^* = 0$ then $j \in S^\perp$. But $j \in S$ and $S \cap S^\perp = (0)$. This shows that (4) implies (3).

Assume (3). Let j be a minimal idempotent, $I = jR$. Since $jj^* \neq 0, jj^*R = I$. There exists $u \in R, jj^*u = j$. As noted above j^* is a minimal idempotent. By (3), $0 \neq j^*j$. Then $0 \neq (u^*jj^*)(jj^*u) = u^*(jj^*)^2u$. Therefore $(jj^*)^2 \neq 0$. Set $h = jj^*$. Since I is minimal, $I = hI$. As in the proof of [17, Lemma 2.1] there exists $u \in I$ such that $h = hu$. Set $e = uu^*$. As in that proof, e is a s.a. idempotent and it remains only to check that $e \neq 0$ to obtain (2) from (3). If $e = 0$ then $0 = uu^* = hu u^*h = h^2$ which is impossible.

5. Normed algebras with minimal ideals. We are concerned here with $*$ -representations of semi-simple normed algebras B with an involution

where B has minimal one-sided ideals. B may be incomplete.

5.1. LEMMA. *Let B be a complex semi-simple normed algebra with minimal one-sided ideals. Let e_1, e_2 be minimal idempotents of B . Then the following statements are equivalent.*

- (1) $e_1Be_2 \neq (0)$.
- (2) $e_2Be_1 \neq (0)$,
- (3) e_1Be_2 is one-dimensional.
- (4) e_2Be_1 is one-dimensional.

Suppose (1). There exists $u \in B, e_1ue_2 \neq 0$. Since $e_1ue_2B = e_1B$, there exists $v \in B$ where $e_1ue_2v = e_1$. Then $e_2ve_1 \neq 0$ and (1) implies (2). Let $E = \{\lambda e_2ve_1 \mid \lambda \text{ complex}\}$. Clearly $e_2Be_1 \supset E$. Let $e_2xe_1 \in e_2Be_1$. Then $e_2xe_1 = e_2x(e_1ue_2ve_1) = (e_2xe_1ue_2)e_2ve_2$, a scalar multiple of e_2 by the Gelfand-Mazur Theorem. Thus (1) implies (4). The remainder of the argument is trivial.

For the remainder of § 5, B denotes a semi-simple complex normed algebra with an involution and with minimal one-sided ideals.

5.2. THEOREM. *The following statements concerning B are equivalent.*

- (1) *Every minimal one-sided ideal is generated by a s.a. idempotent.*
- (2) *There exists a $*$ -representation with kernel S^\perp .*
- (3) *There exists a $*$ -representation with kernel contained in S^\perp .*
- (4) *$j - j^*$ is quasi-regular for every minimal idempotent j .*
- (5) *$jBj^* \neq (0)$ for every minimal idempotent j and $xx^* = 0$ implies $x^*x \in S^\perp, x \in B$.*

Suppose that (1) holds. Let Q be the set of all s.a. minimal idempotents of B and let $j \in Q$. By the Gelfand-Mazur Theorem, $jBj = \{\lambda j \mid \lambda \text{ complex}\}$. Suppose $jx^*xj = \lambda j$. Taking adjoints, $\lambda = \bar{\lambda}$ so λ is real. We show that $jx^*xj = -j$ is impossible. For suppose $jx^*xj = -j$. Now $jxj = \alpha j$ for some scalar $\alpha = a + bi$, where a, b are real. Set $c = a + (a^2 + 1)^{1/2}$. By the use of $jx^*xj = -j$ one obtains $(jx^* - cj)(jx^* - cj)^* = 0$. From Lemma 4.3 we have $jx^* - cj = 0$. Then $(a - bi)j = jx^*j = cj$. It follows that $c = a$ and $b = 0$. This is impossible.

For $j \in Q$ we define the functional $f_j(x)$ on B by the rule $f_j(x)j = jxj$. By the above, $f_j(x^*x) \geq 0, x \in B, x \in B$ and $f_j(x^*) = \overline{f_j(x)}$. The functional f_j is a positive linear functional on B and is continuous on B .

The following inequality of Kaplansky [9, p. 55] is then available.

$$(5.1) \quad f_j(y^*x^*xy) \leq \nu(x^*x)f_j(y^*y), \quad x, y \in B,$$

where $\nu(x^*x) = \lim ||(x^*x)^n||^{1/n}$. Let $I_j = \{x \mid f_j(x^*x) = 0\}$. Let π be the natural homomorphism of B onto B/I_j . The definition $(\pi(x), \pi(y)) = f_j(y^*x)$ makes B/I_j a pre-Hilbert space. Let \mathfrak{H}_j be its completion. See the discussion of the Gelfand-Neumark procedure in § 3. To each $y \in B$ we correspond

the operator A_y^j defined by $A_y^j[\pi(x)] = \pi(yx)$. Then

$$\|A_y^j[\pi(x)]\|^2 = f_j(x^*y^*yx) \leq \nu(y^*y) \|\pi(x)\|^2$$

by (5.1). Thus A_y^j can be extended to a bounded linear operator T_y^j on \mathfrak{H}_j , and the mapping $y \rightarrow T_y^j$ is a $*$ -representation of B .

Since $\|T_y^j\| \leq \nu(y^*y)^{1/2}$ and the estimate is independent of $j \in Q$ we can take the direct sum \mathfrak{H} of the Hilbert spaces $\mathfrak{H}_j, j \in Q$ and the direct sum $x \rightarrow T_x$ of the representations $x \rightarrow T_x^j$. This gives a $*$ -representation of B with kernel K where

$$K = \{x \in B \mid xy \in \bigcap_{j \in Q} I_j, \text{ for all } y \in B\} .$$

We show that $K = S^\perp$.

It is clear that $S^* = S$ and therefore $(S^\perp)^* = S^\perp$. Using this and Lemma 4.3 we obtain the following chain of equivalences: $x \in \bigcap I_j \leftrightarrow jx^*xj = 0, \text{ all } j \in Q \leftrightarrow jx^* \in S^\perp, \text{ all } j \in Q \leftrightarrow jx^* = 0, \text{ all } j \in Q \leftrightarrow x^* \in S^\perp \leftrightarrow x \in S^\perp$. Therefore $\bigcap I_j = S^\perp$. Thus $K = \{x \mid xy \in S^\perp, \text{ all } y \in B\}$. If $x \in K$ then $xj \in S^\perp \cap S = (0)$ for all $j \in Q$ and $x \in S^\perp$. Clearly $S^\perp \subset K$. Therefore $K = S^\perp$. Hence (1) implies (2). Clearly (2) implies (3).

Assume (3) and let φ be a $*$ -representation whose kernel $\subset S^\perp$. Let j be a minimal idempotent of B . Let A be the subalgebra of B generated by j and j^* . By the Gelfand-Mazur Theorem, $jj^*j = \lambda j$ for some scalar λ . Thus A is the linear space spanned by j, j^*, jj^* and j^*j . A is finite-dimensional and $A \subset S$. Since $S \cap S^\perp = (0)$, φ is one-to-one on A . Note that $A = A^*$. Let E be the B^* -algebra obtained by taking the closure in the operator algebra on the appropriate Hilbert space of $\varphi(B)$. Clearly $\varphi(A)$ is a closed $*$ -subalgebra of E . The element $\varphi(j - j^*)$ is a skew element of E and therefore quasi-regular in E . By [8, Theorem 4.2] its quasi-inverse in E already lies in $\varphi(A)$. As φ is one-to-one on $A, j - j^*$ has a quasi-inverse in A . Thus (3) implies (4).

Assume (4). Let j be a minimal idempotent of B . There exists $u \in B$ such that $j - j^* + u - (j - j^*)u = 0$. If $jj^* = 0$ then left multiplication by j gives $j = 0$ which is impossible. Therefore $jj^* \neq 0$. By Lemma 4.3, we see that (4) implies (1). Clearly (1) implies (5) by Lemma 4.3. Assume (5). Let j be a minimal idempotent of B . If $j^*j = 0$ then $0 = x^*j^*jx = (jx)^*(jx)$. Also $jxx^*j^* \in S^\perp \cap S = (0)$ for all $x \in B$. Since $jBj^* \neq (0), jBj^*$ is one-dimensional by Lemma 5.1. Hence there exists $u \neq 0$ in B and a linear functional $f(x)$ on B such that $jxj^* = f(x)u$. Then $f(xx^*) = 0$ for all $x \in B$. Expanding $0 = f[(x + y)(x + y)^*] = f[(x + iy)(x + iy)^*]$ we see that $f(xy^*) = 0$ for all $x, y \in B$. Hence f vanishes on B^2 . Take any $z \in B$. We have $f(jz) = 0$ or $jzj^* = 0$. Thus $jBj^* = (0)$ which is impossible. Therefore $j^*j \neq 0$. By Lemma 4.3, (5) implies (1).

Algebras to which Theorem 5.2 can be applied most easily are those for

which $S^\perp = (0)$. Examples are semi-simple annihilator algebras studied by Bonsall and Goldie [3] and primitive algebras (Corollary 5.4).

5.3. COROLLARY. *If B is an Arens*-algebra with non-zero socle then $N \subset S^\perp$.*

Let $x_0 \in N$, $sp(x_0x_0^*) \subset (-\infty, 0]$. Then we can write $x_0x_0^* = -h^2$ where h is s.a. The ideal S^\perp is closed and self-adjoint. Let π be the natural homomorphism of B onto B/S^\perp . An involution can be defined in B/S^\perp by the rule $[\pi(x)]^* = \pi(x^*)$. Since B is semi-simple, B/S^\perp has non-zero socle. Let $\pi(x)$ be a minimal idempotent of B/S^\perp . Then $[\pi(x)]^* - \pi(x) = \pi(x^* - x)$ is quasi-regular in B/S^\perp since $x^* - x$ is quasi-regular in B . By Theorem 5.2 and Lemma 4.1, B/S^\perp has a faithful*-representation. Then, by Theorem 3.4, $\pi(x_0x_0^*) = 0 = \pi(h^2)$. Therefore $x_0x_0^* \in S^\perp$ and $(jx_0)(jx_0)^* = 0$ for each minimal idempotent j of B . Therefore $jx_0 = 0$ for all such j and $x_0 \in S^\perp$.

We call the involution $x \rightarrow x^*$ *proper* if $xx^* = 0$ implies $x = 0$. We call the involution *quasi-proper* if $xx^* = 0$ implies $x^*x = 0$. Not every involution is quasi-proper. For example let B be all 2×2 matrices with the involution defined by

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix}^* = \begin{pmatrix} \bar{a} & -\bar{c} \\ -\bar{b} & \bar{d} \end{pmatrix}.$$

To see that this is not quasi-proper choose x as

$$\begin{pmatrix} 1 & i \\ 0 & 0 \end{pmatrix}.$$

Every proper involution is quasi-proper but the converse is false. Consider, for example $B = C([0, 1])$ and set $x^*(t) = \overline{x(1-t)}$.

5.4. COROLLARY. *Let B be primitive with non-zero socle. Then the following statements are equivalent.*

- (1) *The involution* is proper.*
- (2) *The involution* is quasi-proper.*
- (3) *There exists a faithful*-representation of B .*

Suppose that $S^\perp \neq (0)$. Then by [5, p. 75], $S \subset S^\perp$. Since $S \cap S^\perp = (0)$ this is impossible. Therefore $S^\perp = (0)$. Assume (2). Let j be a minimal idempotent of B . Then $jBj^* \neq (0)$ (see the proof of [16, Theorem 4.4]) and, consequently (5) of Theorem 5.2 is satisfied. Then by Theorem 5.2, (2) implies (3); the remainder of the proof is obvious.

The equivalence of (1) and (3) was noted by Rickart [17, Theorem 3.5]. By Lemma 4.3 and Theorem 5.2 this equivalence of (1) and (3) holds for any B for which $S^\perp = (0)$.

If B is complete the following statements hold. (1) Any*-representation of B is continuous [16, Theorem 6.2]. (2) If B has a faithful*-representation then the involution is continuous [16, Lemma 5.3]. We show that

both these statements can be false for B incomplete. Our discussion is based on work of Kakutani and Mackey [6, p. 56] (see also [7] for the complex case). Let \mathfrak{X} be an infinite-dimensional complex Hilbert space, $(x, x)^{1/2} = \|x\|$. Let $\|x\|$ be any other norm on \mathfrak{X} such that $\|x\| \leq \|x\|, x \in \mathfrak{X}$. Let $\mathfrak{X}_1 = \{y \in \mathfrak{X} | (x, y) \text{ is continuous on } \mathfrak{X} \text{ in the norm } \|x\|\}$ and endow \mathfrak{X}_1 with the norm $\|x\|$. Then [6, p. 56] a linear functional $f(x)$ on \mathfrak{X}_1 has the form $f(x) = (x, y)$. Moreover \mathfrak{X}_1 is dense in \mathfrak{X} in both norms. If there exists $c > 0$ such that $\|x\| \leq c \|x\|, x \in \mathfrak{X}_1$ then $\mathfrak{X} = \mathfrak{X}_1$ and \mathfrak{X}_1 is complete.

Let $\mathfrak{G}(\mathfrak{X}_1)$ be the normed algebra of all bounded linear operators on \mathfrak{X}_1 . As shown in [6, p. 56], $\mathfrak{G}(\mathfrak{X}_1)$ has an involution $T \rightarrow T^*$ where $(T(x), y) = (x, T^*(y)), x, y \in \mathfrak{X}_1$. In these terms we show the following.

5.5. THEOREM. *The following statements are equivalent.*

- (1) \mathfrak{X}_1 is complete.
- (2) The involution in $\mathfrak{G}(\mathfrak{X}_1)$ is continuous.
- (3) The faithful*-representation of Theorem 5.2 for $\mathfrak{G}(\mathfrak{X}_1)$ is continuous.

As already noted (1) implies (2) and (3). Assume (2) and let M be the norm of the involution. By [2] any minimal idempotent of $\mathfrak{G}(\mathfrak{X}_1)$ is one-dimensional and the operator J defined by the rule $J(x) = (x, u)u$ where $(u, u) = 1$ is a minimal idempotent. Since $(J(x), y) = (x, u)(u, y) = (x, J(y))$ we have $J = J^*$. The functional f defined by $f(U)J = JUJ$ is a continuous positive linear functional on $\mathfrak{G}(\mathfrak{X}_1)$. For $z \in \mathfrak{X}_1$ define the operator W_z by the rule $W_z(x) = (x, u)z$. Then we can write the norm of W_z as $C \|z\|$ where C is independent of z . A simple computation gives $JW_z^*W_zJ = (z, z)J$. By formula (5.1), where $\|U\|$ denotes the norm in $\mathfrak{G}(\mathfrak{X}_1)$,

$$\|z\|^2 = (z, z) \leq \nu(W_z^*W_z) \leq \|W_z^*W_z\| \leq C^2M \|z\|^2.$$

This shows that \mathfrak{X}_1 is complete.

Assume (3) and let N be the norm of the faithful*-representation. Let $I_f = \{U \in \mathfrak{G}(\mathfrak{X}_1) | f(U^*U) = 0\}$, π be the natural homomorphism of $\mathfrak{G}(\mathfrak{X}_1)$ onto $\mathfrak{G}(\mathfrak{X}_1)/I_f$ and $(\xi, \eta)_f$ be the inner product for the pre-Hilbert space $\mathfrak{G}(\mathfrak{X}_1)/I_f$. Let $V \rightarrow T_f$ be the partial*-representation induced by f . Its norm cannot exceed N . Now $(\pi(J), \pi(J))_f = 1$ and

$$N^2 \|U\|^2 \geq \|T_f[\pi(J)]\|^2 = (UJ, UJ)_f = f(JU^*UJ) = f(U^*U).$$

Applying this formula to $U = W_z$ we obtain $N^2C^2 \|z\|^2 \geq (z, z)$ and again \mathfrak{X}_1 is complete.

A specific example is suggested in [6, p. 57]. Let $\mathfrak{X} = l^2, \|\{x_n\}\| = \sup |x_n|$. An easy computation gives $\mathfrak{X}_1 = l^2 \cap l^1$ in the sup norm. Here the involution and*-representation are therefore not continuous.

6. Involutions on $\mathfrak{G}(\mathfrak{H})$. Let \mathfrak{H} be a Hilbert space and $\mathfrak{G}(\mathfrak{H})$ the B^* -

algebra of all bounded linear operators on \mathfrak{H} . We determine in Theorem 6.2 all the involutions on $\mathfrak{C}(\mathfrak{H})$ for which there are faithful adjoint-preserving representations.

6.1. LEMMA. *Let T^* be any involution on $\mathfrak{C}(\mathfrak{H})$. Then there exists an invertible s.a. element U in $\mathfrak{C}(\mathfrak{H})$ such that $T^* = U^{-1}T^*U$ for all $T \in \mathfrak{C}(\mathfrak{H})$. Conversely any such mapping is an involution.*

The mapping $T \rightarrow T^{**}$, $T \in \mathfrak{C}(\mathfrak{H})$, is an automorphism of $\mathfrak{C}(\mathfrak{H})$. Thus there exists $V \in \mathfrak{C}(\mathfrak{H})$ where $T^{**} = VTV^{-1}$, $T \in \mathfrak{C}(\mathfrak{H})$. Set $U = V^*$. Then $T^* = U^{-1}T^*U$. Since $T^{**} = T$, $T = (U^{-1}T^*U)^* = U^{-1}U^*T(U^*)^{-1}U$. Thus $U^{-1}U^*$ lies in the center of $\mathfrak{C}(\mathfrak{H})$. Consequently $U = \lambda U^*$ for some scalar λ . Since $U^*U = |\lambda|^2 U^*U$ we see that $|\lambda| = 1$. Set $\lambda = \exp(i\theta)$ and $W = \exp(-i\theta/2)U$. Then $W^* = W$ and $T^* = W^{-1}T^*W$, $T \in \mathfrak{C}(\mathfrak{H})$. The remaining statement is easily verified.

6.2. THEOREM. *An involution $T \rightarrow T^*$ on $\mathfrak{C}(\mathfrak{H})$ is proper if and only if it can be expressed in the form $T^* = U^{-1}T^*U$, $U \in \mathfrak{C}(\mathfrak{H})$ where U is s.a. and $sp(U) \subset (0, \infty)$.*

If $T \rightarrow T^*$ is a proper involution then (see [7]) an inner product can be defined in \mathfrak{H} in terms of which T^* is the adjoint of T . Hence the proper involutions are those for which there is an adjoint preserving faithful representation.

Let W be a one-dimensional operator, $W(x) = (x, z)w$ with $w \neq 0$, $z \neq 0$. Then $W^*(x) = (x, w)z$. By Lemma 6.1 we can write $T^* = U^{-1}T^*U$, $T \in \mathfrak{C}(\mathfrak{H})$, where U is s.a. Then $0 \neq W^*W = U^{-1}W^*UW$. Hence $0 \neq W^*UW$. But $W^*UW(x) = (x, z)W^*U(w) = (x, z)(U(w), w)z$. Therefore $(U(w), w) \neq 0$ for an arbitrary non-zero $w \in \mathfrak{H}$. Hence $(U(w), w) \neq 0$ for an arbitrary non-zero $w \in H$. Hence $(U(w), w)$ has a constant sign and, by changing to $-U$ if necessary, we may suppose that $(U, w), w \geq 0$, $w \in \mathfrak{H}$. Then we can write $U = V^2$ where V is s.a. in $\mathfrak{C}(\mathfrak{H})$.

Suppose conversely that $T^* = V^{-2}T^*V^2$, $T \in \mathfrak{C}(\mathfrak{H})$ where V is s.a. Then $TT^* = (TV^{-1})(TV^{-1})^*V^2$. Thus $TT^* = 0$ implies that $TV^{-1} = 0$ and that $T = 0$.

REFERENCES

1. R. Arens, *Representations of *-algebras*, Duke Math. J. **14** (1947), 269-282.
2. B. H. Arnold, *Rings of operators on vector spaces*, Ann. of Math. **45** (1944), 24-49.
3. F. F. Bonsall and A. W. Goldie, *Annihilator algebras*, Proc. London Math. Soc. **14** (1954), 154-167.
4. P. Civin and B. Yood, *Involutions on Banach algebras*, Pacific J. Math., **9** (1959), 415-436.
5. N. Jacobson, *Structure of rings*, Amer. Math. Soc. Colloq. Publ. vol. **37**, Providence, 1956.
6. S. Kakutani and G. W. Mackey, *Two characterizations of real Hilbert space*, Ann. of Math. **45** (1944), 50-58.

7. S. Kakutani and G. W. Mackey, *Ring and lattice characterization of complex Hilbert space*, Bull. Amer. Math. Soc. **52** (1946), 727-733.
8. I. Kaplansky, *Normed algebras*, Duke Math. J. **16** (1949), 399-418.
9. ———, *Topological algebra*, Department of Mathematics, University of Chicago, 1952, mimeographed notes.
10. J. L. Kelley and R. L. Vaught, *The positive cone in Banach algebras*, Trans. Amer. Math. Soc. **74** (1953), 44-55.
11. V. L. Klee, Jr., *Convex sets in linear spaces*, Duke Math. J. **18** (1951), 443-466.
12. L. H. Loomis, *An introduction to abstract harmonic analysis*, D. Van Nostrand Co., New York, 1953.
13. M. A. Naimark, *Involutive Algebren*, Sowjetische Arbeiten zur Funktionalysis, Berlin (1954), 89-196.
14. M. A. Naimark, *Normed rings*, Moscow, 1956. (Russian).
15. J. D. Newburgh, *The variation of spectra*, Duke. Math. J. **18** (1951), 165-177.
16. C. E. Rickart, *The uniqueness of norm problem in Banach algebras*, Ann. of Math. **51** (1950), 615-628.
17. ———, *Representations of certain Banach algebras on Hilbert space*, Duke Math. J. **18** (1951), 27-39.
18. B. Yood, *Topological properties of homomorphisms between Banach algebras*, Amer. J. Math. **76** (1954), 155-167.

YALE UNIVERSITY AND UNIVERSITY OF OREGON

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

DAVID GILBARG
Stanford University
Stanford, California

F. H. BROWNELL
University of Washington
Seattle 5, Washington

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

L. J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
G. M. CHERRY
D. DERRY

E. HEWITT
A. HORN
L. NACHBIN

M. OHTSUKA
H. L. ROYDEN
M. M. SCHIFFER

E. SPANIER
E. G. STRAUS
F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Printed in Japan by Kokusai Bunken Insatsusha
(International Academic Printing Co., Ltd.), Tokyo, Japan

Pacific Journal of Mathematics

Vol. 10, No. 1

September, 1960

Richard Arens, <i>Extensions of Banach algebras</i>	1
Fred Guenther Brauer, <i>Spectral theory for linear systems of differential equations</i>	17
Herbert Busemann and Ernst Gabor Straus, <i>Area and normality</i>	35
J. H. Case and Richard Eliot Chamberlin, <i>Characterizations of tree-like continua</i>	73
Ralph Boyett Crouch, <i>Characteristic subgroups of monomial groups</i>	85
Richard J. Driscoll, <i>Existence theorems for certain classes of two-point boundary problems by variational methods</i>	91
A. M. Duguid, <i>A class of hyper-FC-groups</i>	117
Adriano Mario Garsia, <i>The calculation of conformal parameters for some imbedded Riemann surfaces</i>	121
Irving Leonard Glicksberg, <i>Homomorphisms of certain algebras of measures</i>	167
Branko Grünbaum, <i>Some applications of expansion constants</i>	193
John Hilzman, <i>Error bounds for an approximate solution to the Volterra integral equation</i>	203
Charles Ray Hobby, <i>The Frattini subgroup of a p-group</i>	209
Milton Lees, <i>von Neumann difference approximation to hyperbolic equations</i>	213
Azriel Lévy, <i>Axiom schemata of strong infinity in axiomatic set theory</i>	223
Benjamin Muckenhoupt, <i>On certain singular integrals</i>	239
Kotaro Oikawa, <i>On the stability of boundary components</i>	263
J. Marshall Osborn, <i>Loops with the weak inverse property</i>	295
Paulo Ribenboim, <i>Un théorème de réalisation de groupes réticulés</i>	305
Daniel Saltz, <i>An inversion theorem for Laplace-Stieltjes transforms</i>	309
Berthold Schweizer and Abe Sklar, <i>Statistical metric spaces</i>	313
Morris Weisfeld, <i>On derivations in division rings</i>	335
Bertram Yood, <i>Faithful $*$-representations of normed algebras</i>	345