

Pacific Journal of Mathematics

TRANSFORMATIONS OF SERIES OF *E*-FUNCTIONS

T. M. MACROBERT

TRANSFORMATIONS OF SERIES OF *E*-FUNCTIONS

T. M. MACROBERT

1. Introductory. The transformation [1, p. 25, 2, p. 369]

$$(1) \quad F\left(\begin{matrix} \alpha, \beta, \gamma, \delta, -l; 1 \\ \alpha-\beta+1, \alpha-\gamma+1, \alpha-\delta+1, \alpha+l+1 \end{matrix}\right) \\ = \frac{(\alpha+1; l)\left(\frac{1}{2}\alpha-\beta+1; l\right)}{\left(\frac{1}{2}\alpha+1; l\right)(\alpha-\beta+1; l)} F\left(\begin{matrix} \alpha-\gamma-\delta+1, \frac{1}{2}\alpha, \beta, -l; 1 \\ \alpha-\gamma+1, \alpha-\delta+1, \beta-\frac{1}{2}\alpha-l \end{matrix}\right),$$

where l is a positive integer, is a special case of a formula of Whipple's. It, and other transformations of the same kind, can be employed to obtain transformations of series of *E*-functions. Two such transformations are:

$$\sum_{n=0}^l \frac{(\alpha; n)(\beta; n)(-l; n)}{n!(\alpha-\beta+l; n)(\alpha+l+1; n)} E\left\{ \frac{p; \alpha_r}{\Delta(m; \rho-n), \Delta(m; \sigma-n), \Delta(m; \alpha+\rho+n)} : z \right\}$$

$$(2) \quad = \frac{(\alpha+1; l)\left(\frac{1}{2}\alpha-\beta+1; l\right)}{\left(\frac{1}{2}\alpha+1; l\right)(\alpha-\beta+1; l)} \sum_{n=0}^l \frac{\left(\frac{1}{2}\alpha; n\right)(\beta; n)(-l; n)}{n!\left(\beta-\frac{1}{2}\alpha-l; n\right)} \left(\frac{2}{m}\right)^n \\ \times E\left\{ \frac{\Delta(2m; \alpha+\rho+\sigma+n-1), \alpha_1, \alpha_2, \dots, \alpha_p}{\Delta(2m; \alpha+\rho+\sigma-1), \Delta(m; \rho), \Delta(m; \sigma), \Delta(m; \alpha+\rho+n), \Delta(m; \alpha+\sigma+n), \rho_1, \rho_2, \dots, \rho_q} : z \right\};$$

$$\sum_{n=0}^l \frac{(\alpha; n)(\beta; n)(-l; n)}{n!(\alpha-\beta+1; n)(\alpha+l+1; n)} E\left\{ \frac{\Delta(m; \gamma+n), \Delta(m; \gamma-\alpha-n), \alpha_1, \dots, \alpha_p}{\Delta(m; \sigma+n), \Delta(m; \sigma-\alpha-n), \rho_1, \dots, \rho_q} : z \right\} \\ (3) \quad = \frac{(\alpha+1; l)\left(\frac{1}{2}\alpha-\beta+1; l\right)}{\left(\frac{1}{2}\alpha+1; l\right)(\alpha-\beta+1; l)} \sum_{n=0}^l \frac{(\sigma-\gamma; n)\left(\frac{1}{2}\alpha; n\right)(\beta; n)(-l; n)}{n!\left(\beta-\frac{1}{2}\alpha-l; n\right)(-m^2)^n} \\ \times E\left\{ \frac{\Delta(m; \gamma), \Delta(m; \gamma-\alpha-n), \alpha_1, \dots, \alpha_p}{\Delta(m; \sigma-\alpha), \Delta(m; \sigma+n), \rho_1, \dots, \rho_q} : z \right\}.$$

In these formulae m is a positive integer,

$$(4) \quad (\alpha; 0) = 1, \quad (\alpha; m) = \alpha(\alpha+1) \cdots (\alpha+m-1),$$

and $\Delta(m; \alpha)$ denotes the set of parameters

$$\frac{\alpha}{m}, \frac{\alpha+1}{m}, \dots, \frac{\alpha+m-1}{m}.$$

The proofs of (2) and (3) are given in § 2. The following formulae are required.

If m is a positive integer

$$(5) \quad \Gamma(mz) = (2\pi)^{[(1/2) - (1/2)m]} m^{[mz - (1/2)]} \Gamma(z) \Gamma\left(z + \frac{1}{m}\right) \cdots \Gamma\left(z + \frac{m-1}{m}\right).$$

From this it follows that, if m and n are positive integers,

$$(6) \quad \begin{aligned} \Gamma\left(\frac{\alpha+n}{m}\right) \Gamma\left(\frac{\alpha+1+n}{m}\right) \cdots \Gamma\left(\frac{\alpha+m-1+n}{m}\right) \\ = \Gamma\left(\frac{\alpha}{m}\right) \Gamma\left(\frac{\alpha+1}{m}\right) \cdots \Gamma\left(\frac{\alpha+m-1}{m}\right) m^{-n}(\alpha; n), \end{aligned}$$

and

$$(7) \quad \begin{aligned} \Gamma\left(\frac{\alpha-n}{m}\right) \Gamma\left(\frac{\alpha+1-n}{m}\right) \cdots \Gamma\left(\frac{\alpha+m-1-n}{m}\right) \\ = \Gamma\left(\frac{\alpha}{m}\right) \Gamma\left(\frac{\alpha+1}{m}\right) \cdots \Gamma\left(\frac{\alpha+m-1}{m}\right) (-m)^n/(1-\alpha; n). \end{aligned}$$

The Barnes' integral for the E -function is [2, p. 374]

$$(8) \quad E(p; \alpha_r; q; \rho_s; z) = \frac{1}{2\pi i} \int \frac{\Gamma(\xi) \prod \Gamma(\alpha_r - \xi)}{\prod \Gamma(\rho_s - \xi)} z^\xi d\xi$$

where $|\operatorname{amp} z| < \pi$ and the integral is taken up the η -axis with loops, if necessary, to ensure that the origin lies to the left of the contour and the points $\alpha_1, \alpha_2, \dots, \alpha_p$ to the right of the contour. Zero and negative integral values of the parameters are excluded, and the α 's must not differ by integral values. When $p < q + 1$ the contour is bent to the left at both ends. When $p > q + 1$ the formula is valid for $|\operatorname{amp} z| < \frac{1}{2}(p - q + 1)\pi$.

2. Proofs of the transformations. Using (8), (6) and (7), the left-hand side of (2), with $p = q = 0$, can be written

$$\begin{aligned} & \sum_{n=0}^l \frac{(\alpha; n)(\beta; n)(-l; n)}{n!(\alpha - \beta + 1; n)(\alpha + l + 1; n)} \\ & \times \frac{1}{2\pi i} \int \frac{\Gamma(\xi) z^\xi (1 - \rho + m\xi; n)(1 - \sigma + m\xi; n) d\xi}{\prod_{\rho, \sigma} \left[\prod_{u=0}^{m-1} \left\{ \Gamma\left(\frac{\rho+u}{m} - \xi\right) \Gamma\left(\frac{\alpha+\rho+u}{m} - \xi\right) \right\} (\alpha + \rho - m\xi; n) \right]} \\ & = \frac{1}{2\pi i} \int \frac{\Gamma(\xi) z^\xi}{\prod_{\rho, \sigma} \left[\prod_{u=0}^{m-1} \left\{ \Gamma\left(\frac{\rho+u}{m} - \xi\right) \Gamma\left(\frac{\alpha+\rho+u}{m} - \xi\right) \right\} \right]} Id\xi, \end{aligned}$$

where

$$I = F \begin{pmatrix} \alpha, & \beta, & 1 - \rho + m\xi, & 1 - \sigma + m\xi, & -l; & 1 \\ \alpha - \beta + 1, & \alpha + \rho - m\xi, & \alpha + \sigma - m\xi, & \alpha + l + 1 \end{pmatrix},$$

From (1) it follows that this can be put in the form

$$\frac{(\alpha + 1; l) \left(\frac{1}{2} \alpha - \beta + 1; l \right)}{\left(\frac{1}{2} \alpha + 1; l \right) (\alpha - \beta + 1; 1)} \times \frac{1}{2\pi i} \int \frac{\Gamma(\xi) z^\xi}{\prod_{\rho, \sigma} \prod_{u=0}^{m-1} \left\{ \Gamma \left(\frac{\rho + u}{m} - \xi \right) \Gamma \left(\frac{\alpha + \rho + u}{m} - \xi \right) \right\}} J d\xi,$$

where

$$J = F \begin{pmatrix} \alpha + \rho + \sigma - 1 - 2m\xi, & \frac{1}{2} \alpha, & \beta, & -l; & 1 \\ \alpha + \rho - m\xi, & \alpha + \sigma - m\xi, & \beta - \frac{1}{2} \alpha - l \end{pmatrix}.$$

Now the integral is equal to

$$\sum_{n=0}^l \frac{\left(\frac{1}{2} \alpha; n \right) (\beta; n) (-l; n)}{n! \left(\beta - \frac{1}{2} \alpha - l; n \right) 2\pi i} \times \int \frac{\Gamma(\xi) z^\xi \Gamma(\alpha + \rho + \sigma - 1 + n - 2m\xi)}{\prod_{\rho, \sigma} \prod_{u=0}^{m-1} \left\{ \Gamma \left(\frac{\rho + u}{m} - \xi \right) \Gamma \left(\frac{\alpha + \rho + u}{m} - \xi \right) \right\}} \left| \frac{\Gamma(\alpha + \rho - m\xi) \Gamma(\alpha + \sigma - m\xi) d\xi}{\Gamma(\alpha + \rho + n - m\xi)} \right\} \Gamma(\alpha + \rho + \sigma - 1 - 2m\xi),$$

and, on applying (5) to the gamma functions whose arguments contain $-2m\xi$ or $-m\xi$, the right-hand side of (2) with $p = q = 0$ is obtained. Formula (2) can then be derived by generalising.

Formula (3) can be proved in the same way. It should be noted that

$$(\alpha - \gamma + 1 + m\xi; n) = (-1)^n (\gamma - \alpha - n - m\xi; n).$$

The restrictions on amp z and on the parameters can be removed by analytical continuation, provided that the functions exist.

REFERENCES

1. W. N. Bailey, *Generalized Hypergeometric Series*, Cambridge 1935.
2. T. M. MacRobert, *Functions of a Complex Variable*, London 1954.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

F. H. BROWNELL

University of Washington
Seattle 5, Washington

L. J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
T. M. CHERRY

D. DERRY
M. OHTSUKA

H. L. ROYDEN
E. SPANIER

E. G. STRAUS
F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE,
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Printed in Japan by International Academic Printing Co., Ltd., Tokyo, Japan

Reprinted 1966 in the United States of America

Pacific Journal of Mathematics

Vol. 11, No. 1

November, 1961

A. A. Albert, <i>Generalized twisted fields</i>	1
Richard Arens, <i>Operational calculus of linear relations</i>	9
John Herbert Barrett, <i>Disconjugacy of a self-adjoint differential equation of the fourth order</i>	25
Paul Richard Beesack, <i>Hardy's inequality and its extensions</i>	39
Julius Rubin Blum and David Lee Hanson, <i>On invariant probability measures. II</i>	63
Robert Allen Bonic, <i>Symmetry in group algebras of discrete groups</i>	73
R. Creighton Buck, <i>Multiplication operators</i>	95
Jack Gary Ceder, <i>Some generalizations of metric spaces</i>	105
Meyer Dwass, <i>Random crossings of cumulative distribution functions</i>	127
Albert Edrei, Wolfgang H. J. Fuchs and Simon Hellerstein, <i>Radial distribution and deficiencies of the values of a meromorphic function</i>	135
William Cassidy Fox, <i>Harmonic functions with arbitrary local singularities</i>	153
Theodore Thomas Frankel, <i>Manifolds with positive curvature</i>	165
Avner Friedman, <i>A strong maximum principle for weakly subparabolic functions</i>	175
Watson Bryan Fulks and J. O. Sather, <i>Asymptotics. II. Laplace's method for multiple integrals</i>	185
Adriano Mario Garsia and Eugene Richard Rodemich, <i>An embedding of Riemann surfaces of genus one</i>	193
Irving Leonard Glicksberg, <i>Weak compactness and separate continuity</i>	205
Branko Grünbaum, <i>On a conjecture of H. Hadwiger</i>	215
Frank J. Hahn, <i>On the action of a locally compact group on E_n</i>	221
Magnus R. Hestenes, <i>Relative hermitian matrices</i>	225
G. K. Kalisch, <i>On similarity invariants of certain operators in L_p</i>	247
Yitzhak Katznelson and Walter Rudin, <i>The Stone-Weierstrass property in Banach algebras</i>	253
Samir A. Khabbaz, <i>The subgroups of a divisible group G which can be represented as intersections of divisible subgroups of G</i>	267
Marvin Isadore Knopp, <i>Construction of a class of modular functions and forms</i>	275
Charles Alan McCarthy, <i>Commuting Boolean algebras of projections</i>	295
T. M. MacRobert, <i>Transformations of series of E-functions</i>	309
Heinz Renggli, <i>An inequality for logarithmic capacities</i>	313
M. S. Robertson, <i>Applications of the subordination principle to univalent functions</i>	315
David Sachs, <i>Partition and modulated lattices</i>	325
Frank S. Scalora, <i>Abstract martingale convergence theorems</i>	347
Elbert A. Walker, <i>Torsion endomorphic images of mixed Abelian groups</i>	375
Morgan Ward, <i>The prime divisors of Fibonacci numbers</i>	379
Charles R. B. Wright, <i>On the nilpotency class of a group of exponent four</i>	387