

Pacific Journal of Mathematics

CONVERGENT SEQUENCES OF FINITELY ADDITIVE MEASURES

TSUYOSHI ANDÔ

CONVERGENT SEQUENCES OF FINITELY ADDITIVE MEASURES

TSUYOSHI ANDÔ

1. Introduction and Main theorem. Though sequences of countably additive measures have been investigated by many authors, comparatively little attention has been paid to finitely additive measures in general. The main purpose of this paper is to give a generalization of a classical convergence theorem to the case of finitely additive measures and its improvement.

Let \mathcal{B} be a σ -complete (infinite) Boolean algebra with the unit I . \mathfrak{M} is the class of finitely additive measures on \mathcal{B} with bounded variations, that is, real valued functions μ on \mathcal{B} with the following properties:

$$\sup_{E \in \mathcal{B}} |\mu(E)| < \infty, \quad \mu(O) = 0$$

and

$$\mu(E \cup F) + \mu(E \cap F) = \mu(E) + \mu(F)$$

for every $E, F \in \mathcal{B}$. We shall call elements of \mathfrak{M} simply *measures*. Under the ordinary addition and scalar multiplication, \mathfrak{M} is a linear space. Moreover it is a universally continuous semi-ordered linear space [6] (=a conditionally complete vector lattice [2]) under the order relation: $\nu \geq \mu$ means $\nu(E) \geq \mu(E)$ for every $E \in \mathcal{B}$. The symbols \vee and \wedge will denote supremum and infimum in \mathfrak{M} respectively. We shall write $\mu^+ = \mu \vee 0$, $\mu^- = (-\mu) \vee 0$ and $|\mu| = \mu \vee (-\mu)$, then $\mu = \mu^+ - \mu^-$ and $|\mu| = \mu^+ + \mu^-$. For each subset \mathcal{C} of \mathfrak{M} its orthogonal complement, i.e., $\{\mu \mid |\mu| \wedge |\nu| = 0 \text{ for every } \nu \in \mathcal{C}\}$ will be denoted by \mathcal{C}^\perp . We quote some results from the theory of vector lattices (see [6] chap. I). \mathcal{C} is called *normal*, if $\mathcal{C} = (\mathcal{C}^\perp)^\perp$. Any orthogonal complement is normal. Every normal subset is a direct summand, that is, $\mathfrak{M} = \mathcal{C} \oplus \mathcal{C}^\perp$ (in linear order sense). Thus each normal subset \mathcal{C} determines a linear lattice homomorphism of \mathfrak{M} onto \mathcal{C} which makes \mathcal{C} invariant. Following [6] §5 this homomorphism will be denoted by $[\mathcal{C}]$, that is,

$$\begin{aligned} [\mathcal{C}](\alpha\mu + \beta\nu) &= \alpha[\mathcal{C}]\mu + \beta[\mathcal{C}]\nu \quad (\alpha, \beta \text{ real}), \\ [\mathcal{C}](\mu \vee \nu) &= [\mathcal{C}]\mu \vee [\mathcal{C}]\nu, \end{aligned}$$

and $[\mathcal{C}]\mu = \mu$ is equivalent to $\mu \in \mathcal{C}$. When $\mathcal{C} = (\{\nu\}^\perp)^\perp$ where $\{\nu\}$ consists of a single measure ν , the linear operator $[\mathcal{C}]$ will be denoted simply by $[\nu]$. μ is said to be *absolutely continuous with respect to* ν ,

Received June 2, 1960.

if $|\nu|(E) \rightarrow 0$ implies $\mu(E) \rightarrow 0$. It is known that this is equivalent to the relation $[\nu]\mu = \mu$. A measure μ is called *countably additive*, if $|\mu|(\bigcup_{k=1}^{\infty} E_k) \leq \sum_{k=1}^{\infty} |\mu|(E_k)$ for every sequence $\{E_k\}$. The set of countably additive measures is normal. It will be denoted by \mathfrak{L} . Following [8] we shall call measures in \mathfrak{L}^\perp *purely finitely additive*.

A classical convergence theorem of countably additive measures can be formulated as follows (see [7]):

Let $\{\mu_k\}$ be a sequence of measures such that $\lim_{k \rightarrow \infty} \mu_k(E)$ exists and is finite for every $E \in \mathcal{B}$. If every μ_k is absolutely continuous with respect to a fixed countably additive measure ν , then the function $\mu(E) \stackrel{\text{def}}{=} \lim_{k \rightarrow \infty} \mu_k(E)$ is also a measure absolutely continuous with respect to ν , and the sequence has the following uniform absolute continuity:

$$|\nu|(E) \rightarrow 0 \text{ implies } \sup_k |\mu_k|(E) \rightarrow 0 .$$

We shall prove the theorem without assumption of countable additivity. Since, as stated before, absolute continuity can be expressed in terms of $[\mathfrak{C}]$ the following theorem will give a more complete answer.

MAIN THEOREM. *If $\lim_{k \rightarrow \infty} \mu_k(E)$ exists and is finite for every $E \in \mathcal{B}$, then the function $\mu(E) \stackrel{\text{def}}{=} \lim_{k \rightarrow \infty} \mu_k(E)$ is a measure, and for each normal \mathfrak{C}*

$$\lim_{k \rightarrow \infty} [\mathfrak{C}]\mu_k(E) = [\mathfrak{C}]\mu(E) \quad \text{for every } E \in \mathcal{B} .$$

Moreover the sequence $\{[\mathfrak{C}]\mu_k\}$ has the following uniform absolute continuity:

$$|\nu|(E) \rightarrow 0 \text{ (for every } \nu \in \mathfrak{C}) \text{ implies } \sup_k |[\mathfrak{C}]\mu_k|(E) \rightarrow 0 .$$

2. Proof of Main theorem. In connection with uniform absolute continuity we begin with some lemmas.

LEMMA 1. *Let $\{\mu_k\}$ be a sequence of measures with the property:*

$$(*) \quad \lim_{k, j \rightarrow \infty} \rho(E_k - E_j) = 0$$

for every monotone sequence $\{E_k\}$ where

$$\rho(E) = \sup_k |\mu_k|(E) .$$

Then for each sequence $\{F_k\}$ and $\varepsilon > 0$ there exist two sequences $\{G_k\}$ and $\{H_k\}$ such that

$$(1) \quad G_k = F_k \cup F_{k+1} \cup \dots \cup F_{j(k)} \quad \text{for some } j(k) ,$$

$$\begin{aligned}
 (2) \quad & G_k \supset H_k \supset H_{k+1} , \\
 (3) \quad & \rho(G_k - H_k) \leq \varepsilon \qquad k = 1, 2, \dots
 \end{aligned}$$

Proof. Since the sequence $F_{k,j} \stackrel{\text{def}}{=} \bigcup_{i=k}^j F_i$ ($j \geq k$) is monotone (for each fixed k), by property (*) there exists a sequence $\{j(k)\}$ of positive integers such that $j(k) \leq j(k+1)$ and

$$\rho(F_{k,i} - F_{k,j(k)}) \leq \varepsilon/2^k \quad \text{for } i \geq j(k) .$$

We define the desired sequences by

$$G_k = F_{k,j(k)} \quad \text{and} \quad H_k = \bigcap_{i=1}^k G_i \qquad k = 1, 2, \dots .$$

Then (1) and (2) are trivially satisfied. As to (3)

$$\begin{aligned}
 \rho(G_k - H_k) &\leq \rho\left(\bigcup_{i=1}^{k-1} (G_{i+1} - G_i)\right) \\
 &\leq \sum_{i=1}^{\infty} \rho(G_{i+1} - G_i) = \sum_{i=1}^{\infty} \rho(F_{i+1,j(i+1)} - F_{i,j(i)}) \\
 &\leq \sum_{i=1}^{\infty} \rho(F_{i,j(i+1)} - F_{i,j(i)}) \leq \sum_{i=1}^{\infty} \varepsilon/2^i = \varepsilon .
 \end{aligned}$$

This completes the proof.

REMARK. Likewise we can choose $\{G_k\}$ and $\{H_k\}$ as follows:

$$\begin{aligned}
 (1') \quad & G_k = F_k \cap F_{k+1} \cap \dots \cap F_{j(k)} \quad \text{for some } j(k) , \\
 (2') \quad & G_k \subset H_k \subset H_{k+1} , \\
 (3') \quad & \rho(H_k - G_k) \leq \varepsilon \qquad k = 1, 2, \dots .
 \end{aligned}$$

LEMMA 2. For any non negative measures ν and μ

$$(4) \quad [\nu]\mu(E) = \inf \left\{ \sup_k \mu(E_k) \right\}$$

where the infimum is taken over all the sequences $\{E_k\}$ such that $\bigcup_k E_k \subset E$ and $\lim_{k \rightarrow \infty} \nu(E - E_k) = 0$.

Proof. First remark that $\lim_{k \rightarrow \infty} [\nu]\mu(E_k) = [\nu]\mu(E)$ for every such sequence, because $[\nu]\mu$ is absolutely continuous with respect to ν . Since

$$[\nu]\mu(E) = \inf \left\{ \sup_k [\nu]\mu(E_k) \right\} \leq \inf \left\{ \sup_k \mu(E_k) \right\} \leq \mu(E) ,$$

(because $0 \leq [\nu]\mu \leq \mu$), the function $\mu_1(E)$ defined by the right side of (4), is a measure with the property: $[\nu]\mu \leq \mu_1 \leq \mu$. If it is proved that μ_1 itself is absolutely continuous with respect to ν (i.e. $[\nu]\mu_1 = \mu_1$), by

the order-preserving property of $[\nu]$ we have

$$[\nu]\mu = [\nu]([\nu]\mu) \leq [\nu]\mu_1 = \mu_1 \leq [\nu]\mu ,$$

that is, $[\nu]\mu = \mu_1$. Now suppose that μ_1 is not absolutely continuous with respect to ν , then there exist a sequence $\{F_k\}$ and $\varepsilon > 0$ such that

$$(5) \quad \nu(F_k) \leq 1/2^k \quad \text{and} \quad \mu_1(F_k) > 3\varepsilon \quad k = 1, 2, \dots .$$

Since condition (*) is evidently satisfied for single μ_1 , Lemma 1 guarantees the existence of $\{G_k\}$ and $\{H_k\}$ with the properties (1), (2) and (3) (with μ_1 instead of ρ). From (1), (2), (3) and (5) it follows

$$(6) \quad \begin{aligned} \mu_1(H_k) = \mu_1(G_k) - \mu_1(G_k - H_k) &\geq \mu_1(F_k) - \mu_1(G_k - H_k) \geq 2\varepsilon \quad \text{i.e.} \\ \inf_k \mu_1(H_k) &\geq 2\varepsilon . \end{aligned}$$

By the definition of μ_1 there can be chosen a double sequence $\{E_{kj}\}$ such that

$$(7) \quad \begin{aligned} \bigcup_j E_{kj} \subset H_k - H_{k+1} , \quad \lim_{j \rightarrow \infty} \nu(H_k - H_{k+1} - E_{kj}) &= 0 , \\ \sup_j \mu(E_{kj}) \leq \mu_1(H_k - H_{k+1}) + \varepsilon/2^k & \quad k = 1, 2, \dots . \end{aligned}$$

Writing $D_j = \bigcup_{k=1}^j E_{kj}$, it follows

$$\nu(H_1 - D_j) \leq \sum_{k=1}^{j-1} \nu(H_k - H_{k+1} - E_{kj}) + \nu(H_j) \xrightarrow{j \rightarrow \infty} \nu(H_1) \quad i = 1, 2, \dots$$

consequently we have

$$(8) \quad \lim_{j \rightarrow \infty} \nu(H_1 - D_j) = 0 ,$$

because by (5) and (1)

$$\nu(H_i) \leq \nu(G_i) \leq \sum_{j=i}^{\infty} \nu(F_j) \leq \sum_{j=i}^{\infty} 1/2_j = 1/2^{i-1} .$$

On the other hand, on account of (2), (6) and (7)

$$\begin{aligned} \mu(D_j) &= \sum_{k=1}^j \mu(E_{kj}) \leq \sum_{k=1}^j \mu_1(H_k - H_{k+1}) + \sum_{k=1}^{\infty} \varepsilon/2^k \\ &= \mu_1(H_1) - \mu_1(H_j) + \varepsilon \leq \mu_1(H_1) - \varepsilon , \end{aligned}$$

that is,

$$\sup_j \mu(D_j) \leq \mu_1(H_1) - \varepsilon .$$

Taking (8) into consideration, by the definition of μ_1 .

$$\mu_1(H_1) \leq \sup_j \mu(D_j) \leq \mu_1(H_1) - \varepsilon .$$

This contradiction establishes the assertion.

We shall reduce a proof of Main theorem to the case of a concrete Boolean algebra. The simplest σ -complete (infinite) Boolean algebra is the class \mathcal{N} of all subsets of natural numbers. Phillips proved a special case of our Main theorem when $\mathcal{B} = \mathcal{N}$. The following Lemma due to him is essential.

LEMMA 3 (Phillips). *Let $\{\varphi_k\}$ be a sequence of measures on \mathcal{N} . If $\lim_{k \rightarrow \infty} \varphi_k(A)$ exists and is finite for every $A \in \mathcal{N}$, then*

$$\lim_{k \rightarrow \infty} \varphi_k(\{k\}) = 0$$

where $\{k\}$ is the set consisting of single k .

This is a slight modification of [1] p. 32 Lemma.

Up to this point the σ -completeness of \mathcal{B} has not been used, however in the following Lemma it plays a decisive rôle.

LEMMA 4. *If $\lim_{k \rightarrow \infty} \mu_k(E)$ exists and is finite for every $E \in \mathcal{B}$, then $\sup_k |\mu_k|(I) < \infty$ and (*) is satisfied.*

Proof. As proofs of two assertions are similar, we confine ourselves to the proof of (*). Supposing that (*) is not satisfied, we can choose a sequence $\{E_k\}$ and $\varepsilon > 0$ such that $E_1 \subset E_2 \subset \dots$ and

$$|\mu_k|(E_{k+1} - E_k) > 2\varepsilon \quad k = 1, 2, \dots$$

(taking a subsequence of $\{\mu_k\}$ if necessary). Since in general (see [8])

$$|\mu|(E) = \sup_{F \subset E} |\mu(F) - \mu(E - F)|,$$

there exists a sequence $\{F_k\}$ such that

$$F_k \subset E_{k+1} - E_k \quad \text{and} \quad |\mu_k(F_k)| > \varepsilon \quad k = 1, 2, \dots$$

Writing $\varphi_k(A) = \mu_k(\bigcup_{i \in A} F_i)$ for $A \in \mathcal{N}$ (here the σ -completeness of \mathcal{B} is necessary), we obtain a sequence of measures on \mathcal{N} with the property: $\lim_{k \rightarrow \infty} \varphi_k(A)$ exists and is finite for every $A \in \mathcal{N}$. Then Lemma 3 shows that $\mu_k(F_k) = \varphi_k(\{k\}) \xrightarrow{k \rightarrow \infty} 0$. This contradiction establishes the assertion.

With these preparations we are now in position to prove Main theorem.

Proof of Main theorem. Since $\sup_k |\mu_k|(I) < \infty$ by Lemma 4, the function $\mu(E) \stackrel{\text{def}}{=} \lim_{k \rightarrow \infty} \mu_k(E)$ is a measure. Considering the sequence $\{\mu_k - \mu\}$ instead, we may assume $\mu = 0$. Define

$$\nu(E) = \sum_{k=1}^{\infty} \frac{[\mathcal{C}]\mu_k(E)}{2^k [\mu_k(I)} \quad E \in \mathcal{B} .$$

It is not difficult to see that ν is a measure and $[\mathcal{C}]\mu_k = [\nu]\mu_k$ $k = 1, 2, \dots$ (see [6] § 5), hence we may also assume $[\mathcal{C}] = [\nu]$, and we shall prove

$$\lim_{k \rightarrow \infty} [\nu]\mu_k(E) = 0 \quad \text{for every } E \in \mathcal{B} .$$

By Lemma 2 for each E there exist sequences $\{C_{kj}\}$ and $\{D_{kj}\}$ such that

$$\begin{aligned} \bigcup_j C_{kj} &\subset E, & \bigcup_j D_{kj} &\subset E, \\ \lim_{j \rightarrow \infty} \nu(E - C_{kj}) &= \lim_{j \rightarrow \infty} \nu(E - D_{kj}) = 0, \\ \lim_{j \rightarrow \infty} \mu_k^+(C_{kj}) &= [\nu]\mu_k^+(E), \end{aligned}$$

and

$$\lim_{j \rightarrow \infty} \mu_k^-(D_{kj}) = [\nu]\mu_k^-(E) \quad k = 1, 2, \dots .$$

Since

$$\nu(E - C_{kj} \cap D_{kj}) \leq \nu(E - C_{kj}) + \nu(E - D_{kj}) \xrightarrow{j \rightarrow \infty} 0$$

by Lemma 2 we obtain

$$[\nu]\mu_k^+(E) \leq \overline{\lim}_{j \rightarrow \infty} \mu_k^+(E_{kj}) \leq \lim_{j \rightarrow \infty} \mu_k^+(C_{kj}) = [\nu]\mu_k^+(E),$$

where $E_{kj} = C_{kj} \cap D_{kj}$, similarly

$$\lim_{j \rightarrow \infty} \mu_k^-(E_{kj}) = [\nu]\mu_k^-(E) \quad k = 1, 2, \dots .$$

Writing $F_{kj} = \bigcap_{i=1}^k E_{ij}$, the similar arguments show

$$\lim_{j \rightarrow \infty} \nu(E - F_{kj}) = 0$$

and

$$\lim_{j \rightarrow \infty} \mu_i^+(F_{kj}) = [\nu]\mu_i^+(E) \quad i = 1, 2, \dots, k ,$$

and similarly for $\{\mu_k^-\}$. By a diagonal method we can find a subsequence $\{F_i\}$ of $\{F_{kj}\}$ such that

$$\nu(E - F_i) < 1/2^i \quad i = 1, 2, \dots$$

and

$$\lim_{i \rightarrow \infty} \mu_k^+(F_i) = [\nu]\mu_k^+(E) \quad k = 1, 2, \dots$$

and similarly for $\{\mu_k^-\}$. Since condition (*) is satisfied by Lemma 4, there exist sequences $\{G_k\}$ and $\{H_k\}$ with the properties (1'), (2') and (3').

On account of (1'), by the similar way as above, it is not difficult to see that

$$\lim_{j \rightarrow \infty} \mu_k^+(G_j) = [\nu]\mu_k^+(E) \quad k = 1, 2, \dots$$

and similarly for $\{\mu_k^-\}$, hence by subtraction

$$(9) \quad \lim_{j \rightarrow \infty} \mu_k(G_j) = [\nu]\mu_k(E) \quad k = 1, 2, \dots$$

On the other hand, for each $\varepsilon > 0$ (*) and (2') guarantee the existence of $n = n(\varepsilon)$ such that

$$\rho(H_j - H_n) \leq \varepsilon \quad \text{for } j \geq n,$$

consequently by (3')

$$(10) \quad |\mu_k(H_n) - \mu_k(G_j)| \leq \rho(H_j - H_n) + \rho(H_j - G_j) \leq 2\varepsilon \quad \text{for } j \geq n.$$

Then from (9) and (10) it follows

$$|\mu_k(H_n) - [\nu]\mu_k(E)| = \lim_{j \rightarrow \infty} |\mu_k(H_n) - \mu_k(G_j)| \leq 2\varepsilon \quad k = 1, 2, \dots$$

Since $\lim_{k \rightarrow \infty} \mu_k(H_n) = 0$ by hypothesis, combining this with the above, we obtain $\bar{\lim}_{k \rightarrow \infty} [\nu]\mu_k(E) \leq 2\varepsilon$. The arbitrariness of ε implies $\lim_{k \rightarrow \infty} [\nu]\mu_k(E) = 0$.

Next we shall turn to a proof of the second assertion. Here we may again assume $[\mathfrak{S}] = [\nu]$. First remark that the sequence $\{[\nu]\mu_k\}$ satisfies (*). Given a sequence $\{F_k\}$ with the property: $\nu(F_k) < 1/2^k$ $k = 1, 2, \dots$, applying Lemma 1 to this sequence and $\{[\nu]\mu_k\}$, for any $\varepsilon > 0$ we can find sequences $\{G_k\}$ and $\{H_k\}$ with properties (1), (2) and (3) (with $\rho(E) = \sup_k |[\nu]\mu_k|(E)$). Since (*) and the absolute continuity of every $[\nu]\mu_k$ (with respect to ν) imply $\rho(H_k) \xrightarrow{k \rightarrow \infty} 0$, we obtain $\bar{\lim}_{k \rightarrow \infty} \rho(F_k) \leq 2\varepsilon$, because

$$\rho(F_k) \leq \rho(G_k) \leq \rho(H_k) + \rho(G_k - H_k) \leq \rho(H_k) + 2\varepsilon \quad k = 1, 2, \dots$$

The arbitrariness of ε establishes the assertion.

REMARK. When \mathfrak{B} is moreover *complete*, our Main theorem can be deduced from a result of Grothendieck [3] by the following way. By means of the theory of Boolean algebra, \mathfrak{B} can be represented by the class of open-closed subsets of a compact Stonian space Ω . Then by the natural way \mathfrak{M} may be considered as the dual of the Banach space $C(\Omega)$ (the space of continuous functions on Ω with the supremum norm). Grothendieck proved that if μ_k is $\sigma(\mathfrak{M}, C(\Omega))$ -convergent, then it is also $\sigma(\mathfrak{M}, \mathfrak{M}')$ -convergent, where \mathfrak{M}' is the dual of \mathfrak{M} and $\sigma(\dots)$ denotes the weak topology. Since every operator $[\mathfrak{S}]$ is $\sigma(\mathfrak{M}, \mathfrak{M}')$ -continuous,

the first part of our Main theorem follows immediately. The second part is also stated there.

3. Corollaries. An immediate corollary (which is a direct generalization of the classical theorem in question) is as follows.

COROLLARY 1. *Every normal set of \mathfrak{M} is sequentially complete under the weak topology defined by \mathcal{B} .*

Let $\{E_\lambda\}$ be the set of all atoms in \mathcal{B} and

$$\mathfrak{F} = \{\mu \in \mathfrak{L} \mid |\mu|(E_\lambda) = 0 \text{ for every } E_\lambda\}.$$

Then \mathfrak{F} is normal. A modification of a recent result of Kaplan [5] § 9 shows that the closure of \mathfrak{L}^\perp with respect to the topology in question coincides with $\mathfrak{F} \oplus \mathfrak{L}^\perp$. In this regards, the following special case of Corollary 1 is of some interest.

COROLLARY 2. *The set of purely finitely additive measures is sequentially complete under the weak topology defined by \mathcal{B} .*

4. Sequences of purely finitely additive measures. In connection with Main theorem a question arises whether $\mu_k \xrightarrow{k \rightarrow \infty} 0$ implies $|\mu_k| \xrightarrow{k \rightarrow \infty} 0$. When $\sup_{\mu \in \mathfrak{L}} \mathfrak{B}\mu(E) > 0$, for every $E \in \mathcal{B}$, Halperin and Nakano [4] proved that the property “ $\mu_k \in \mathfrak{L}, \mu_k \xrightarrow{k \rightarrow \infty} 0$ implies $|\mu_k| \xrightarrow{k \rightarrow \infty} 0$ ” is equivalent to the atomicity of \mathcal{B} . We shall treat this problem in \mathfrak{L}^\perp , namely, does $\mu_k \xrightarrow{k \rightarrow \infty} 0$ (all μ_k being purely finitely additive) imply $|\mu_k| \xrightarrow{k \rightarrow \infty} 0$? The answer is negative.

THEOREM. *There exists a sequence $\{\mu_k\}$ of purely finitely additive measures such that*

$$\lim_{k \rightarrow \infty} \mu_k(E) = 0 \qquad \text{for every } E \in \mathcal{B}$$

but

$$|\mu_k|(I) = 1 \qquad k = 1, 2, \dots$$

Proof. As in § 2 we shall reduce the proof to the case $\mathcal{B} = \mathcal{N}$. Banach (see [1] p. 83) proved that there exists a positive measure φ_0 on \mathcal{N} invariant under translation, that is, $\varphi_0(I_0) = 1$ and

$$0 \leq \varphi_0(A) = \varphi_0(\tau A) \qquad \text{for every } A \in \mathcal{N}$$

where I_0 is the unit of \mathcal{N} and $\tau A = \{j + 1 \mid j \in A\}$. We define a sequence $\{\varphi_k\}$ recurrently by the formula

$$\varphi_{k+1}(A) = \varphi_k\left(A \cap \bigcup_{j=1}^{2^k} B_{k+1,j}\right) - \varphi_k\left(A \cap \bigcup_{j=2^{k+1}}^{2^{k+1}} B_{k+1,j}\right)$$

where

$$B_{k,j} = \{i \mid i \equiv j \pmod{2^k}\}.$$

From the arguments in [4] it results

$$(11) \quad \lim_{k \rightarrow \infty} \varphi_k(A) = 0 \quad \text{for every } A \in \mathcal{N}$$

but

$$|\varphi_k| = \varphi_0 \quad k = 1, 2, \dots$$

Let $\{F_k\}$ be a sequence in \mathcal{B} with the property

$$(12) \quad \bigcup_k F_k = I \quad \text{and} \quad F_k \cap F_j = \emptyset \quad (k \neq j).$$

On account of the representation theorem of Boolean algebra (see [6] § 8; [2] Chap. X) there exists a sequence $\{\nu_k\}$ of two-valued (say 1 and 0) measures on \mathcal{B} such that

$$\nu_k(F_j) = \delta_{kj} \quad k, j = 1, 2, \dots$$

We construct the desired sequence $\{\mu_k\}$ from $\{\varphi_k\}$ and $\{\nu_k\}$ by the formula:

$$\mu_k(E) = \varphi_k(A)$$

where $A = \{j \mid \nu_j(E) = 1\}$. From (11) and (12) it results

$$\lim_{k \rightarrow \infty} \mu_k(E) = 0 \quad \text{for every } E \in \mathcal{B}.$$

but $|\mu_k| = \mu_0$ $k = 1, 2, \dots$. There remains to prove pure finite additivity of μ_k . For this purpose it is enough to prove it for μ_0 only. Invariance of φ_0 under translation shows

$$\mu_0(F_j) = \varphi_0(\{j\}) = \varphi_0(\{i\}) = \mu_0(F_i) \quad i, j = 1, 2, \dots$$

hence

$$k\mu_0(F_j) = \sum_{i=1}^k \mu_0(F_i) \leq \mu_0(I) = 1 \quad j, k = 1, 2, \dots$$

finally

$$\mu_0(F_j) = 0 \quad j = 1, 2, \dots$$

Since $\mu_0(I_j) = 1$, this implies pure finite additivity (cf. [8] § 4).

REFERENCES

1. M. M. Day, *Normed linear spaces*, Berlin, 1958.
2. G. Birkoff, *Lattice theory*, New York, 1948.
3. A. Grothendieck, *Sur les applications linéaires faiblement compactes d'espace du type $C(K)$* , *Canad. J. Math.*, **5** (1953), 129-173.

4. I. Halperin, and H. Nakano, *Discrete semi-ordered linear spaces*, Canad. J. Math., **3** (1951), 293-298.
5. S. Kaplan, *The second dual of the space of continuous functions*, II, Trans. Amer. Math. Soc., **93** (1959), 329-350.
6. H. Nakano, *Modulared semi-ordered linear spaces*, Tokyo, 1950.
7. O. Nikodym, *Sur les suites convergentes de fonctions parfaitement additives d'ensemble abstrait*, Monat. für Math. Phys., **40** (1933), 427-432.
8. K. Yosida and E. Hewitt, *Finitely additive measures*, Trans. Amer. Math. Soc., **72** (1952), 46-66.

RESEARCH INSTITUTE OF APPLIED ELECTRICITY,
HOKKAIDÔ UNIVERSITY, SAPPORO, JAPAN

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS
Stanford University
Stanford, California

A. L. WHITEMAN
University of Southern California
Los Angeles 7, California

F. H. BROWNELL
University of Washington
Seattle 5, Washington

L. J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
T. M. CHERRY

D. DERRY
M. OHTSUKA

H. L. ROYDEN
E. SPANIER

E. G. STRAUS
F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE COLLEGE
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE COLLEGE
UNIVERSITY OF WASHINGTON
* * *

AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
HUGHES AIRCRAFT COMPANY
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Tsuyoshi Andô, <i>Convergent sequences of finitely additive measures</i>	395
Richard Arens, <i>The analytic-functional calculus in commutative topological algebras</i>	405
Michel L. Balinski, <i>On the graph structure of convex polyhedra in n-space</i>	431
R. H. Bing, <i>Tame Cantor sets in E^3</i>	435
Cecil Edmund Burgess, <i>Collections and sequences of continua in the plane. II</i>	447
J. H. Case, <i>Another 1-dimensional homogeneous continuum which contains an arc</i>	455
Lester Eli Dubins, <i>On plane curves with curvature</i>	471
A. M. Duguid, <i>Feasible flows and possible connections</i>	483
Lincoln Kearney Durst, <i>Exceptional real Lucas sequences</i>	489
Gertrude I. Heller, <i>On certain non-linear operators and partial differential equations</i>	495
Calvin Virgil Holmes, <i>Automorphisms of monomial groups</i>	531
Wu-Chung Hsiang and Wu-Yi Hsiang, <i>Those abelian groups characterized by their completely decomposable subgroups of finite rank</i>	547
Bert Hubbard, <i>Bounds for eigenvalues of the free and fixed membrane by finite difference methods</i>	559
D. H. Hyers, <i>Transformations with bounded mth differences</i>	591
Richard Eugene Isaac, <i>Some generalizations of Doeblin's decomposition</i>	603
John Rolfe Isbell, <i>Uniform neighborhood retracts</i>	609
Jack Carl Kiefer, <i>On large deviations of the empiric $D. F.$ of vector chance variables and a law of the iterated logarithm</i>	649
Marvin Isadore Knopp, <i>Construction of a class of modular functions and forms. II</i>	661
Gunter Lumer and R. S. Phillips, <i>Dissipative operators in a Banach space</i>	679
Nathaniel F. G. Martin, <i>Lebesgue density as a set function</i>	699
Shu-Teh Chen Moy, <i>Generalizations of Shannon-McMillan theorem</i>	705
Lucien W. Neustadt, <i>The moment problem and weak convergence in L^2</i>	715
Kenneth Allen Ross, <i>The structure of certain measure algebras</i>	723
James F. Smith and P. P. Saworotnow, <i>On some classes of scalar-product algebras</i>	739
Dale E. Varberg, <i>On equivalence of Gaussian measures</i>	751
Avrum Israel Weinzweig, <i>The fundamental group of a union of spaces</i>	763