

# Pacific Journal of Mathematics

**APPLICATIONS OF THE TOPOLOGICAL METHOD OF  
WAŻEWSKI TO CERTAIN PROBLEMS OF ASYMPTOTIC  
BEHAVIOR IN ORDINARY DIFFERENTIAL EQUATIONS**

NELSON ONUCHIC

APPLICATIONS OF THE TOPOLOGICAL METHOD OF  
WAZEWSKI TO CERTAIN PROBLEMS OF  
ASYMPTOTIC BEHAVIOR IN ORDINARY  
DIFFERENTIAL EQUATIONS

NELSON ONUCHIC

**Introduction.** The main objective of this paper is to present some results concerning the asymptotic behavior of the integrals of some systems of ordinary differential equations.

As Wazewski's theorem, used in our work, is not very well known, we state it here, giving first some definitions and notations.

**HYPOTHESIS H.** (a) *The real-valued functions  $f_i(t, x_1, \dots, x_n)$ ,  $i = 1, \dots, n$ , of the real variables  $t, x_1, \dots, x_n$ , are continuous in an open set  $\Omega \subset R^{n+1}$ .*

(b) *Through every point of  $\Omega$  passes only one integral of the system*

$$\dot{x} = f(t, x) \quad \left( \cdot = \frac{d}{dt} \right) \quad \text{where}$$

$$x = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}, \quad f(t, x) = \begin{pmatrix} f_1(t, x_1, \dots, x_n) \\ \vdots \\ f_n(t, x_1, \dots, x_n) \end{pmatrix} \quad \text{and } (t, x) \in \Omega .$$

Let  $\omega$  be an open set of  $R^{n+1}$ ,  $\omega \subset \Omega$  and let us denote by  $B(\omega, \Omega)$  the boundary of  $\omega$  in  $\Omega$ .

Let  $P_0: (t_0, x_0) \in \Omega$ . We write  $I(t, P_0) = (t, x(t, P_0))$ , where  $x(t, P_0)$  is the integral of the system  $\dot{x} = f(t, x)$  passing through the point  $P_0$ .

Let  $(\alpha(P_0), \beta(P_0))$  be the maximal open interval in which the integral passing through  $P_0$  exists. We write

$$I(\mathcal{A}, P_0) = \{(t, x(t, P_0)) \mid t \in \mathcal{A}\}$$

for every set  $\mathcal{A}$  contained in  $(\alpha(P_0), \beta(P_0))$ .

We say that the point  $P_0: (t_0, x_0) \in B(\omega, \Omega)$  is a *point of egress* from  $\omega$  (with respect to the system  $\dot{x} = f(t, x)$  and the set  $\Omega$ ) if there exists a positive number  $\delta$  such that  $I([t_0 - \delta, t_0), P_0) \subset \omega$ ;  $P_0$  is a *point of strict egress* from  $\omega$  if  $P_0$  is a point of egress and if there exists a positive number  $\delta$  such that  $I((t_0, t_0 + \delta], P_0) \subset \Omega - \bar{\omega}$ . The set of all points of egress (strict egress) is denoted by  $S(S^*)$ .

If  $A \subset B$  are any two sets of a topological space and  $K: B \rightarrow A$  is

a continuous mapping from  $B$  onto  $A$  such that  $K(P) = P$  for every  $P \in A$ , then  $K$  is said to be a *retraction* from  $B$  into  $A$  and  $A$  a *retract* of  $B$ .

**THEOREM OF WAŻEWSKI.** *Suppose that the system  $\dot{x} = f(t, x)$  and the open sets  $\omega \subset \Omega \subset R^{n+1}$  satisfy the following hypotheses:*

- (1) *Hypothesis H.*
- (2)  $S = S^*$ .
- (3) *There exists a set  $Z \subset \omega \cup S$  such that  $Z \cap S$  is a retract of  $S$  but is not a retract of  $Z$ .*

*Then there is at least one point  $P_0: (t_0, x_0) \in Z - S$  such that  $I(t, P_0) \subset \omega$  for every  $t_0 \leq t < \beta(P_0)$ .*

The theorem of Ważewski [6, Théorème 1, p. 299] is actually more general than the one stated above.

If  $f_i(t, x_1, \dots, x_n)$ ,  $i = 1, \dots, n$ , are complex-valued functions of the real variable  $t$  and of the complex variables  $x_1, \dots, x_n$ , the  $n$ -dimensional complex system  $\dot{x} = f(t, x)$  can be considered as a  $2n$ -dimensional real system, so that the theorem of Ważewski is also extensible, in a natural way, to complex systems [5, p. 19. § 1 and p. 21, § 2].

The most difficult part in the applications of the method of Ważewski is, in general, to verify that  $S = S^*$ . To accomplish this Ważewski introduced the concept of a regular polyfacial set [6, § 14 p. 307 and § 15, p. 309]. However the distinction established by Ważewski between positive and negative faces has certain inconveniences. In some applications of the method of Ważewski there appear sets  $\omega$  such that  $S = S^*$  but whose faces are only "almost positive" and "almost negative". We thus have to work sometimes with sets  $\omega$  that are similar, in some sense, to the regular polyfacial sets and that satisfy the condition  $S = S^*$ .

In the first part of our work we give a generalization of polyfacial regular sets eliminating the distinction between positive and negative faces and such that the main theorem concerning the polyfacial regular sets [6, Théorème 5, p. 310] remains valid. We observe that the sets  $\omega$  considered in Z. Szmydtówna's paper [5, § 4, Théorème 1, p. 24]<sup>1</sup>, in our Theorem II-1 and in Barbālat's paper [1, Théorème 1, p. 303; Théorème 2, p. 305] are generalized regular polyfacial sets, in our sense, but are not regular polyfacial sets.

Szmydtówna [5, Corollaire 1-Remarque 2, p. 30] proves a theorem

<sup>1</sup> Szmydtówna's Theorem 1 is false. We observed that the proof is wrong because the statement: "La frontière de  $\omega$  touchant celle de  $\Omega$  exclusivement sur le plan  $t = \infty$  ..." [5, p. 28] is false.

J. Lewowics [3], developing a counter-example suggested by J. L. Massera, has shown that the theorem is actually false. Nevertheless, Theorems 2 and 3 deduced from Theorem 1 are correct because, in the particular case of linear systems  $\dot{x} = A(t)x$ , with  $A(t)$  defined for  $T \leq t < \infty$ , the solutions are defined for all  $T \leq t < \infty$ .

which generalizes a theorem of Perron. In part II of our work (Theorem II-1) we obtain the same conclusion but starting from hypotheses different from those of Szmydtówna.

*Note*<sup>2</sup>. Our Theorem II-1 improves a result of N. I. Gavrilov. I. M. Rapoport in his book "On some asymptotic methods in the theory of differential equations", Kiev (1954) has also studied problems of this type. For some reference to their work to see "Forty years of Soviet Mathematics", Moscow (1959), Vol. i., pp. 520-521.

Our Theorem II-2 follows the same line of ideas.

Theorem II-3, due to Professor J. L. Massera, shows that in the case  $n = 2$  the asymptotic behavior can be described more completely.

Consider two systems

$$\begin{aligned} (1) \quad & \dot{y} = A(t)y \\ (2) \quad & \dot{x} = A(t)x + g(t, x) \end{aligned}$$

where  $A(t)$  is a continuous matrix for  $t \geq T$  and  $g(t, x)$  a continuous vector-function in  $\Omega = [T, \infty) \times R^{2n}$ .

Suppose that  $g(t, x)$  satisfies some condition ensuring the uniqueness of the solution through each point  $P_0 \in \Omega$  and that all solutions are defined for  $T \leq t < \infty$ . We say that (1) and (2) are asymptotically equivalent if there exists a homeomorphism  $\phi$  from the plane  $t = T$  onto itself such that if  $Q_0 = \phi(P_0)$  then  $\lim_{t \rightarrow \infty} [x(t, P_0) - y(t, Q_0)] = 0$  [4, Cap. IX, § 4, p. 634].

In part III of our work the main result is the establishment of a condition that implies the asymptotic equivalence between two linear systems (Theorem III-3).

The author is deeply indebted to Professor J. L. Massera for his constant guidance and invaluable help during the preparatation of this paper, the result of work done at the Instituto de Matemática y Estadística, Montevideo, Uruguay

### PART I

Let the real-valued functions

$$f_i(t, x_1, \dots, x_n), \quad i = 1, \dots, n,$$

of real variables  $t, x_1, \dots, x_n$  belong to  $C^p, p \geq 1$ , on an open set  $\Omega \subset R^{n+1}$ , i.e., all partial derivatives

$$\frac{\partial^k f_i}{\partial t^{p_0} \partial x_1^{p_1} \dots \partial x_n^{p_n}} \quad (p_0 + p_1 + \dots + p_n = k \leq p)$$

---

<sup>2</sup> The information given in this Note is due to the referee. We have not had access to the above works. We are indebted to him for this.

exist and are continuous on  $\Omega$ .

Consider the differential system

$$(I) \quad \dot{x} = f(t, x)$$

where

$$x = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} \text{ and } f(t, x) = \begin{pmatrix} f_1(t, x_1, \dots, x_n) \\ \vdots \\ f_n(t, x_1, \dots, x_n) \end{pmatrix}$$

with  $(t, x) \in \Omega$ .

Let  $g(t, x)$  be a real-valued function belonging to  $C^{p+1}$  on  $\Omega$ , let  $P_0: (t_0, x_0) \in \Omega$  and let  $x(t)$  be the integral of system (I) passing through the point  $P_0$ . We set  $\varphi(t) = g(t, x(t))$ ; since  $f(t, x) \in C^p$  and  $g(t, x) \in C^{p+1}$  it follows  $\varphi(t) \in C^{p+1}$  on  $(\alpha(P_0), \beta(P_0))$ .

The  $q$ th derivative,  $q \leq p + 1$ , of  $g(t, x)$  at the point  $P_0: (t_0, x_0)$  with respect to the system (I), is by definition

$$\left[ \frac{d^q}{dt^q} \varphi(t) \right]_{t_0} \text{ and is denoted by } [D^q_{(t)}g(P)]_{P_0}.$$

Let  $H_i(P) = H_i(t, x)$ ,  $i = 1, \dots, m$ , be functions  $\in C^{p+1}$  on the open set  $\Omega \subset R^{n+1}$ .

Let

$$\omega = \{P \in \Omega \mid H_i(P) < 0, i = 1, \dots, m\}$$

$$\Gamma_i = \{P \in \Omega \mid H_i(P) = 0, H_j(P) \leq 0, j = 1, \dots, m\}$$

The  $\Gamma_i$  are called *faces* of  $\omega$ .

Such a set  $\omega$  will be called a *generalized regular polyfacial set* relative to (I) if, for each  $i = 1, \dots, m$  and each  $P_0: (t_0, x_0) \in \Gamma_i$ , the following alternative holds:

(1) The smallest index  $q \leq p + 1$  such that  $[D^q_{(t)}H_i(P)]_{P_0} \neq 0$  is odd and the corresponding derivative is positive;

(2)  $P_0$  is not a point of egress.

Let  $L_i, M_i$  be the corresponding sets of points. Useful criteria to verify  $P_0 \in M_i$  are:

(a) the smallest index  $q \leq p + 1$  such that  $[D^q_{(t)}H_i(p)]_{P_0} \neq 0$  is either odd with a negative value of the derivative or even with a positive value of the derivative;

(b) There exists  $[a, b] \subset (\alpha(P_0), \beta(P_0))$  such that  $a < t_0 \leq b$  and  $I([a, b], P_0) \subset \Gamma_i$ .

LEMMA 1. *If  $\omega$  is a generalized regular polyfacial set relative to (I),*

$$S = S^* = \bigcup_{i=1}^m L_i - \bigcup_{i=1}^m M_i .$$

*Proof.* Since  $\Gamma_i = L_i \cup M_i$ ,  $B(\omega, \Omega) \subset \bigcup_{i=1}^m \Gamma_i$ ,

$$S^* \subset S \subset \bigcup_{i=1}^m L_i - \bigcup_{i=1}^m M_i ,$$

it is enough to show that any point  $P_0$  belonging to this last set is a point of strict egress. For such a  $P_0$ ,  $J = \{j \mid P_0 \in L_j\} \neq \phi$ . If  $j \in J$ ,  $H_j(P_0) = 0$  and there exists a  $\delta > 0$  such that  $H_j(t, I(t, P_0)) < 0$  in  $[t_0 - \delta, t_0)$  and  $H_j(t, I(t, P_0)) > 0$  in  $(t_0, t_0 + \delta]$ . If  $j \notin J$ ,  $P_0 \notin \Gamma_j$  whence  $H_j(P_0) < 0$  and there exists a  $\delta > 0$  such that  $H_j(t, I(t, P_0)) < 0$  in  $[t_0 - \delta, t_0)$ . There exists therefore a  $\delta > 0$  such that  $H_j(t, I(t, P_0)) < 0, j = 1, \dots, m, t \in [t_0 - \delta, t_0)$ , and, for at least one  $j(\in J), H_j(t, P_0) > 0, t \in (t_0, t_0 + \delta]$ , so that  $P_0 \in S^*$ .

PART II

Consider the linear differential system

$$\dot{y}_i = f_i(t)y_i + \sum_{j=1}^n g_{ij}(t)y_j , \quad i = 1, \dots, n$$

where the coefficients  $f_i, g_{ij}, T \leqq t < \infty$ , are continuous functions (in general complex-valued) of the real variable  $t$ .

By using Wazewski's method Z. Szmydtówna proved that if

$$R(f_k - f_{k+1}) > 0 , \quad \int_T^\infty R(f_k - f_{k+1})dt = \infty , \quad k = 1, \dots, n - 1 ,$$

and

$$\lim_{t \rightarrow \infty} \frac{g_{ij}}{R(f_k - f_{k+1})} = 0 , \quad i, j = 1, \dots, n , \quad k = 1, \dots, n - 1 ,$$

then there is a system of  $n$  linearly independent solutions  $(y_{1k}, \dots, y_{nk}), k = 1, \dots, n$ , with  $\lim_{t \rightarrow \infty} y_{ik}/y_{kk} = 0$  for  $i \neq k$  [5, Corollaire 1, Remarque 2, p. 30]. This theorem generalizes a theorem of Perron who obtains the same result requiring the existence of a constant  $c > 0$  such that  $R(f_k) > R(f_{k+1}) + c, k = 1, \dots, n - 1$ , and  $\lim_{t \rightarrow \infty} g_{ij} = 0$ .

We notice that Szmydtówna allows the  $f_i, i = 1, \dots, n$ , to be large and the  $g_{ij}$  to be small in some sense. In the following theorem we obtain the same result allowing also the  $f_i$  to be large and the  $g_{ij}$  to be small but in a sense completely different from Szmydtówna's.

THEOREM II-1. *Suppose that the system*

$$(II) \quad \dot{x}_i = f_i(t)x_i + \sum_{j=1}^n g_{ij}(t)x_j, \quad i = 1, \dots, n,$$

satisfies the following hypotheses:

(1) *The coefficients  $f_i, g_{ij}, T \leq t < \infty$ , are continuous functions (in general complex-valued) of the real variable  $t$ .*

(2) *There exists a real-valued continuous function  $h(t), T \leq t < \infty$ , such that for all  $i \neq j$  we have*

$$\begin{aligned} |R(f_i - f_j)| &\leq h(t), \\ \int_T^\infty |g_{ij}(t)| e^{H(t)} dt &< \infty \end{aligned}$$

and

$$\int_T^\infty |R(g_{ii} - g_{jj})| e^{H(t)} dt < \infty,$$

where  $H(t) = \int_T^t h(s) ds$

Then there is a system of  $n$  linearly independent solutions

$$(x_1(t), \dots, x_n(t)) = \begin{pmatrix} x_{11}(t), \dots, x_{1n}(t) \\ \dots\dots\dots\dots\dots\dots \\ x_{n1}(t), \dots, x_{nn}(t) \end{pmatrix}$$

with  $\lim_{t \rightarrow \infty} x_{ik}/x_{kk} = 0$  for all  $i \neq k$ .

*Proof.*

For every fixed integer  $p, 0 < p \leq n$ , we set

$$\omega_p = \{P: (t, x) \mid |x_i|^2 - |x_p|^2 \varphi^2(t) < 0, i \neq p, t > t_0 \geq T\}$$

where  $\varphi(t)$  and  $t_0$  will be conveniently chosen so that, for every  $t \geq t_0$ ,  $\varphi(t) > 0$ ,  $\varphi$  is differentiable,  $\lim_{t \rightarrow \infty} \varphi(t) = 0$  and  $\omega_p$  is a generalized regular polyfacial set.

Let

$$\begin{aligned} H_i(P) &= |x_i|^2 - |x_p|^2 \varphi^2(t), & i \neq p, \\ H_p(P) &= t_0 - t, \end{aligned}$$

it follows that  $\omega_p = \{P \mid H_i(P) < 0, i = 1, \dots, n\}$ .

Set, for  $q \neq p$ ,

$$\begin{aligned} \tilde{\Gamma}_p &= \Gamma_p - \{Q: (t, x) \mid x = 0\} \\ &= \{P \mid |x_q| = |x_p| \varphi(t), |x_i| \leq |x_p| \varphi(t) \text{ for } i \neq p, t \geq t_0, x_p \neq \mathbf{0}\}. \end{aligned}$$

An easy computation shows that

$$\begin{aligned} & \frac{1}{2}[D_{(11)}H_q(P)]P \in \tilde{\Gamma}_q \geq |x_p|^2 \varphi^2(t)[R(f_q - f_p + g_{qq} - g_{pp})] \\ & - |x_p|^2 \varphi(t) \dot{\varphi}(t) - |x_p|^2 \varphi^2(t) \sum_{j \neq p} |g_{pj}| \frac{|x_j|}{|x_p|} \\ & - |x_p|^2 \sum_{j \neq q} |g_{qj}| \frac{|x_j|}{|x_q|} \cdot \frac{|x_q|}{|x_p|} . \end{aligned}$$

Since  $|x_q| = |x_p| \varphi(t) \geq |x_j|$  for  $j \neq p$  it follows that  $|x_j|/|x_p| \leq \varphi(t)$ . As we want  $\varphi(t) > 0$  and  $\lim_{t \rightarrow \infty} \varphi(t) = 0$  we can take  $t_0$  such that  $\varphi(t) < 1$  for  $t \geq t_0$ . Then

$$\begin{aligned} & \frac{1}{2}[D_{(11)}H_q(P)]P \in \tilde{\Gamma} \geq |x_p|^2 \varphi^2(t) R(f_q - f_p + g_{qq} - g_{pp}) \\ & - |x_p|^2 \varphi(t) \dot{\varphi}(t) - |x_p|^2 \varphi^2(t) \sum_{j \neq p} |g_{pj}| - |x_p|^2 \varphi(t) \sum_{j \neq q} |g_{qj}| . \end{aligned}$$

since

$$\begin{aligned} \varphi(t) R(f_q - f_p + g_{qq} - g_{pp}) - \dot{\varphi}(t) - \varphi(t) \sum_{j \neq p} |g_{pj}| - \sum_{j \neq q} |g_{qj}| > \\ - \dot{\varphi}(t) - \varphi(t) h(t) - g(t) , \end{aligned}$$

where

$$g(t) = \{ \sum_{i \neq j} |R(g_{ii} - g_{jj})| + |g_{ij}| \} + e^{-H(t)-t} ,$$

in order to have, for  $q \neq p$ ,  $[D_{(11)}H_q(P)]P \in \tilde{\Gamma}_q > 0$ , it is sufficient to choose  $\varphi(t)$  such that

(A)  $\dot{\varphi}(t) + \varphi(t)h(t) + g(t) = 0$ .

$\varphi(t) = e^{-H(t)} \int_t^\infty g(s)e^{H(s)} ds$  is indeed a solution of (A) satisfying the conditions  $\varphi(t) > 0$ ,  $\varphi$  differentiable and  $\lim_{t \rightarrow \infty} \varphi(t) = 0$ .

If  $\omega_p$  is defined in this way, taking into account that  $[D_{(11)}H_p(P)]P \in \Gamma_p = -1$  and that the set  $\{P \in \Gamma_q \mid x_p = 0\} \subset M_q$ , for  $q \neq p$ , it follows that  $\omega_p$  is a generalized regular polyfacial set.

For  $i \neq p$  we have

$$\begin{aligned} L_i &= \tilde{\Gamma}_i \text{ and } L_p = \phi , \\ M_i &= M = \{P: (t, x) \mid t \geq t_0, x = 0\} \text{ and } M_p = \Gamma_p . \end{aligned}$$

By Lemma 1

$$S = S = \bigcup_{i \neq p} \tilde{\Gamma}_i - \Gamma_p - M .$$

We choose

$$Z_p = \{P: (t, x) \mid t = \tau > t_0, x_p = x_p^0 \neq 0, |x_i| \leq |x_p^0| \varphi(\tau), i \neq p\} = \prod_{j \neq p} B_j^2 ,$$

where  $B_j^2$  is a solid sphere in  $R^2$ . We have



$$Z_p \cap S = Z_p \cap \left[ \bigcup_{i \neq p} \tilde{\Gamma}_i - \Gamma_p - M \right] = \bigcup_{i \neq p} Z_p \cap [\tilde{\Gamma}_i - \Gamma_p - M].$$

For  $i \neq p$

$$Z_p \cap [\tilde{\Gamma}_i - \Gamma_p - M] = \{P: (t, x) \mid t = \tau, x_p = x_p^0, |x_i| = |x_p^0| \varphi(\tau), |x_j| \leq |x_p^0| \varphi(\tau), j \neq p\} = B_1^2 \times \cdots \times B_{i-1}^2 \times S_i^1 \times B_{i+1}^2 \times \cdots \times B_n^2$$

(in the cartesian product above  $B_p^2$  is excluded) where  $S_i^1$  is the boundary of  $B_i^2$  in  $R^2$ .

Modulo homeomorphisms we have therefore  $Z_p = B^{2n-2}$  (solid sphere in  $R^{2n-2}$ ) and  $Z_p \cap S = S^{2n-3} =$  Boundary of  $B^{2n-2}$  in  $R^{2n-2}$ , so that  $Z_p \cap S$  is not a retract of  $Z_p$ .

There is however a retraction  $\phi: S \rightarrow Z_p \cap S$  given by  $\phi(P) = P^*$ , with  $t^* = \tau, x_p^* = x_p^0, x_i^* = \varphi(\tau)/\varphi(t) |x_p^0|/|x_p| \cdot x_i, i \neq p$ . The verification is trivial.

By using the theorem of Ważewski we can conclude the existence of at least one point  $P_0: (\tau, x_0) \in Z_p - S$  with  $I(t, P_0) \subset \omega_p$  for every  $t \geq \tau$ . This means that the solution  $x_p(t) = (x_{1p}(t), \dots, x_{np}(t))$  of (II) passing through  $P_0$  satisfies

$$\frac{|x_{ip}(t)|}{|x_{pp}(t)|} < \varphi(t) \text{ for } t \geq \tau \text{ and } i \neq p.$$

Letting  $p = 1, \dots, n$  we find  $n$  solutions  $(x_1(t), \dots, x_n(t))$  with the required property. Let us show that these solutions can be taken linearly independent.

By choosing  $Z_p$  with sufficiently large  $\tau$  and  $x_{pp}^0 = 1$  the absolute values of the coordinates  $x_{ip}, i \neq p$ , of the points of  $Z_p$  can be made arbitrarily small. We then have

$$\begin{pmatrix} x_1(\tau) \\ x_2(\tau) \\ \vdots \\ x_n(\tau) \end{pmatrix} = \begin{pmatrix} \varepsilon_{11} \cdots \varepsilon_{1n} \\ \varepsilon_{21} \cdots \varepsilon_{2n} \\ \dots\dots\dots \\ \varepsilon_{n1} \cdots \varepsilon_{nn} \end{pmatrix}$$

where  $\varepsilon_{ii} = 1$  and the  $|\varepsilon_{ij}|$  are smaller than any given positive number for all  $i \neq j$ . This completes the proof

In the following theorem we will look for linearly independent solutions of (II) with similar properties to those of Theorem II-I but not necessarily requiring that they form a fundamental set of solutions of (II).

**THEOREM II-2.** *Suppose that the system (II) satisfies the following hypotheses:*

(1) *The coefficients  $f_i, g_{ij}, T \leq t < \infty$ , are continuous functions (in general complex-valued) for the real variable  $t$ .*

(2) *There exists a natural number  $r \leq n$  such that  $R(f_r) = \dots = R(f_n)$ ,  $R(f_i) \geq R(f_r)$  for all  $i < r$ ,  $\int_T^\infty |g_{ij}(t)| dt < \infty$  for all  $i \neq j$  and  $\int_T^\infty |R(f_{ii} - g_{jj})| dt < \infty$ .*

*Then there exists  $s + 1$  ( $r + s = n$ ) linearly independent solutions*

$$(x_r(t), \dots, x_n(t)) = \begin{pmatrix} x_{1r}(t) \dots x_{1n}(t) \\ \dots\dots\dots\dots\dots\dots \\ x_{nr}(t) \dots x_{nn}(t) \end{pmatrix}$$

*such that  $\lim_{t \rightarrow \infty} x_{ik}/x_{kk} = 0$  for all  $i \neq k, k = r, \dots, n$ .*

*Proof.* Given an integer  $p, r \leq p \leq n$ , we proceed exactly as in Theorem II-1 up to the point where we got the expression:

$$\varphi(t)R(f_q - f_p + g_{qq} - g_{pp}) - \dot{\varphi}(t) - \varphi(t) \sum_{j \neq p} |g_{pj}| - \sum_{j \neq q} |g_{qj}|$$

which we denote by  $B_q$ .

As we have  $R(f_q - f_p) \geq 0$  for all  $q, 0 < q \leq n$ , it follows ( $\varphi(t) < 1$ )

$$B_q \geq -\dot{\varphi}(t) - g(t) \text{ where } \int_{t_0}^\infty g(t)dt < \infty.$$

Making  $\varphi(t) = \int_t^\infty [g(s) + e^{-s}]ds$  it follows that  $\varphi(t) > 0$ ,  $\varphi$  is differentiable,  $\lim_{t \rightarrow \infty} \varphi(t) = 0$  and  $B_q > 0$ .

Proceeding as in Theorem II-1 we find a set of  $(s + 1)$  solutions  $(x_r(t), \dots, x_n(t))$ . Still by a similar reasoning we may show that these solutions can be so chosen that for a sufficiently large  $\tau$  we have

$$(x_r(\tau), \dots, x_n(\tau)) = \begin{pmatrix} \varepsilon_{1r}, \dots, \varepsilon_{1n} \\ \dots\dots\dots\dots\dots\dots \\ \varepsilon_{rr}, \dots, \varepsilon_{rn} \end{pmatrix}$$

with  $\varepsilon_{ii} = 1$  and the  $|\varepsilon_{ij}|, i \neq j$ , smaller than any given positive number, so that, they are linearly independent.

If  $n = 2$  Theorem II-2, with some supplementary hypotheses, leads us to a deeper result. As already mentioned in the Introduction the following theorem is due to Professor J. L. Massera with whose permission it is reproduced here.

**THEOREM II-3.** *Suppose that the system*

$$\dot{x} = f_1(t)x + g_{11}(t)x + g_{12}(t)y$$

$$\dot{y} = f_2(t)y + g_{21}(t)x + g_{22}(t)y$$

satisfies the following hypotheses:

(1) The coefficients  $f_i, g_{ij}, T \leqq t < \infty$ , are continuous real-valued functions of the real variable  $t$ .

(2)  $f_1(t) \geqq f_2(t)$ ,  $\int_T^\infty (f_1(t) - f_2(t))dt = \infty$ ,  $\int_T^\infty |g_{ij}(t)| dt < \infty$  for  $i \neq j$  and  $\int_T^\infty |g_{11}(t) - g_{22}(t)| dt < \infty$ .

Then there exists a solution  $(x_1(t), y_1(t))$  satisfying  $\lim_{t \rightarrow \infty} x_1(t)/y_1(t) = 0$  and, for any other solution  $(x(t), y(t))$  which is not proportional to  $(x_1(t), y_1(t))$ , we have  $\lim_{t \rightarrow \infty} y(t)/x(t) = 0$ .

*Proof.* The existence of a solution  $(x_1(t), y_1(t))$  with the required property follows from Theorem II-2.

Without loss of generality we may assume  $g_{11} = g_{22} = 0$ . Choose  $t_0 \geqq T$  so large that  $\int_{t_0}^\infty (|g_{12}| + |g_{21}|)dt < \pi/4$ . Let  $(x_2(t), y_2(t))$  be the solution which satisfies  $x_2(t_0) = 1, y_2(t_0) = 0$ . Setting  $\theta(t) = \arg(x_2(t), y_2(t))$ , we claim, in the first place, that  $|\theta(t)| < \pi/4$  for  $t \geqq t_0$ . Assume that this were not the case. It then follows that there exists an interval  $(t_1, t_2), t_1 \geqq t_0$ , such that  $\theta(t_1) = 0, |\theta(t_2)| = \pi/4, 0 < |\theta(t)| < \pi/4$  for  $t_1 < t < t_2$ , say,  $\theta(t_2) = \pi/4, 0 < \theta(t) < \pi/4$  for  $t_1 < t < t_2$ , whence  $x_2(t) \cdot y_2(t) > 0$  in  $(t_1, t_2)$ . Since

$$\dot{\theta} = \frac{\dot{y}_2 x_2 - \dot{x}_2 y_2}{x_2^2 + y_2^2} = \frac{g_{21} x_2^2 - g_{12} y_2^2 + (f_2 - f_1) x_2 y_2}{x_2^2 + y_2^2},$$

we arrive to the contradiction  $\pi/4 = \theta(t_2) - \theta(t_1) = \int_{t_1}^{t_2} \dot{\theta} dt < \pi/4$ .

We next prove that  $\lim_{t \rightarrow \infty} y_2(t)/x_2(t) = 0$ , or equivalently  $\lim_{t \rightarrow \infty} \theta(t) = 0$ . There exists a sequence  $t_n \rightarrow \infty$  with  $\theta(t_n) \rightarrow 0$ , otherwise  $\theta(t) > \theta_0 > 0$ , say, which leads to the contradiction

$$\begin{aligned} \theta(t) - \theta(t_0) &\leqq - \int_{t_0}^t (f_1(t) - f_2(t)) \sin \theta(t) \cos \theta(t) dt \\ &\quad + \int_{t_0}^t (|g_{12}| + |g_{21}|) dt \rightarrow -\infty. \end{aligned}$$

Now, given  $\varepsilon > 0$ , choose  $t_n$  such that  $|\theta(t_n)| < \varepsilon/2, \int_{t_n}^\infty (|g_{12}| + |g_{21}|) dt < \varepsilon/2$ . An argument similar to the one used to prove  $|\theta(t)| < \pi/4$  then shows that  $|\theta(t)| < \varepsilon$  for  $t \geqq t_n$ .

Assume  $t_0$  large enough so that  $|x_1(t)|/|y_1(t)| < 1, |y_2(t)|/|x_2(t)| < 1$  for  $t \geqq t_0$  and, say,  $y_1(t) > 0, x_2(t) > 0$ ; then

$$\dot{y}_1(t) \leqq (f_2(t) + |g_{21}(t)|) \cdot y_1(t),$$

$$\dot{x}_2(t) \geq (f_1(t) - |g_{12}(t)|) \cdot x_2(t) ,$$

whence

$$y_1(t) \leq y_1(t_0) \cdot \exp \left( \int_{t_0}^t (f_2(t) + |g_{21}(t)|) dt \right) ,$$

$$x_2(t) \geq x_2(t_0) \cdot \exp \left( \int_{t_0}^t (f_1(t) - |g_{12}(t)|) dt \right)$$

and

$$\frac{y_1(t)}{x_2(t)} \leq \frac{y_1(t_0)}{x_2(t_0)} \cdot \exp \left( \int_{t_0}^t (f_2(t) - f_1(t) + |g_{12}(t)| + |g_{21}(t)|) dt \right) \rightarrow 0 .$$

Finally, any solution  $(x(t), y(t))$  which is not proportional to  $(x_1(t), y_1(t))$  satisfies, for a certain constant value  $k$ ,

$$\frac{y(t)}{x(t)} = \frac{y_2(t) + ky_1(t)}{x_2(t) + ky_2(t)} = \frac{(y_2(t)/x_2(t)) + k(y_1(t)/x_2(t))}{1 + k(x_1(t)/y_1(t))(y_1(t)/x_2(t))} \rightarrow 0 .$$

### PART III

Consider the linear differential systems

$$(III) \quad \dot{x} = A(t)x + B(t)x$$

$$(III') \quad \dot{y} = A(t)y$$

where  $A(t), B(t), T \leq t < \infty$ , are continuous complex matrix functions.

Conti [2, Theorem I, p. 589] proved that: *if  $\int_0^\infty |B(t)| dt < \infty$  where  $B(t) = (b_{ij}^i(t))$  and  $|B(t)| = \sum_{i,j} |b_{ij}^i(t)|$  and if (III') is uniformly stable, then the system (III) and (III') are asymptotically equivalent<sup>3</sup>.*

The theorem of Wintner [7, 7-i, p. 423] stating that:

*If  $B(t) = (b_{ij}^i(t))$ ,  $T \leq t < \infty$ ,  $i, j = 1, \dots, n$ , is a matrix of  $n^2$  continuous functions satisfying  $\int_0^\infty |B(t)| dt < \infty$ , then every solution of  $\dot{x} = B(t)x$  tends to a finite limit as  $t \rightarrow \infty$ , is a particular case of Conti's result ( $A(t) = 0$ ).*

Our Theorem III-3, is also a generalization of Wintner's theorem but different from that of Conti.

Theorems III-1 and III-2, which are preliminary to Theorem III-3, give us some information, though less than asymptotic equivalence, concerning the behavior of two systems, one of which not necessarily linear.

**THEOREM III-1.** *Suppose that the systems*

---

<sup>3</sup> The theorem of Conti is actually more general. We have considered the theorem applied to linear systems only.

$$\begin{aligned} \text{(III-1)} \quad & \dot{x}_i = f_i(t)x_i + g_i(t, x) , \\ \text{(III-2)} \quad & \dot{y}_i = f_i(t)y_i , \\ & i = 1, \dots, n , \quad g(t, x) = (g_i(t, x)) \end{aligned}$$

satisfy the following hypotheses:

(1)  $f_i(t)$ ,  $T \leq t < \infty$ , are continuous functions (in general complex-valued) of the real variable  $t$ ;  $g_i(t, x)$  are functions (in general complex-valued) continuous in

$$\Omega = \{(t, x) \mid t > T, |x| = \sum_{i=1}^n |x_i| < \infty\}$$

and satisfy some condition which implies the existence of only one integral passing through each point of  $\Omega$ .

$$(2) \quad |g(t, x)| \leq |x| F(t) \text{ on } \Omega.$$

(3) There exists a negative constant  $K$  such that

$$K \leq \int_t^v R(f_i(\tau)) d\tau$$

for all  $v \geq t > T$  and

$$\int_t^\infty F(t) \exp \left[ \int_t^t R(f_i(s)) ds \right] dt < \infty$$

for all  $i = 1, \dots, n$ .

Then for every solution  $y(t)$  of III-2 there is a solution  $x(t)$  of III-1 such that  $\lim_{t \rightarrow \infty} [x(t) - y(t)] = 0$ .

*Proof.* We define  $\omega = \{P \in \Omega \mid |x_i - y_i(t)| < \varphi_i(t), t > t_0 \geq T\}$  where the  $\varphi_i(t)$  and  $t_0$  will be adequately chosen so that for all  $t \geq t_0$ ,  $i = 1, \dots, n$ , we have:  $\varphi_i(t) > 0$ ,  $\varphi_i$  differentiable,  $\lim_{t \rightarrow \infty} \varphi_i(t) = 0$  and  $\omega$  a generalized regular polyfacial set.

If we put

$$\begin{aligned} H_i(P) &= |x_i - y_i(t)|^2 - \varphi_i^2(t) , & i = 1, \dots, n \\ H_{n+1}(P) &= t_0 - t \end{aligned}$$

it follows that  $\omega = \{P \mid H_i(P) < 0, i = 1, \dots, n + 1\}$ .

For all  $i$ ,  $1 \leq i \leq n$ ,

$$\Gamma_i = \{P \in \Omega \mid |x_i - y_i(t)| = \varphi_i(t), |x_j - y_j(t)| \leq \varphi_j(t), j = 1, \dots, n, t \geq t_0\} .$$

An easy computation shows that

$$\frac{1}{2} [D_{(III-1)} H_i(P)] P \in \Gamma_i \geq \varphi_i^2(t) R[f_i(t)] - \varphi_i(t) \dot{\varphi}_i(t) - \varphi_i(t) F(t) \left[ \sum_{k=1}^n |x_i - y_k| + |y_k| \right].$$

As we want  $\varphi_i(t) > 0$  and  $\lim_{t \rightarrow \infty} \varphi_i(t) = 0$ , we can take  $t_0$  such that  $\sum_{i=1}^n \varphi_i(t) < 1$  for all  $t \geq t_0$ . Then,

$$\begin{aligned} \frac{1}{2} [D_{(III-1)} H_i(P)] P \in \Gamma_i &\geq \varphi_i^2(t) R[f_i(t)] - \varphi_i(t) \dot{\varphi}_i(t) - \varphi_i(t) F(t) \\ &- \varphi_i(t) F(t) \sum_{k=1}^n |c_k| \exp \int_{t_0}^t R[f_k(s)] ds \geq \varphi_i^2(t) R[f_i(t)] - \varphi_i(t) \dot{\varphi}_i(t) \\ &- \varphi_i(t) F(t) \sum_{k=1}^n |d_k| \exp \int_{t_0}^t R[f_k(s)] ds \geq \varphi_i^2(t) R[f_i(t)] - \varphi_i(t) \dot{\varphi}_i(t) \\ &- \varphi_i(t) h(t) \end{aligned}$$

where we can assume  $\int_{t_0}^{\infty} h(t) dt < \infty$  and, without loss of generality,  $h(t) > 0$  for all  $t \geq t_0$ .

In order to have, for all  $i = 1, \dots, n$ ,  $[D_{(III-1)} H_i(P)] P \in \Gamma_i > 0$  it is sufficient to choose  $\varphi(t)$  such that

$$-\dot{\varphi}_i(t) + R[f_i(t)]\varphi_i(t) - h(t) > 0.$$

The problem is then to look for a solution  $z(t)$  of  $\dot{z} < \sigma(t)z - \gamma(t)$  satisfying  $z(t) > 0$  for all  $t \geq t_0$ ,  $\lim_{t \rightarrow \infty} z(t) = 0$ , knowing that  $\gamma(t) > 0$  for all  $t \geq t_0$ ,  $\int_{t_0}^{\infty} \gamma(t) dt < \infty$  and  $\int_{t_0}^v \sigma(s) ds \geq K$  for some constant  $K$  and all  $v \geq t \geq t_0$ . If  $W(t)$  is a solution of  $\dot{W} = \sigma(t)W - \gamma(t)$  it follows that  $z(t) = 2W(t)$  is a solution of  $\dot{z} < \sigma(t)z - \gamma(t)$ . It is then sufficient to find a solution  $W(t)$  satisfying  $W(t) > 0$  for all  $t \geq t_0$  and  $\lim_{t \rightarrow \infty} W(t) = 0$ . The solution  $W(t) = \exp \left( \int_{t_0}^t \sigma(s) ds \right) \cdot \int_t^{\infty} \gamma(v) \exp \left( - \int_{t_0}^v \sigma(s) ds \right) dv$  exists and indeed  $W(t) \rightarrow 0$  as  $t \rightarrow \infty$  because

$$W(t) = \int_t^{\infty} \gamma(v) \exp \left( - \int_t^v \sigma(s) ds \right) dv \leq e^{-K} \int_t^{\infty} \gamma(v) dv.$$

Since  $[D_{(III-1)} H_{n+1}(P)] = -1$  it follows that  $\omega$  is a generalized regular polyfacial set and  $S = S^* = \bigcup_{i=1}^n \Gamma_i - \Gamma_{n+1}$ .

If we choose

$$Z = \{(t, x) \mid t = \tau > t_0, |x_j - y_j(\tau)| \leq \varphi_j(\tau), j = 1, \dots, n\}$$

it follows that  $S \cap Z = \bigcup_{i=1}^n \Gamma_i \cap Z - \Gamma_{n+1}$

$$\begin{aligned} \Gamma_i \cap Z &= \{(t, x) \mid t = \tau, |x_i - y_i(\tau)| \\ &= \varphi_i(\tau), |x_j - y_j(\tau)| \leq \varphi_j(\tau), j = 1, \dots, n\}. \end{aligned}$$

Then  $Z = \prod_{j=1}^n B_j^2$

$$Z \cap S = \bigcup_{j=1}^n B_1^3 \times \cdots \times B_{j-1}^3 \times S_j^1 \times B_{j+1}^3 \times \cdots \times B_n^3$$

and, modulo homomorphisms,  $Z = B^{2n}$ ,  $Z \cap S = S^{2n-1}$  so that  $Z \cap S$  is not a retract of  $Z$ . However, it is easily seen that  $\phi: S \rightarrow S \cap Z$  given by  $\phi(P) = P^*$ , with  $t^* = \tau$ ,  $x_i^* = y_i(\tau) + [x_i - y_i(t)]\varphi_i(\tau)/\varphi_i(t)$ , is a retraction.

Using the theorem of Ważewski we can conclude the existence of at least one point  $P_0: (\tau, x_0) \in Z - S$  such that  $(t, x(t, P_0)) = I(t, P_0) \subset \omega$  for all  $t \geq t_0$ .

Since  $x(t, P_0)$  is defined in the future, i.e.,  $\beta(P_0) = \infty$  (because  $\beta(P_0) < \infty$  implies  $\{I(t, P_0) \mid t_0 \leq t < \beta(P_0)\}$  bounded, which is not possible), it follows that  $\lim_{t \rightarrow \infty} [x(t, P_0) - y(t)] = 0$ .

**COROLLARY 1.** *Suppose that the systems*

$$(III-1') \quad \dot{x}_i = f_i(t)x_i + \sum_{j=1}^n g_{ij}(t)x_j$$

$$(III-2') \quad \dot{y}_i = f_i(t)y_i$$

$$i = 1, \dots, n, \quad g(t) = (g_{ij}(t))$$

*satisfy the following hypotheses:*

(1) *The coefficients  $f_i, g_{ij}, T \leq t < \infty$ , are continuous functions (in general complex-valued) of the real variable  $t$ .*

(2) *There exists a constant  $K$  such that*

$$K \leq \int_t^v R[f_i(s)]ds \text{ for all } v \geq t \geq T \text{ and}$$

$$\int_x^t |g(t)| \exp \left\{ \int_x^t R[f_i(s)]ds \right\} dt < \infty, \quad i = 1, \dots, n.$$

*Then for every solution  $y(t)$  of (III-2') there exists a solution  $x(t)$  of (III-1') such that  $\lim_{t \rightarrow \infty} [x(t) - y(t)] = 0$ .*

The theorem of Wintner mentioned before follows a once from Corollary 1.

**THEOREM III-2.** *Suppose that the systems*

$$(III-A) \quad \dot{x}_i = \sum_{j=1}^n f_{ij}(t)x_j + g_i(t, x)$$

$$(III-B) \quad \dot{y}_i = \sum_{j=1}^n f_{ij}(t)y_j$$

$$i, j = 1, \dots, n, \quad g(t, x) = (g_j(t, x))$$

*satisfy the following hypotheses:*

(1)  $f_{ij}(t), T \leq t < \infty$ , are continuous functions (in general complex-valued) of real variable  $t$ ;  $g_i(t, x)$  are functions (in general complex-valued) continuous in

$$\Omega = \{(t, x) \mid t < T, |x| < \infty\}$$

and satisfy some condition which implies the existence of only one integral passing through each point of  $\Omega$ .

(2)  $|g(t, x)| \leq |x| F(t)$  in  $\Omega$ .

(3) There exists a constant  $K$  such that

$$K \leq \int_t^v R[f_{ii}(s)]ds \text{ for all } v \geq t \geq T \text{ and}$$

$$\int_r^\infty F(t) \exp \left\{ \int_r^t R[f_{ii}(s)]ds \right\} dt < \infty, \quad i = 1, \dots, n$$

$$\int_r^\infty |f_{ij}(t)| \exp \left\{ \int_r^t R[f_{kk}(s)]ds \right\} dt < \infty, \quad k = 1, \dots, n, i \neq j.$$

Then for every solution  $y(t)$  of (III-B) there is a solution  $x(t)$  of (III-A) such that  $\lim_{t \rightarrow \infty} [x(t) - y(t)] = 0$

*Proof.* Consider the systems

$$(III-A) \quad \dot{x}_i = f_{ii}(t)x_i + \tilde{g}_i(t, x) \text{ where } \tilde{g}_i(t, x) = g_i(t, x) + \sum_{j \neq i} f_{ij}(t)x_j$$

$$(III-C) \quad \dot{z}_i = f_{ii}(t)z_i.$$

These systems satisfy the condition of Theorem III-1. Hence for every solution  $z(t)$  of (III-C) there is a solution  $x(t)$  of

$$(III-A) \quad \text{such that } \lim_{t \rightarrow \infty} [z(t) - x(t)] = 0$$

Consider now the systems

$$(III-B) \quad \dot{y}_i = \sum_{j=1}^n f_{ij}(t)y_j$$

$$(III-C) \quad \dot{z}_i = f_{ii}(t)z_i.$$

It is easy to see that they also satisfy the hypotheses of Theorem III-1. Hence for every solution  $z(t)$  of (III-C) there is a solution  $y(t)$  of (III-B) such that  $\lim_{t \rightarrow \infty} [y(t) - z(t)] = 0$ . But we can also prove that for every solution  $y(t)$  of (III-B) there is a solution  $z(t)$  of (III-C) such that  $y(t) - z(t) \rightarrow 0$  as  $t \rightarrow \infty$ . For that purpose it is enough to show that there is a fundamental set  $z^1(t), \dots, z^n(t)$  of solutions of (III-C) such that the solutions  $y^1(t), \dots, y^n(t)$  satisfying  $y^i(t) - z^i(t) \rightarrow 0$  as  $t \rightarrow \infty$ , for all  $i = 1, \dots, n$ , are a fundamental set of solutions of (III B).



Let us take  $z^i(t) = \begin{pmatrix} z_1^i(t) \\ \vdots \\ z_n^i(t) \end{pmatrix}$  such that  $z_j^i(t) = 0$  for all  $j \neq i$  and  $z_j^i(t) = \exp\left(\int_T^t f_{ii}(s) ds\right)$  for all  $i = 1, \dots, n$ .

The corresponding  $y^i(t)$ ,  $i = 1, \dots, n$ , satisfy  $\lim_{t \rightarrow \infty} y_j^i(t) = 0$  if  $j \neq i$  and  $\lim_{t \rightarrow \infty} |y^i(t) - \exp\left[\int_T^t f_{ii}(s) ds\right]| = 0$ . Hence, there exists  $t_0$  such that  $t \geq t_0$  implies

$$|y^i(t) - \exp\left[\int_T^t f_{ii}(s) ds\right]| < \frac{1}{2}e^\kappa.$$

Whence

$$|y^i(t)| > \exp\left\{\int_T^t R[f_{ii}(s)] ds\right\} - \frac{1}{2}e^\kappa \geq \frac{1}{2}e^\kappa.$$

Therefore, for any  $\varepsilon > 0$  there is a  $t(\varepsilon)$  such that  $t \geq t(\varepsilon)$  implies  $|y^i(t)| > 1/2 e^\kappa$ ,  $i = 1, \dots, n$ , and  $|y_j^i(t)| < \varepsilon$  for all  $i \neq j$ . This implies the existence of a  $\tau \geq T$  with  $\det(y^1(\tau), \dots, y^n(\tau)) \neq 0$  and  $(y^1(t), \dots, y^n(t))$  is a fundamental set of solutions of (III-B).

From the results concerning the systems (III-A), (III-C) and (III-B), (III-C) we conclude that for every solution  $y(t)$  of (III-B) there is a solution  $x(t)$  of (III-A) such that  $\lim_{t \rightarrow \infty} [x(t) - y(t)] = 0$ .

**THEOREM III-3.** *Suppose that the systems*

$$(III-\alpha) \quad \dot{x}_i = \sum_{j=1}^n f_{ij}(t)x_j(t)x_j + \sum_{j=1}^n g_{ij}(t)x_j$$

$$(III-\beta) \quad \dot{y}_i = \sum_{j=1}^n f_{ij}(t)y_j \quad i, j = 1, \dots, n$$

*satisfy the following hypotheses:*

(1) *The coefficients  $f_{ij}, g_{ij}$ ,  $T \leq t < \infty$ , are continuous functions (in general complex-valued) of the real variable  $t$ .*

(2) *There exists constant  $K$  such that  $K \leq \int_t^v R[f_{ii}(s)] ds$  for all  $v \geq t \geq T$ ,  $i = 1, \dots, n$ , and*

$$\int_t^\infty |g_{ij}(t)| \exp\left\{\int_T^t R[f_{kk}(s)] ds\right\} dt < \infty, \quad i, j, k = 1, \dots, n$$

$$\int_t^\infty |f_{ij}(t)| \exp\left\{\int_T^t R[f_{kk}(s)] ds\right\} dt < \infty, \quad i, j, k = 1, \dots, n, i \neq j.$$

*Then the systems (III- $\alpha$ ) and (III- $\beta$ ) are asymptotically equivalent.*

*Proof.* By Theorem III-2 for every solution  $y(t)$  of (III- $\beta$ ) there is a solution  $x(t)$  of (III- $\alpha$ ) such that  $\lim_{t \rightarrow \infty} [x(t) - y(t)] = 0$ .

Let us show that given a fundamental set  $(y^1(t), \dots, y^n(t))$  of solutions of (III- $\beta$ ) the corresponding solutions  $(x^1(t), \dots, x^n(t))$  of (III- $\alpha$ ) satisfying  $\lim_{t \rightarrow \infty} [x^i(t) - y^i(t)] = 0, i = 1, \dots, n$ , also form a fundamental set of solutions.

Consider the auxiliary system

$$(III-\gamma) \quad \dot{z}_i = f_{ii}(t)z_i, \quad i = 1, \dots, n.$$

Applying the argument used in Theorem III-2 to the systems (III- $\beta$ ), (III- $\gamma$ ) we conclude that there exists a fundamental set  $(y^1(t), \dots, y^n(t))$  of solutions of (III- $\beta$ ) and a  $t_0$  such that  $t \geq t_0$  implies

$$|y_i^i(t)| \geq \frac{1}{2}e^\kappa \text{ and } y_j^i(t) \rightarrow 0 \text{ as } t \rightarrow \infty \text{ for all } i \neq j.$$

Let  $(x^1(t), \dots, x^n(t))$  be the solutions of (III- $\alpha$ ) such that  $\lim_{t \rightarrow \infty} [x^i(t) - y^i(t)] = 0$  (the existence of which follows from Theorem III-2). Then  $\lim_{t \rightarrow \infty} x_j^i(t) = 0$  for all  $i \neq j$  and there exists  $\tau \geq t_0$  such that  $t \geq \tau$  implies  $|x_i^i(t)| > 1/4e^\kappa$ .

For sufficiently large  $t$  we have therefore

$$\det(x^1(t), \dots, x^n(t)) \neq 0$$

and this means that  $(x^1(t), \dots, x^n(t))$  is a fundamental set of solutions of (III- $\alpha$ ).

The systems (III- $\alpha$ ) and (III- $\beta$ ) being linear this implies that they are asymptotically equivalent.

## BIBLIOGRAPHY

1. I. Barbălat, *Applications du principe topologique de T. Wazewski aux équations différentielles du second ordre*, Ann. Polon. Math., **3** (1958-1959), 303-317.
2. R. Conti, *Sistemi differenziali asintoticamente equivalenti*, Rendiconti dell'Accademia Nazionale dei Lincei. ser. VIII, vol. XXII, fasc. 5, (1957), 588-592.
3. J. Lewowics, *Sobre um teorema de Szmidtówna*. Publicaciones del Instituto de Matemática y Estadística de la Facultad de Ingeniería y Agrimensura, Montevideo, Uruguay, III, (1960), 125-134.
4. G. R. Sansone, e Conti, *Equazioni differenziali non lineari*, Consiglio Nazionale delle Ricerche, Monografie Matematiche, 3 Edizioni Cremonese, Roma, (1956).
5. Z. Szmydtówna, *Sur l'allure asymptotique des intégrales des équations différentielles ordinaires*, Ann. Soc. Polon. Math. Tath. Tome 24, Fasc. II, (1951), 17-34.
6. T. Wazewski, *Sur un principe topologique de l'examen de l'allure asymptotique des intégrales des équations différentielles ordinaires*, Ann. Soc. Polon. Math. Tome 20, (1947), 279-313.
7. A. Wintner, *Small perturbations*, Amer. J. Math., **67**, (1945), 417-430.



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

RALPH S. PHILLIPS

Stanford University  
Stanford, California

F. H. BROWNELL

University of Washington  
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California  
Los Angeles 7, California

L. J. PAIGE

University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

E. F. BECKENBACH  
T. M. CHERRY

D. DERRY  
M. OHTSUKA

H. L. ROYDEN  
E. SPANIER

E. G. STRAUS  
F. WOLF

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE COLLEGE  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE COLLEGE  
UNIVERSITY OF WASHINGTON  
\* \* \*

AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
HUGHES AIRCRAFT COMPANY  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. The price per volume (4 numbers) is \$12.00; single issues, \$3.50. Back numbers are available. Special price to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues, \$1.25.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Reprinted 1966 in the United States of America

# Pacific Journal of Mathematics

Vol. 11, No. 4

, 1961

A. V. Balakrishnan, <i>Prediction theory for Markoff processes</i> . . . . .	1171
Dallas O. Banks, <i>Upper bounds for the eigenvalues of some vibrating systems</i> . . . . .	1183
A. Białynicki-Birula, <i>On the field of rational functions of algebraic groups</i> . . . . .	1205
Thomas Andrew Brown, <i>Simple paths on convex polyhedra</i> . . . . .	1211
L. Carlitz, <i>Some congruences for the Bell polynomials</i> . . . . .	1215
Paul Civin, <i>Extensions of homomorphisms</i> . . . . .	1223
Paul Joseph Cohen and Milton Lees, <i>Asymptotic decay of solutions of differential inequalities</i> . . . . .	1235
István Fáry, <i>Self-intersection of a sphere on a complex quadric</i> . . . . .	1251
Walter Feit and John Griggs Thompson, <i>Groups which have a faithful representation of degree less than <math>(p - 1/2)</math></i> . . . . .	1257
William James Firey, <i>Mean cross-section measures of harmonic means of convex bodies</i> . . . . .	1263
Avner Friedman, <i>The wave equation for differential forms</i> . . . . .	1267
Bernard Russel Gelbaum and Jesus Gil De Lamadrid, <i>Bases of tensor products of Banach spaces</i> . . . . .	1281
Ronald Kay Gettoor, <i>Infinitely divisible probabilities on the hyperbolic plane</i> . . . . .	1287
Basil Gordon, <i>Sequences in groups with distinct partial products</i> . . . . .	1309
Magnus R. Hestenes, <i>Relative self-adjoint operators in Hilbert space</i> . . . . .	1315
Fu Cheng Hsiang, <i>On a theorem of Fejér</i> . . . . .	1359
John McCormick Irwin and Elbert A. Walker, <i>On <math>N</math>-high subgroups of Abelian groups</i> . . . . .	1363
John McCormick Irwin, <i>High subgroups of Abelian torsion groups</i> . . . . .	1375
R. E. Johnson, <i>Quotient rings of rings with zero singular ideal</i> . . . . .	1385
David G. Kendall and John Leonard Mott, <i>The asymptotic distribution of the time-to-escape for comets strongly bound to the solar system</i> . . . . .	1393
Kurt Kreith, <i>The spectrum of singular self-adjoint elliptic operators</i> . . . . .	1401
Lionello Lombardi, <i>The semicontinuity of the most general integral of the calculus of variations in non-parametric form</i> . . . . .	1407
Albert W. Marshall and Ingram Olkin, <i>Game theoretic proof that Chebyshev inequalities are sharp</i> . . . . .	1421
Wallace Smith Martindale, III, <i>Primitive algebras with involution</i> . . . . .	1431
William H. Mills, <i>Decomposition of holomorphs</i> . . . . .	1443
James Donald Monk, <i>On the representation theory for cylindric algebras</i> . . . . .	1447
Shu-Teh Chen Moy, <i>A note on generalizations of Shannon-McMillan theorem</i> . . . . .	1459
Donald Earl Myers, <i>An imbedding space for Schwartz distributions</i> . . . . .	1467
John R. Myhill, <i>Category methods in recursion theory</i> . . . . .	1479
Paul Adrian Nickel, <i>On extremal properties for annular radial and circular slit mappings of bordered Riemann surfaces</i> . . . . .	1487
Edward Scott O'Keefe, <i>Primal clusters of two-element algebras</i> . . . . .	1505
Nelson Onuchic, <i>Applications of the topological method of Ważewski to certain problems of asymptotic behavior in ordinary differential equations</i> . . . . .	1511
Peter Perkins, <i>A theorem on regular matrices</i> . . . . .	1529
Clinton M. Petty, <i>Centroid surfaces</i> . . . . .	1535
Charles Andrew Swanson, <i>Asymptotic estimates for limit circle problems</i> . . . . .	1549
Robert James Thompson, <i>On essential absolute continuity</i> . . . . .	1561
Harold H. Johnson, <i>Correction to "Terminating prolongation procedures"</i> . . . . .	1571