

Pacific Journal of Mathematics

CONVERGENCE OF INVERSE SYSTEMS

JACK SEGAL

CONVERGENCE OF INVERSE SYSTEMS

JACK SEGAL

1. Introduction. Let X_i be a metrizable continuum for each positive integer i , and let f_i be a mapping of X_{i+1} onto X_i . Let X be the inverse limit space of the inverse system $(\{X_i\}, \{f_i\})$; in notation $X = \lim (\{X_i\}, \{f_i\})$, and let π_i be the projection mapping of X onto X_i . In [2] it is proved that if metrics d_i for X_i are properly chosen then the inverse limit space $X = \lim (\{X_i\}, \{f_i\})$ is locally connected if and only if the collection $\{(X_i, d_i) \mid i \text{ a positive integer}\}$ of metric spaces is equi-uniformly locally connected. Also the X_i were embedded in their cartesian product in such a way that X is locally connected if and only if the sequence X_1, X_2, X_3, \dots converges 0-regularly to X .

In this paper similar relations between semi-local connectedness, equi-uniform semi-local connectedness and 0-coregular convergence are established. These results are then combined with known results about 0-regular and 0-coregular convergence to obtain properties of certain inverse limit spaces. For example, if each X_i is a simple closed curve and X is semi-locally connected and cyclic (i.e., without cut points), then X is a simple closed curve. Similar theorems for 2-spheres and 2-cells are also obtained.

For definitions and results on semi-local connectedness see [5] or [6], equi-uniform local connectedness [2], inverse limits [1], 0-regular and 0-coregular convergence [3] and [4]. Throughout this paper $S_\varepsilon(p)$ will denote ε -neighborhood of p .

REMARK 1. Note the following relationship for the case where X is cyclic between semi-local connectedness and local connectedness:

If X is cyclic then it is $\left\{ \begin{array}{l} \text{locally connected} \\ \text{semi-locally connected} \end{array} \right\}$ at a point p if and only if for each $\varepsilon > 0$ there exists a $\delta > 0$ such that any two points $\left\{ \begin{array}{l} \text{inside } S_\delta(p) \\ \text{outside } S_\varepsilon(p) \end{array} \right\}$ can be joined by a connected set lying $\left\{ \begin{array}{l} \text{inside } S_\varepsilon(p) \\ \text{outside } S_\delta(p) \end{array} \right\}$. Note also that since X is compact it is locally connected (semi-locally connected) if and only if it is uniformly locally connected (uniformly semi-locally connected).

2. Equi-uniform semi-local connectedness.

DEFINITION 1. A collection $\{(Y_\alpha, \rho_\alpha) \mid \alpha \in A\}$ of metric spaces is *equi-uniformly locally connected* if and only if corresponding to each $\varepsilon > 0$

there is a $\delta > 0$ such that if $\alpha \in A$ and s and t are members of Y_α for which $\rho_\alpha(s, t) < \delta$ then s and t lie in a connected subset of Y_α of diameter less than ϵ .

DEFINITION 2. A collection $\{(Y_\alpha, \rho_\alpha) | \alpha \in A\}$ of metric spaces is *equi-uniformly semi-locally connected* if and only if corresponding to each $\epsilon > 0$ there is a $\delta > 0$ such that if $\alpha \in A$ then for any p in Y_α and s and t in $Y_\alpha - S_\delta(p)$ we have s and t lie in a connected subset of $Y_\alpha - S_\delta(p)$.

DEFINITION 3. Let d_i be a metric for X_i for each positive integer i . The sequence d_1, d_2, d_3, \dots is *admissible* if there exists a metric d for X such that

$$\lim_{i \rightarrow \infty} d_i(\pi_i(u), \pi_i(v)) = d(u, v)$$

uniformly on $X \times X$.

REMARK 2. Let D_i be a metric for X_i such that $D_i(x, y) \leq 1$ for all x and y in X_i . If $i > j$ let f_{ij} denote the composite mapping $f_j \cdots f_{i-2} f_{i-1}$. Define

$$d_i(x, y) = \sum_{j=1}^i 2^{-j} D_j(f_{ij}(x), f_{ij}(y))$$

for each positive integer i and all x and y in X_i . Also define

$$d(u, v) = \sum_{j=1}^{\infty} 2^{-j} D_j(\pi_j(u), \pi_j(v)) .$$

In ([7], Theorem 1) it is shown that $\lim_{i \rightarrow \infty} d_i(\pi_i(u), \pi_i(v)) = d(u, v)$ uniformly on $X \times X$ and hence the sequence d_1, d_2, \dots is admissible.

THEOREM 1. *Suppose d_i is a metric for X_i for each i and the sequence d_1, d_2, d_3, \dots is admissible. Then if X is cyclic it is semi-locally connected if and only if the collection $\{(X_i, d_i) | i \geq n\}$ is equi-uniformly semi-locally connected for some positive integer n .*

Proof. Suppose $\{(X_i, d_i) | i \geq n\}$ is equi-uniformly semi-locally connected. For any $\epsilon > 0$, there exists a $\delta > 0$ such that for any p in X_i and x and y in $X_i - S_{\epsilon/3}(p)$ we have x and y are contained in a connected subset of $X_i - S_\delta(p)$. Now suppose u and v are in $X - S_{\epsilon/3}(q)$ where $\pi_i(q) = p$. There exists a positive integer m such that if $i \geq m$ then $\pi_i(u)$ and $\pi_i(v)$ are in $X_i - S_{\epsilon/3}(p)$. For $i \geq \max(m, n)$ let K_i be a connected subset of $X_i - S_\delta(p)$ containing $\pi_i(u)$ and $\pi_i(v)$. Let H_i denote the closure of $\bigcup_{j \geq i} f_{ji}[K_j]$. Each H_i is a continuum, $f_{ji}[H_j] \subset H_i$ for $j > i$, so $H = \lim(\{H_i\}, \{f_i | H_{i+1}\})$ is a subcontinuum of $X - S_\delta(q)$ containing u and v . Hence since X is cyclic it is semi-locally connected.

Now assume X is semi-locally connected and cyclic. Suppose $\varepsilon > 0$. Then there exists a $\delta^* > 0$ such that for any p in X if u and v are in $X - S_{\varepsilon/2}(p)$ they lie in a connected subset of $X - S_{\delta^*/2}(p)$. There is a positive integer n such that if $i \geq n$ and u and v are in X then $|d_i(\pi_i(u), \pi_i(v)) - d(u, v)| < \min(\delta^*/6, \varepsilon/2)$. Now suppose $i \geq n$ and x and y are in X_i for which x and y are in $X_i - S_\varepsilon(p_i)$ where $\pi_i(p) = p_i$. There exists u and v in X such that $\pi_i(u) = x$ and $\pi_i(v) = y$. It follows u and v are in $X - S_{\varepsilon/2}(p)$ and hence u and v lie in a connected subset K of $X - S_{\delta^*/2}(p)$. Since $\text{diam}(\pi_i^{-1}(p_i)) < \delta^*/6$ we have $K \cap \pi_i^{-1}(p_i) = \phi$. Therefore $\pi_i(K)$ is a connected subset of X_i which contains x and y and is contained in $X_i - S_{\delta^*/3}(p_i)$. Let $\delta = \delta^*/3$. Then the collection $\{(X_i, d_i) | i \geq n\}$ is equi-uniformly semi-locally connected.

3. 0-coregular convergence. Let P be the cartesian product of the sequence X_1, X_2, \dots . Let D_i, d_i and d be metrics defined as in Remark 2. If we define a metric d^* for P by

$$d^*(a, b) = \sum_{i=1}^{\infty} 2^{-j} D_j(a_j, b_j) \quad \text{for } a = (a_1, a_2, \dots) \text{ and } b = (b_1, b_2, \dots),$$

then the inclusion map is an isometry of (X, d) into (P, d^*) . Choose a point $p = (p_1, p_2, \dots)$ in P , and define for each positive integer i an isometry h_i on (X_i, d_i) into (P, d^*) by letting

$$h_i(x) = (f_{i1}(x), \dots, f_{ii}(x), p_{i+1}, p_{i+2}, \dots)$$

where f_{ii} is the identity map. In the following denote $h_i[X_i]$ by X_i^* .

THEOREM 2. *The sequence X_1^*, X_2^*, \dots converges 0-coregularly to X if and only if X is semi-locally connected and cyclic.*

Proof. It is obvious that X_1^*, X_2^*, \dots converges to X . Suppose X is semi-locally connected and cyclic, then it follows from Theorem 1 that the collection $\{(X_i, d_i) | i \geq n\}$ of metric spaces is equi-uniformly semi-locally connected for some positive integer n . Since each h_i is an isometry, it follows at once that X_1^*, X_2^*, \dots must converge 0-coregularly to X .

Now suppose the sequence X_1^*, X_2^*, \dots converges 0-coregularly to X . Then by ([4], Theorem 2.1) X must be semi-locally connected and cyclic. White proved the following lemma in [4].

LEMMA 1. *If $A_i \rightarrow A$ 0-coregularly and if each A_i is a*

$$\left\{ \begin{array}{l} \text{simple closed curve} \\ \text{2-sphere and } A_i \rightarrow A \text{ 0-regularly} \\ \text{2-cell and } A_i \rightarrow A \text{ 0-regularly} \end{array} \right\}$$

then A is a $\left\{ \begin{array}{l} \text{simple closed curve or a point} \\ \text{2-sphere or a point} \\ \text{2-cell, 2-sphere or a point} \end{array} \right\}$.

THEOREM 3. *If X is semi-locally connected and cyclic and each X_i is a simple closed curve, then X is a simple closed curve.*

Proof. Since X is semi-locally connected and cyclic, by Theorem 2 $X_i^* \rightarrow X$ 0-coregularly where X_i^* is isometric to X_i . So by Lemma 1 X is a simple closed curve.

LEMMA 2. ([2], Theorem 3) *The sequence X_1^*, X_2^*, \dots converges 0-regularly to X if and only if X is locally connected.*

THEOREM 4. *If X is locally connected and cyclic and each X_i is a 2-sphere then X is a 2-sphere.*

Proof. For compact spaces local connectedness implies semi-local connectedness. So X is semi-locally connected and cyclic. Therefore by Theorem 2 $X_i^* \rightarrow X$ 0-coregularly and by Lemma 2 $X_i^* \rightarrow X$ 0-regularly where X_i^* is isometric to X_i . So by Lemma 1, X is a 2-sphere or a point and since π_i maps X onto X_i , we have X is a 2-sphere.

THEOREM 5. *If X is locally connected and cyclic and each X_i is a 2-cell then X is a 2-cell.*

Proof. Since X is locally connected and cyclic $X_i^* \rightarrow X$ 0-coregularly and by Lemma 2 $X_i^* \rightarrow X$ 0-regularly where X_i^* is isometric to X_i . So by Lemma 1 X is a 2-cell, 2-sphere or a point. The third possibility is ruled out since π_i maps X onto X_i . Since the second Čech homology group of each X_i is zero so is that of X . Hence the second possibility is ruled out also. So X is a 2-cell.

REMARK 3. To see the need for requiring X to be cyclic in Theorem 5 consider Example 1 of [2]. Each X_i is a 2-cell but X is an arc. So since X is locally connected $X_i^* \rightarrow X$ 0-regularly but not 0-coregularly since X has a cut point.

BIBLIOGRAPHY

1. C. E. Capel, *Inverse limit spaces*, Duke Math. J., **26** (1954), 233-245.
2. M. K. Fort and J. Segal, *Local connectedness of inverse limit spaces*, Duke Math. J., **28** (1961), 253-260.
3. P. A. White, *Regular convergence*, Bull. Amer. Math. Soc., **60** (1954), 431-443.
4. ———, *New types of regular convergence*, Duke Math. J., **12** (1945), 305-315.
5. G. T. Whyburn, *Semi-locally connected sets*, Amer. J. of Math., **61** (1939), 733-749.
6. ———, *Analytic Topology*, Amer. Math. Soc. Colloquium Publications, Vol. 28, 1942.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

M. OHTSUKA

H. L. ROYDEN

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Jonathan L. Alperin, <i>Groups with finitely many automorphisms</i>	1
Martin Arthur Arkowitz, <i>The generalized Whitehead product</i>	7
John D. Baum, <i>Instability and asymptoticity in topological dynamics</i>	25
William Aaron Beyer, <i>Hausdorff dimension of level sets of some Rademacher series</i>	35
Frank Herbert Brownell, III, <i>A note on Cook's wave-matrix theorem</i>	47
Gulbank D. Chakerian, <i>An inequality for closed space curves</i>	53
Inge Futtrup Christensen, <i>Some further extensions of a theorem of Marcinkiewicz</i>	59
Charles Vernon Coffman, <i>Linear differential equations on cones in Banach spaces</i>	69
Eckford Cohen, <i>Arithmetical notes. III. Certain equally distributed sets of integers</i>	77
John Irving Derr and Angus E. Taylor, <i>Operators of meromorphic type with multiple poles of the resolvent</i>	85
Jacob Feldman, <i>On measurability of stochastic processes in products space</i>	113
Robert S. Freeman, <i>Closed extensions of the Laplace operator determined by a general class of boundary conditions, for unbounded regions</i>	121
Robert E. Fullerton, <i>Geometric structure of absolute basis systems in a linear topological space</i>	137
Dieter Gaijer, <i>On conformal mapping of nearly circular regions</i>	149
Andrew Mattei Gleason and Hassler Whitney, <i>The extension of linear functionals defined on H^∞</i>	163
Seymour Goldberg, <i>Closed linear operators and associated continuous linear operators</i>	183
Basil Gordon, Aviezri Siegmund Fraenkel and Ernst Gabor Straus, <i>On the determination of sets by the sets of sums of a certain order</i>	187
Branko Grünbaum, <i>The dimension of intersections of convex sets</i>	197
Paul Daniel Hill, <i>On the number of pure subgroups</i>	203
Robert Peter Holten, <i>Generalized Goursat problem</i>	207
Alfred Horn, <i>Eigenvalues of sums of Hermitian matrices</i>	225
Henry C. Howard, <i>Oscillation and nonoscillation criteria for $y''(x) + f(y(x))p(x) = 0$</i>	243
Taqdir Husain, <i>S-spaces and the open mapping theorem</i>	253
Richard Eugene Isaac, <i>Markov processes and unique stationary probability measures</i>	273
John Rolfe Isbell, <i>Supercomplete spaces</i>	287
John Rolfe Isbell, <i>On finite-dimensional uniform spaces. II</i>	291
N. Jacobson, <i>A note on automorphisms of Lie algebras</i>	303
Antoni A. Kosinski, <i>A theorem on families of acyclic sets and its applications</i>	317
Marvin David Marcus and H. Minc, <i>The invariance of symmetric functions of singular values</i>	327
Ralph David McWilliams, <i>A note on weak sequential convergence</i>	333
John W. Milnor, <i>On axiomatic homology theory</i>	337
Victor Julius Mizel and Malempati Madhusudana Rao, <i>Nonsymmetric projections in Hilbert space</i>	343
Calvin Cooper Moore, <i>On the Frobenius reciprocity theorem for locally compact groups</i>	359
Donald J. Newman, <i>The Gibbs phenomenon for Hausdorff means</i>	367
Jack Segal, <i>Convergence of inverse systems</i>	371
Józef Siciak, <i>On function families with boundary</i>	375
Hyman Joseph Zimmerberg, <i>Two-point boundary conditions linear in a parameter</i>	385