

Pacific Journal of Mathematics

CONTINUOUSLY INVERTIBLE SPACES

P. H. DOYLE, III AND JOHN GILBERT HOCKING

CONTINUOUSLY INVERTIBLE SPACES

P. H. DOYLE AND J. G. HOCKING

In [4] we introduced the concept of an invertible space. (A topological space S is *invertible* if, for each open set U in S , there is an *invertting homeomorphism for U* , $h: S \rightarrow S$, of S onto itself such that $h(S - U)$ lies in U .) This paper is a continuation of the investigation of invertibility in which we concentrate upon that aspect contained in the following definition: A topological space S is *continuously invertible* if it is invertible and if, for each open set in S , there is an invertting homeomorphism that is isotopic to the identity mapping.

It is easy to see that the n -sphere S^n enjoys this property for, given an open set U in S^n , there is an invertting homeomorphism for U in each component of the space of homeomorphisms of S^n onto itself. It was this observation which suggested our present investigation. We acknowledge a debt of gratitude to the unknown referee of our paper [4] who brought to our attention the papers by Dancer [2], L. Whyburn [6] and Wilder [7]. While they were of little assistance, these papers do contain results allied to our own and hence provide us with a link to the past.

Let S be a continuously invertible space and let $\mathcal{G}(S)$ denote the group of homeomorphisms of S onto itself. (Note that each element of $\mathcal{G}(S)$ is an invertting homeomorphism for *some* open set, the identity mapping included.) Let $\mathcal{I}(S)$ be the subgroup of $\mathcal{G}(S)$ consisting of all homeomorphisms that are isotopic to the identity mapping. If x is a point of S , we let

$$O_x = \{y | y = g(x), \quad g \in \mathcal{G}(S)\}$$

denote the *total orbit* of x and we let

$$P_x = \{y | y = h(x), \quad h \in \mathcal{I}(S)\}$$

denote the *continuous orbit* of x .

By Theorem 8 of [4], each total orbit O_x is dense in S and an obvious modification of the same argument proves that each continuous orbit P_x is also dense in S . We note that each continuous orbit is connected. For if y is any point of the continuous orbit P_x , then the isotopy path of x during an isotopy carrying x onto y is a continuum. Therefore P_x is a union of continua having the point x in common.

THEOREM 1. *If S is a continuously invertible space, then every continuous orbit (and every total orbit) is connected and dense in S .*

Also, S itself is connected.

THEOREM 2. *If S is a continuously invertible T_1 space and if c is a cut point of S , then the continuous orbit P_c of c is S itself.*

Proof. Suppose there were a point x in $S - P_c$. Since $S - c = A \cup B$ set, we may assume that x lies in A . The set B is open under the hypotheses and hence there is an inverting homeomorphism h , isotopic to the identity, which carries x into B . But then the isotopy path of x must pass through the cut point c and hence the continuous orbits P_x and P_c intersect. This is impossible.

COROLLARY. *No continuously invertible Hausdorff continuum has a cut point.*

Proof. If such a continuum is degenerate, then it has no cut points. If it is nondegenerate, then it has at least two noncut points and the existence of a cut point would contradict Theorem 2.

THEOREM 3. *In a (nondegenerate) continuously invertible Hausdorff space, each continuous orbit is arcwise connected and each point in a total orbit lies on an arc in the orbit.*

Proof. The isotopy path of a point is a continuous image of the unit interval in a Hausdorff space and hence is a Peano continuum. Then by Theorem 10 of [4], each orbit is homogeneous.

COROLLARY. *No orbit in a (nondegenerate) continuously invertible Hausdorff space is degenerate.*

COROLLARY. *Every continuously invertible (nondegenerate) Hausdorff space is a union of nondegenerate, dense, disjoint, homogeneous, arcwise connected, continuously invertible subspaces.*

Proof. Each such space is the union of its continuous orbits.

If a continuously invertible Hausdorff space S contains no simple closed curve, then every Peano continuum in S is a dendrite and if S itself is a Peano continuum, then S is a dendrite. But by virtue of the corollary to Theorem 2, a dendrite is not continuously invertible.

THEOREM 4. *Each nondegenerate continuously invertible Peano continuum contains a simple closed curve.*

THEOREM 5. *If the invertible space S contains a separating proper subcontinuum C , then each open set in S contains a separating continuum imbedded in S as is C .*

Proof. This is an application of Theorem 6 of [4].

THEOREM 6. *If an invertible space S is separated by a proper closed set C which is irreducible with respect to separating S , then C contains no open set of S .*

Proof. This is an application of Theorem 5 above.

Next we have an extrinsic characterization of the n -sphere which may be compared with the intrinsic characterization given in [3].

THEOREM 7. *Let M be a set in E^{n+1} which is continuously invertible and which contains an n -sphere, S . Then M is the n -sphere S .*

Proof. Suppose that $M - S$ is not empty. There is no loss of generality in assuming that there are points of M in the bounded component A of $E^{n+1} - S = A \cup B$. By Theorem 5, there is an n -sphere S' in $A \cap M$ and, in particular, there is an isotopy H_t which carries S onto S' where $H_t(S)$ lies in M for each t , $0 \leq t \leq 1$.

Now M cannot contain an $(n + 1)$ -cell for if it were locally Euclidean of dimension $n + 1$ at any point, then Theorem 1 of [3] would imply that M is an $(n + 1)$ -sphere imbedded in E^{n+1} which is impossible. Thus there must be a point p lying in the annular region between S and S' such that p is not in M . Similarly, there is a point q in the unbounded domain B such that q is not in M . Clearly, the continuous cycles S and $p \cup q$ are linked, whence the isotopic cycle S' must be linked with $p \cup q$. This is contradictory.

THEOREM 8. *The only continuously invertible Peano continua in the plane are the simple closed curves.*

Proof. By Theorem 4, each such continuum contains a simple closed curve and then Theorem 7 applies.

We conclude this report with a few results on continuously invertible plane continua which serve only to indicate a direction in which further study may be fruitful.

THEOREM 9. *Let C be a continuously invertible plane continuum that is not a simple closed curve. If x and y are two points in the same continuous orbit in C , then there is a unique arc in C having*

x and y as endpoints.

Proof. By Theorem 3 there is at least one arc joining x and y . If there were another, then C would contain (and hence be) a simple closed curve.

THEOREM 10. *Let C be a continuously invertible plane continuum that is not a simple closed curve. Then every Peano continuum in C is a simple arc.*

Proof. From a previous remark and Theorem 7 we know that every Peano continuum in C is a dendrite. If there is a dendrite in C other than a simple arc, then there would be a simple triod T in C . Now $C - T$ is not empty and hence there is an isotopy carrying T into its complement. The isotopy path of T is a Peano continuum in C and hence is a dendrite. But then the isotopy path of the branch point b of T contains uncountably many branch points in the isotopy path of T . This is impossible because no dendrite contains uncountably many branch points.

THEOREM 11. *No proper subcontinuum of a continuously invertible plane continuum separates the plane.*

Proof. If some proper subcontinuum separates the plane, then by Theorem 5 there is a separating subcontinuum in every open set of the continuum. A construction as in the proof of Theorem 7 using Theorem 7, Chap. 1 of [5] will then yield a contradiction.

THEOREM 12. *Let C be a continuously invertible plane continuum that is not a simple closed curve. Then every proper subcontinuum of C is arcwise connected.*

Proof. Let C' be a proper subcontinuum of C and suppose there are points x, y in C' which lie on no arc in C' . Let U be an open set of C such that \bar{U} lies in $C - C'$. Then there is an isotopy of C with terminal mapping h such that $h(C') = C''$ lies in U . Letting $h(x) = x'$ and $h(y) = y'$, we note that there are arcs xx', yy' from C' to C'' and that these arcs must pass into $C - C'$, as we go from x to x' and y to y' . It follows that the continuum $D = C' \cup xx' \cup yy' \cup C''$ separates the plane. Thus by Theorem 11, we must have $D = C$. But this, too, is impossible since D obviously contains an open arc as an open set and hence Theorem 1 of [3] concludes that D is a simple closed curve, contrary to hypothesis.

THEOREM 13. *Let C be a continuously invertible plane continuum that is not a simple closed curve. Then every proper subcontinuum of C is an arc in some continuous orbit of C .*

Proof. Suppose there were a continuum C' in C which did not lie entirely in a continuous orbit. Let x and y be points of C' such that y is not in P_x . Then the arc from x to y given by Theorem 12 together with an arc in P_x having x as an interior point will contain a triod T . Then the same argument as in Theorem 10 yields a contradiction. The fact that a proper subcontinuum is an arc easily follows from Theorem 9.

THEOREM 14. *The only decomposable, continuously invertible plane continua are the simple closed curves.*

Proof. Suppose there is a decomposable continuously invertible plane continuum C that is not a simple closed curve. By definition, $C = C_1 \cup C_2$ where C_1 and C_2 are proper subcontinua. Then by Theorem 13, both C_1 and C_2 are simple arcs. Hence the open set $C_1 - C_2$ contains an open arc, whence Theorem 1 of [3] proves that C is a simple closed curve after all.

In [1], R. H. Bing has shown that a homogeneous plane continuum that contains an arc is necessarily a simple closed curve. Since a continuously invertible continuum contains an arc, we have the following result.

COROLLARY. *The only homogeneous, continuously invertible plane continua are the simple closed curves.*

The existence of an indecomposable, continuously invertible plane continuum remains an open question. In §8 of [1], Bing gives an example which may enjoy these properties but this has not been established.

REFERENCES

1. R. H. Bing, *A simple closed curve is the only homogeneous bounded plane continuum that contains an arc*, Can. J. Math., **12** (1960), 209-230.
2. W. Dancer, *Symmetrical cut sets*, Fund. Math., **27** (1936), 123-135.
3. P. H. Doyle, and J. G. Hocking, *A characterization of Euclidean n -space*, Mich. Math. J., **7** (1960), 199-200.
4. ———, *Invertible spaces*, Amer. Math. Monthly **68** (1961), 959-965.
5. S. Eilenberg, *Topologie du plan*, Fund. Math., **26** (1936), 61-112.
6. L. Whyburn, *Rotation groups about a set of fixed points*, Fund. Math., **28** (1937), 124-130.
7. R. L. Wilder, *The strong symmetrical cut set of closed Euclidean n -space*, Fund. Math., **27** (1936), 136-139.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

M. OHTSUKA

H. L. ROYDEN

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

CALIFORNIA RESEARCH CORPORATION

SPACE TECHNOLOGY LABORATORIES

NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insa'susha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

William George Bade and Robert S. Freeman, <i>Closed extensions of the Laplace operator determined by a general class of boundary conditions</i>	395
William Browder and Edwin Spanier, <i>H-spaces and duality</i>	411
Stewart S. Cairns, <i>On permutations induced by linear value functions</i>	415
Frank Sydney Cater, <i>On Hilbert space operators and operator roots of polynomials</i>	429
Stephen Urban Chase, <i>Torsion-free modules over $K[x, y]$</i>	437
Heron S. Collins, <i>Remarks on affine semigroups</i>	449
Peter Crawley, <i>Direct decompositions with finite dimensional factors</i>	457
Richard Brian Darst, <i>A continuity property for vector valued measurable functions</i>	469
R. P. Dilworth, <i>Abstract commutative ideal theory</i>	481
P. H. Doyle, III and John Gilbert Hocking, <i>Continuously invertible spaces</i>	499
Shaul Foguel, <i>Markov processes with stationary measure</i>	505
Andrew Mattei Gleason, <i>The abstract theorem of Cauchy-Weil</i>	511
Allan Brasted Gray, Jr., <i>Normal subgroups of monomial groups</i>	527
Melvin Henriksen and John Rolfe Isbell, <i>Lattice-ordered rings and function rings</i>	533
Amnon Jakimovski, <i>Tauberian constants for the $[J, f(x)]$ transformations</i>	567
Hubert Collins Kennedy, <i>Group membership in semigroups</i>	577
Eleanor Killam, <i>The spectrum and the radical in locally m-convex algebras</i>	581
Arthur H. Kruse, <i>Completion of mathematical systems</i>	589
Magnus Lindberg, <i>On two Tauberian remainder theorems</i>	607
Lionello A. Lombardi, <i>A general solution of Tonelli's problem of the calculus of variations</i>	617
Marvin David Marcus and Morris Newman, <i>The sum of the elements of the powers of a matrix</i>	627
Michael Bahir Maschler, <i>Derivatives of the harmonic measures in multiply-connected domains</i>	637
Deane Montgomery and Hans Samelson, <i>On the action of $SO(3)$ on S^n</i>	649
J. Barros-Neto, <i>Analytic composition kernels on Lie groups</i>	661
Mario Petrich, <i>Semicharacters of the Cartesian product of two semigroups</i>	679
John Sydney Pym, <i>Idempotent measures on semigroups</i>	685
K. Rogers and Ernst Gabor Straus, <i>A special class of matrices</i>	699
U. Shukla, <i>On the projective cover of a module and related results</i>	709
Don Harrell Tucker, <i>An existence theorem for a Goursat problem</i>	719
George Gustave Weill, <i>Reproducing kernels and orthogonal kernels for analytic differentials on Riemann surfaces</i>	729
George Gustave Weill, <i>Capacity differentials on open Riemann surfaces</i>	769
G. K. White, <i>Iterations of generalized Euler functions</i>	777
Adil Mohamed Yaqub, <i>On certain finite rings and ring-logics</i>	785