

Pacific Journal of Mathematics

**ON THE GREEN'S FUNCTION OF AN N -POINT BOUNDARY
VALUE PROBLEM**

PAUL RICHARD BEESACK

ON THE GREEN'S FUNCTION OF AN N -POINT BOUNDARY VALUE PROBLEM

PAUL R. BEESACK

1. Introduction. In a recent paper [3], D. V. V. Wend made use of the Green's functions $g_2(x, s)$, $g_3(x, s)$ for the boundary value problems

$$\begin{aligned} u'' &= 0; u(a_1) = u(a_2) = 0, & (a_1 < a_2), \\ u''' &= 0; u(a_1) = u(a_2) = u(a_3) = 0, & (a_1 < a_2 < a_3). \end{aligned}$$

In particular, he showed that if $a_1 \geq 0$, then

$$|g_2(x, s)| < a_2, \quad |g_3(x, s)| < a_3^2$$

for $a_1 \leq x, s \leq a_2$ or $a_1 \leq x, s \leq a_3$ respectively. He conjectured that if $g_n(x, s)$ is the Green's function for the boundary value problem

$$(1.1) \quad u^{(n)} = 0; u(a_1) = \dots = u(a_n) = 0, \quad (a_1 < a_2 < \dots < a_n),$$

then

$$|g_n(x, s)| < a_n^{n-1}, \quad a_1 \leq x, s \leq a_n,$$

(if $a_1 \geq 0$) and states in a footnote that this conjecture has been verified for $n < 6$. Assuming this conjecture valid he uses the inequality to obtain a lower bound for the m th positive zero of a solution of the differential equation

$$(1.2) \quad y^{(n)} + f(x)y = 0$$

where $f(x)$ is continuous and complex-valued on $0 \leq x < \infty$. In this proof, all zeros of the solution are *counted* as though they were *simple* zeros.

In this paper, we consider a more general boundary value problem allowing for multiple zeros of $y(x)$. Let $g_n(x, s)$ now denote the Green's function of the differential system

$$(1.3) \quad \begin{cases} y^{(n)} = 0, \\ y(a_i) = y'(a_i) = y''(a_i) = \dots = y^{(k_i)}(a_i) = 0. \end{cases} \quad 1 \leq i \leq r,$$

where $a_1 < a_2 < \dots < a_r$, $0 \leq k_i$, $k_1 + k_2 + \dots + k_r + r = n$. In §2, we shall prove that

$$(1.4) \quad |g_n(x, s)| \leq \frac{\prod_{i=1}^r |x - a_i|^{k_i+1}}{(n-1)!(a_r - a_1)} \leq \left(\frac{n-1}{n}\right)^{n-1} \frac{(a_r - a_1)^{n-1}}{n!}$$

for $a_1 < x, s < a_r$. In the case $r = n$, Wend's conjecture is thus verified,

Received September 28, 1961.

and improved. In §3, we apply this inequality to differential equations of the form (1.2), and more generally to nonlinear differential equations

$$(1.5) \quad y^{(n)} + f(x, y, y', \dots, y^{(n-1)}) = 0,$$

to obtain lower bounds for the m th zero of solutions. The inequality (1.4) also leads to an extension of an oscillation criterion of Liapounoff for the case $n = 2$.

2. The Green's function. If $g_n(x, s)$ denotes the Green's function of the system (1.3), then g_n satisfies—and in fact, is defined by—the three conditions:

1°. $g_n, g'_n, \dots, g_n^{(n-2)}$ are continuous functions of (x, s) on the square $a_1 \leq x, s \leq a_r$, while $g_n^{(n-1)}$ is a continuous function of (x, s) in each of the two triangles $a_1 \leq x \leq s \leq a_r$ and $a_1 \leq s \leq x \leq a_r$ with

$$g_n^{(n-1)}(s+, s) - g_n^{(n-1)}(s-, s) \equiv -1, \quad a_1 < s < a_r.$$

2°. $g_n^{(n)}(x, s) \equiv 0$ in each of the two triangles above.

3°. For each s , with $a_1 < s < a_r$, $g_n(x, s)$ satisfies the n boundary conditions of the system (1.3).

In the above statements (and throughout this paper), all derivatives are taken with respect to x . For a thorough discussion of Green's functions for much more general systems than (1.3), see Toyoda [2]. The existence of g_n depends on the fact that the system (1.3) is incompatible. We need not verify this directly since the result will follow from our method of proof which is by induction.

Although the conditions 1°–3° define g_n on the square $a_1 \leq x, s \leq a_r$, we want to extend the domain of definition of g_n to the entire plane. We assert that this can be done in such a way that

(I_n) $g_n, g'_n, \dots, g_n^{(n-2)}$ are continuous for all (x, s) , while $g_n^{(n-1)}$ is continuous in each of the half-planes $x \leq s$ and $s \leq x$, with $g_n^{(n-1)}(s+, s) - g_n^{(n-1)}(s-, s) \equiv -1, -\infty < s < \infty$.

(II_n) $g_n^{(n)}(x, s) \equiv 0$ in each of the above half-planes.

(III_n) For each $s, (-\infty < s < \infty)$, $g_n(x, s)$ satisfies the n boundary conditions of the system (1.3).

(IV_n) $g_n(x, s) \equiv 0$ if $s \leq \min(a_1, x)$, or $s \geq \max(a_r, x)$.

We proceed to prove these assertions by induction. Suppose they are valid for any system of the form (1.3). If a_j is any zero of a boundary value problem of this form for the equation $y^{(n+1)} = 0$, the corresponding set of boundary conditions is either of the form (1.3) with k_j replaced by $k_j + 1$ (in case a_j is not a simple zero for the new system), or is of the form

$$(2.1) \quad \begin{cases} y(a_i) = y'(a_i) = \dots = y^{(k_i)}(a_i) = 0, & 1 \leq i \leq r, i \neq j, \\ y(a_j) = 0, \end{cases}$$

where now $k_1 + \dots + k_{j-1} + k_{j+1} + \dots + k_r + r = n + 1$. Let $g_{n+1}(x, s)$ be the Green's function for this new system. We assert that

$$(2.2) \quad g_{n+1}(x, s) = \frac{1}{n} \left\{ (x - s)g_n(x, s) - \frac{(a_j - s)g_n^{(k_j+1)}(a_j, s)}{(k_j + 1)!} (x - a_j)^{k_j+1} \prod_{\substack{i=1 \\ i \neq j}}^r \left(\frac{x - a_i}{a_j - a_i} \right)^{k_i+1} \right\}$$

in the first case noted above, while

$$(2.3) \quad g_{n+1}(x, s) = \frac{1}{n} \left\{ (x - s)g_n(x, s) - (a_j - s)g_n(a_j, s) \prod_{\substack{i=1 \\ i \neq j}}^r \left(\frac{x - a_i}{a_j - a_i} \right)^{k_i+1} \right\}$$

in the second case. Note that (2.3) is formally included in (2.2) by setting $k_j = -1$ in (2.2). In the sequel we work with (2.2) only; (2.3) will follow by making use of this formal identity. We remark that g_n is defined by the conditions 1°-3° in $2(r - 1)$ "pieces", an explicit determination of any "piece" requiring the solution of n nonhomogeneous linear equations. For this reason, the recursion relations (2.2), (2.3) may be of some interest in themselves.

For brevity, set

$$P(x, s) = \frac{(a_j - s)g_n^{(k_j+1)}(a_j, s)}{(k_j + 1)!} (x - a_j)^{k_j+1} \prod_{i=1, i \neq j}^r \left(\frac{x - a_i}{a_j - a_i} \right)^{k_i+1}.$$

For each s , $P(x, s)$ is a polynomial of degree n in x . If $k_j < n - 2$ it follows from our induction assumptions that P , as well as all its derivatives with respect to x , is a continuous function of (x, s) in the entire plane. This also holds if $k_j = n - 2$ because of the factor $(a_j - s)$, provided we define $P(x, a_j) \equiv 0$. We also note that

$$(2.4) \quad \begin{aligned} P^{(m)}(a_i, s) &\equiv 0, & 0 \leq m \leq k_i, & i \neq j, \\ P^{(m)}(a_j, s) &\equiv 0, & 0 \leq m \leq k_j, \\ P^{(k_j+1)}(a_j, s) &\equiv (a_j - s)g_n^{(k_j+1)}(a_j, s). \end{aligned}$$

(In the case $k_j = -1$, the second of the identities (2.4) does not appear.) Differentiating (2.2) partially with respect to x , we obtain

$$\begin{aligned} g'_{n+1}(x, s) &= \frac{1}{n} \{ (x - s)g'_n(x, s) + g_n(x, s) - P'(x, s) \}, \\ g''_{n+1}(x, s) &= \frac{1}{n} \{ (x - s)g''_n(x, s) + 2g'_n(x, s) - P''(x, s) \}, \\ &\vdots \\ g^{(m)}_{n+1}(x, s) &= \frac{1}{n} \{ (x - s)g_n^{(m)}(x, s) + mg_n^{(m-1)}(x, s) - P^{(m)}(x, s) \}, \end{aligned}$$

$$1 \leq m \leq n + 1.$$

By our induction assumptions, together with the preceding remarks concerning $P(x, s)$, it follows that $g_{n+1}, g'_{n+1}, \dots, g_{n+1}^{(n-2)}$ are continuous in the entire plane. The same is true of $g_{n+1}^{(n-1)}$, because of the factor $(x - s)$. Finally, for $x \neq s$,

$$g_{n+1}^{(n)}(x, s) = \frac{1}{n} \{ (x - s)g_n^{(n)}(x, s) + ng_n^{(n-1)}(x, s) - P^{(n)}(x, s) \},$$

so that

$$g_{n+1}^{(n)}(s+, s) - g_{n+1}^{(n)}(s-, s) = \frac{1}{n} \cdot n \{ g_n^{(n-1)}(s+, s) - g_n^{(n-1)}(s-, s) \} \equiv -1,$$

and condition (I_{n+1}) is thus satisfied. Condition (II_{n+1}) is also satisfied since $g_n^{(n)} \equiv 0$ and $P^{(n+1)} \equiv 0$ in each of the two half-planes $x \leq s$ and $s \leq x$. For the boundary conditions we have

$$g_{n+1}^{(m)}(a_i, s) = \frac{1}{n} \{ (a_i - s)g_n^{(m)}(a_i, s) + mg_n^{(m-1)}(a_i, s) - P^{(m)}(a_i, s) \} \equiv 0$$

for $1 \leq i \leq r$ and $0 \leq m \leq k_i$, using the first two of (2.4). Using the last of (2.4), we also see that $g_{n+1}^{(k_j+1)}(a_j, s) \equiv 0$, and (III_{n+1}) is satisfied. Finally, suppose $s \leq \min(a_1, x)$ so that $g_n(x, s) \equiv 0$, and hence also $g_n^{(k_j+1)}(a_j, s) \equiv 0$ since $s \leq a_1 \leq a_j$. Thus, by (2.2), the first of conditions (IV_{n+1}) is satisfied, and similarly, so is the second.

For $n = 2$, we have explicitly

$$(2.5) \quad g_2(x, s) = \begin{cases} x - s, & x \leq s, & -\infty < s \leq a_1, \\ 0, & s \leq x, & \\ \frac{(x - a_1)(a_2 - s)}{a_2 - a_1}, & x \leq s, & a_1 \leq s \leq a_2, \\ \frac{(s - a_1)(a_2 - x)}{a_2 - a_1}, & s \leq x, & \\ 0, & x \leq s, & a_2 \leq s < \infty, \\ s - x, & s \leq x, & \end{cases}$$

from which one easily verifies that the conditions (I_2) – (IV_2) are satisfied, thus completing the induction for all $n \geq 2$.

Our goal now is to obtain an upper bound for $|g_n(x, s)|$. It will, however, be easier to work with the related function $G_n(x, s)$ defined by

$$(2.6) \quad \begin{cases} g_n(x, s) = G_n(x, s) \prod_{i=1}^r (x - a_i)^{k_i+1} \text{ for } x \neq a_i. \\ g_n^{(k_i+1)}(a_i, s) = (k_i + 1)! \prod_{m \neq i} (a_i - a_m)^{k_m+1} G_n(a_i, s) \text{ for } x = a_i. \end{cases}$$

We note that for each fixed $s \neq a_i$, $G_n(x, s)$ is continuous for all x . If

$k_i < n - 2$, $G_n(x, a_i)$ is also continuous for all x , while if $k_i = n - 2$, $G_n(x, a_i)$ has a finite jump at $x = a_i$ and is otherwise continuous. (The case $k_j = -1$ is again included in (2.6), the factor $(x - a_j)^{k_j+1}$ becoming unity in this case.) We now have

$$g_{n+1}(x, s) = G_{n+1}(x, s) (x - a_j) \prod_{i=1}^r (x - a_i)^{k_i+1}, \quad x \neq a_i.$$

Using (2.2) and (2.6), we also have

$$\begin{aligned} g_{n+1}(x, s) &= \frac{1}{n} \left\{ (x - s) \prod_{i=1}^r (x - a_i)^{k_i+1} G_n(x, s) - (a_j - s) \prod_{i=1}^r (x - a_i)^{k_i+1} G_n(a_j, s) \right\} \\ &= \frac{1}{n} \prod_{i=1}^r (x - a_i)^{k_i+1} \{ (x - s) G_n(x, s) - (a_j - s) G_n(a_j, s) \}, \end{aligned}$$

so that

$$(2.7) \quad (x - a_j) G_{n+1}(x, s) = \frac{1}{n} \{ (x - s) G_n(x, s) - (a_j - s) G_n(a_j, s) \}.$$

We note that for all $n \geq 2$, (IV _{n}) gives

$$(2.8) \quad G_n(x, s) \equiv 0 \text{ for } s \leq \min(a_1, x) \text{ and } s \geq \max(a_r, x).$$

We now prove by induction that

$$(2.9) \quad |G_n(x, s)| \leq \begin{cases} \frac{1}{(n-1)!} \frac{1}{a_r - x}, & -\infty < s < a_1, \quad x \leq s, \\ \frac{1}{(n-1)!} \frac{1}{a_r - a_1}, & a_1 < s < a_r, \quad -\infty < x < \infty, \\ \frac{1}{(n-1)!} \frac{1}{x - a_1}, & a_r < s < \infty, \quad s \leq x. \end{cases}$$

For $n = 2$, (2.5) gives

$$|G_2(x, s)| = \begin{cases} \frac{(s-x)}{(a_1-x)(a_2-x)}, & -\infty < s < a_1, \quad x \leq s, \\ \frac{(a_2-s)}{(a_2-x)(a_2-a_1)}, & x \leq s, \\ \frac{(s-a_1)}{(x-a_1)(a_2-a_1)}, & s \leq x, \quad a_1 < s < a_2, \\ \frac{(x-s)}{(x-a_1)(x-a_2)}, & a_2 < s < \infty, \quad s \leq x, \end{cases}$$

from which (2.9) is immediately verified for $n = 2$.

We will first prove (2.9) under the assumption that g_{n+1} has at

least three distinct zeros. Taking $j = 1$ in (2.2) and (2.7), we have for $-\infty < s < a_1$, $x \leq s$,

$$\begin{aligned} G_{n+1}(x, s) &= \frac{1}{n} \left\{ \frac{x-s}{x-a_1} G_n(x, s) - \frac{a_1-s}{x-a_1} G_n(a_1, s) \right\} \\ &= \frac{1}{n} \frac{x-s}{x-a_1} G_n(x, s), \end{aligned}$$

by (2.8); hence

$$|G_{n+1}(x, s)| = \frac{1}{n} \frac{s-x}{a_1-x} |G_n(x, s)| \leq \frac{1}{n!} \frac{1}{a_r-x}.$$

Similarly, taking $j = r$ in (2.2) and (2.7) we obtain for $a_r < s < \infty$, $s \leq x$,

$$|G_{n+1}(x, s)| = \frac{1}{n} \frac{x-s}{x-a_r} |G_n(x, s)| \leq \frac{1}{n!} \frac{1}{x-a_1}.$$

Note that the above work is valid whether a_1 or a_r are simple zeros of g_{n+1} or not. Also, the first inequality is valid even when $r = 2$ provided a_1 is *not* a simple zero of g_{n+1} , and similarly for the second inequality provided $a_r = a_2$ is not a simple zero of g_{n+1} .

In order to complete the induction on the middle inequality of (2.9), we suppose first that $a_2 < s < a_r$ and $s \leq x$. Taking $j = 2$ in (2.2) and (2.7), we obtain

$$\begin{aligned} |G_{n+1}(x, s)| &\leq \frac{1}{n} \left\{ \frac{x-s}{x-a_2} |G_n(x, s)| + \frac{s-a_2}{x-a_2} |G_n(a_2, s)| \right\} \\ &\leq \frac{1}{n!} \frac{1}{a_r-a_1}. \end{aligned}$$

If, however, $a_1 < s \leq a_2$ and $s \leq x$, we again take $j = 1$, whence

$$\begin{aligned} (2.10) \quad |G_{n+1}(x, s)| &\leq \frac{1}{n} \left\{ \frac{x-s}{x-a_1} |G_n(x, s)| + \frac{s-a_1}{x-a_1} |G_n(a_1, s)| \right\} \\ &\leq \frac{1}{n!} \frac{1}{a_r-a_1}, \end{aligned}$$

if a_1 is not a simple zero of g_{n+1} , or

$$|G_{n+1}(x, s)| = \frac{1}{n} \frac{s-a_1}{x-a_1} |G_n(a_1, s)| \leq \frac{1}{n!} \frac{1}{a_r-a_1}$$

if a_1 is a simple zero of g_{n+1} . (In this latter case, we used (2.8) and the first of inequalities (2.9).)

Similarly, if $a_1 < s < a_{r-1}$ and $x \leq s$, we take $j = r-1$ in (2.7) to obtain

$$\begin{aligned}
 |G_{n+1}(x, s)| &\leq \frac{1}{n} \left\{ \frac{s-x}{a_{r-1}-x} |G_n(x, s)| + \frac{a_{r-1}-s}{a_{r-1}-x} |G_n(a_{r-1}, s)| \right\} \\
 &\leq \frac{1}{n!} \frac{1}{a_r - a_1}.
 \end{aligned}$$

If $a_{r-1} \leq s < a_r$ and $x \leq s$, we again take $j = r$, whence

$$\begin{aligned}
 (2.11) \quad |G_{n+1}(x, s)| &\leq \frac{1}{n} \left\{ \frac{s-x}{a_r-x} |G_n(x, s)| + \frac{a_r-s}{a_r-x} |G_n(a_r, s)| \right\} \\
 &\leq \frac{1}{n!} \frac{1}{a_r - a_1}
 \end{aligned}$$

if a_r is not a simple zero of g_{n+1} , or

$$|G_{n+1}(x, s)| = \frac{1}{n} \frac{a_r - s}{a_r - x} |G_n(a_r, s)| \leq \frac{1}{n!} \frac{1}{a_r - a_1}$$

if a_r is a simple zero of g_{n+1} .

We now complete the induction in the case that g_{n+1} has only two distinct zeros. For $n \geq 2$, at least one of a_1, a_2 must be a multiple zero of g_{n+1} . Suppose a_2 is a multiple zero of g_{n+1} . Let $g_n(x, s)$ denote the Green's function for the boundary conditions

$$\begin{aligned}
 y(a_1) = y'(a_1) = \dots = y^{(k_1)}(a_1) = 0, \\
 y(a_2) = y'(a_2) = \dots = y^{(k_2)}(a_2) = 0,
 \end{aligned}$$

and $g_{n+1}(x, s)$ the Green's function for these boundary conditions with k_2 replaced by $k_2 + 1$. For any α with $a_1 < \alpha < a_2$, let $g_{n+1}(x, s; \alpha)$ denote the Green's function for the boundary conditions of g_n together with the condition $y(\alpha) = 0$. Let $G_n(x, s), G_{n+1}(x, s), G_{n+1}(x, s; \alpha)$ denote the related functions defined by (2.6). By (2.7)

$$\begin{aligned}
 G_{n+1}(x, s) &= \frac{1}{n} \left\{ \frac{x-s}{x-a_2} G_n(x, s) - \frac{a_2-s}{x-a_2} G_n(a_2, s) \right\}, \quad x \neq a_2, \\
 G_{n+1}(x, s; \alpha) &= \frac{1}{n} \left\{ \frac{x-s}{x-\alpha} G_n(x, s) - \frac{\alpha-s}{x-\alpha} G_n(\alpha, s) \right\}, \quad x \neq \alpha.
 \end{aligned}$$

For each $s \neq a_2$ and $x \neq a_2$, we have

$$\lim_{\alpha \rightarrow a_2} G_{n+1}(x, s; \alpha) = G_{n+1}(x, s)$$

since $G_n(\alpha, s)$ is a continuous function of α for any $s \neq a_2$. Since we have already established

$$|G_{n+1}(x, s; \alpha)| \leq \begin{cases} \frac{1}{n!} \frac{1}{a_2 - x}, & -\infty < s < a_1, x \leq s, \\ \frac{1}{n!} \frac{1}{a_2 - a_1}, & a_1 < s < a_2, -\infty < x < \infty, \\ \frac{1}{n!} \frac{1}{x - a_1}, & a_2 < s < \infty, s \leq x, \end{cases}$$

the same result holds for $|G_{n+1}(x, s)|$. The proof of (2.9) is now complete. From (2.6) and (2.9) it follows that, for each $n \geq 2$, we have

$$(2.12) \quad |g_n(x, s)| \leq \frac{\prod_{i=1}^r |x - a_i|^{k_i+1}}{(n-1)!(a_r - a_1)}, \quad a_1 < s < a_r, \quad -\infty < x < \infty.$$

We now prove that

$$(2.13) \quad \prod_{i=1}^r |x - a_i|^{k_i+1} \leq \left(\frac{n-1}{n}\right)^{n-1} \frac{(a_r - a_1)^n}{n} \quad \text{for } a_1 \leq x \leq a_r,$$

which will complete the proof of (1.4). Here, $r \geq 2, 0 \leq k_i, a_1 < a_2 < \dots < a_r$, and $k_1 + k_2 + \dots + k_r + r = n$. We note that equality is attained in (2.13) for $r = 2, k_1 = 0, k_2 = n - 2, x = [(n - 1)a_1 + a_2]/n$, or for $r = 2, k_1 = n - 2, k_2 = 0$, and $x = [(n - 1)a_2 + a_1]/n$.

Instead of proving (2.13) in the form stated, we will prove

$$(2.13_1) \quad \left| \prod_{i=1}^n (x - a_i) \right| \leq \left(\frac{n-1}{n}\right)^{n-1} \frac{(a_n - a_1)^n}{n} \quad \text{for } a_1 \leq x \leq a_n,$$

where $a_1 \leq a_2 \leq \dots \leq a_n$. As a first step, we prove

$$(2.14) \quad \left| \prod_{i=1}^n (x - a_i) \right| \leq \max \left\{ \begin{aligned} &(x - a_1)(a_n - x)^{n-1}, \\ &(x - a_1)^{n-1}(a_n - x), \end{aligned} \right. \quad a_1 \leq x \leq a_n.$$

To this end, suppose $a_j < x < a_{j+1}$. If $x - a_1 \geq a_n - x$, then

$$\left| \prod_{i=1}^n (x - a_i) \right| \leq (x - a_1)^j (a_n - x)^{n-j} \leq (x - a_1)^{n-1} (a_n - x);$$

if $x - a_1 \leq a_n - x$ (and $a_j < x < a_{j+1}$), then

$$\left| \prod_{i=1}^n (x - a_i) \right| \leq (x - a_1)^j (a_n - x)^{n-j} \leq (x - a_1) (a_n - x)^{n-1},$$

proving (2.14). Now, setting $f_1(x) = (x - a_1)(a_n - x)^{n-1}$, we see that $f_1(x)$ has an absolute maximum on $a_1 < x < a_n$ when $x = [(n - 1)a_1 + a_n]/n$. Similarly, $f_2(x) = (x - a_1)^{n-1}(a_n - x)$ has an absolute maximum on $a_1 < x < a_n$ when $x = [(n - 1)a_n + a_1]/n$. The inequalities (2.13₁) and (2.13) now follow by computation.

There remains the question as to whether the above inequalities are best possible. The inequality (2.9), or rather the restriction

$$(2.15) \quad |G_n(x, s)| \leq \frac{1}{(n-1)!(a_r - a_1)}, \quad a_1 < s < a_r, \quad -\infty < x < \infty,$$

is best possible. Indeed, equality holds in (2.15) for precisely the cases in which equality was attained in (2.13), that is, for $r = 2, k_1 = 0, k_2 = n - 1$ and for $r = 2, k_1 = n - 2, k_2 = 0$. For the first of these cases we shall show that

$$(2.16) \quad \lim_{x \rightarrow a_2^+} |G_n(x, a_2)| = \frac{1}{(n-1)!(a_2 - a_1)}$$

which will prove that (2.15) is best possible. Indeed, taking $j = 2$ and $s = a_2$ in (2.7), we have

$$G_{n+1}(x, a_2) = \frac{1}{n} G_n(x, a_2), \quad x \neq a_2,$$

so that (2.16) holds for $(n + 1)$ if it holds for n . One easily verifies that (2.16) is valid for $n = 2$. Similarly, for the second case noted above, we have

$$\lim_{x \rightarrow a_1^-} |G_n(x, a_1)| = \frac{1}{(n-1)!(a_2 - a_1)}.$$

It seems likely that equality is possible in (2.15) *only* in these two cases.

Nevertheless, the inequality (1.4) is *not the best possible, even in the simple case* $r = 2, k_1 = 0, k_2 = 1$, when (2.15) is best possible. We leave it to the reader to verify that in this case

$$|g_3(x, s)| \leq \frac{5\sqrt{5} - 11}{4} (a_2 - a_1)^2,$$

with equality holding for $s = 1/2 \{(3 - \sqrt{5})a_1 + (\sqrt{5} - 1)a_2\} = s_0$, and $x = (a_2 s_0 - a_1^2)/(a_2 + s_0 - 2a_1)$. This is an improvement over our estimate (1.4) which, for this case, is

$$|g_3(x, s)| \leq \frac{2}{27} (a_2 - a_1)^2.$$

3. Applications. Consider the ordinary differential equation

$$(3.1) \quad y^{(n)} + f(x, y, y', \dots, y^{(n-1)}) = 0,$$

where we assume that f is a continuous, complex-valued function for $a_1 \leq x \leq a_r$, and for all $y, y', \dots, y^{(n-1)}$, and

$$(3.2) \quad |f(x, y, \dots, y^{(n-1)})| \leq h(x) |y|$$

in this domain, where $h(x)$ is a nonnegative continuous function with $h(x) \neq 0$ on $a_1 \leq x \leq a_r$. Suppose (3.1) has a nontrivial solution $y(x)$ satisfying the boundary conditions

$$(3.3) \quad y(a_i) = y'(a_i) = \dots = y^{(k_i)}(a_i) = 0, \quad 1 \leq i \leq r,$$

where $a_1 < a_2 < \dots < a_r$, $0 \leq k_i$, $k_1 + k_2 + \dots + k_r + r = n$. Then $y(x)$ is a solution of the linear nonhomogeneous equation

$$y^{(n)} = -f[x, y(x), y'(x), \dots, y^{(n-1)}(x)]$$

which satisfies the linear homogeneous boundary conditions (3.3). By Theorem 1 of [2] it follows that $y(x)$ satisfies the integral equation

$$(3.4) \quad y(x) = \int_{a_1}^{a_r} g_n(x, s) f[s, y(s), \dots, y^{(n-1)}(s)] ds, \quad a_1 \leq x \leq a_r,$$

where $g_n(x, s)$ is the Green's function of the system (1.3). Taking x to be the point—or one of the points—at which $|y(x)|$ assumes its maximum value on $a_1 \leq x \leq a_r$, we obtain

$$(3.5) \quad 1 < \int_{a_1}^{a_r} |g_n(x, s)| h(s) ds,$$

by (3.2). Hence, using the inequality (1.4),

$$(3.6) \quad 1 < \left(\frac{n-1}{n} \right)^{n-1} \frac{(a_r - a_1)^{n-1}}{n!} \int_{a_1}^{a_r} h(s) ds.$$

The inequality (3.6) is thus a *necessary condition* for the existence of a solution of the boundary value problem (3.1), (3.3). If the system (1.3) is self-adjoint, we may improve this necessary condition. The system (1.3) is self-adjoint if $n = 2m$, $r = 2$, and the boundary conditions are

$$(3.7) \quad \begin{cases} y(a_1) = y'(a_1) = \dots = y^{(m-1)}(a_1) = 0, \\ y(a_2) = y'(a_2) = \dots = y^{(m-1)}(a_2) = 0. \end{cases}$$

The Green's function is now symmetric, and by (2.12) we have

$$|g_n(x, s)| = |g_n(s, x)| \leq \frac{(s - a_1)^m (a_2 - s)^m}{(2m - 1)! (a_2 - a_1)}, \quad a_1 < s < a_2.$$

On substituting this in (3.5), we obtain

$$(3.8) \quad 1 < \frac{1}{(2m - 1)! (a_2 - a_1)} \int_{a_1}^{a_2} (s - a_1)^m (a_2 - s)^m h(s) ds$$

as a necessary condition for the existence of a solution of the boundary value problem consisting of (3.1)—with $n = 2m$ —and (3.7).

We may adopt a different point of view and use (3.6) or (3.8) to obtain an extension of the following oscillation criterion due originally to Liapounoff (cf. [1]): *If $y''(x)$ and $y''(x)y^{-1}(x)$ are continuous for $a_1 \leq x \leq a_2$, with $y(a_1) = y(a_2) = 0$, then*

$$(3.9) \quad \int_{a_1}^{a_2} |y''y^{-1}| dx > \frac{4}{a_2 - a_1} .$$

By taking $f \equiv -y^{(n)}(x)y^{-1}(x)y$ in (3.1), $h(x) = |y^{(n)}(x)y^{-1}(x)|$ in (3.2), (3.6) leads to the following extension: *If $y^{(n)}(x)$ and $y^{(n)}(x)y^{-1}(x)$ are continuous for $a_1 \leq x \leq a_r$, and $y(x)$ has n zeros (counting multiplicity) including a_1 and a_r , on $a_1 \leq x \leq a_r$, then*

$$(3.10) \quad \int_{a_1}^{a_r} |y^{(n)}(x)y^{-1}(x)| dx > \left(\frac{n}{n-1}\right)^{n-1} \frac{n!}{(a_r - a_1)^{n-1}} .$$

This reduces to (3.9) when $n = 2$. Similarly, using (3.8) in the self-adjoint case: *If $y^{(2m)}(x)$ and $y^{(2m)}(x)y^{-1}(x)$ are continuous for $a_1 \leq x \leq a_2$, with $y^{(k)}(a_1) = y^{(k)}(a_2) = 0$ for $0 \leq k \leq m - 1$, then*

$$(3.11) \quad \int_{a_1}^{a_2} (x - a_1)^m (a_2 - x)^m |y^{(2m)}(x)y^{-1}(x)| dx > (2m - 1)! (a_2 - a_1) ,$$

$$(3.12) \quad \int_{a_1}^{a_2} |y^{(2m)}(x)y^{-1}(x)| dx > \frac{(2m - 1)! 2^{2m}}{(a_2 - a_1)^{2m-1}} .$$

The inequality (3.12) also reduces to (3.9) when $m = 1$, but is better than (3.10) for $n = 2m \geq 4$.

Next we turn to the question of obtaining a lower bound for the m th zero of solutions of the linear equation

$$(3.13) \quad y^{(n)} + h(x)y = 0$$

on an interval $I: x_0 \leq x < \infty$. cf. [3, Theorem 5]. *We suppose that $h(x)$ is continuous, complex-valued, with $h(x) \neq 0$ on I , and*

$$(3.14) \quad \int_{x_0}^{\infty} |h(x)| dx = K .$$

If $a_1 \leq a_2 \leq \dots \leq a_m$ are m consecutive zeros of any solution of (3.13) on the interval I , then for $m \geq n$

$$(3.15) \quad a_m > a_1 + \frac{n}{n-1} \sqrt[n-1]{\frac{(m-n+1) [(n-1)!]}{K}} .$$

To prove this, we first note that for the equation (3.13)—but not necessarily for (3.1)—no solution can have a zero of multiplicity greater than $(n - 1)$ at any point of I . Hence, if $a_i \leq a_{i+1} \leq \dots \leq a_{i+n-1}$ are n consecutive zeros of a solution of (3.13) on I then $a_i < a_{i+n-1}$, and (3.6) applies to give

$$(3.16) \quad \left(\frac{n}{n-1}\right)^{n-1} n! < (a_{i+n-1} - a_i)^{n-1} \int_{a_i}^{a_{i+n-1}} |h(x)| dx.$$

Suppose $m = qn + s$, where $q \geq 1$, $0 \leq s \leq n - 1$, so $a_m \geq a_{qn}$. Taking $i = 1, n + 1, \dots, (q - 1)n + 1$ in (3.16) and adding these inequalities gives

$$\left(\frac{n}{n-1}\right)^{n-1} qn! < \sum_{i=1}^q [a_{i_n} - a_{(i-1)n+1}]^{n-1} \int_{a_{(i-1)n+1}}^{a_{i_n}} |h(x)| dx,$$

whence, since $qn = m - s \geq m - n + 1$,

$$(3.17) \quad \left(\frac{n}{n-1}\right)^{n-1} (m - n + 1) [(n - 1)!] < (a_m - a_1)^{n-1} \int_{a_1}^{a_m} |h(x)| dx.$$

The inequality (3.15) follows at once from (3.17). As in [3], (3.17) can be used to obtain a lower bound for a_m even when $K = \infty$.

In case $m > 2n - 1$, these inequalities can be improved slightly, as follows. If $m = qn + s$, with $q \geq 1$, $0 \leq s \leq n - 1$, there exists precisely one integer $r \geq 1$ such that

$$rn - (r - 1) \leq m < (r + 1)n - r.$$

Now taking $i = 1, n, 2n - 1, \dots, (r - 1)n - (r - 2)$ in (3.16), and proceeding as above, we obtain

$$\left(\frac{n}{n-1}\right)^{n-1} rn! < [a_{rn-(r-1)} - a_1]^{n-1} \int_{a_1}^{a_{rn-(r-1)}} |h(x)| dx.$$

Since $r(n - 1) + n > m$ and $a_m \geq a_{rn-(r-1)}$, we have

$$\left(\frac{n}{n-1}\right)^{n-1} \frac{m - n}{n - 1} n! < (a_m - a_1)^{n-1} \int_{a_1}^{a_m} |h(x)| dx;$$

this yields the estimate

$$(3.18) \quad a_m > a_1 + \frac{n}{n-1} \sqrt[n-1]{\frac{(m-n)n!}{(n-1)K}}$$

which is a slight improvement on (3.15) for $m > 2n - 1$.

REFERENCES

1. G. Borg, *On a Liapounoff criterion of stability*, Amer. J. of Math., **71** (1949), 67-70.
2. K. Toyoda, *Green's function in space of one dimension*, Tôhoku Math. J., **38** (1933), 343-355.
3. D. V. V. Wend, *On the zeros of solutions of some linear complex differential equations*, Pacific J. Math., **10** (1960), 713-722.

CARLETON UNIVERSITY,
OTTAWA, CANADA,

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

T. M. CHERRY

D. DERRY

M. OHTSUKA

H. L. ROYDEN

E. SPANIER

E. G. STRAUS

F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF CALIFORNIA

MONTANA STATE UNIVERSITY

UNIVERSITY OF NEVADA

NEW MEXICO STATE UNIVERSITY

OREGON STATE UNIVERSITY

UNIVERSITY OF OREGON

OSAKA UNIVERSITY

UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY

UNIVERSITY OF TOKYO

UNIVERSITY OF UTAH

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF WASHINGTON

* * *

AMERICAN MATHEMATICAL SOCIETY

CALIFORNIA RESEARCH CORPORATION

SPACE TECHNOLOGY LABORATORIES

NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 12, No. 3

March, 1962

Alfred Aeppli, <i>Some exact sequences in cohomology theory for Kähler manifolds</i>	791
Paul Richard Beesack, <i>On the Green's function of an N-point boundary value problem</i>	801
James Robert Boen, <i>On p-automorphic p-groups</i>	813
James Robert Boen, Oscar S. Rothaus and John Griggs Thompson, <i>Further results on p-automorphic p-groups</i>	817
James Henry Bramble and Lawrence Edward Payne, <i>Bounds in the Neumann problem for second order uniformly elliptic operators</i>	823
Chen Chung Chang and H. Jerome (Howard) Keisler, <i>Applications of ultraproducts of pairs of cardinals to the theory of models</i>	835
Stephen Urban Chase, <i>On direct sums and products of modules</i>	847
Paul Civin, <i>Annihilators in the second conjugate algebra of a group algebra</i>	855
J. H. Curtiss, <i>Polynomial interpolation in points equidistributed on the unit circle</i>	863
Marion K. Fort, Jr., <i>Homogeneity of infinite products of manifolds with boundary</i>	879
James G. Glimm, <i>Families of induced representations</i>	885
Daniel E. Gorenstein, Reuben Sandler and William H. Mills, <i>On almost-commuting permutations</i>	913
Vincent C. Harris and M. V. Subba Rao, <i>Congruence properties of $\sigma_r(N)$</i>	925
Harry Hochstadt, <i>Fourier series with linearly dependent coefficients</i>	929
Kenneth Myron Hoffman and John Wermer, <i>A characterization of $C(X)$</i>	941
Robert Weldon Hunt, <i>The behavior of solutions of ordinary, self-adjoint differential equations of arbitrary even order</i>	945
Edward Takashi Kobayashi, <i>A remark on the Nijenhuis tensor</i>	963
David London, <i>On the zeros of the solutions of $w''(z) + p(z)w(z) = 0$</i>	979
Gerald R. Mac Lane and Frank Beall Ryan, <i>On the radial limits of Blaschke products</i>	993
T. M. MacRobert, <i>Evaluation of an E-function when three of its upper parameters differ by integral values</i>	999
Robert W. McKelvey, <i>The spectra of minimal self-adjoint extensions of a symmetric operator</i>	1003
Adegoke Olubummo, <i>Operators of finite rank in a reflexive Banach space</i>	1023
David Alexander Pope, <i>On the approximation of function spaces in the calculus of variations</i>	1029
Bernard W. Roos and Ward C. Sangren, <i>Three spectral theorems for a pair of singular first-order differential equations</i>	1047
Arthur Argyle Sagle, <i>Simple Malcev algebras over fields of characteristic zero</i>	1057
Leo Sario, <i>Meromorphic functions and conformal metrics on Riemann surfaces</i>	1079
Richard Gordon Swan, <i>Factorization of polynomials over finite fields</i>	1099
S. C. Tang, <i>Some theorems on the ratio of empirical distribution to the theoretical distribution</i>	1107
Robert Charles Thompson, <i>Normal matrices and the normal basis in abelian number fields</i>	1115
Howard Gregory Tucker, <i>Absolute continuity of infinitely divisible distributions</i>	1125
Elliot Carl Weinberg, <i>Completely distributed lattice-ordered groups</i>	1131
James Howard Wells, <i>A note on the primes in a Banach algebra of measures</i>	1139
Horace C. Wisner, <i>Decomposition and homogeneity of continua on a 2-manifold</i>	1145