

Pacific Journal of Mathematics

**SOME THEOREMS ON THE RATIO OF EMPIRICAL
DISTRIBUTION TO THE THEORETICAL DISTRIBUTION**

S. C. TANG

SOME THEOREMS ON THE RATIO OF EMPIRICAL DISTRIBUTION TO THE THEORETICAL DISTRIBUTION

S. C. TANG

1. Introduction. Let X_1, X_2, \dots, X_n be mutually independent random variables with the common cumulative distribution function $F(x)$. Let $X_1^*, X_2^*, \dots, X_n^*$ be the same set of variables rearranged in increasing order of magnitude. In statistical language X_1, X_2, \dots, X_n form a sample of n drawn from the distribution with distribution function $F(x)$. The empirical distribution of the sample X_1, \dots, X_n is the step function $F_n(x)$ defined by

$$(1) \quad F_n(x) = \begin{cases} 0 & \text{for } x \leq X_1^* \\ \frac{k}{n} & \text{for } X_k^* < x \leq X_{k+1}^* \\ 1 & \text{for } x > X_n^* . \end{cases}$$

A. Kolmogorov developed a well-known limit distribution law for the difference between the empirical distribution and the corresponding theoretical distribution, assuming $F(x)$ continuous:

$$\lim_{n \rightarrow \infty} P\left\{ \sqrt{n} \sup_{-\infty < x < \infty} |F_n(x) - F(x)| < z \right\} = \begin{cases} 0, & \text{for } z \leq 0, \\ \sum_{k=-\infty}^{\infty} (-1)^k e^{-2k^2z^2}, & \text{for } z > 0. \end{cases}$$

Equally interesting is Smirnov's theorem:

$$\lim_{n \rightarrow \infty} P\left\{ \sqrt{n} \sup_{-\infty < x < \infty} [F_n(x) - F(x)] < z \right\} = \begin{cases} 0, & \text{for } z \leq 0, \\ 1 - e^{-2z^2}, & \text{for } z > 0. \end{cases}$$

In this paper we shall study the ratio of the empirical distribution to the theoretical distribution, and evaluate the distribution functions of the upper and lower bounds of the ratio. We shall prove the following four theorems.

THEOREM 1. *If F is everywhere continuous then*

$$P\left\{ \sup_{0 \leq F(x) \leq 1} \frac{F_n(x)}{F(x)} < z \right\} = s(z) = \begin{cases} 0 & \text{for } z \leq 1 \\ 1 - \frac{1}{z} & \text{for } z > 1. \end{cases}$$

Received July 28, 1961. I am deeply indebted to the referee for his corrections of my English writing. Without his aid, the present paper would not be read clearly, although the responsibility of errors rests with me.

THEOREM 2. Let c be a positive number no larger than n and suppose that $F(x)$ is continuous in the range $0 \leq F(x) \leq c/n$. Then

$$P\left\{\sup_{0 < F(x) \leq \frac{c}{n}} \frac{F_n(x)}{F(x)} < z\right\} = \begin{cases} \sum_{r=0}^{\lfloor cz \rfloor} \binom{n}{r} \left(1 - \frac{c}{n}\right)^{n-r} \left(\frac{c}{n}\right)^r \\ \quad - \sum_{k=1}^{\lfloor cz \rfloor} \binom{n}{k} \frac{k^{k-1}(nz - k)^{n-k}}{(nz)^n} \\ \quad \cdot \sum_{r=k}^{\lfloor cz \rfloor} \binom{n-k}{r-k} \left(\frac{cz - k}{nz - k}\right)^{r-k} \left(1 - \frac{cz - k}{nz - k}\right)^{n-r} & \text{for } 0 < z \leq \frac{n}{c} \\ 0 & \text{for } z \leq 0 \\ 1 - \frac{1}{z} & \text{for } z > \frac{n}{c}. \end{cases}$$

THEOREM 3. Under the assumption of Theorem 1,

$$P\left\{\inf_{0 < F_n(x) \leq 1} \frac{F_n(x)}{F(x)} \leq z\right\} = I_n(z) = \begin{cases} 0 & \text{for } z < \frac{1}{n} \\ \left(1 - \frac{1}{nz}\right)^n \\ \quad + \sum_{k=1}^{\lfloor nz \rfloor} \binom{n}{k} \frac{(k-1)^{k-1}(nz - k)^{n-k}}{(nz)^n} & \text{for } \frac{1}{n} \leq z \leq 1 \\ 1 & \text{for } z > 1. \end{cases}$$

THEOREM 4. Under the assumptions of Theorem 2, if $1 \leq c \leq n$ then

$$P\left\{\inf_{0 < F_n(x) \leq \frac{c}{n}} \frac{F_n(x)}{F(x)} \leq z\right\} = \begin{cases} 0 & \text{for } z < \frac{1}{n} \\ \left(1 - \frac{1}{nz}\right)^n + \sum_{k=1}^{\lfloor nz \rfloor} \binom{n}{k} \frac{(k-1)^{k-1}(nz - k)^{n-k}}{(nz)^n} & \text{for } \frac{1}{n} \leq z \leq \frac{c}{n} \\ \left(1 - \frac{1}{nz}\right)^n + \sum_{k=1}^{\lfloor c \rfloor} \binom{n}{k} \frac{(k-1)^{k-1}(nz - k)^{n-k}}{(nz)^n} & \text{for } z > \frac{c}{n}. \end{cases}$$

2. An elementary lemma. We shall use the following lemma.

LEMMA. Let k be a positive integer and α an arbitrary real number. Then

$$(2) \quad \sum_{r=1}^k \frac{r^{r-1}}{r!} \frac{(\alpha - r)^{k-r}}{(k-r)!} = \frac{\alpha^{k-1}}{(k-1)!},$$

$$(3) \quad \frac{(\alpha - 1)^k}{k!} + \sum_{r=1}^k \frac{(r-1)^{r-1}}{r!} \frac{(\alpha - r)^{k-r}}{(k-r)!} = \frac{\alpha^k}{k!}.$$

If $\alpha = k$ (a special case) then

$$(4) \quad \sum_{r=1}^k \frac{r^{r-1}}{r!} \frac{(k-r)^{k-r}}{(k-r)!} = \frac{k^{k-1}}{(k-1)!} = \frac{k^k}{k!},$$

$$(5) \quad \frac{(k-1)^k}{k!} + \sum_{r=1}^k \frac{(r-1)^{r-1}}{r!} \frac{(k-r)^{k-r}}{(k-r)!} = \frac{k^k}{k!}.$$

Proof. Let $f(x)$ be a polynomial whose degree is less than k . Then

$$\sum_{r=0}^k \binom{k}{r} (-1)^r f(r) = (-1)^n [\mathcal{A}_n f(x)]_{x=0} = 0$$

i.e.
$$\sum_{r=1}^k \binom{k}{r} (-1)^r f(r) = -f(0).$$

Now we can directly obtain (2) and (3):

$$\begin{aligned} \sum_{r=1}^k \frac{r^{r-1}}{r!} \frac{(\alpha - r)^{k-r}}{(k-r)!} &= \sum_{r=1}^k \frac{r^{r-1}}{r!} \sum_{s=0}^{k-r} \frac{\alpha^s (-r)^{k-r-s}}{s! (k-r-s)!} \\ &= \sum_{s=0}^{k-1} \frac{(-1)^{k-s} \alpha^s}{s!} \sum_{r=1}^{k-s} \frac{(-1)^r r^{k-s-1}}{r! (k-r-s)!} \\ &= \frac{(-1)^{k-(k-1)} \alpha^{k-1}}{(k-1)!} (-1) = \frac{\alpha^{k-1}}{(k-1)!}. \\ \frac{(\alpha - 1)^k}{k!} + \sum_{r=1}^k \frac{(r-1)^{r-1}}{r!} \frac{(\alpha - r)^{k-r}}{(k-r)!} &= \frac{(\alpha - 1)^k}{k!} + \sum_{r=1}^k \frac{(r-1)^{r-1}}{r!} \sum_{s=0}^{k-r} \frac{(\alpha - 1)^s (-r + 1)^{k-r-s}}{s! (k-r-s)!} \\ &= \frac{(\alpha - 1)^k}{k!} + \sum_{s=0}^{k-1} \frac{(-1)^{k-s} (\alpha - 1)^s}{s!} \sum_{r=1}^{k-s} \frac{(-1)^r (r-1)^{k-s-1}}{r! (k-r-s)!} \\ &= \frac{(\alpha - 1)^k}{k!} + \sum_{s=0}^{k-1} \frac{(-1)^{k-s} (\alpha - 1)^s (-1)^{k-s}}{s! (k-s)!} \\ &= \sum_{s=0}^k \frac{(\alpha - 1)^s}{s! (k-s)!} = \frac{\alpha^k}{k!}. \end{aligned}$$

3. Proof of Theorem 1. First we consider the case $z \geq 1$. Let y_k be the largest root of the equation

$$F(x) = \frac{k}{nz} \quad (1 \leq k \leq n).$$

Since F is continuous, y_k is well defined. Now we evaluate probability of the inequality

$$(6) \quad \sup_{0 \leq F(x) \leq 1} \frac{F_n(x)}{F(x)} \geq z$$

that is, the probability that there exists an x such that

$$(7) \quad \frac{F_n(x)}{F(x)} \geq z.$$

If (7) is true, then, since $F(x)$ is nondecreasing and $\lim_{x \rightarrow \infty} (F_n(x)/F(x)) = 1$, there exists an x_0 such that

$$\frac{F_n(x_0)}{F(x_0)} = z.$$

By the definition of F_n , $F_n(x_0) = k/n$ for some k , so that $F(x_0) = k/nz$. Therefore we can take x_0 as one of y_k . In other words, for one value of y_k we have

$$F_n(y_k) = \frac{k}{n}$$

and the inequality

$$(8) \quad X_k^* < y_k \leq X_{k+1}^*$$

is true. Now let A_k be the event that this inequality holds.

Clearly (6) occurs if, and only if, at least one among the events

$$A_1, A_2, A_3, \dots, A_n$$

occurs. Generally the A_k are not mutually exclusive events so that the additive law is of no use. We may, with the help of an associated set of mutually exclusive events, deal with this situation. Put

$$u_k = \bar{A}_1 \bar{A}_2 \cdots \bar{A}_{k-1} A_k \quad (k = 1, 2, \dots, n).$$

where \bar{A} denotes the complementary event of A . We have

$$(9) \quad P\left\{\sup_{0 \leq F(x) \leq 1} \frac{F_n(x)}{F(x)} \geq z\right\} = P\left\{\sum_{k=1}^n A_k\right\} = P\left\{\sum_{k=1}^n u_k\right\} = \sum_{k=1}^n P(u_k).$$

If we employ the following conditional probability formula, we can evaluate $P(u_k)$.

$$(10) \quad P(A_k) = \sum_{r=1}^k P(u_r)P\{A_k | u_r\} = \sum_{r=1}^k P(u_r)P\{A_k | A_r\}.$$

Now A_k occurs if k of the n observations fall to the left of y_k , and $n - k$ to the right; hence

$$P(A_k) = \binom{n}{k} \left(\frac{k}{nz} \right)^k \left(1 - \frac{k}{nz} \right)^{n-k}.$$

And $P(A_k | A_r)$ is the probability that of $(n - r)$ observations, known to lie to the right of y_r , $(k - r)$ lie to the left of y_k and $n - k$ to the right. The probability of occurrence of an event, whose value is greater than or equal to y_r and is on the interval $y_r \leq x \leq y_k$ is

$$\frac{k/nz - r/nz}{1 - r/nz} = \frac{k - r}{nz - r}.$$

Therefore

$$P(A_k | A_r) = \binom{n-r}{k-r} \left(\frac{k-r}{nz-r} \right)^{k-r} \left(1 - \frac{k-r}{nz-r} \right)^{n-k}.$$

If we employ the notation

$$p_k(z) = \frac{(nz - n + k)^k}{k!},$$

we get

$$P(A_k) = \frac{p_k(1)p_{n-k}(z)}{p_n(z)}, \quad P(A_k | A_r) = \frac{p_{k-r}(1)p_{n-k}(z)}{p_{n-r}(z)}.$$

Equation (10) can be reduced to

$$\begin{aligned} \frac{p_k(1)p_{n-k}(z)}{p_n(z)} &= \sum_{r=1}^k P(u_r) \frac{p_{k-r}(1)p_{n-k}(z)}{p_{n-r}(z)}, \\ p_k(1) &= \sum_{r=1}^k P(u_r) \frac{p_n(z)}{p_{n-r}(z)} p_{k-r}(1). \end{aligned}$$

Hence $P_k(1) = k^k/k!$ and $P_{k-r}(1) = (k - r)^{k-r}/(k - r)!$. By (3) of our lemma we know that

$$P(u_r) \frac{p_n(z)}{p_{n-r}(z)} = \frac{r^{r-1}}{r!}, \quad P(u_r) = \frac{r^{r-1}}{r!} \frac{p_{n-r}(z)}{p_n(z)} = \frac{n!}{(nz)^n} \frac{r^{r-1}}{r!} \frac{(nz - r)^{n-r}}{(n - r)!}.$$

And (9) and the (2) of our lemma tell us that

$$\begin{aligned} P\left\{\sup_{0 < F(x) \leq 1} \frac{F_n(x)}{F(x)} < z\right\} &= 1 - P\left\{\sup_{0 \leq F(x) \leq 1} \frac{F_n(x)}{F(x)} \geq z\right\} = 1 - \sum_{k=1}^n P(u_k) \\ &= 1 - \frac{n!}{(nz)^n} \sum_{k=1}^n \frac{k^{k-1}}{k!} \frac{(nz - k)^{n-k}}{(n - k)!} \\ &= 1 - \frac{n!}{(nz)^n} \frac{(nz)^{n-1}}{(n - 1)!} = 1 - \frac{1}{z}, \quad (z \geq 1). \end{aligned}$$

This completes the proof of Theorem 1.

Proof of Theorem 2. Since the ratio is nonnegative, the probability of the inequality is zero if $z \leq 0$. Under the condition $z > n/c$, we know that the event $\{\sup_{0 < F(x) \leq c/n} F_n(x)/F(x) \geq z\}$ is still equal to $\sum_{k=1}^n A_k$ by the result of Theorem 1. Suppose $0 < z \leq n/c$. Then

$$P\left\{\sup_{0 < F(x) \leq \frac{c}{n}} \frac{F_n(x)}{F(x)} \geq z\right\} = P\left\{\sum_{k=1}^{[cz]} A_k + A_{cz}^*\right\} = \sum_{k=1}^{[cz]} P(u_k) + P(u_{cz}^*) ,$$

where

$$\begin{aligned} A_{cz}^* &= \{X_{[cz]+1}^* < y_{cz}\} , \quad y_{cz} = \sup \left\{x : F(x) = \frac{c}{n}\right\} , \\ u_{cz}^* &= \bar{A}_1 \bar{A}_2 \cdots \bar{A}_{[cz]} A_{cz}^* . \end{aligned}$$

Since $P(U_k)$ has been computed, we need only evaluate $P(U_{cz}^*)$. We have

$$\begin{aligned} P(A_{cz}^*) &= \sum_{k=1}^{[cz]} P(u_k) P\{A_{cz}^* | A_k\} + P(u_{cz}^*) P\{A_{cz}^* | A_{cz}^*\} , \\ P(u_{cz}^*) &= P(A_{cz}^*) - \sum_{k=1}^{[cz]} P\{A_{cz}^* | A_k\} . \end{aligned}$$

From this we obtain

$$\begin{aligned} P\left\{\sup_{0 \leq F(x) \leq \frac{c}{n}} \frac{F_n(x)}{F(x)} < z\right\} &= 1 - \sum_{k=1}^{[cz]} P(u_k) - P(u_{cz}^*) \\ &= 1 - P(A_{cz}^*) - \sum_{k=1}^{[cz]} P(u_k)[1 - P\{A_{cz}^* | A_k\}] \\ &= P(\bar{A}_{cz}^*) - \sum_{k=1}^{[cz]} P(u_k) P\{\bar{A}_{cz}^* | A_k\} \\ &= \sum_{r=0}^{[cz]} \binom{n}{r} \left(\frac{c}{n}\right)^r \left(1 - \frac{c}{n}\right)^{n-r} - \sum_{k=1}^{[cz]} \binom{n}{k} \frac{k^{k-1}(nz-k)^{n-k}}{(nz)^n} \\ &\quad \cdot \sum_{r=k}^{[cz]} \binom{n-k}{r-k} \left(\frac{cz-k}{nz-k}\right)^{r-k} \left(1 - \frac{cz-k}{nz-k}\right)^{n-r} . \end{aligned}$$

Since

$$\begin{aligned} P(\bar{A}_{cz}^*) &= P\{y_{cz} \leq X_{[cz]+1}^*\} = \sum_{r=0}^{[cz]} P\{X_r^* < y_{cz} \leq X_{r+1}^*\} \\ &= \sum_{r=0}^{[cz]} \binom{n}{r} \left(\frac{c}{n}\right)^r \left(1 - \frac{c}{n}\right)^{n-r} \end{aligned}$$

we have

$$P(\bar{A}_{cz}^* | A_k) = \sum_{r=k}^{[cz]} \binom{n-k}{r-k} \left(\frac{cz-k}{nz-k}\right)^{r-k} \left(1 - \frac{cz-k}{nz-k}\right)^{n-r}$$

which completes the proof. The distribution of Theorem 1 is continuous; the distribution of Theorem 2 is not continuous on the interval $0 \leq z \leq n/c$.

Theorem 3 is a special case of Theorem 4 under the condition $c = n$. In fact, by (4) of our lemma, setting $z = 1$, we deduce Theorem 3 as follows:

$$\begin{aligned} P\left\{\inf_{0 < F_n(x) \leq 1} \frac{F_n(x)}{F(x)} \leq 1\right\} &= \left(1 - \frac{1}{n}\right)^n + \sum_{k=1}^n \binom{n}{k} \frac{(k-1)^{k-1}(n-k)^{n-k}}{n^n} \\ &= \frac{n!}{n^n} \left[\frac{(n-1)^n}{n!} + \sum_{k=1}^n \frac{(k-1)^{k-1}}{k!} \frac{(n-k)^{n-k}}{(n-k)!} \right] \\ &= \frac{n!}{n^n} \frac{n^n}{n!} = 1. \end{aligned}$$

Thus we need only prove Theorem 4. The distribution function of Theorem 3 has a discontinuity corresponding to $z = 1$; the distribution function of Theorem 4 is continuous on the interval $1 \leq c \leq n$.

5. Proof of Theorem 4. On the internal $F_n(x) > 0$ the maximum of $F(x)$ is 1; the minimum of $F_n(x)$ is $1/n$. From these results it follows that the lower bound of the ratio is no smaller than $1/n$. Therefore

$$P\left\{\inf_{0 < F_n(x) \leq \frac{c}{n}} \frac{F_n(x)}{F(x)} \leq z\right\} = 0$$

if $z < 1/n$. Let z_k be the least root of the equation

$$F(x) = \frac{k}{nz} \quad \left(z \geq \frac{1}{n}, \quad 1 \leq k \leq [nz]\right).$$

Define events

$$\begin{aligned} B_0: \quad &X_1 \geq z_1, \\ B_k: \quad &X_k^* < z_k \leq X_{k+1}^*. \end{aligned}$$

If $1/n \leq z \leq c/n$, the event

$$\inf_{0 < F_n(x) \leq \frac{c}{n}} \frac{F_n(x)}{F(x)} \leq z$$

is the union of the events

$$B_0, B_1, B_2, \dots, B_{[nz]}.$$

For the purpose of evaluating the probability of $\sum_k B_k$ we put

$$V_0 = B_0, \quad V_k = \bar{B}_0 \bar{B}^1 \cdots \bar{B}_{k-1} B_k \quad (k = 1, 2, \dots, [nz]).$$

We have

$$(11) \quad P(B_k) = \sum_{r=0}^k P(V_r)P(B_k | B_r), \quad (k = 0, 1, \dots, [nz]).$$

Here

$$P(V_0) = P(B_0) = \left(1 - \frac{1}{nz}\right)^n,$$

$$P\{B_k | B_0\} = \binom{n}{k} \left(\frac{k-1}{nz-1}\right)^k \left(\frac{nz-k}{nz-1}\right)^{n-k}.$$

Now we transform (11) into the following form:

$$\frac{p_k(1)p_{n-k}(z)}{p_n(z)} = \frac{(k-1)^k p_{n-k}(z)}{k! p_n(z)} + \sum_{r=1}^k P(V_r) \frac{p_{k-r}(1)p_{n-k}(z)}{p_{n-r}(z)},$$

$$p_k(1) = \frac{(k-1)^k}{k!} + \sum_{r=1}^k P(V_r) \frac{p_n(z)}{p_{n-r}(z)} p_{k-r}(1).$$

By (4) of our lemma we obtain

$$P(V_r) = \frac{(r-1)^{r-1}}{r!} \frac{p_{n-r}(z)}{p_n(z)} = \binom{n}{r} \frac{(r-1)^{r-1}(nz-r)^{n-r}}{(nz)^n}, \quad (1 \leq r \leq [nz]).$$

It follows that if $1/n \leq z \leq c/n$,

$$P\left\{\inf_{0 < F_n(x) \leq \frac{c}{n}} \frac{F_n(x)}{F(x)} \leq z\right\} = P\left\{\sum_{k=0}^{[nz]} B_k\right\} = \sum_{k=0}^{[nz]} P(V_k)$$

$$= \left(1 - \frac{1}{nz}\right)^n + \sum_{k=1}^{[nz]} \binom{n}{k} \frac{(k-1)^{k-1}(nz-k)^{n-k}}{(nz)^n}.$$

If $z > c/n$,

$$P\left\{\inf_{0 < F_n(x) \leq \frac{c}{n}} \frac{F_n(x)}{F(x)} \leq z\right\} = P\left\{\sum_{k=0}^{[c]} B_k\right\} = \sum_{k=0}^{[c]} P(V_k)$$

$$= \left(1 - \frac{1}{nz}\right)^n + \sum_{k=1}^{[c]} \binom{n}{k} \frac{(k-1)^{k-1}(nz-k)^{n-k}}{(nz)^n}.$$

Theorem 4 is thus proved.

6. Conclusions. Theorems 1 and 3 can be applied to test whether statistical data correspond with the theoretical distribution or not. In the process of testing, z_s and z_i are separately representative of the ratio's upper and lower bounds. If $S(z_s)$ is small and $I_n(z_i)$ large, we conclude that the empirical data in two extreme tails correspond with the theoretical distribution; conversely, if $S(z_s)$ is very large and $I_n(z_i)$ very small, the statistical data do not correspond with the theoretical distribution. Our test is more sensitive in the two tails, but less sensitive in the central part, than Smirnov's test.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

RALPH S. PHILLIPS

Stanford University
Stanford, California

M. G. ARSOVE

University of Washington
Seattle 5, Washington

A. L. WHITEMAN

University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE

University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH
T. M. CHERRY

D. DERRY
M. OHTSUKA

H. L. ROYDEN
E. SPANIER

E. G. STRAUS
F. WOLF

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA

STANFORD UNIVERSITY

CALIFORNIA INSTITUTE OF TECHNOLOGY

UNIVERSITY OF TOKYO

UNIVERSITY OF CALIFORNIA

UNIVERSITY OF UTAH

MONTANA STATE UNIVERSITY

WASHINGTON STATE UNIVERSITY

UNIVERSITY OF NEVADA

UNIVERSITY OF WASHINGTON

NEW MEXICO STATE UNIVERSITY

* * *

OREGON STATE UNIVERSITY

AMERICAN MATHEMATICAL SOCIETY

UNIVERSITY OF OREGON

CALIFORNIA RESEARCH CORPORATION

OSAKA UNIVERSITY

SPACE TECHNOLOGY LABORATORIES

UNIVERSITY OF SOUTHERN CALIFORNIA

NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and the author should keep a complete copy. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunkenshuppan (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 12, No. 3

March, 1962

Alfred Aeppli, <i>Some exact sequences in cohomology theory for Kähler manifolds</i>	791
Paul Richard Beesack, <i>On the Green's function of an N-point boundary value problem</i>	801
James Robert Boen, <i>On p-automorphic p-groups</i>	813
James Robert Boen, Oscar S. Rothaus and John Griggs Thompson, <i>Further results on p-automorphic p-groups</i>	817
James Henry Bramble and Lawrence Edward Payne, <i>Bounds in the Neumann problem for second order uniformly elliptic operators</i>	823
Chen Chung Chang and H. Jerome (Howard) Keisler, <i>Applications of ultraproducts of pairs of cardinals to the theory of models</i>	835
Stephen Urban Chase, <i>On direct sums and products of modules</i>	847
Paul Civin, <i>Annihilators in the second conjugate algebra of a group algebra</i>	855
J. H. Curtiss, <i>Polynomial interpolation in points equidistributed on the unit circle</i>	863
Marion K. Fort, Jr., <i>Homogeneity of infinite products of manifolds with boundary</i>	879
James G. Glimm, <i>Families of induced representations</i>	885
Daniel E. Gorenstein, Reuben Sandler and William H. Mills, <i>On almost-commuting permutations</i>	913
Vincent C. Harris and M. V. Subba Rao, <i>Congruence properties of $\sigma_r(N)$</i>	925
Harry Hochstadt, <i>Fourier series with linearly dependent coefficients</i>	929
Kenneth Myron Hoffman and John Wermer, <i>A characterization of $C(X)$</i>	941
Robert Weldon Hunt, <i>The behavior of solutions of ordinary, self-adjoint differential equations of arbitrary even order</i>	945
Edward Takashi Kobayashi, <i>A remark on the Nijenhuis tensor</i>	963
David London, <i>On the zeros of the solutions of $w''(z) + p(z)w(z) = 0$</i>	979
Gerald R. Mac Lane and Frank Beall Ryan, <i>On the radial limits of Blaschke products</i>	993
T. M. MacRobert, <i>Evaluation of an E-function when three of its upper parameters differ by integral values</i>	999
Robert W. McKelvey, <i>The spectra of minimal self-adjoint extensions of a symmetric operator</i>	1003
Adegoke Olubummo, <i>Operators of finite rank in a reflexive Banach space</i>	1023
David Alexander Pope, <i>On the approximation of function spaces in the calculus of variations</i>	1029
Bernard W. Roos and Ward C. Sangren, <i>Three spectral theorems for a pair of singular first-order differential equations</i>	1047
Arthur Argyle Sagle, <i>Simple Malcev algebras over fields of characteristic zero</i>	1057
Leo Sario, <i>Meromorphic functions and conformal metrics on Riemann surfaces</i>	1079
Richard Gordon Swan, <i>Factorization of polynomials over finite fields</i>	1099
S. C. Tang, <i>Some theorems on the ratio of empirical distribution to the theoretical distribution</i>	1107
Robert Charles Thompson, <i>Normal matrices and the normal basis in abelian number fields</i>	1115
Howard Gregory Tucker, <i>Absolute continuity of infinitely divisible distributions</i>	1125
Elliot Carl Weinberg, <i>Completely distributed lattice-ordered groups</i>	1131
James Howard Wells, <i>A note on the primes in a Banach algebra of measures</i>	1139
Horace C. Wiser, <i>Decomposition and homogeneity of continua on a 2-manifold</i>	1145