

Pacific Journal of Mathematics

SOME APPLICATIONS OF MEANS OF CONVEX BODIES

WILLIAM JAMES FIREY

SOME APPLICATIONS OF MEANS OF CONVEX BODIES

WILLIAM J. FIREY

Let A be a real, positive definite, $n \times n$ matrix; with A we associate, in the Euclidean n -space R_n , the ellipsoid $E(A)$ of points x for which

$$(x, Ax) \leq 1$$

where (x, y) denotes the usual inner product. In references [5], [6], [7] certain means of convex bodies were studied. It will be shown here that two particular means of ellipsoids of the type $E(A)$ correspond to two simple combinations of the corresponding matrices A . The applications mentioned in the title rest upon this correspondence. The first two give results about positive definite matrices, including a refinement of a determinant inequality of Minkowski; the third application shows the existence of a set of unique ellipsoids related to a convex body by a set of similar extremal problems, the classical Loewner ellipsoid being a particular instance.

Throughout this paper the letters A and B , sometimes with distinguishing marks, denote real, positive definite, $n \times n$ matrices. The distance from x to the origin is written $\|x\|$.

1. The distance and support functions of $E(A)$ are:

$$F(x) = \sqrt{(x, Ax)}, \quad H(x) = \sqrt{(x, A^{-1}x)}.$$

In the first case, if $x \neq 0$, we have $F(x) = \|x\|/\|z\|$ where $x/\|x\| = z/\|z\|$ and $(z, Az) = 1$, and so

$$\begin{aligned} \|x\|/\|z\| &= \|x\| \sqrt{(z/\|z\|, Az/\|z\|)} \\ &= \|x\| \sqrt{(x/\|x\|, Ax/\|x\|)} = \sqrt{(x, Ax)}. \end{aligned}$$

In the second case

$$H(x) = \max_y (x, y) \quad \text{where} \quad (y, Ay) = 1.$$

We represent y in the form $\lambda A^{-1}x + v$ where $(x, v) = 0$. Then

$$(y, Ay) = \lambda^2(x, A^{-1}x) + (v, Av),$$

whence

Received April 4, 1963. This work was supported in part by a grant from the National Science Foundation, NSF-G19838. The author is grateful to the referee for a constructive comment on the last application.

$$(x, y) = \lambda(x, A^{-1}x) = \sqrt{(x, A^{-1}x)}\sqrt{1 - (v, Av)},$$

and the maximum is attained for $v = 0$.

The polar reciprocal $\hat{E}(A)$ of $E(A)$ with respect to the unit sphere $E(I)$ has $H(x)$ as its distance function, $F(x)$ as its support function. Consequently

$$\hat{E}(A) = E(A^{-1}).$$

In [5] the p -dot mean of two convex bodies K_0, K_1 in R_n , which have the origin as a common interior point, was defined for $p \geq 1$ to be the convex body $\dot{M}_p(K_0, K_1; \vartheta)$ whose distance function is

$$[(1 - \vartheta)F_0^p(x) + \vartheta F_1^p(x)]^{1/p}$$

where F_i is the distance function of K_i and $0 \leq \vartheta \leq 1$. From this it follows that $\dot{M}_2(E(A_0), E(A_1); \vartheta)$ has the distance function

$$\sqrt{[(1 - \vartheta)(x, A_0x) + \vartheta(x, A_1x)]} = \sqrt{(x, [(1 - \vartheta)A_0 + \vartheta A_1]x)}.$$

Thus

$$(2) \quad \dot{M}_2(E(A_0), E(A_1); \vartheta) = E((1 - \vartheta)A_0 + \vartheta A_1).$$

In [7] the p -mean $M_p(K_0, K_1; \vartheta)$ was defined for $p \geq 1$ to be the convex body whose support function is

$$[(1 - \vartheta)H_0^p(x) + \vartheta H_1^p(x)]^{1/p}$$

where H_i is the support function of K_i . Therefore, by reasoning similar to the preceding, we have

$$(3) \quad M_2(E(A_0), E(A_1); \vartheta) = E([(1 - \vartheta)A_0^{-1} + \vartheta A_1^{-1}]^{-1}).$$

2. Our first application is based on the inclusion

$$(4) \quad \dot{M}_2(K_0, K_1; \vartheta) \subseteq M_2(K_0, K_1; \vartheta),$$

established in [5] and [7]¹ with equality if and only if $K_0 = K_1$, and the observation that

$$E(A) \subseteq E(B)$$

if and only if $A - B$ is positive semi-definite. For the latter we write $A \geq B$; we call such an inequality strict if $A - B$ is not a zero matrix. From (2), (3) and (4) we have

$$(5) \quad E((1 - \vartheta)A_0 + \vartheta A_1) \subseteq E([(1 - \vartheta)A_0^{-1} + \vartheta A_1^{-1}]^{-1}).$$

¹ The inclusion is not specifically mentioned, but in [7] it is proved that $M_1 \subseteq M_p$ for $p > 1$ and in [5] that $\dot{M}_p \subseteq \dot{M}_1$ and $\dot{M}_1 \subseteq M_1$.

Hence, from (5) we obtain an "inequality of arithmetic and harmonic means" for positive definite matrices.

THEOREM 1. *If A_0, A_1 are any two real, positive definite, $n \times n$ matrices, then*

$$(1 - \vartheta)A_0 + \vartheta A_1 \geq [(1 - \vartheta)A_0^{-1} + \vartheta A_1^{-1}]^{-1}$$

for $0 \leq \vartheta \leq 1$. The inequality is strict except in the trivial cases $A_0 = A_1$ or $\vartheta = 0, 1$.

3. The next application is a refinement of the following determinant inequality of Minkowski, cf [1], p. 70.

$$\det^{1/n}(A_0 + A_1) \geq \det^{1/n} A_0 + \det^{1/n} A_1.$$

Let V be the volume functional. In [5] it was shown that

$$(6) \quad V(\dot{M}_p(K_0, K_1; \vartheta)) \leq [(1 - \vartheta)V^{-p/n}(K_0) + \vartheta V^{-p/n}(K_1)]^{-n/p}$$

with equality if and only if $K_0 = \lambda K_1$ for some $\lambda > 0$. Since

$$V(E(A)) = \pi^{n/2} \Gamma(1 + n/2) \sqrt{\det A},$$

we have, with $p = 2$ in (6),

$$(7) \quad \det [(1 - \vartheta)A_0 + \vartheta A_1] \geq [(1 - \vartheta) \det^{1/n} A_0 + \vartheta \det^{1/n} A_1]^n$$

with equality if and only if $A_0 = \lambda A_1$ for some $\lambda > 0$. With a slight change in notation, this is Minkowski's determinant inequality.

If L is any k -dimensional linear subspace of R_n , then

$$\dot{M}_2(E(A_0) \cap L, E(A_1) \cap L; \vartheta) = \dot{M}_2(E(A_0), E(A_1); \vartheta) \cap L.$$

Consequently, by letting A' be the $k \times k$, positive definite matrix associated with $E(A) \cap L$, we obtain

$$E((1 - \vartheta)A'_0 + \vartheta A'_1) = E([(1 - \vartheta)A_0 + \vartheta A_1]') .$$

To this we apply (7), with $n = k$, to get

$$(8) \quad \det [(1 - \vartheta)A'_0 + \vartheta A'_1] \geq [(1 - \vartheta) \det^{1/k} A'_0 + \vartheta \det^{1/k} A'_1]^k .$$

Let us define $|A|_k$ to be the product of the k least eigenvalues of A , repeated eigenvalues being counted according to their multiplicity. The inequality

$$\det A' \geq |A|_k$$

with equality if and only if L is the k -dimensional space spanned by the eigenvectors corresponding to the k least eigenvalues of A , is

essentially Theorem 20, p. 74 of [1].

In (8) choose L to be the linear subspace spanned by those eigenvectors of $(1 - \vartheta)A_0 + \vartheta A_1$, which correspond to the k smallest eigenvalues of $(1 - \vartheta)A_0 + \vartheta A_1$. By (9):

$$(10) \quad \det A'_0 \geq |A_0|_k, \quad \det A'_1 \geq |A_1|_k,$$

and so (8) becomes

$$(11) \quad |(1 - \vartheta)A_0 + \vartheta A_1|_k \geq [(1 - \vartheta)|A_0|_k^{1/k} + \vartheta|A_1|_k^{1/k}]^k.$$

There is equality in (8) if and only if, for some $\lambda > 0$,

$$A'_0 = \lambda A'_1$$

and equality in (10) if and only if the subspaces L appropriate to $|A_0|_k$, $|A_1|_k$ are the same. Hence, in (11), there is equality if and only if the following conditions are met. Let x_1, \dots, x_k be eigenvectors of A_0 corresponding to the k smallest eigenvalues $\lambda_1 \leq \dots \leq \lambda_k$. These are eigenvectors of A_1 corresponding to the k smallest eigenvalues of A_1 which are of the form $\lambda\lambda_1 \leq \dots \leq \lambda\lambda_k$ for some $\lambda > 0$.

Inequality (11), which includes (7) when $k = n$, is an improvement of a result of Ky Fan, cf. [1], Theorem 21, p. 74, in which the right side of (11) is replaced by the geometric mean $|A_0|_k^{1-\vartheta}|A_1|_k^\vartheta$ since the power mean of order $1/k$ appearing on the right side of (11) exceeds this geometric mean.

If we define $|A|_k$ to be the product of the k greatest eigenvalues of A , then

$$(12) \quad |A^{-1}|_k = 1/|A|_k.$$

We apply (11) to $(1 - \vartheta)A_0^{-1} + \vartheta A_1^{-1}$ and obtain, after taking reciprocals,

$$1/|(1 - \vartheta)A_0^{-1} + \vartheta A_1^{-1}|_k \leq [(1 - \vartheta)|A_0|_k^{-1/k} + \vartheta|A_1|_k^{-1/k}]^{-k}.$$

With the use of (12) on the left side, we have finally

$$k|[(1 - \vartheta)A_0^{-1} + \vartheta A_1^{-1}]^{-1}| \leq [(1 - \vartheta)|A_0|_k^{-1/k} + \vartheta|A_1|_k^{-1/k}]^{-k}$$

as a "dual" result to (11). The cases of equality are given by the conditions for equality in (11) with the word "smallest" replaced by "greatest" throughout.

The last application concerns a generalization of the Loewner ellipsoid of a convex body K . Let x be an interior point of K . The classical Loewner ellipsoid is that *unique* ellipsoid, centred at x and containing K , which has minimum volume, cf. [3]. Let us take the point x to be the origin and denote the mean cross-sectional measures W_ν , $\nu = 0, 1, \dots, n-1$, of $E(A)$ by $W_\nu(A)$; for their definition see

[2]. In particular $W_0(A) = V(E(A))$. We will show that, for each ν there is a unique ellipsoid $E(A)$ containing K for which $W_\nu(A)$ is a minimum.

It is clear that $W_\nu(A)$ depends continuously on the entries a_{ij} of A . Moreover, when we restrict the ellipsoids $E(A)$ not only to contain K , but also to be contained in the sphere $E(I/\rho^2)$, the domain of definition of the functions $W_\nu(A)$ is closed and bounded. Consequently each of the functions $W_\nu(A)$ attains a minimum. Furthermore, if the radius of the bounding sphere $E(I/\rho^2)$ is chosen to be sufficiently large, the minimum of $W_\nu(A)$ and the matrix or matrices for which it is attained will be independent of ρ . Thus the uniqueness is the only point in question.

In [6] inequality (6) was extended to read

$$(13) \quad W_\nu^{1/(n-\nu)}[\dot{M}_p(K_0, K_1; \vartheta)] \leq [(1 - \vartheta) W_\nu^{-p/(n-\nu)}(K_0) + \vartheta W_\nu^{-p/(n-\nu)}(K_1)]^{-1/p}$$

for $p = 1$, with equality if and only if $K_0 = \lambda K_1$ for some $\lambda > 0$. Inequality (13) is true for all $p \geq 1$ however. This can be shown from the special case $p = 1$ in the following fashion. We make the usual type of reduction to the special case in which $W_\nu(K_i) = 1$, $i = 0, 1$, by setting:

$$\begin{aligned} \lambda_i &= W_\nu^{1/(n-\nu)}(K_i), & K_i &= \lambda_i K'_i, \\ \vartheta' &= \vartheta \lambda_1^{-p} / [(1 - \vartheta) \lambda_0^{-p} + \vartheta \lambda_1^{-p}]. \end{aligned}$$

Then

$$\dot{M}_p(K'_0, K'_1; \vartheta') = \dot{M}_p(K_0, K_1; \vartheta) / \mu$$

where

$$\mu = [(1 - \vartheta) \lambda_0^{-p} + \vartheta \lambda_1^{-p}]^{-1/p}.$$

Since $W_\nu(K'_i) = 1$, in order to prove (13) it is enough to prove

$$W_\nu(\dot{M}_p(K'_0, K'_1; \vartheta')) \leq 1.$$

This has been shown to be true for $p = 1$. By Theorem 2 of [5]

$$\dot{M}_p(K'_0, K'_1; \vartheta') \subseteq \dot{M}_1(K'_0, K'_1; \vartheta')$$

with equality if and only if $K'_0 = K'_1$. These assertions, together with the monotonic character of W_ν , cf. [2], p. 50, prove (13) and establish the cases of equality. Naturally we will use (13) for $p = 2$.

Let A_ν be a matrix which is a solution of the minimum problem:

$$K \subseteq E(A), \quad W_\nu(A) = \text{minimum.}$$

Suppose A'_ν is a second solution. From

$$K \subseteq E(A_\nu), \quad K \subseteq E(A'_\nu)$$

we have

$$K \subseteq E((1 - \vartheta)A_\nu + \vartheta A'_\nu);$$

from (13) we have

$$W_\nu((1 - \vartheta)A_\nu + \vartheta A'_\nu) \leq W_\nu(A_\nu) = W_\nu(A'_\nu)$$

with equality in the inequality if and only if $A_\nu = \lambda A'_\nu$. The last equality shows that we must have $\lambda = 1$ and so A_ν is unique.

In a similar way we can establish that, given K and an interior point of K which we take as the origin, there is a unique ellipsoid $E(B_\nu)$ which is contained in K for which is a maximum. The only difference is the use of Theorem 2 of [7] in lieu of inequality (13).

We summarize:

Theorem 2. *Given a convex body K in Euclidean n -space and an interior point of K which we take as the origin, there are positive definite $n \times n$ matrices $A_\nu, B_\nu, \nu = 0, 1, \dots, n - 1$ such that, among the ellipsoids $E(A)$ which contain K , $E(A_\nu)$ is the unique, outer, Loewner ellipsoid minimizing W_ν , and among the ellipsoids $E(B)$ which are contained in K , $E(B_\nu)$ is the unique inner, Loewner ellipsoid maximizing W_ν .*

We close with several observations. Suppose \hat{K} is the polar reciprocal of K with respect to $E(I)$, then, in the notation of Theorem 2, $E(B_\nu^{-1})$ is the ν th outer Loewner ellipsoid of \hat{K} while $E(A_\nu^{-1})$ is the ν th inner Loewner ellipsoid. To prove this, we denote the outer and inner Loewner ellipsoids of \hat{K} with respect to the origin by $E(\hat{A}_\nu)$, $E(\hat{B}_\nu)$ respectively. If $K_0 \subseteq K_1$, then $\hat{K}_0 \supseteq \hat{K}_1$. Consequently, by (1),

$$\hat{E}(A_\nu) = E(A_\nu^{-1}) \subseteq \hat{K}, \quad \hat{E}(B_\nu) = E(B_\nu^{-1}) \supseteq \hat{K}.$$

Therefore

$$E(A_\nu^{-1}) \subseteq E(\hat{B}_\nu), \quad E(B_\nu^{-1}) \supseteq E(\hat{A}_\nu).$$

Applying the same argument to \hat{A}_ν and \hat{B}_ν , we get

$$E(\hat{A}_\nu^{-1}) \subseteq E(B_\nu), \quad E(\hat{B}_\nu^{-1}) \supseteq E(A_\nu).$$

In terms of the ordering of positive definite matrices, these inclusions become

$$(14) \quad A_\nu^{-1} \geq \hat{B}_\nu, \quad \hat{A}_\nu \geq B_\nu^{-1}, \quad \hat{A}_\nu^{-1} \geq B_\nu, \quad A_\nu \geq \hat{B}_\nu^{-1}.$$

Now when $B \supseteq A$, then $A^{-1} \supseteq B^{-1}$ since, from the first condition we have

$$E(A) \supseteq E(B)$$

and, by taking polar reciprocals, we obtain

$$E(A^{-1}) \subseteq E(B^{-1}).$$

Apply this to the last inequality of (14). Taken together with the first inequality of (14), this yields

$$A^{-1} \supseteq \hat{B}_\nu \supseteq A_\nu^{-1}.$$

Thus $\hat{B}_\nu - A_\nu^{-1}$ is both positive and negative semi-definite. Hence

$$A_\nu^{-1} = \hat{B}_\nu.$$

By a similar argument it is shown that

$$B_\nu^{-1} = \hat{A}_\nu.$$

Part of Theorem 2 remains true even if the centre of the ellipsoids to be considered does not lie within K . We give this as a corollary.

COROLLARY TO THEOREM 2. *Given a convex body K , not necessarily containing the origin, there are positive definite matrices A_ν , $\nu = 0, 1, \dots, n-1$, such that, among the ellipsoids $E(A)$ which contain K , $E(A_\nu)$ is the unique outer Loewner ellipsoid minimizing W_ν .*

Suppose $E(A)$ contains K ; since $E(A)$ is centred at the origin it also contains a sufficiently small sphere $E(\rho I)$ and so, by the convexity of $E(A)$, $E(A)$ contains

$$K' = \overline{K \cup E(\rho I)}$$

where the bar denotes the convex closure. Conversely, if $E(A)$ contains K' it contains the subset K . We claim as proof of the corollary that the outer Loewner ellipsoid $E(A_\nu)$ of K' is also that of K . Indeed $E(A_\nu)$ contains K and if an ellipsoid $E(A'_\nu)$ contains K and is such that

$$W_\nu(A'_\nu) \leq W_\nu(A_\nu)$$

then $E(A'_\nu)$ must contain K' and so, by Theorem 2, $A'_\nu = A_\nu$.

Let x be the interior point mentioned in Theorem 2 and let $E(A_\nu(x))$, $E(B_\nu(x))$ be the ν th outer and inner Loewner ellipsoids of K which are centred at x . We allow x to vary and so generate two collections of ellipsoids $\{E(A_\nu(x))\}$ and $\{E(B_\nu(x))\}$. For $\nu = 0$ Danzer, Laugwitz and Lenz in [4] have shown that in the first collection there is a unique

ellipsoid for which the volume W_0 is a minimum and in the second collection there is a unique ellipsoid for which the volume is a maximum. We have not been able to decide if this is also true for $\nu = 1, 2, \dots, n - 1$ with W_ν in place of the volume.

REFERENCES

1. E. Beckenbach and R. Bellman, *Inequalities*, Berlin, Göttingen, Heidelberg, 1961.
2. T. Bonnesen and W. Fenchel, *Theorie der konvexen Körper*, Berlin, 1934
3. H. Busemann, *The Geometry of Geodesics*, New York, 1955.
4. L. Danzer, D. Laugwitz and H. Lenz, *Über das Löwnersche Ellipsoid und sein Analogon unter den Eikörper einbeschriebenen Ellipsoiden*, Archiv der Mathematik, **8** (1957), 214-218.
5. W. Firey, *Polar means of convex bodies and a dual to the Brunn-Minkowski theorem*, Canadian J. Math., **13** (1961) 444-453.
6. ———, *Mean cross-section measures of harmonic means of convex bodies*, Pacific J. Math., **11** (1961), 1263-1266.
7. ———, *p-means of convex bodies*, Mathematica Scandinavica, **10** (1962), 17-24.

OREGON STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

ROBERT OSSERMAN
Stanford University
Stanford, California

M. G. ARSOVE
University of Washington
Seattle 5, Washington

J. DUGUNDJI
University of Southern California
Los Angeles 7, California

LOWELL J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Richard Arens, <i>Normal form for a Pfaffian</i>	1
Charles Vernon Coffman, <i>Non-linear differential equations on cones in Banach spaces</i>	9
Ralph DeMarr, <i>Order convergence in linear topological spaces</i>	17
Peter Larkin Duren, <i>On the spectrum of a Toeplitz operator</i>	21
Robert E. Edwards, <i>Endomorphisms of function-spaces which leave stable all translation-invariant manifolds</i>	31
Erik Maurice Ellentuck, <i>Infinite products of isolos</i>	49
William James Firey, <i>Some applications of means of convex bodies</i>	53
Haim Gaifman, <i>Concerning measures on Boolean algebras</i>	61
Richard Carl Gilbert, <i>Extremal spectral functions of a symmetric operator</i>	75
Ronald Lewis Graham, <i>On finite sums of reciprocals of distinct nth powers</i>	85
Hwa Suk Hahn, <i>On the relative growth of differences of partition functions</i>	93
Isidore Isaac Hirschman, Jr., <i>Extreme eigen values of Toeplitz forms associated with Jacobi polynomials</i>	107
Chen-jung Hsu, <i>Remarks on certain almost product spaces</i>	163
George Seth Innis, Jr., <i>Some reproducing kernels for the unit disk</i>	177
Ronald Jacobowitz, <i>Multiplicativity of the local Hilbert symbol</i>	187
Paul Joseph Kelly, <i>On some mappings related to graphs</i>	191
William A. Kirk, <i>On curvature of a metric space at a point</i>	195
G. J. Kurowski, <i>On the convergence of semi-discrete analytic functions</i>	199
Richard George Laatsch, <i>Extensions of subadditive functions</i>	209
V. Marić, <i>On some properties of solutions of $\Delta\psi + A(r^2)X\nabla\psi + C(r^2)\psi = 0$</i> ...	217
William H. Mills, <i>Polynomials with minimal value sets</i>	225
George James Minty, Jr., <i>On the monotonicity of the gradient of a convex function</i>	243
George James Minty, Jr., <i>On the solvability of nonlinear functional equations of 'monotonic' type</i>	249
J. B. Muskat, <i>On the solvability of $x^e \equiv e \pmod{p}$</i>	257
Zeev Nehari, <i>On an inequality of P. R. Bessack</i>	261
Raymond Moos Redheffer and Ernst Gabor Straus, <i>Degenerate elliptic equations</i>	265
Abraham Robinson, <i>On generalized limits and linear functionals</i>	269
Bernard W. Roos, <i>On a class of singular second order differential equations with a non linear parameter</i>	285
Tôru Saitô, <i>Ordered completely regular semigroups</i>	295
Edward Silverman, <i>A problem of least area</i>	309
Robert C. Sine, <i>Spectral decomposition of a class of operators</i>	333
Jonathan Dean Swift, <i>Chains and graphs of Ostrom planes</i>	353
John Griggs Thompson, <i>2-signalizers of finite groups</i>	363
Harold Widom, <i>On the spectrum of a Toeplitz operator</i>	365