

# Pacific Journal of Mathematics

**POLYNOMIALS WITH MINIMAL VALUE SETS**

WILLIAM H. MILLS

# POLYNOMIALS WITH MINIMAL VALUE SETS

W. H. MILLS

Let  $\mathcal{K}$  be a finite field of characteristic  $p$  that contains exactly  $q$  elements. Let  $F(x)$  be a polynomial over  $\mathcal{K}$  of degree  $f, f > 0$ , and let  $r + 1$  denote the number of distinct values  $F(\tau)$  as  $\tau$  ranges over  $\mathcal{K}$ . Carlitz, Lewis, Mills, and Straus [1] pointed out that  $r \geq [(q - 1)/f]$ , and raised the question of determining all polynomials for which  $r = [(q - 1)/f]$ . The cases  $r = 0$  and  $r = 1$  are special cases that do not fit into the general pattern. These are treated in [1], and do not concern us here. Thus we arrive at the statement of our main problem: For what polynomials  $F(x)$  do we have

$$(I) \quad r = [(q - 1)/f] \geq 2?$$

Carlitz, Lewis, Mills, and Straus [1] determined all polynomials with  $f < 2p + 2$  for which (I) holds. In the present paper this result is extended—all polynomials with  $f \leq \sqrt{q}$  for which (I) holds are determined. These are polynomials of the form

$$F(x) = \alpha L^v + \gamma,$$

where  $L$  is a polynomial that factors into distinct linear factors over  $\mathcal{K}$  and that has the form

$$L = \beta + \sum_i \rho_i x^{p^{ki}},$$

and where  $v$  and  $k$  are integers such that  $v | (p^k - 1)$  and  $q$  is a power of  $p^k$ . Regardless of the size of  $f$  our present methods give a great deal of information about  $F(x)$ . Furthermore many of the proofs of [1] can be shortened and simplified by using the results of §1 of the present paper.

The results of [1] provide a complete answer for the case  $q = p$ . In the present paper the problem is completely solved for the case  $q = p^2$ .

**1. Preliminaries.** Let  $\mathcal{K}$  be a finite field with  $q$  elements and characteristic  $p$ . We use Greek letters for elements of  $\mathcal{K}$ , and small Latin letters, other than  $x$ , for nonnegative integers. We use capital letters for polynomials in one variable over  $\mathcal{K}$ . The polynomials denoted by  $A, B, C, D, E$  and the integers denoted by  $a, b, c, d, e$

---

Received May 1, 1963. Presented to the American Mathematical Society March 4, 1963. This work was partially supported by the National Science Foundation under NSF Grant 18916.

vary from proof to proof. The polynomials and integers denoted by other letters, except  $i$  and  $j$ , remain the same throughout the paper.

Let  $F = F(x)$  be a polynomial over  $\mathcal{K}$  of degree  $f, f > 0$ . Let  $\gamma_0, \gamma_1, \dots, \gamma_r$  denote the distinct values assumed by  $F(\tau)$  as  $\tau$  ranges over  $\mathcal{K}$ . It follows easily from the fact that a polynomial of degree  $f$  has at most  $f$  roots, that  $r + 1 \geq q/f$ . This is equivalent to  $r \geq [(q - 1)/f]$ . We intend to study the question raised in [1] of characterizing those polynomials for which  $r = [(q - 1)/f]$ . The cases  $r = 0$  and  $r = 1$  were fully treated in [1]. Hence we make the assumption that

$$(1) \quad r = [(q - 1)/f] \geq 2.$$

Subtracting the constant  $\gamma_0$  from  $F$  does not change the value of  $r$ . Thus it is sufficient to consider the case  $\gamma_0 = 0$ . In the first two sections of this paper, we assume that

$$\gamma_0 = 0.$$

Then  $\gamma_i \neq 0$  for  $i > 0$ . We now set

$$F_i = F - \gamma_i, \quad 0 \leq i \leq r.$$

The polynomials  $F_i$  are relatively prime in pairs, and each of them has at least one root in  $\mathcal{K}$ . Let  $\pi_{i1}, \pi_{i2}, \dots, \pi_{il_i}$  be the distinct roots of  $F_i$  that lie in  $\mathcal{K}$  and set

$$L_i = \prod_{j=1}^{l_i} (x - \pi_{ij}), \quad 0 \leq i \leq r.$$

Then  $l_i = \deg L_i \geq 1, 0 \leq i \leq r$ , and<sup>1</sup>

$$(2) \quad x^q - x = \prod_{i=0}^r L_i.$$

Now set  $F_i = L_i U_i, 0 \leq i \leq r$ , and

$$(3) \quad G = \prod_{i=0}^r U_i.$$

Then the  $L_i$ , the  $U_i$  and  $G$  are polynomials over  $\mathcal{K}$ , and

$$(4) \quad (x^q - x)G = \prod_{i=0}^r F_i.$$

Now (4) and (1) give us an upper bound on the degree of  $G$ , namely

$$\deg G = (r+1)f - q \leq q - 1 + f - q = f - 1.$$

---

<sup>1</sup> The relations (2), (3), (4), (5), (6), and (7) can all be found in [1] under the assumption that the leading coefficient of  $F$  is 1.

Thus we have

$$(5) \quad \deg G < f.$$

Set  $u_i = \deg U_i$ ,  $0 \leq i \leq r$ . We already have  $F' = F'_0$  by the assumption  $\gamma_0 = 0$ . We set  $L = L_0$ ,  $U = U_0$ ,  $l = l_0$ , and  $u = u_0$ .

We now differentiate both sides of (2) and obtain  $-1 \equiv L'L^* \pmod{L}$ , where  $L^* = L_1 L_2 \cdots L_r$ . Hence  $G \equiv -L'L^*G \pmod{LG}$ . Since  $F = LU$  and  $U|G$ , it follows that  $F'|LG$  and thus

$$G \equiv -L'L^*G \pmod{F}.$$

Now

$$L^*G = U \prod_{i=1}^r (L_i U_i) = U \prod_{i=1}^r (F' - \gamma_i) \equiv -\zeta U \pmod{F},$$

where

$$\zeta = -\prod_{i=1}^r (-\gamma_i) \neq 0.$$

Hence  $G \equiv \zeta L'U \pmod{F}$ . Since  $\deg(\zeta L'U) < \deg(LU) = f$  and  $\deg G < f$ , we must have

$$(6) \quad G = \zeta L'U.$$

By symmetry it follows that

$$(7) \quad G = \zeta_i L'_i U_i, \quad 0 \leq i \leq r,$$

for suitable nonzero elements  $\zeta_i$  of  $\mathcal{K}$ .

We next derive another expression for  $G$ .

**LEMMA 1.** *There exists a nonzero element  $\theta$  in  $\mathcal{K}$  such that  $G = \theta F'$ .*

*Proof.* Since  $F' = F'_i = L'_i U_i + L_i U'_i$ , it follows from (7) that

$$L_i U'_i = F' - G/\zeta_i, \quad 0 \leq i \leq r.$$

Therefore  $L_0 U'_0 = LU'$ ,  $L_1 U'_1$ , and  $L_2 U'_2$  are linearly dependent. Thus there exist  $\lambda$ ,  $\lambda_1$ , and  $\lambda_2$  in  $\mathcal{K}$ , not all zero, such that

$$\lambda LU' + \lambda_1 L_1 U'_1 + \lambda_2 L_2 U'_2 = 0.$$

Multiplying this relation by  $UU_1 U_2$  and noting that  $LU = F$ ,  $L_1 U_1 = F - \gamma_1$ ,  $L_2 U_2 = F - \gamma_2$ , we obtain

$$(8) \quad (\lambda U' U_1 U_2 + \lambda_1 U U'_1 U_2 + \lambda_2 U U_1 U'_2) F = \lambda_1 \gamma_1 U U'_1 U_2 + \lambda_2 \gamma_2 U U_1 U'_2.$$

Now the degree of the right side of (8) is less than  $u + u_1 + u_2$  and

$$u + u_1 + u_2 \leq \deg G < f = \deg F'.$$

This is possible only if we have

$$(9) \quad \lambda U' U_1 U_2 + \lambda_1 U U_1' U_2 + \lambda_2 U U_1 U_2' = 0.$$

The constants  $\lambda$ ,  $\lambda_1$ , and  $\lambda_2$  are not all zero. Without loss of generality we suppose  $\lambda_2 \neq 0$ . Then (9) gives us  $U_2 | U U_1 U_2'$ . Since  $U_2 | F_2$ ,  $U_2$  must be relatively prime to both  $F$  and  $F_1$ . Hence  $U_2$  is relatively prime to  $U U_1$ , and  $U_2 | U_2'$ . This implies that  $U_2' = 0$ . Hence

$$F' = F_2' = L_2' U_2 + L_2 U_2' = L_2' U_2 = G / \zeta_2.$$

Thus  $G = \zeta_2 F'$ , which completes this proof.

Lemma 1 is false for  $r \leq 1$ —counter examples can be readily constructed.

LEMMA 2. For each  $j$ ,  $0 \leq j \leq r$ ,  $U_j$  is of the form

$$U_j = L_j^{w_j} H_j^p,$$

where  $w_j$  is a nonnegative integer,  $H_j$  is a polynomial over  $\mathcal{K}$ , and  $L_j \nmid H_j$ .

*Proof.* By symmetry it is sufficient to prove the lemma for the case  $j = 0$ . Combining (6) with Lemma 1 we obtain

$$\zeta L' U = G = \theta F' = \theta L' U + \theta L U'.$$

Thus

$$(10) \quad \theta L U' = (\zeta - \theta) L' U.$$

We set  $U = L^w A$ , where  $L \nmid A$  and  $w \geq 0$ . Then substitution in (10) gives us

$$\theta w L^w L' A + \theta L^{w+1} A' = (\zeta - \theta) L' L^w A.$$

This reduces to

$$\theta L A' = (\zeta - \theta - w\theta) L' A.$$

Thus  $L | (\zeta - \theta - w\theta) L' A$ . Since  $L$  is the product of distinct linear factors, it follows that  $L$  and  $L'$  are relatively prime. Since  $L \nmid A$ , this implies that  $\zeta - \theta - w\theta = 0$ . Therefore  $\theta L A' = 0$ . It follows that  $A' = 0$ . Hence  $A = H^p$  for some polynomial  $H$ . Then we have  $L \nmid H$  and  $U = L^w H^p$ , which completes this proof.

We now suppose, without loss of generality, that

$$(11) \quad l \leq l_j, \quad 0 \leq j \leq r.$$

LEMMA 3. Under the assumption (11), the constants  $w_j$  of Lemma 2 satisfy

$$w_1 = w_2 = \dots = w_r = 0 .$$

*Proof.* Combining (3) and (6) we obtain

$$\zeta L' U = G = U U_1 U_2 \dots U_r .$$

Now suppose  $1 \leq j \leq r$ . Then  $U_j \mid L'$ , and hence

$$u_j \leq \deg L' < l \leq l_j .$$

Therefore  $L_j \nmid U_j$ , so that we have  $w_j = 0$ . This completes the proof.

Set  $H = H_0$  and  $v = w_0 + 1$ . Then from Lemmas 2 and 3 we obtain

$$(12) \quad F = LU = L^v H^p ,$$

and

$$(13) \quad F_i = L_i U_i = L_i H_i^p , \quad 1 \leq i \leq r ,$$

where  $L \nmid H$ ,  $L_i \nmid H_i$ . Moreover

$$\zeta L' = G/U = U_1 U_2 \dots U_r = (H_1 H_2 \dots H_r)^p .$$

Thus  $L' = S^p$ , where  $S = \zeta^{-1/p} H_1 H_2 \dots H_r$ . Therefore  $L$  is of the form

$$(14) \quad L = x S^p + T^p ,$$

where  $T$ , as well as  $S$ , is a polynomial over  $\mathcal{K}$ .

2. The polynomial  $R(x)$ . Set

$$R(x) = \prod_{i=1}^r (x - \gamma_i) = \sum_{j=0}^r \rho_j x^j ,$$

where  $\rho_j \in \mathcal{K}$ ,  $0 \leq j \leq r$ ,  $\rho_r = 1$ . From (4) and (6) we obtain

$$LUR(F) = FR(F) = \prod_{i=0}^r F_i = (x^q - x)G = \zeta(x^q - x)L'U .$$

These identities and (12) give us

$$(15) \quad \sum_{j=0}^r \rho_j L^{1+vj} H^{pj} = LR(F) = \zeta(x^q - x)L' .$$

Differentiating both sides of (15) and noting that  $L'' = 0$  by (14), we get the congruence

$$\rho_0 L' \equiv -\zeta L' \pmod{L}.$$

Since  $L' \neq 0$ , we obtain

$$(16) \quad \rho_0 = -\zeta.$$

By Lemma 1 we have  $F' = G/\theta \neq 0$ . Hence  $p \nmid v$ .

Let  $k$  be the smallest positive integer such that  $v \mid (p^k - 1)$ . The main objective of this section is to show that  $1 + vj$  is a power of  $p^k$  for every nonzero coefficient  $\rho_j$  of  $R(x)$ .

In the proof of the following lemma the notation  $A \parallel B$  means that  $A \mid B$  and  $(A, B/A) = 1$ .

**LEMMA 4.** *Let  $d$  be a nonnegative integer such that  $L'$  is a  $p^d$ th power and  $1 + vr > p^{d-1}$ . If  $j$  is an integer such that  $\rho_j \neq 0$ , then either (i)  $1 + vj$  is a power of  $p^k$ , or (ii)  $p^d \mid (1 + vj)$ . Moreover  $H$  is a  $p^{d-1}$ st power.*

*Proof by induction on  $d$ .* The desired result is trivial for  $d = 0$ . We suppose that it is true for an integer  $d$  and show that this implies that it is true for  $d + 1$ . Thus we assume that  $L'$  is a  $p^{d+1}$ st power and  $1 + vr > p^d$ . Then the induction hypothesis applies so that  $R(x)$  is of the form

$$(17) \quad R(x) = \sum_{i=0}^c \omega_i x^{(p^{kt}-1)/v} + \Sigma' \rho_j x^j,$$

where  $\omega_i \in \mathcal{K}$ ,  $0 \leq i \leq c$ ,  $c = [d/k]$ , and the second summation  $\Sigma'$  is over all  $j$  such that

$$p^d \mid (1 + vj), \quad p^d < 1 + vj, \quad j \leq r.$$

Moreover  $H$  is a  $p^{d-1}$ st power. Thus

$$H = A^{p^{d-1}} \quad \text{and} \quad F = L^v A^{p^d}$$

for some polynomial  $A$  over  $\mathcal{K}$ . Substitution in (15) gives us

$$(18) \quad \Sigma' \rho_j L^{1+vj} A^{jp^d} = \zeta x^q L' + B,$$

where

$$B = -\zeta x L' - \sum_{i=0}^c \omega_i L^{p^{kt}} A^{p^d(p^{kt}-1)/v}.$$

The left side of (18) is a  $p^d$ th power. Since

$$q \geq 1 + fr \geq 1 + vr > p^d$$

and  $q$  is a power of  $p$ , it follows that  $p^{d+1} \mid q$ . Hence  $\zeta x^q L'$  is a  $p^{d+1}$ st power. Therefore  $B$  is a  $p^d$ th power. Thus we can set

$$\zeta x^q L' = C^{p^{a+1}} \quad \text{and} \quad B = D^{p^a}.$$

Since  $1 + vr > p^a$  and  $\rho_r \neq 0$ , it follows that the left side of (18) does not vanish identically. Let the term corresponding to  $j = a$  be the nonzero term of lowest degree in the left side of (18). Thus  $a$  is the least integer such that  $\rho_a \neq 0$  and  $1 + va > p^a$ . Then  $p^a | (1 + va)$ , and hence  $1 + va \geq 2p^a$ . Because of the way  $a$  was chosen we have

$$(19) \quad L^{1+va} A^{ap^a} | (\zeta x^q L' + B).$$

Extracting the  $p^a$ th roots of both sides of (19) we get

$$L^{(1+va)p^{-a}} A^a | (C^p + D).$$

Since  $1 + va \geq 2p^a$  this gives us  $L^2 A^a | (C^p + D)$ . By differentiation we obtain

$$(20) \quad LA^{a-1} | D'.$$

Now

$$\deg D' < p^{-a} \deg B \leq p^{-a} \deg \{L^{p^{kc}} A^{p^a(p^{kc}-1)/v}\} \leq \deg \{LA^{(p^{kc}-1)/v}\}.$$

Since

$$a > (p^a - 1)/v \geq (p^{kc} - 1)/v,$$

we have  $(p^{kc} - 1)/v \leq a - 1$ , and

$$\deg D' < \deg (LA^{a-1}).$$

Combining this with (20) we get  $D' = 0$ . Thus  $D$  must be a  $p$ th power, and  $B$  a  $p^{a+1}$ st power. Thus the right side of (19) is a  $p^{a+1}$ st power. Hence the left side of (19) is also a  $p^{a+1}$ st power. Now  $L \nmid H$ . Since  $L$  is the product of distinct linear factors we have  $L \nmid A$ ,  $p^{a+1} | (1 + va)$ , and  $A^a$  is a  $p$ th power. Hence  $p \nmid a$ , and  $A$  itself is a  $p$ th power. It follows that  $H$  is a  $p^a$ th power. Suppose there is a  $b$  such that  $\rho_b \neq 0$ ,  $1 + vb$  is not a power of  $p^k$ , and  $p^{a+1} \nmid (1 + vb)$ . Without loss of generality suppose that  $b$  is the smallest integer with these properties. By (17) we have  $1 + vb > p^a$ , and by (18) we have

$$(21) \quad L^{1+vb} A^{bp^a} | \{\zeta x^q L' + B - \Sigma'' \rho_j L^{1+vj} A^{jp^a}\},$$

where  $\Sigma''$  is over those  $j$  such that  $j < b$ ,  $p^{a+1} | (1 + vj)$ . The right side of (21) is a  $p^{a+1}$ st power. Hence the left side of (21) is also a  $p^{a+1}$ st power. Therefore  $p^{a+1} | (1 + vb)$ , a contradiction. It follows that for every  $j$  such that  $\rho_j \neq 0$ , either  $1 + vj$  is a power of  $p^k$  or  $p^{a+1} | (1 + vj)$ . This establishes the desired result for  $d + 1$ , and



completes this proof.

**LEMMA 5.** *Suppose there exists an integer  $d$  such that  $L'$  is a  $p^d$ th power but not a  $p^{d+1}$ st power, and  $1 + vr > p^d$ . Then  $v = 1$  and  $p^{d+1} \nmid (1 + r)$ .*

*Proof.* Since  $L'$  is a  $p$ th power by (14), we have  $d \geq 1$ . By Lemma 4 we have

$$R(x) = \sum_{i=0}^c \omega_i x^{(p^{ki}-1)/v} + \Sigma^* \rho_j x^j + x^r,$$

where the  $\omega_i$  are elements of  $\mathcal{K}$ ,  $c = [d/k]$ , and the summation  $\Sigma^*$  is over all  $j$  such that  $p^d \mid (1 + vj)$ ,  $p^d < 1 + vj$ ,  $j < r$ . Moreover since  $1 + vr > p^d$  and  $\rho_r \neq 0$ , we have  $p^d \mid (1 + vr)$ . Furthermore  $H$  is a  $p^{d-1}$ st power. Since  $\zeta \in \mathcal{K}$ , it follows that  $\zeta L'$  is a  $p^d$ th power but not a  $p^{d+1}$ st power. Thus we can set

$$H = A^{p^{d-1}} \quad \text{and} \quad \zeta L' = C^{p^d},$$

where  $C$  is not a  $p$ th power. Substitution in (15) gives us

$$(22) \quad L^{1+vr} A^{rp^d} = x^q C^{p^d} + B,$$

where

$$\begin{aligned} B &= -\zeta x L' - L R(F) + L F^r \\ &= -\zeta x L' - \sum_{i=0}^c \omega_i L^{p^{ki}} A^{p^d(p^{ki}-1)/v} - \Sigma^* \rho_j L^{1+vj} A^{jp^d}. \end{aligned}$$

Now the left side of (22) is a  $p^d$ th power. Moreover

$$q \geq 1 + fr \geq 1 + vr > p^d,$$

so that  $p^{d+1} \mid q$ . Therefore  $B$  is a  $p^d$ th power, say  $B = D^{p^d}$ . Extracting the  $p^d$ th roots of both sides of (22) we obtain

$$(23) \quad L^{(1+vr)p^{-d}} A^r = x^{qp^{-d}} C + D.$$

Differentiation now yields

$$(24) \quad L^{-1+(1+vr)p^{-d}} A^{r-1} \{(1 + vr)p^{-d} L' A + r L A'\} = x^{qp^{-d}} C' + D'.$$

since  $p^{d+1} \mid q$ . Multiplying (24) by  $C$ , (23) by  $C'$ , and subtracting, we get

$$(25) \quad L^{-1+(1+vr)p^{-d}} A^{r-1} E = C D' - C' D,$$

where

$$E = (1 + vr)p^{-d} L' A C + r L A' C - L A C'.$$

Now  $A \mid H$  and therefore  $LA \mid F$ . Moreover

$$C \mid L' = G/(\zeta U) = \zeta^{-1}U_1U_2 \cdots U_r \mid F_1F_2 \cdots F_r.$$

Hence  $C$  is relatively prime to  $LA$ . Since  $C$  is not a  $p$ th power we have  $C' \neq 0$ . Hence  $C \nmid LAC'$ . It follows that  $E \neq 0$ . From (25) we obtain  $CD' \neq C'D$  and

$$(26) \quad L^{-e+(1+vr)p^{-a}}A^{r-1} \mid (CD' - C'D),$$

where

$$e = \begin{cases} 0 & \text{if } p^{a+1} \mid (1 + vr), \\ 1 & \text{if } p^{a+1} \nmid (1 + vr). \end{cases}$$

Comparing degrees in (26) we obtain

$$(27) \quad (1 + vr - ep^a)l + p^a(r - 1) \deg A < p^a \deg(CD) = \deg(L'B).$$

Now the leading term of  $R(x)$  is  $x^r$  and  $R(x) \neq x^r$ . Set  $b = \deg\{R(x) - x^r\}$ . Then we have  $0 \leq b < r$  and

$$\begin{aligned} \deg B &\leq \deg(LF^b) \\ &= (1 + vb)l + bp^a \deg A \leq (1 + vb)l + (r - 1)p^a \deg A. \end{aligned}$$

Substitution in (27) gives us, after simplification,

$$v(r - b)l < ep^al + \deg L' < (ep^a + 1)l.$$

Hence  $v(r - b) \leq ep^a$ . Therefore  $e \neq 0$ . Hence  $e = 1$  and

$$v(r - b) \leq p^a.$$

Since  $p^a \mid (1 + vr)$  and  $1 + vr > p^a$ , we have  $1 + vr \geq 2p^a$  and

$$1 + vb = 1 + vr - v(r - b) \geq p^a.$$

Since  $\rho_b \neq 0$ , this gives us  $p^a \mid (1 + vb)$ . Since  $p^a \mid (1 + vr)$ , it follows that  $p^a \mid v(r - b)$  and  $p \nmid v$ . Hence  $v(r - b) = p^a$  and  $v = 1$ . Finally since  $e = 1$  we have

$$p^{a+1} \nmid (1 + vr) = 1 + r,$$

which completes this proof.

**LEMMA 6.** *If  $d$  is an integer such that  $p^a < 1 + vr$ , then  $L'$  is a  $p^{a+1}$ st power.*

*Proof.* Suppose the result is false. Then  $L'$  is not a  $p^{a+1}$ st power and  $p^a < 1 + vr$ . Without loss of generality we suppose that  $L'$  is a  $p^a$ th power. By Lemma 5 we have  $v = 1$  and  $p^{a+1} \nmid (1 + r)$ .

Therefore  $k = 1$  and  $p^a < 1 + r$ . It follows from Lemma 4 that  $R(x)$  is of the form

$$R(x) = \sum_{i=0}^{d-1} \omega_i x^{p^i-1} + \Sigma^+ \rho_j x^j ,$$

where the summation  $\Sigma^+$  is over all  $j$  such that  $p^a | (1 + j)$ ,  $j \leq r$ . Moreover  $H$  is a  $p^{a-1}$ -st power and  $p^a | (1 + r)$ . Now

$$FR(F) = \prod_{i=0}^r (F - \gamma_i) = \prod_{i=0}^r F_i = (x^q - x)G$$

by (4), so that

$$(28) \quad \Sigma^+ \rho_j F^{j+1} = x^q G + B ,$$

where  $\deg B \leq p^{a-1}f$ . The left side of (28) is a  $p^a$ th power. Moreover  $q \geq 1 + fr \geq 1 + r > p^a$ , so that  $x^q$  is a  $p^{a+1}$ -st power. Since  $G = \zeta L'U$  and  $U = L^{v-1}H^p = H^p$ , it follows that  $G$  is a  $p^a$ th power. Hence  $B$  is also a  $p^a$ th Power. We set

$$G = C^{p^a} \quad \text{and} \quad B = D^{p^a} .$$

Then, extracting the  $p^a$ th roots of both sides of (28), we get

$$(29) \quad \sum_{j=1}^a \xi_j F^j = x^{qp-a} C + D ,$$

where  $a = (r + 1)p^{-a} \geq 2$ , the  $\xi_j$  are in  $\mathcal{K}$ ,  $\xi_a = 1$ , and  $\deg D \leq f/p$ . Now  $p \nmid a$  since  $p^{a+1} \nmid (r + 1)$ . We set  $\bar{F} = F + \xi_{a-1}/a$ . Then (29) becomes

$$(30) \quad \sum_{j=0}^a \eta_j \bar{F}^j = x^{qp-a} C + D ,$$

where the  $\eta_j$  are in  $\mathcal{K}$ ,  $\eta_a = 1$ , and  $\eta_{a-1} = 0$ . Differentiating (30) we obtain

$$(31) \quad \sum_{j=1}^a j \eta_j \bar{F}^{j-1} \bar{F}' = x^{qp-a} C' + D' .$$

Eliminating  $x^{qp-a}$  from (30) and (31) we get

$$\eta_0 C' + \sum_{j=1}^a \eta_j \bar{F}^{j-1} (C' \bar{F} - j C \bar{F}') = C' D - C D' .$$

Since  $\eta_{a-1} = 0$ , it follows that

$$(32) \quad \bar{F}^{a-1} (C' \bar{F} - a C \bar{F}') = C' D - C D' - E ,$$

where

$$\deg E < (a - 2)f + \deg C .$$

Now

$$\deg C = p^{-a} \deg G < p^{-a} f \leq f/p$$

by (5). Hence  $\deg E < (a-1)f$ , and

$$\deg (C'D - CD') < \deg (CD) < 2f/p \leq (a-1)f.$$

Therefore

$$\deg (C'D - CD' - E) < (a-1)f = \deg \bar{F}^{a-1},$$

and (32) yields

$$C'\bar{F} = aC\bar{F}'.$$

Now  $\bar{F}' = F' = \theta^{-1}G \neq 0$  by Lemma 1. Therefore  $aC\bar{F}' \neq 0$ . Hence  $C' \neq 0$  and thus  $C \nmid C'$ . It follows that  $(\bar{F}, C) \neq 1$ . Since

$$C^{p^a} = G = \prod_{i=0}^r U_i$$

we have  $(\bar{F}, U_b) \neq 1$  for some  $b, 0 \leq b \leq r$ . Hence  $(\bar{F}, F_b) \neq 1$ . Since  $\bar{F} - F_b \in \mathcal{K}$ , we must have  $\bar{F} = F_b$ . Therefore

$$C'F_b = aCF'_b.$$

Since  $v = 1$ , we have  $F_b = L_b H_b^p$ , whether or not  $b = 0$ . Hence

$$C'L_b H_b^p = aCL'_b H_b^p,$$

and  $C'L_b = aCL'_b$ . Now  $L_b$  is relatively prime to  $L'_b$ . Therefore  $L_b \mid C$ . Since  $v = 1$  we have

$$C^{p^a} = G = \prod_{i=0}^r U_i = \prod_{i=0}^r H_i^p.$$

It follows that  $L_b \mid H_0 H_1 \cdots H_r$ . On the other hand  $L_b \nmid H_b$ , while for  $i \neq b$  we have  $(L_b, H_i) = 1$ . Therefore  $L_b \nmid H_0 H_1 \cdots H_r$ , a contradiction. This completes the proof of this lemma.

We are now in a position to prove the most general theorem of this paper. We drop the assumption  $\gamma_0 = 0$ .

**THEOREM 1.** *Let  $\mathcal{K}$  be a finite field of characteristic  $p$  that contains exactly  $q$  elements. Let  $F(x)$  be a polynomial over  $\mathcal{K}$  of degree  $f, f > 0$ . Let  $\gamma_0, \gamma_1, \dots, \gamma_r$  be the distinct values  $F(\tau)$  as  $\tau$  ranges over  $\mathcal{K}$ , and let  $l_i$  denote the number of distinct roots in  $\mathcal{K}$  of the polynomial  $F(x) - \gamma_i$ . Let the  $\gamma_i$  be arranged in such a way that  $l_0 \leq l_i, 1 \leq i \leq r$ . Set  $L = \prod (x - \pi)$ , where the product is over the distinct roots  $\pi$  of  $F(x) - \gamma_0$  that lie in  $\mathcal{K}$ . Suppose that*

$r = [(q-1)/f] \geq 2$ . Then there exist positive integers  $v, k, m$ ; a polynomial  $N$  over  $\mathcal{K}$ ; and  $\omega_0, \omega_1, \dots, \omega_m$  in  $\mathcal{K}$  such that  $L \nmid N$ ,  $v \mid (p^k - 1)$ ,  $1 + vr = p^{mk}$ ,  $L'$  is a  $p^{mk}$ th power,  $\omega_0 \neq 0$ ,  $\omega_m = 1$ ,

$$F(x) = L^v N^{p^{mk}} + \gamma_0,$$

$$(33) \quad \prod_{i=1}^r (x - \gamma_i + \gamma_0) = \sum_{i=0}^m \omega_i x^{(p^{ki}-1)/v},$$

and

$$(34) \quad \sum_{i=0}^m \omega_i L^{p^{ki}} N^{p^{km}(p^{ki}-1)/v} = -\omega_0 (x^q - x) L'.$$

*Proof.* Without loss of generality we can suppose that  $\gamma_0 = 0$ , so that our previous discussion applies. Let  $d$  be the integer such that

$$p^d \geq 1 + vr > p^{d-1}.$$

It follows from Lemma 6 that  $L'$  is a  $p^d$ th power. We now apply Lemma 4 to conclude that either  $1 + vr$  is a power of  $p^k$  or  $p^a \mid (1 + vr)$ . In either case we must have  $p^a = 1 + vr$ . Since  $k$  is the smallest positive integer such that  $v \mid (p^k - 1)$ , it follows that  $k \mid d$ . We put  $m = d/k$ . Then  $L'$  is a  $p^{mk}$ th power and  $1 + vr = p^{mk}$ . Applying Lemma 4 again we find that  $R(x)$  is of the form

$$R(x) = \sum_{i=0}^m \omega_i x^{(p^{ki}-1)/v},$$

so that (33) holds. Moreover  $H$  is a  $p^{d-1}$ st power by Lemma 4, and therefore  $H^p$  is a  $p^{mk}$ th power. Thus there is a polynomial  $N$  over  $\mathcal{K}$  such that

$$F = L^v H^p = L^v N^{p^{mk}}.$$

Furthermore since  $L \nmid H$ , it follows that  $L \nmid N$ . Using (16) we obtain  $\omega_0 = \rho_0 = -\zeta \neq 0$ . It follows at once from (33) that  $\omega_m = 1$ . Finally we substitute in (15) to obtain (34). This completes the proof of the theorem.

In the next two sections we apply Theorem 1 to a number of special cases.

**3. A special case.** There are two general types of polynomials known for which (1) holds [1, § 5]. For every polynomial of the first type both  $L'$  and  $N$  are constants. Thus this case is of special interest. Here we have the following result:

LEMMA 7. *Suppose that  $L'$  and  $N$  are both constants. Then  $q$  is a power of  $p^k$ , and  $F$  is of the form*

$$(35) \quad F = \alpha L^v + \gamma, \quad L = \beta + \sum_{j=0}^d \varphi_j x^{p^{kj}},$$

where  $L$  factors into distinct linear factors over  $\mathcal{K}$  and  $v \mid (p^k - 1)$ .

*Proof.* Since  $N$  is a constant it follows from Theorem 1 that  $F = \alpha L^v + \gamma$ , where  $\alpha \in \mathcal{K}$  and  $\gamma = \gamma_0 \in \mathcal{K}$ . Suppose that  $L$  is not of the form given in (35). Then, since  $L'$  is a constant, we can write

$$(36) \quad L = \beta + \sum_{j=0}^c \varphi_j x^{p^{kj}} + \sum_{j=a}^{l/p} \delta_j x^{p^j}$$

where  $a$  and  $c$  are integers such that

$$p^{k(c+1)} > pa > p^{kc}, \quad l \geq pa,$$

and  $\delta_a \neq 0$ . Moreover  $L' = \varphi_0 \neq 0$ . Now (34) becomes

$$(37) \quad \sum_{i=0}^m \chi_i L^{p^{ki}} = -\omega_0 \mathcal{P}_0(x^q - x),$$

where the  $\chi_i$  are in  $\mathcal{K}$ ,  $\chi_0 = \omega_0 \neq 0$ , and  $\chi_m \neq 0$ . Substituting (36) in (37) we get

$$\psi + \sum_{j=0}^c \psi_j x^{p^{kj}} + \chi_0 \delta_a x^{pa} + \sum_{j=pa+1}^{lp^{km}} \sigma_j x^j = -\omega_0 \mathcal{P}_0(x^q - x),$$

for suitable  $\psi$ ,  $\psi_j$ ,  $\sigma_j$  in  $\mathcal{K}$ . Since  $\chi_0 \delta_a \neq 0$ , this implies that either  $pa = 1$  or  $pa = q$ . Comparing degrees we obtain

$$q = lp^{km} > l \geq pa.$$

Clearly  $pa \neq 1$ . This contradiction implies that  $L$  is of the desired form, which completes this proof.

The converse of Lemma 7 is already known [1]: *If  $q$  is a power of  $p^k$ , and if  $F$  is of the form (35), then the polynomial  $F$  satisfies the equality  $r = [(q-1)/f]$ . This was proved in [1] as follows: Let  $\pi$  be a root of  $L$ . Replacing  $x$  by  $x + \pi$  we can assume that  $\beta = 0$ . Let  $l = \deg L$  as before, and set  $L(x) = L$ . Because of the form of  $L$  the values assumed by  $L(\tau)$  as  $\tau$  ranges over  $\mathcal{K}$  form a vector space over the subfield  $GF(p^k)$ . Since we have assumed that  $L$  factors into distinct linear factors over  $\mathcal{K}$ , it follows that  $L$  has exactly  $l$  distinct roots in  $\mathcal{K}$ . Therefore this vector space contains exactly  $q/l$  distinct elements. Then since  $F = \alpha L^v + \gamma$ , where  $v \mid (p^k - 1)$ , it follows that the number of values assumed by  $F(\tau)$  as*

$\tau$  ranges over  $\mathcal{K}$  is exactly

$$1 + (-1 + q/l)/v = 1 + (q - l)/f = 1 + [(q - 1)/f] .$$

Hence  $r = [(q - 1)/f]$ .

Thus we have a complete characterization of those polynomials for which  $r = [(q - 1)/f] \geq 2$ , subject to the condition that  $L'$  and  $N$  are both constants. One significance of this result can be seen from the following lemma:

**LEMMA 8.** *If  $f \leq \sqrt{q}$ , and  $r = [(q - 1)/f] \geq 2$ , then  $L'$  and  $N$  are both constants.*

*Proof.* Theorem 1 applies so that we have  $1 + rv = p^{m_k}$ , and  $f = vl + p^{m_k} \deg N$ . Moreover  $f^2 \leq q$  and  $r = [(q - 1)/f]$  so that

$$f \leq q/f \leq r + 1 = 1 + (p^{m_k} - 1)/v \leq p^{m_k} .$$

Thus  $p^{m_k} \deg N < f \leq p^{m_k}$ ,  $\deg N = 0$ , and  $N$  is a constant. Furthermore  $L'$  is a  $p^{m_k}$ th power by Theorem 1 and  $\deg L' < l \leq f \leq p^{m_k}$ . Hence  $L'$  is also a constant, and the proof of this lemma is complete.

The above results give us a complete characterization of those polynomials  $F$  for which  $r = [(q - 1)/f] \geq 2$  and  $0 < f \leq \sqrt{q}$ . Now suppose that  $r = [(q - 1)/f] < 2$  and  $0 < f \leq \sqrt{q}$ . Then

$$2 > (q - 1)/f \geq (f^2 - 1)/f ,$$

$f^2 - 2f - 1 < 0$ , and thus  $f = 1$  or  $f = 2$ . Now  $q$  is a prime power and  $f^2 \leq q < 2f + 1$ . Hence we have either (i)  $f = 1$  and  $q = 2$ , or (ii)  $f = 2$  and  $q = 4$ . If  $f = 1$ , then  $F$  is clearly of the form (35) with  $v = k = 1$  and  $d = 0$ . If  $f = 2$  and  $q = 4$ , then  $r = 1$ , and since  $F_0$  and  $F_1$  together have 4 distinct roots in  $\mathcal{K}$ , it follows that  $F_0$  has two distinct roots in  $\mathcal{K}$ , so that  $F$  is still of the form (35), this time with  $p = 2$  and  $v = k = d = 1$ . Thus we see that the condition  $r \geq 2$  can be dropped here. Combining all these results we obtain one of our major results:

**THEOREM 2.** *Let  $F(x)$  be a polynomial over the finite field  $\mathcal{K}$  of characteristic  $p$  and let  $q$  denote the number of elements of  $\mathcal{K}$ . Let  $r + 1$  denote the number of distinct values assumed by  $F(\tau)$  as  $\tau$  ranges over  $\mathcal{K}$ , and let  $f$  be the degree of  $F(x)$ . Suppose that  $0 < f \leq \sqrt{q}$ . Then*

$$r = [(q - 1)/f]$$

*if and only if  $F$  is of the form*

$$F = \alpha L^v + \gamma ,$$

where  $L$  is a polynomial that factors into distinct linear factors over  $\mathcal{K}$  and that has the form

$$L = \beta + \sum_{i=0}^d \varphi_i x^{p^{ki}} .$$

and where  $v$  and  $k$  are integers such that  $v \mid (p^k - 1)$ ,  $q$  is a power of  $p^k$ , and  $\alpha, \beta, \gamma$ , and the  $\varphi_i$  are elements of  $\mathcal{K}$ .

**4. The cases  $q = p$  and  $q = p^2$ .** The results of §1 enable us to treat the case  $q = p$  quickly.

Suppose  $q = p$  and  $r = [(q - 1)/f] \geq 2$ . If  $\gamma_0 = 0$ , then the results of §1 apply, so that

$$F = L^v H^v, \quad L = xS^p + T^p$$

by (12) and (14). Since

$$\deg F = f \leq \frac{1}{2}(q - 1) = \frac{1}{2}(p - 1) < p ,$$

the polynomials  $H, S$ , and  $T$  are all constants. Thus  $F$  is of the form  $\alpha(x + \beta)^v$  and  $v = f$ . It is easily shown that  $v \mid (q - 1)$  here. Dropping the assumption  $\gamma_0 = 0$ , we see that if  $q = p$  and  $r = [(q - 1)/f] \geq 2$ , then  $f \mid (q - 1)$  and  $F$  is of the form

$$F = \alpha(x + \beta)^f + \gamma .$$

We note that in this case  $L'$  and  $N$  must both be constants, so that we could have obtained this result from Lemma 7.

Let us now consider the case  $q = p^2$ . Comparing the degrees of the two sides of (34) we obtain

$$p^{mk}l + rp^{mk} \deg N = q + \deg L' \leq q + l - 1 = p^2 + l - 1 .$$

Therefore

$$(38) \quad pl + p \deg N \leq p^2 + l - 1 .$$

Thus  $pl \leq p^2 + l - 1$  or  $l \leq p + 1$ . Since  $L'$  is a  $p$ th power, it follows that  $l \equiv 0$  or  $1 \pmod{p}$ . Therefore  $l = 1, p$ , or  $p + 1$ . If  $l = p$  or  $p + 1$ , the inequality (38) gives us

$$p \deg N \leq p^2 - l(p - 1) - 1 \leq p - 1 ,$$

$\deg N = 0$  and  $N$  is a constant. If  $l = 1$ , then  $L$  is of the form  $x + \beta$ ,  $L' = 1$ , and (34) gives us



$$N|(-\omega_0 x^q + w_0 x - \omega_0 L) = -\omega_0(x^q + \beta) = -\omega_0 L^q .$$

Thus in case  $l = 1$ , we see that  $N$  is a constant times a power of  $L$ . Since  $L \nmid N$ , this implies that  $N$  is a constant. Thus  $N$  is a constant in all three cases.

If  $L'$  is also a constant then Lemma 7 applies, and  $F$  is of the form (35) with either (i)  $l = 1$ ,  $d = 0$ , and  $v \mid (p^2 - 1)$ , or (ii)  $l = p$ ,  $k = d = 1$ , and  $v \mid (p - 1)$ .

Now suppose that  $L'$  is not a constant. Since  $L'$  is a  $p^{m/k}$ th power by Theorem 1, we must have  $l = p + 1$  and  $m = k = 1$ . Since  $N$  is a constant we have  $F = \alpha L^p + \gamma$ , where  $\alpha \in \mathcal{K}$  and  $\gamma = \gamma_0 \in \mathcal{K}$ . Moreover  $L$  is of the form  $L = xS^p + T^p$  by (14). Since  $L$  has leading coefficient 1,  $S$  is of the form  $S = x + \varphi$ . Moreover  $T$  is of the form  $T = \mu x + \nu$ . Now (34) becomes

$$\omega_0 L + \chi L^p = -\omega_0(x^q - x)S^p ,$$

where  $\chi \in \mathcal{K}$ . Comparing leading coefficients we see that  $\chi = -\omega_0$ . Therefore

$$L^p = (x^q - x)S^p + L = x^{p^2}S^p + T^p .$$

Extracting  $p$ th roots we obtain  $L = x^p S + T$ . Thus

$$xS^p + T^p = x^p S + T ,$$

or

$$(39) \quad x^{p+1} + \mu^p x^p + \varphi^p x + \nu^p = x^{p+1} + \varphi x^p + \mu x + \nu .$$

Comparing the coefficients of  $x$  in (39) we obtain  $\mu = \varphi^p$ . Therefore

$$L = x^p S + T = x^{p+1} + \varphi x^p + \varphi^p x + \nu = (x + \varphi)^{p+1} + \beta ,$$

where  $\beta = \nu - \varphi^{p+1}$ . Comparing the constant terms of (39) we get  $\nu^p = \nu$ . Therefore  $\nu \in GF(p)$ , the prime field of  $\mathcal{K}$ . Now  $\varphi^{p+1} \in GF(p)$ . Hence  $\beta \in GF(p)$ . Since  $L$  has distinct roots we have  $\beta \neq 0$ . Now if  $v = 1$ , then  $F = \alpha L + \gamma$ , and  $F - \gamma - \alpha\beta$  has exactly one distinct root in  $\mathcal{K}$ , contradicting (11). Thus  $v \geq 2$ . We have shown that if  $q = p^2$ ,  $r = [(q - 1)/f] \geq 2$  and  $L'$  is not constant, then  $F$  is of the form  $\alpha L^p + \gamma$ , where  $L$  is of the form

$$L = (x + \varphi)^{p+1} + \beta ,$$

where  $\beta \in GF(p)$ ,  $\beta \neq 0$ ,  $v \mid (p - 1)$ ,  $v \geq 2$ .

Conversely if  $q = p^2$  and  $F$  has this form, then  $L(\tau) \in GF(p)$  for all  $\tau \in \mathcal{K}$ , and thus  $F$  assumes at most

$$1 + (p - 1)/v = 1 + (q - 1)/f = 1 + [(q - 1)/f]$$

distinct values. Since we always have  $r \geq [(q-1)/f]$ , this implies that  $r = [(q-1)/f]$ .

We have completed the discussion of the case  $q = p^2$ . We sum up our results for this case in our final theorem:

**THEOREM 3.** *Let  $\mathcal{K}$  be a field of characteristic  $p$  that contains exactly  $p^2$  elements. Let  $F(x)$  be a polynomial over  $\mathcal{K}$  of degree  $f$ ,  $f > 0$ . Let  $F(\tau)$  assume exactly  $r+1$  distinct values as  $\tau$  ranges over  $\mathcal{K}$ . If  $r = [(p^2-1)/f] \geq 2$ , then  $F(x)$  has one of the following three forms:*

- (i)  $F(x) = \alpha(x + \beta)^v + \gamma$ , where  $v \mid (p^2 - 1)$ ,  $\alpha \neq 0$ ,
- (ii)  $F(x) = \alpha(x^p + \varphi x + \beta)^v + \gamma$ , where  $x^p + \varphi x + \beta$  has  $p$  distinct roots in  $\mathcal{K}$ ,  $v \mid (p-1)$ ,  $\alpha \neq 0$ ,
- (iii)  $F(x) = \alpha\{(x + \varphi)^{p+1} + \beta\}^v + \gamma$ , where  $\beta \in GF(p)$ ,  $\beta \neq 0$ ,  $v \geq 2$ ,  $v \mid (p-1)$ , and  $\alpha \neq 0$ .

*Conversely if  $F(x)$  has one of these three forms, then  $r = [(q-1)/f]$ .*

For  $q > p^2$ , the question of the characterization of all polynomials  $F$  for which (1) holds, remains open. The most general types of polynomials known for which (1) holds are described in [1, §5]. At present it seems unlikely that there are any more.

#### REFERENCE

1. L. Carlitz, D. J. Lewis, W. H. Mills and E. G. Straus, *Polynomials over finite fields with minimal value sets*, *Mathematika* **8** (1961), 121-130.

YALE UNIVERSITY



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

ROBERT OSSERMAN  
Stanford University  
Stanford, California

M. G. ARSOVE  
University of Washington  
Seattle 5, Washington

J. DUGUNDJI  
University of Southern California  
Los Angeles 7, California

LOWELL J. PAIGE  
University of California  
Los Angeles 24, California

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

Richard Arens, <i>Normal form for a Pfaffian</i> .....	1
Charles Vernon Coffman, <i>Non-linear differential equations on cones in Banach spaces</i> .....	9
Ralph DeMarr, <i>Order convergence in linear topological spaces</i> .....	17
Peter Larkin Duren, <i>On the spectrum of a Toeplitz operator</i> .....	21
Robert E. Edwards, <i>Endomorphisms of function-spaces which leave stable all translation-invariant manifolds</i> .....	31
Erik Maurice Ellentuck, <i>Infinite products of isols</i> .....	49
William James Firey, <i>Some applications of means of convex bodies</i> .....	53
Haim Gaifman, <i>Concerning measures on Boolean algebras</i> .....	61
Richard Carl Gilbert, <i>Extremal spectral functions of a symmetric operator</i> .....	75
Ronald Lewis Graham, <i>On finite sums of reciprocals of distinct <math>n</math>th powers</i> .....	85
Hwa Suk Hahn, <i>On the relative growth of differences of partition functions</i> .....	93
Isidore Isaac Hirschman, Jr., <i>Extreme eigen values of Toeplitz forms associated with Jacobi polynomials</i> .....	107
Chen-jung Hsu, <i>Remarks on certain almost product spaces</i> .....	163
George Seth Innis, Jr., <i>Some reproducing kernels for the unit disk</i> .....	177
Ronald Jacobowitz, <i>Multiplicativity of the local Hilbert symbol</i> .....	187
Paul Joseph Kelly, <i>On some mappings related to graphs</i> .....	191
William A. Kirk, <i>On curvature of a metric space at a point</i> .....	195
G. J. Kurowski, <i>On the convergence of semi-discrete analytic functions</i> .....	199
Richard George Laatsch, <i>Extensions of subadditive functions</i> .....	209
V. Marić, <i>On some properties of solutions of <math>\Delta\psi + A(r^2)X\nabla\psi + C(r^2)\psi = 0</math></i> ...	217
William H. Mills, <i>Polynomials with minimal value sets</i> .....	225
George James Minty, Jr., <i>On the monotonicity of the gradient of a convex function</i> .....	243
George James Minty, Jr., <i>On the solvability of nonlinear functional equations of 'monotonic' type</i> .....	249
J. B. Muskat, <i>On the solvability of <math>x^e \equiv e \pmod{p}</math></i> .....	257
Zeev Nehari, <i>On an inequality of P. R. Bessack</i> .....	261
Raymond Moos Redheffer and Ernst Gabor Straus, <i>Degenerate elliptic equations</i> .....	265
Abraham Robinson, <i>On generalized limits and linear functionals</i> .....	269
Bernard W. Roos, <i>On a class of singular second order differential equations with a non linear parameter</i> .....	285
Tôru Saitô, <i>Ordered completely regular semigroups</i> .....	295
Edward Silverman, <i>A problem of least area</i> .....	309
Robert C. Sine, <i>Spectral decomposition of a class of operators</i> .....	333
Jonathan Dean Swift, <i>Chains and graphs of Ostrom planes</i> .....	353
John Griggs Thompson, <i>2-signalizers of finite groups</i> .....	363
Harold Widom, <i>On the spectrum of a Toeplitz operator</i> .....	365