

Pacific Journal of Mathematics

CLIFFORD VECTORS

CURTIS M. FULTON

CLIFFORD VECTORS

CURTIS M. FULTON

In this paper we present a generalization of parallel vector fields in a Riemannian space. As it turns out, such fields exist in spaces of constant positive curvature.

Restricting ourselves to a Riemannian 3-space throughout, we need the oriented third-order tensor [3, p. 249]

$$\eta_{ijh} = [\text{sgn}(g)g]^{1/2}\varepsilon_{ijh} .$$

whose covariant derivative vanishes [3, pp. 251-252]. The latter fact is best ascertained by the use of geodesic coordinates. If we write the determinant of the metric tensor with the aid of permutation symbols we also find without difficulty

$$(1) \quad g^{pq}\eta_{ijp}\eta_{kha} = g_{hj}g_{ik} - g_{hi}g_{jk} .$$

DEFINITION. *Let the direction of a vector field at any point be that of the unit vector V . The field is said to consist of Clifford vectors if*

$$(2) \quad V_{i,j} = L^n{}_{ijh} V^h , \quad L \neq 0 .$$

THEOREM. *If the Riemannian curvature K is constant and equal to L^2 , the system of equations (2) is completely integrable. If, at any point, solutions of (2) exist in all directions, then $K = L^2 = \text{const}$.*

It is known that integrability conditions for (2) are obtained using covariant differentiation. Hence, on account of a Ricci identity [3, p. 83] and (1) we have

$$(3) \quad L_{,k}\eta_{ijh} V^h - L_{,j}\eta_{ikh} V^h + L^2(g_{hj}g_{ik} - g_{hk}g_{ij}) V^h = R_{hijk} V^h .$$

If the Riemannian curvature is constant [3, p. 112],

$$(4) \quad R_{hijk} = K(g_{hj}g_{ik} - g_{hk}g_{ij})$$

and conditions (3) are identically satisfied. This proves the first part of our theorem.

For proof of the second part we multiply (3) by $W^i V^j W^k$ and get

$$L^2(g_{hj}g_{ik} - g_{hk}g_{ij}) V^h W^i V^j W^k = R_{hijk} V^h W^i V^j W^k .$$

Thus L^2 is the Riemannian curvature associated with the unit vectors

Received August 29, 1963.

V , W [3, p. 95]. Assume now that W is a solution of (2) and M the corresponding scalar factor. Then the above curvature is also equal to M^2 . Continuing this process we conclude from Schur's theorem [3, p. 112] that the curvature is constant and because of (4) that $K = L^2$.

To conclude, we demonstrate a geometric property of Clifford vectors justifying the name chosen for them. Let t be the unit tangent to a geodesic and U a unit vector which undergoes a parallel displacement along the geodesic. Hence $U^i{}_{,j}t^j = 0$ and U remains in a plane passing through the geodesic [1, p. 161]. On the other hand, because of (2), $V_{i,j}t^i t^j = 0$ which shows that a Clifford vector, propagated along the geodesic, is inclined at a constant angle to it. Letting $\cos \theta = U^i V_i$, we see that

$$-\sin \theta d_s \theta = L \eta_{ijh} U^i t^j V^h .$$

We now make the simplifying assumption that both U and V are perpendicular to t . In this case the vector $\eta_{ijh} U^i V^h$ has the direction of t and using (1) we find its length to be $\sin \theta$. Thus $d_s \theta = \pm L$ and the Clifford vector rotates about the geodesic in either sense through an angle proportional to the displacement. This property may be used to define the Clifford parallels or paratactic lines in elliptic 3-space [2, p. 108].

REFERENCES

1. A. Duschek, W. Mayer, *Lehrbuch der Differentialgeometrie*, Band II, Teubner, Leipzig und Berlin, 1930.
2. D. M. Y. Sommerville, *The elements of non-Euclidean geometry*, Dover, New York, 1958.
3. J. L. Synge, A. Schild, *Tensor calculus*, University of Toronto Press, Toronto, 1959.

UNIVERSITY OF CALIFORNIA, DAVIS

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

ROBERT OSSERMAN
Stanford University
Stanford, California

J. DUGUNDJI
University of Southern California
Los Angeles 7, California

M. G. ARSOVE
University of Washington
Seattle 5, Washington

LOWELL J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and on submission, must be accompanied by a separate author's résumé. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 14, No. 3

July, 1964

Erik Balslev and Theodore William Gamelin, <i>The essential spectrum of a class of ordinary differential operators</i>	755
James Henry Bramble and Lawrence Edward Payne, <i>Bounds for derivatives in elliptic boundary value problems</i>	777
Hugh D. Brunk, <i>Integral inequalities for functions with nondecreasing increments</i>	783
William Edward Christilles, <i>A result concerning integral binary quadratic forms</i>	795
Peter Crawley and Bjarni Jónsson, <i>Refinements for infinite direct decompositions of algebraic systems</i>	797
Don Deckard and Carl Mark Percy, <i>On continuous matrix-valued functions on a Stonian space</i>	857
Raymond Frank Dickman, Leonard Rubin and P. M. Swingle, <i>Another characterization of the n-sphere and related results</i>	871
Edgar Earle Enochs, <i>A note on reflexive modules</i>	879
Vladimir Filippenko, <i>On the reflection of harmonic functions and of solutions of the wave equation</i>	883
Derek Joseph Haggard Fuller, <i>Mappings of bounded characteristic into arbitrary Riemann surfaces</i>	895
Curtis M. Fulton, <i>Clifford vectors</i>	917
Irving Leonard Glicksberg, <i>Maximal algebras and a theorem of Radó</i>	919
Kyong Taik Hahn, <i>Minimum problems of Plateau type in the Bergman metric space</i>	943
A. Hayes, <i>A representation theory for a class of partially ordered rings</i>	957
J. M. C. Joshi, <i>On a generalized Stieltjes transform</i>	969
J. M. C. Joshi, <i>Inversion and representation theorems for a generalized Laplace transform</i>	977
Eugene Kay McLachlan, <i>Extremal elements of the convex cone B_n of functions</i>	987
Robert Alan Melter, <i>Contributions to Boolean geometry of p-rings</i>	995
James Ronald Retherford, <i>Basic sequences and the Paley-Wiener criterion</i>	1019
Dallas W. Sasser, <i>Quasi-positive operators</i>	1029
Oved Shisha, <i>On the structure of infrapolynomials with prescribed coefficients</i>	1039
Oved Shisha and Gerald Thomas Cargo, <i>On comparable means</i>	1053
Maurice Sion, <i>A characterization of weak* convergence</i>	1059
Morton Lincoln Slater and Robert James Thompson, <i>A permanent inequality for positive functions on the unit square</i>	1069
David A. Smith, <i>On fixed points of automorphisms of classical Lie algebras</i>	1079
Sherman K. Stein, <i>Homogeneous quasigroups</i>	1091
J. L. Walsh and Oved Shisha, <i>On the location of the zeros of some infrapolynomials with prescribed coefficients</i>	1103
Ronson Joseph Warne, <i>Homomorphisms of d-simple inverse semigroups with identity</i>	1111
Roy Westwick, <i>Linear transformations on Grassman spaces</i>	1123