

Pacific Journal of Mathematics

**MINIMUM PROBLEMS OF PLATEAU TYPE IN THE
BERGMAN METRIC SPACE**

KYONG TAIK HAHN

MINIMUM PROBLEMS OF PLATEAU TYPE IN THE BERGMAN METRIC SPACE

KYONG T. HAHN

Dedicated to my teacher Professor C. Loewner on his seventieth birthday

1. **Introduction.** In this paper we are concerned with the existence of minimal surfaces with respect to the B -area (see §4) and related problems in a bounded domain D in the space C^2 of two complex variables z_1, z_2 .

Let $K_D(z, \bar{z})$, $z = (z_1, \dots, z_n)$, be the Bergman kernel function of a bounded domain D in the space C^n of n complex variables. Throughout this paper, we assume $K_D(z, \bar{z})$ has the boundary value infinity at every point on the boundary of D . The kernel $K_D(z, \bar{z})$ enables us to define the Bergman metric

$$(1.1) \quad ds_D^2(z) = \sum_{\mu, \nu=1}^n T_{\mu\bar{\nu}}(z, \bar{z}) dz_\mu d\bar{z}_\nu, \quad T_{\mu\bar{\nu}} = \frac{\partial^2 \log K_D}{\partial z_\mu \partial \bar{z}_\nu},$$

which is invariant with respect to pseudo-conformal mappings [4, pp. 51–53]. Using (1.1) we construct (see §2) the complete Bergman metric space (D, d) over D and state a theorem for complete Riemannian spaces that for any two points in D , there exists a minimal curve with respect to d which connects the two points.

In §3 we show that, if D is a plane domain bounded by finitely many boundary components b_1, b_2, \dots, b_n , then there exists a minimal closed curve with respect to d among those curves which are homotopic to a fixed inner boundary component, say b_1 , in $\overline{D(b_1)}$ (see §3 for notation). If D is doubly connected, there exists a unique minimal closed curve in D . Furthermore, we prove a distortion theorem which gives bounds for the Bergman lengths of the minimal closed curves.

Analogous results are obtained in the case of two complex variables replacing the length by the B -area.

For a closed Jordan curve Γ in a complete metric space (D, d) , we ask whether there exists a minimal surface with respect to the B -area which spans Γ . Answers to this question which constitute the main result of this paper are given in §4.

As a generalization of §3, we consider a domain D which is topologically equivalent to a product domain of the form $D_1 \times D_2$,

Received July 5, 1963. This paper represents the essential contents of the author's doctor thesis at Stanford University. The author wishes to express his thanks to Professor S. Bergman and Professor C. Loewner for their pertinent criticism in the preparation of this paper. This research was supported by Navy Contract Nonr-225(11) and NSF Grant 21344.

where D_k is a bounded domain as considered in § 3. When does there exist a minimal closed surface with respect to the B -area among those surfaces which are homotopic to T_1 in $\overline{D}(T_1)$ (see § 5 for notation)?

Answers are given in § 5. Distortion theorems for the minimal surfaces are given in § 6.

2. The Bergman metric space. A (continuous) curve c in D is said to be *regular* if it admits a regular (parametric) representation, i.e., there exists a continuously differentiable representation

$$(2.1) \quad G|I: z_k = G_k(t), k = 1, 2, \dots, n, t \in I = [a, b],$$

and dG_k/dt never vanish simultaneously at any $t \in I$. A curve c in D is said to be *piecewise regular* if it admits a piecewise regular representation, i.e., there exists a partition $\Delta: a = t_0 < t_1 < \dots < t_{m-1} < t_m = b$ such that $G| [t_{k-1}, t_k]$ is regular for $k = 1, 2, \dots, m$.

For a piecewise regular curve c given by (2.1), we define

$$(2.2) \quad L_D(c) = \int_a^b \left[\sum_{\mu, \nu=1}^n T_{\mu\nu}(G(t), \overline{G(t)}) \frac{dG_\mu}{dt} \frac{d\overline{G}_\nu}{dt} \right]^{1/2} dt.$$

$L_D(c)$ is independent of the choice of piecewise regular representations of c . $L_D(c)$ will be called the *Bergman length* of c .

For any two points z^1 and z^2 in D , we define a distance function d by

$$(2.3) \quad d(z^1, z^2) = \inf_c L_D(c),$$

where c runs over all piecewise regular curves which connect z^1 and z^2 . Then the following theorem holds [15, § 16].

THEOREM 2.1. *d satisfies all the axioms for a metric and the metric space (D, d) is topologically equivalent to the metric space (D, ρ) with the Euclidean metric ρ . Moreover, the metric space (D, d) is finitely connected in the sense that every pair of points in D can be connected by a curve of finite Bergman length.*

The metric space (D, d) will be called the *Bergman metric space* over D . The significance of this metric space is that all metric properties are invariant under pseudo-conformal mappings.

We define the length (generalized) of a continuous curve c in D in the following way: For a partition $\Delta(I) = \{I_1, I_2, \dots, I_m\}$, $I_k = [t_{k-1}, t_k]$, $k = 1, 2, \dots, m$, of I , we define

$$\sigma(G; \Delta(I)) = \sum_{k=1}^m \sigma(G; I_k), \quad \sigma(G; I_k) = d(G(t_k), G(t_{k-1})).$$

Further, we define

$$\mathcal{L}_D(c) = \sup_{\Delta} \sigma(G; \Delta(I)),$$

where Δ runs over all possible partitions of I . Then $\mathcal{L}_D(c)$ is independent of the choice of continuous representations of c . Clearly, the functional \mathcal{L}_D is lower semi-continuous, i.e.,

$$\mathcal{L}_D(c) \leq \liminf_{k \rightarrow \infty} \mathcal{L}_D(c_k), \text{ if } c_k \rightarrow c.$$

Further, for every piecewise regular curve c , $L_D(c) = \mathcal{L}_D(c)$ [15, § 16].

If $\mathcal{L}_D(c) < \infty$, c is said to be *rectifiable*. A curve is said to be *completely degenerated* if there is a representation $G|I$ such that G is constant on I . A representation $G|I$ is said to be *normal* if $\mathcal{L}_D(G; [t, t']) = t' - t$, for $t, t' \in I, t < t'$.

Let c be a rectifiable curve which is not completely degenerated. Then c admits a normal representation $G|[0, \mathcal{L}_D(c)]$. If we set $F(t) = G(t, \mathcal{L}_D(c)), t \in I_0, I_0 = [0, 1]$, then $F|I_0$ is also a representation of c . Such a representation $F|I_0$ is called a *reduced representation* of c . For a closed curve, F is defined on $(-\infty, \infty)$ and is periodic of period 1. It is, therefore, enough to consider F on I_0 . If $F|I_0$ is a reduced representation of a curve c , then the inequality

$$(2.4) \quad d(F(t), F(t')) \leq \mathcal{L}_D(c) |t - t'|$$

holds for every $t, t' \in I_0$.

A metric space is called *complete* if every bounded infinite subset contains a limit point in the metric space. If D is homogeneous, (D, d) is always complete. Further, for every bounded generalized analytic polyhedron $D, (D, d)$ is complete. This is a result of S. Kobayashi (see [11] for details). For domains D in the space $C^n, n \leq 2$, Bergman has shown that the distance from a point in D to the boundary becomes infinite under certain hypothesis on the boundary of D [1], [6, Chap. III]. It is clear, in this case, that the metric space (D, d) is complete. Without going into great details in this direction, we shall assume in the sequel that the metric space (D, d) is always complete.

A curve K in (D, d) which connects z^1 and z^2 in D is called a *minimal curve* between z^1 and z^2 if $\mathcal{L}_D(K) \leq \mathcal{L}_D(c)$ for all curves c connecting z^1 and z^2 .

THEOREM 2.2. *For any two points z^1 and $z^2, z^1 \neq z^2$, in (D, d) , there exists a minimal curve K between z^1 and z^2 . Further, the*

minimal curves are analytic (see [10] or [15, § 17]).

3. The existence of a minimal closed curve in a plane domain and its distortion theorem. We consider a multiply connected bounded domain D in the space C^1 bounded by N Jordan closed curves b_1, b_2, \dots, b_n , where b_n is the outer boundary component. Let (D, d) be the Bergman metric space derived from the Bergman metric

$$(3.1) \quad ds_D^2(z) = K_D(z, \bar{z}) |dz|^2 .$$

It is assumed that (D, d) is complete. Then all the previous considerations, lemmas and theorems can be carried over to this case. We fix an inner boundary component of D , say b_1 . Without loss of generality, we may assume b_1 to be a circle.

Let $\mathfrak{R}(D; b_1)$ be the class of all closed continuous curves c in D which are homotopic to b_1 in $\overline{D(b_1)}$, where $D(b_1)$ is a ring domain bounded by b_1 and b_n which contains the domain D , and $\overline{D(b_1)}$ is the closure of $D(b_1)$. A curve $K(D; b_1)$ in $\mathfrak{R}(D; b_1)$ which satisfies the condition $\mathcal{L}_D(K(D; b_1)) \leq \mathcal{L}_D(c)$ for all $c \in \mathfrak{R}(D; b_1)$, will be called a *minimal closed curve* of D with respect to b_1 . Due to the completeness of $(D; d)$ and the behavior of $K_D(z, \bar{z})$ (described on page 943) on the boundary of D , we have

THEOREM 3.1. *There exists a minimal closed curve $K(D; b_1)$ of the domain D with respect to b_1 . Further, it is analytic.*

Proof. Let $\gamma = \inf_c \mathcal{L}_D(c)$, where c runs over the class $\mathfrak{R}(D; b_1)$. Then $0 < \gamma < \infty$. There exists a minimizing sequence $\{c_k\}$ of rectifiable curves in $\mathfrak{R}(D; b_1)$. Let $G_k|_{I_0}$ be the reduced representation of c_k . By (2.4), we have

$$d(G_k(t), G_k(t')) \leq \mathcal{L}_D(c_k) |t - t'| \text{ for each } k ,$$

and $\{\mathcal{L}_D(c_k)\}$ has an upper bound δ which is finite. We choose an M such that $M^{1/2} > \delta/l(b_1)$, $l(b_1)$ is the Euclidean length of b_1 . Then no c_k lies completely in $D - D_M$, $D_M = [z | K_D(z, \bar{z}) \leq M]$. Let

$$\rho = \max_{z_1, z_2 \in D_M} d(z_1, z_2) ,$$

then for every pair of positive integers p and q , we have $d(G_p(t), G_q(t)) < \rho + 2\delta, 0 \leq t \leq 1$. Hence, we can select a subsequence $\{G_{k_i}\}$ of $\{G_k\}$ which converges uniformly to a continuous function G^0 on I_0 . Let K be the closed curve whose representation is given by $G^0|_{I_0}$. Since $c_{k_i} \rightarrow K$, and by the lower semi-continuity of \mathcal{L}_D , we obtain $\mathcal{L}_D(K) = \gamma$. The analyticity of K is obvious.

THEOREM 3.2. *Every doubly connected domain has a unique minimal closed curve. It is analytic.*

Proof. We shall show first that annulus $Q = [z \mid r < |z| < 1]$ has a unique minimal closed curve given by $c_0 = [z \mid |z| = r^{1/2}]$. Let $P_1 = [z \mid r < |z| < r^{1/2}]$, $P_2 = [z \mid r^{1/2} < |z| < 1]$. If $c \cap P_1 = \phi$, it is immediate that $L_Q(c_0) \leq L_Q(c)$, since the kernel function $K_Q(z, \bar{z})^1$ assumes its minimum on c_0 . If $c \cap P_2 = \phi$, by the conformal mapping $\zeta = r/z$, we have $\tilde{c} \cap P_1 = \phi$, where \tilde{c} is the image curve of c under $\zeta = r/z$. Since $L_Q(\tilde{c}) = L_Q(c)$, $L_Q(c_0) \leq L_Q(c)$ follows. If $c \cap P_1 \neq \phi$ and $c \cap P_2 \neq \phi$, we obtain two closed curves c_1, c_2 consisting of the subarcs of c and c_0 and such that $c_1 \cap P_2 = \phi$, $c_2 \cap P_1 = \phi$. By the previous arguments, $L_Q(c_i) \geq L_Q(c_0)$, $i = 1, 2$. Since $L_Q(c_1) + L_Q(c_2) = L_Q(c) + L_Q(c_0)$, we have $L_Q(c_0) \leq L_Q(c)$. Let D be a doubly connected domain. Then D can be mapped by a univalent analytic function $f(z)$ onto Q . It is clear that $f^{-1}(c_0)$ is the unique minimal closed curve of D with respect to the inner boundary component by the univalence of $f(z)$.

We consider a domain D in the z -plane which is bounded by $b_1 = [z \mid |z| = r]$, $b_N = [z \mid |z| = 1]$, and $(N - 2)$ closed Jordan curves b_2, \dots, b_{N-1} . The curves b_2, \dots, b_{n-1} lie in the domain bounded by b_1 and b_N .

Let $A_1 = [z \mid r < |z| < 1]$, $A_2 = [z \mid |z - a| < \rho, |z| > r]^2$, be exterior and interior domains of comparison for D , respectively, i.e., $A_1 \supset D \supset A_2$. Then

$$(3.2) \quad L_{A_1}(K(A_1)) \leq L_D(K(D)) \leq L_{A_2}(K(A_2)) ,$$

where $K(A_1), K(A_2)$ and $K(D)$ are minimal closed curves of A_1, A_2 and D with respect to b_1 , respectively. It is an immediate consequence of the fact that if $B \subset A$, then $K_B(z, \bar{z}) \geq K_A(z, \bar{z})$ for $z \in B$. The linear transformation

$$(3.3) \quad w = \frac{z - (a + \rho d)}{\rho - d(z - a)} , \quad 0 < |a| \leq \rho - r ,$$

maps A onto $Q_R = [z \mid R < |z| < 1]$, where R is given by

¹ A simple computation shows that the kernel function of Q ,

$$K_Q(z, z) = \frac{1}{\pi |z|^2} \left[\wp(2 \log |z|; -2 \log r, 2\pi i) + \frac{\zeta(\pi i; -2 \log r, 2\pi i)}{\pi i} \right]$$

(see [9], [18]), where \wp and ζ are the Weierstrass elliptic functions, assumes its minimum on c_0 .

² Here we choose a and ρ in such a way that $|z - a| < \rho$ contains b_1 but no other $b_k, k = 2, \dots, N$, and A_2 to be the largest among such domains.

$$(3.4) \quad R = \left[\frac{r^2 - (a + \rho d)^2}{(\rho + ad)^2 - r^2 d^2} \right]^{1/2},$$

$$d = \frac{r^2 - a^2 - \rho^2 + [(r^2 - a^2 - \rho^2)^2 - 4a^2 \rho^2]^{1/2}}{2a\rho}.$$

Since $L_{A_2}(K(A_2)) = L_{Q_R}(K(Q_R))$, using (3.2), we obtain

THEOREM 3.3. $E(r) \leq (1/2)L_D(K(D)) \leq E(R)$,
 where R is given by (3.4) and

$$E(r) = [\pi\wp(\log r; -2 \log r, 2\pi i) - i\zeta(\pi i; -2 \log r, 2\pi i)]^{1/2},$$

\wp and ζ are the Weierstrass elliptic functions.

The estimation of the bounds for the Bergman lengths of the minimal closed curves in Theorem 3.3 seems to be done only for a special domain. However, every multiply connected domain can always be mapped onto such a domain by a conformal mapping. Therefore, if we know the geometry of a given domain D , combining the various distortion theorems in the theory of conformal mappings and the result in Theorem 3.3, we can obtain various bounds for the Bergman lengths of the minimal curves for quite general domains.

4. The existence of a minimal surface which spans a given closed curve in (D, d) . A surface S in the space C^2 is said to be *continuously differentiable* if it admits a continuously differentiable representation

$$G | Q_0 : z_k = G_k(u_1, u_2), k = 1, 2, (u_1, u_2) \in Q_0 = [0 \leq u_1, u_2 \leq 1].$$

A surface S is said to be *piecewise continuously differentiable* if it admits a piecewise continuously differentiable representation $G | Q_0$, i.e., there exists a partition $\Delta = \{A_1, A_2, \dots, A_m\}$ of Q_0 by rectilinear triangles A_k such that $G | A_k$ is continuously differentiable, $k = 1, 2, \dots, m$. The *ordinary B-area element* at a point (z_1, z_2) on a piecewise continuously differentiable surface S is defined by the equation [6, Chap. XI]

$$(4.1) \quad db_S(z) = \left| \frac{\partial(G_1, G_2)}{\partial(u_1, u_2)} \right| du_1 du_2.$$

The ordinary area element of S is given by the equation

$$(4.2) \quad da_S(z) = [g_{11}g_{22} - (Re g_{12})^2]^{1/2} du_1 du_2,$$

$$g_{\alpha\beta} = \sum_{i=1}^2 \frac{\partial G_i}{\partial u_\alpha} \frac{\partial G_i}{\partial u_\beta}, \alpha, \beta = 1, 2.$$

Further (4.1) can also be written in the following form,

$$(4.1)' \quad db_s(z) = [g_{11}g_{22} - |g_{12}|^2]^{1/2} du_1 du_2 .$$

Therefore, $da_s(z) \geq db_s(z)$ at every point $z \in S$; the equality holds if and only if $Im g_{12} = 0$.

For a piecewise continuously differentiable surface S , the ordinary B -area is defined and given by the equation

$$(4.3) \quad b(S) = \iint_{Q_0} \left| \frac{\partial(G_1, G_2)}{\partial(u_1, u_2)} \right| du_1 du_2 .$$

$b(S)$ is independent of the choice of piecewise continuously differentiable representations $G|Q_0$ of S . A surface S is said to be *analytic* if it admits an analytic representation $G|Q_0$, i.e., $\partial G_k / \partial \bar{w} = 0, k = 1, 2, w = u_1 + iu_2$.

For an analytic or an anti-analytic surface $S, b(S) = 0$. It is also clear that $b(S) = 0$ if and only if the tangent plane of S at every point is an analytic plane. A simple computation shows the following lemma:

LEMMA 4.1. *The following three conditions are equivalent:*

- 1) $b(S) = a(S),$
- 2) $\frac{\partial(G_1, \bar{G}_1)}{\partial(u_1, u_2)} + \frac{\partial(G_2, \bar{G}_2)}{\partial(u_1, u_2)} = 0$ at each point on $S,$
- 3) $\oint_c \bar{G}_1 dG_1 + \bar{G}_2 dG_2 = 0$ for every closed curve c on S .

Let D be a bounded domain in the space C^2 on which (D, d) is complete. The quantity

$$(4.4) \quad dB_D(z) = [K_D(z, \bar{z})]^{1/2} db_s(z), z = (z_1, z_2),$$

is invariant with respect to pseudo-conformal mappings and a monotone decreasing functional of D [6]. $dB_D(z)$ is called the *invariant B-area element* of S . For a piecewise continuously differentiable surface S in D , the invariant B -area of S is defined and given by the equation

$$(4.5) \quad B_D(S) = \iint_{Q_0} [K_D(G, \bar{G})]^{1/2} \left| \frac{\partial(G_1, G_2)}{\partial(u_1, u_2)} \right| du_1 du_2$$

and is independent of the choice of piecewise continuously differentiable representations $G|Q_0$ of S .

A surface S in D is said to satisfy the *condition (L) with respect to the metric d* if there exists a representation $G|Q_0$ of S for which there exists a constant $L(S) > 0$ depending only on S and satisfying the inequality

$$(4.6) \quad \mathcal{L}_D(G; \sigma(w_1, w_2)) \leq L(S) |w_1 - w_2|$$

for every pair of points w_1, w_2 in Q_0 ; here $\sigma(w_1, w_2)$ is the line segment that joins w_1 and w_2 in Q_0 , $w_k = u_1^{(k)} + iu_2^{(k)}$, $k = 1, 2$.

It is clear that $G | \partial(Q_0)$, where $\partial(Q_0)$ is the boundary of Q_0 , is a representation of the boundary curve Γ of S and that Γ is rectifiable. It is also clear that every continuously differentiable surface S satisfies the condition (L) with respect to d .

We shall say that a surface S is of class $C'\mathfrak{R}(L, N, \Gamma)$ if S admits a continuously differentiable representation

$$G | Q_0 : z_k = G_k(w), \quad k = 1, 2, w \in Q_0,$$

which satisfies the following conditions:

- (a) for a fixed positive constant L , $L(S) \leq L$,
- (b) for a fixed positive constant N ,

$$\left| \frac{\partial G(w_1)}{\partial u_j} - \frac{\partial G(w_2)}{\partial u_j} \right| \leq N |w_1 - w_2|, \quad j = 1, 2, G = (G_1, G_2),$$

for every pair of points w_1, w_2 in Q_0 ,

- (c) S spans a preassigned closed Jordan curve Γ in D in such a way that G is a one-to-one mapping on $\partial(Q_0)$.

A surface S_m is called *minimal surface* of the class $C'\mathfrak{R}(L, N, \Gamma)$ if $B_D(S_m) \leq B_D(S)$ for all $S \in C'\mathfrak{R}(L, N, \Gamma)$.

THEOREM 4.1.³ *For each L and N for which the class $C'\mathfrak{R}(L, N, \Gamma)$ is not empty, there exists a minimal surface S_m in the class.*

Proof. Let $\inf_S B_D(S) = \gamma$, where S runs over all surfaces in $C'\mathfrak{R}(L, N, \Gamma)$. Then $0 \leq \gamma < \infty$. Hence, there exists a minimizing sequence $\{S_n\}$. Let $G^n | Q_0$ be a representation of S_n which satisfies conditions (a), (b) and (c). From (a) it follows that for any pair of positive integers p, q ,

$$d(G^p(w), G^q(w)) \leq 2 \cdot 2^{p/2} L.$$

Therefore, $\{G^n(w)\}$ is equi-bounded. The equi-continuity of $\{G^n(w)\}$ follows from the inequality

$$(4.7) \quad d(G^n(w), G^n(w')) \leq L |w - w'| \quad \text{for any } w, w' \in Q_0 \text{ and all } n.$$

Hence, we can select a subsequence $\{G^m(w)\}$ of $\{G^n(w)\}$ which converges uniformly to a continuous function $G^0(w)$ defined in Q_0 . Let $G^0 | Q_0$ define a surface S_0 . Then it is clear that S_0 spans Γ in such a way that G^0 is a one-to-one mapping on $\partial(Q_0)$. The family $\{\partial G^m / \partial u_j\}$

³ Replacing (a) by the condition (a') on page 951, a result similar to Theorem 4.1 can be given (see Corollary 2).

of continuous functions $\partial G^m/\partial u_j$ is equi-bounded and equi-continuous by (b) for $j = 1, 2$. Therefore, we can select a subsequence $\{G^{m_i}(w)\}$ of $\{G^m(w)\}$ which converges uniformly to $G^0(w)$ and such that $\{\partial G^{m_i}/\partial u_j\}$ converges uniformly to a continuous function $\partial G^0/\partial u_j$ for $j = 1, 2$. This implies that S_0 is a continuously differentiable surface. In order to show $S_0 \in C'\mathfrak{R}(L, N, \Gamma)$, let c_{m_i} and c_0 be the image curves of a line segment $\sigma(w_1, w_2)$ which connects two points w_1 and w_2 in Q_0 under $G^{m_i}(w)$ and $G^0(w)$, respectively. Then c_{m_i} converges to c_0 and, hence, $\liminf_{i \rightarrow \infty} L_D(c_{m_i}) \geq L_D(c_0)$ by the lower semi-continuity of L_D . Since $L_D(c_{m_i}) \leq L|w_1 - w_2|$ for all m_i , $L_D(c_0) \leq L|w_1 - w_2|$. It is clear that $G^0(w)$ satisfies (b). Since the functional B_D is lower semi-continuous in $C'\mathfrak{R}(L, N, \Gamma)$ and $S_0 \in C'\mathfrak{R}(L, N, \Gamma)$, we have $B_D(S_0) = \gamma$. Thus S_0 is a minimal surface in the class $C'\mathfrak{R}(L, N, \Gamma)$.

REMARK. In the case that Γ lies on an analytic plane π and the portion $\tilde{\pi}$ of $\pi \cap D$ enclosed by Γ is simply connected, $\tilde{\pi}$ is a minimal surface of $C'\mathfrak{R}(L, N, \Gamma)$ with some L and N , and $B_D(S_0) = 0$. In general, if there exists an analytic surface S in D which spans Γ , then S is a minimal surface with some L and N , and $B_0(S) = 0$.

Let $C'\mathfrak{R}(N, \Gamma)$ be the class of continuously differentiable surfaces in the space C^2 which span a preassigned Jordan closed curve Γ in C^2 and satisfy the condition (b). Then (b) implies condition (a) with respect to the Euclidean metric ρ for every surface in $C'\mathfrak{R}(N, \Gamma)$. Since C^2 is complete with respect to ρ , the following corollary follows by the same procedure as in Theorem 4.1.

COROLLARY 1. *In the class $C'\mathfrak{R}(N, \Gamma)$, there exists a minimal surface S'_m in the sense that*

$$b(S'_m) \leq b(S) \text{ for all } S \in C'\mathfrak{R}(N, \Gamma).$$

Let $C'\mathfrak{R}_\alpha(N, \Gamma)$ be the class of continuously differentiable surfaces S in D which satisfy conditions (b), (c) and

$$(a') \text{ for a preassigned real number } \alpha, 0 \leq \alpha \leq 1,$$

$$(4.8) \quad \frac{db_s(z)}{da_s(z)} \geq \alpha \text{ at every point } z \in S.$$

We notice that the class $C'\mathfrak{R}_\alpha(N, \Gamma)$ is motone decreasing with respect to α .

COROLLARY 2. *For a fixed $\alpha > 0$ and N for which $C'\mathfrak{R}_\alpha(N, \Gamma)$ is not empty there exists a minimal surface in the class.*

Proof. The B -areas $B_D(S_\alpha)$ of S_α which belong to a minimizing

sequence $\{S_n\}$ have a fixed upper bound. Therefore, condition (a') ensures the existence of an $M > 0$ such that every S_n lies completely in $D_M, D_M = [z \mid K_D(z, \bar{z}) \leq M]$. This implies condition (a) with some L , which depends on α and N . Hence, the corollary follows from the theorem.

5. **The existence of minimal closed surfaces in (D, d) .** Let D_k be a domain in the space of one complex variable z_k bounded by n_k closed curves $b_1^{(k)}, b_2^{(k)}, \dots, b_{n_k}^{(k)}$. Here $b_{n_k}^{(k)}$ is the outer boundary component of D_k and $b_1^{(k)}$ is an inner boundary component, which is a circle, i.e., $b_1^{(k)} = [z_k \mid |z_k| = r_k]$.

Let D be a domain in the space C^2 which is topologically equivalent to the product domain $\tilde{D} = D_1 \times D_2$, and T_1 the topological image of $\tilde{T}_1 = b_1^{(1)} \times b_1^{(2)}$. A surface S in D which is homotopic to T_1 in $\overline{D(T_1)}$, where $D(T_1)$ is the topological image of $\tilde{D}(\tilde{T}_1) = D_1(b_1^{(1)}) \times D_2(b_1^{(2)})$ (see § 3 for notation), is a closed surface of the torus type and, hence, admits a doubly periodic representation

$$G \mid R^2 : z_k = G_k(u_1, u_2), k = 1, 2, (u_1, u_2) \in R^2, \\ R^2 = (-\infty < u_1, u_2 < +\infty),$$

of periods 1. For our purposes, therefore, it is enough to consider G on the unit square Q_0 as a representation of S .

We shall say that a closed surface S is of class $C' \mathfrak{R}_\alpha(N, T_1)$ if S is homotopic to T_1 in $\overline{D(T_1)}$ and admits a continuously differentiable representation $G \mid Q_0$ satisfying condition (a') and (b) in § 4. By the same procedure as in Corollary 2 of Theorem 4.1, we can prove the following theorem for any fixed $\alpha > 0$.

THEOREM 5.1. *For each N for which the class $C' \mathfrak{R}_\alpha(N, T_1)$ is not empty, there exists a minimal closed surface $S_m(D)$ in the class.*

Let $D' \mathfrak{R}(\tilde{D}, \tilde{T}_1)$ be the class of all closed surfaces S of the form $S = c_1 \times c_2$ in D , where c_k is a piecewise continuously differentiable closed curve in D_k which is homotopic to $b_1^{(k)}$ in $\overline{D_k(b_1^{(k)})}$. For each $S \in D' \mathfrak{R}(\tilde{D}, \tilde{T}_1)$, we have $B_{\tilde{D}}(S) = L_{D_1}(c_1) \cdot L_{D_2}(c_2)$. It follows from the fact that $K_{\tilde{D}}(z, \bar{z}) = K_{D_1}(z_1, \bar{z}_1) \cdot K_{D_2}(z_2, \bar{z}_2)$ [7]. Therefore, the following is an immediate consequence of Theorem 3.1.

THEOREM 5.2. *There exists a minimal surface $S_m(\tilde{D})$ of the class $D' \mathfrak{R}(\tilde{D}, \tilde{T}_1)$. It is given by $K(D_1) \times K(D_2)$, where $K(D_k)$ is a minimal closed curve of D_k with respect to $b_1^{(k)}$.*

Let $A = A_1 \times A_2$, where A_k is a doubly connected plane domain in

the z_k -plane. Let $D'\mathfrak{R}_1(A, T)$ be the class of piecewise continuously differentiable closed surfaces S in A which are homotopic to $T = b_1^{(1)} \times b_2^{(2)}$ in \bar{A} , where $b_1^{(k)}$ is the inner boundary component of A_k , and satisfy the condition $da_s(z) = db_s(z)$.⁴ Then the following theorem holds:

THEOREM 5.3. *There exists a unique minimal closed surface in the class $D'\mathfrak{R}_1(A, T)$. It is given by $K(A_1) \times K(A_2)$, where $K(A_k)$ is a minimal closed curve of A_k with respect to $b_1^{(k)}$.*

Proof. Let $A = Q = Q_1 \times Q_2, Q_k = [z_k \mid r_k < |z_k| < 1]$. We shall show $S_m(Q) = K(Q_1) \times K(Q_2)$ is a unique minimal closed surface of $D'\mathfrak{R}_1(Q, T)$. Let $P_{1k} = [z_k \mid r_k < |z_k| < r_k^{1/2}]$, $P_{2k} = [z_k \mid r_k^{1/2} \leq |z_k| < 1]$. If $S \in P_{21} \times P_{22}$, it is immediate that $B_Q(S) \geq B_Q(S_m)$. For any $S \in D'\mathfrak{R}_1(Q, T)$, S can be replaced by a surface $\tilde{S} \in D'\mathfrak{R}_1(Q, T)$ with $B_Q(S) = B_Q(\tilde{S})$ and lying in $P_{21} \times P_{22}$ by the pseudo-conformal mapping $z_k^* = r_k/z_k, k = 1, 2$. Thus, $B_Q(S) \geq B_Q(S_m)$ for every $S \in D'\mathfrak{R}_1(Q, T)$. There exists a univalent analytic function $f_k(z_k)$ which maps A_k onto Q_k . Therefore, the pseudo-conformal mapping $w_k = f_k^{-1}(z_k)$ maps A onto Q and, hence, $S_m(Q)$ onto $S_m(A), S_m(A) = K(A_1) \times K(A_2)$. The uniqueness of $S_m(A)$ is clear.

6. Bounds for the B -areas of minimal closed surfaces in the space (D, d) . Using the method of exterior and interior domains of comparison, various bounds for the B -areas of minimal surfaces can be obtained. As we have considered in § 3, let D_k be bounded by $b_1^{(k)} = [z_k \mid |z_k| = r_k], b_{n_k}^{(k)} = [z_k \mid |z_k| = 1]$, and $(n_k - 2)$ closed Jordan curves $b_2^{(k)}, \dots, b_{n_k-1}^{(k)}$, which lie in the domain bounded by $b_1^{(k)}$ and $b_{n_k}^{(k)}$. Let $A_{1k} = [z_k \mid r_k < |z_k| < 1], A_{2k} = [z_k \mid |z_k - a_k| < \rho_k, |z_k| > r_k], 0 < |a_k| \leq \rho_k - r_k$, be exterior and interior domains of comparison for D_k , respectively. Then $A_j = A_{j1} \times A_{j2}$ can be used as exterior and interior domains of comparison of $D = D_1 \times D_2$, i.e., $A_1 \supset D \supset A_2$. Let $S_m(\tilde{D})$ and $S_m(A_j)$ be minimal surfaces of the classes $D'\mathfrak{R}(\tilde{D}, \tilde{T}_1)$ and $D'\mathfrak{R}(A_j, \tilde{T}_1)$, respectively. Then $B_{A_1}(S_m(A_1)) \leq B_{\tilde{D}}(S_m(D)) \leq B_{A_2}(S_m(A_2))$. Using this inequality, we have the following distortion theorem for minimal surfaces of the class $D'\mathfrak{R}(\tilde{D}, \tilde{T}_1)$.

THEOREM 6.1. $\prod_{k=1}^2 E(r_k) \leq (1/4)B_{\tilde{D}}(S_m(\tilde{D}, \tilde{T}_1)) \leq \prod_{k=1}^2 E(R_k)$, where R_k is given in (3.4) with the corresponding subscript k and $E(r)$ is given in Theorem 3.3.

By a construction of an interior domain of comparison for D in

⁴ This is the case $\alpha = 1$ in (a') (see (4.8) and Lemma 4.1).

Theorem 5.1, we can also obtain a distortion theorem for minimal surfaces $S_m(D)$ in Theorem 5.1 which gives us an upper bound. Suppose an interior domain of comparison for D is given by A_{2k} , then we have

THEOREM 6.2. $B_D(S_m(D)) \leq 4 \prod_{k=1}^2 E(R_k)$, where R_k and $E(r)$ are given as in Theorem 6.1.

REMARK. For the product domain $Q = Q_1 \times Q_2$ of two annuli Q_1 and Q_2 , $K(Q_1) \times K(Q_2)$ is not necessarily a minimal surface for the class $D'\mathfrak{R}_\alpha(Q, T)$ for a fixed α , $0 < \alpha < 1$.

BIBLIOGRAPHY

1. S. Bergman, *Über die Kernfunktion eines Bereiches und ihr Verhalten am Rande*, J. Reine Angew. Math., **169** (1933), 1-42, and **172** (1934), 89-128.
2. ———, *Sur quelques propriétés des transformations par un couple de fonctions de deux variables complexes*, Rend. Della R. Accad. Naz. Del Lincei, **19** (ser. 6) (1934), 474-478.
3. ———, *Zur Theorie von Pseudokonformen Abbildungen*, Recueil Math., nouv., **1** (43), (1936), 76-96.
4. ———, *Sur les fonctions orthogonales de plusieurs variables complexes*, Mém. des Sciences Math., **106** (1947).
5. ———, *Sur la fonction-noyau d'un domaine et ses applications dans la théorie des transformations pseudo-conformes*, Mém. des Sciences Math., **108** (1948).
6. ———, *The kernel function and conformal mapping*, Math. Surveys 5, Amer. Math. Soc., New York, 1950.
7. H. J. Bremermann, *Holomorphic continuation of the kernel function and the Bergman metric in several complex variables*, in Lectures on functions of a complex variable ed. by W. Kaplan, Univ. of Michigan Press, 1955.
8. H. Busemann, and M. Mayer, *On the foundations of calculus of variations*, Trans. Amer. Math. Soc., **49** (1941), 173-198.
9. B. Epstein, *The kernel-function and conformal invariants*, J. Math. and Mech., **7** (1958), 925-936.
10. D. Hilbert, *Über das Dirichletsche Princip*, Jahr. Dtsch. Math.-Verein., **8** (1900), 184-188.
11. S. Kobayashi, *Geometry of Bounded Domains*, Trans. Amer. Math. Soc., **92** (1959), 267-290.
12. C. B. Morrey Jr., *Quasi-convexity and the lower semi-continuity of multiple integrals*, Pacific J. Math., **2**, (1952), 25-53.
13. ———, *The parametric variational problem for double integrals*, Communications on Pure and Applied Math., **14** (1961), 569-575.
14. K. Reinhardt, *Über Abbildungen durch analytische Funktionen zweier Veränderlicher*, Math. Ann., **83** (1921), 211-255.
15. W. Rinow, *Die Innere Geometrie der Metrischen Räume*, Die Grundlehren der Math. Wissenschaften **105** (1961).
16. J. M. Stark, *On distortion in pseudo conformal mappings*, Pacific J. Math., **6** (1956), 565-582.
17. ———, *Minimum problems in the theory of pseudo-conformal transformations*

and their application to estimation of the curvature of the invariant metric, Pacific J. Math., **10** (1960), 1021-1038.

18. K. Zarankiewicz, *Über ein numerisches Verfahren zur konformen Abbildung zweifach zusammenhängender Gebiete*, Zeits. f. ang. Math. u. Mech. **14** (1934), 97-104.

STANFORD UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

ROBERT OSSERMAN
Stanford University
Stanford, California

J. DUGUNDJI
University of Southern California
Los Angeles 7, California

M. G. ARSOVE
University of Washington
Seattle 5, Washington

LOWELL J. PAIGE
University of California
Los Angeles 24, California

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CALIFORNIA RESEARCH CORPORATION
SPACE TECHNOLOGY LABORATORIES
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced), and on submission, must be accompanied by a separate author's résumé. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, L. J. Paige at the University of California, Los Angeles 24, California.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 14, No. 3

July, 1964

Erik Balslev and Theodore William Gamelin, <i>The essential spectrum of a class of ordinary differential operators</i>	755
James Henry Bramble and Lawrence Edward Payne, <i>Bounds for derivatives in elliptic boundary value problems</i>	777
Hugh D. Brunk, <i>Integral inequalities for functions with nondecreasing increments</i>	783
William Edward Christilles, <i>A result concerning integral binary quadratic forms</i>	795
Peter Crawley and Bjarni Jónsson, <i>Refinements for infinite direct decompositions of algebraic systems</i>	797
Don Deckard and Carl Mark Percy, <i>On continuous matrix-valued functions on a Stonian space</i>	857
Raymond Frank Dickman, Leonard Rubin and P. M. Swingle, <i>Another characterization of the n-sphere and related results</i>	871
Edgar Earle Enochs, <i>A note on reflexive modules</i>	879
Vladimir Filippenko, <i>On the reflection of harmonic functions and of solutions of the wave equation</i>	883
Derek Joseph Haggard Fuller, <i>Mappings of bounded characteristic into arbitrary Riemann surfaces</i>	895
Curtis M. Fulton, <i>Clifford vectors</i>	917
Irving Leonard Glicksberg, <i>Maximal algebras and a theorem of Radó</i>	919
Kyong Taik Hahn, <i>Minimum problems of Plateau type in the Bergman metric space</i>	943
A. Hayes, <i>A representation theory for a class of partially ordered rings</i>	957
J. M. C. Joshi, <i>On a generalized Stieltjes transform</i>	969
J. M. C. Joshi, <i>Inversion and representation theorems for a generalized Laplace transform</i>	977
Eugene Kay McLachlan, <i>Extremal elements of the convex cone B_n of functions</i>	987
Robert Alan Melter, <i>Contributions to Boolean geometry of p-rings</i>	995
James Ronald Retherford, <i>Basic sequences and the Paley-Wiener criterion</i>	1019
Dallas W. Sasser, <i>Quasi-positive operators</i>	1029
Oved Shisha, <i>On the structure of infrapolynomials with prescribed coefficients</i>	1039
Oved Shisha and Gerald Thomas Cargo, <i>On comparable means</i>	1053
Maurice Sion, <i>A characterization of weak* convergence</i>	1059
Morton Lincoln Slater and Robert James Thompson, <i>A permanent inequality for positive functions on the unit square</i>	1069
David A. Smith, <i>On fixed points of automorphisms of classical Lie algebras</i>	1079
Sherman K. Stein, <i>Homogeneous quasigroups</i>	1091
J. L. Walsh and Oved Shisha, <i>On the location of the zeros of some infrapolynomials with prescribed coefficients</i>	1103
Ronson Joseph Warne, <i>Homomorphisms of d-simple inverse semigroups with identity</i>	1111
Roy Westwick, <i>Linear transformations on Grassman spaces</i>	1123