

# Pacific Journal of Mathematics

**EUCLIDEAN FIBERINGS OF SOLVMANIFOLDS**

JOHN STUELPNAGEL

## EUCLIDEAN FIBERINGS OF SOLVMANIFOLDS

JOHN STUELPNAGEL

This paper is concerned with the problem of finding conditions on a solvable Lie group  $G$  and a closed subgroup  $H$  which are sufficient for  $G/H$  to have topological structure of a fiber bundle with compact base space and euclidean fiber (if this is the case, we say that  $G/H$  has a euclidean fibering). The main results are the following two theorems.

**THEOREM 5.3.** Let  $G$  be a connected solvable linear Lie group, and  $H$  a closed subgroup which splits in  $G$ . Then  $G/H$  has a euclidean fibering.

**THEOREM 5.4.** Let  $G$  be a connected solvable matrix group, and assume that  $G$  is of finite index in its algebraic group hull. Then for any closed subgroup  $H$  of  $G$ ,  $G/H$  has a euclidean fibering.

To the best of the author's knowledge, these are the first results on existence of such fiberings which do not require that the isotropy subgroup  $H$  have a finite number of connected components.

A solvmanifold is a Hausdorff space  $X$  on which a solvable Lie group  $G$  acts transitively. It is well-known that, if  $x \in X$  and  $H$  is the subgroup of  $G$  leaving  $x$  fixed, then  $X$  is homeomorphic to the coset space  $G/H$ . The problem investigated in this paper is that of finding conditions on  $G$  and  $H$  which are sufficient for  $G/H$  to have the topological structure of a fiber bundle with compact base space and euclidean fiber, and such that the action of the structure group on the fiber is equivalent to the action of a linear group. Such a fibering will be called a euclidean fibering. We are led to suspect the existence of euclidean fiberings by Mostow's result of [4], in which he shows that  $G/H$  is covered a finite number of times by the Cartesian product of a compact solvmanifold and a euclidean space.

Our main results on euclidean fiberings are the two theorems quoted above.

In addition to these results on euclidean fiberings of solvmanifolds, we obtain in § 3 generalizations of some results of Togo on splittable matrix groups (cf. [9], [10], [11]). Togo [10] has proved Theorem 3.1 for the case that  $H$  is a Zariski-connected  $C^\infty$ -group over an algebraically closed perfect field, and has given an example (p. 319

---

Received April 14, 1964. This research was supported in part by the Army Research Office under Contract No. DA-36-034-ORD-3471.

in [11]) to show that the theorem is not generally true if  $H \cap N$  is not connected.

The research in this paper was originally done for the author's doctoral dissertation at the Johns Hopkins University. The author wishes to express his gratitude to Prof. G. D. Mostow for his advice, encouragement and direction.

**2. Notation and preliminaries.** If  $G$  is a Lie group, we shall always mean by  $\mathfrak{G}$  the Lie algebra of  $G$ , and  $Ad$  will denote the adjoint representation of  $G$  by automorphisms of  $\mathfrak{G}$ ; if  $H$  is a subgroup of  $G$ ,  $AdH$  denotes the image of  $H$  in  $AdG$  under  $Ad$ .

If  $G$  is a connected Lie group (we shall also say, equivalently, that  $G$  is an analytic group), and  $H$  is a subgroup,  $H$  is said to be full in  $G$  if no proper analytic subgroup of  $G$  contains  $H$ .  $H$  is said to be uniform in  $G$  if  $G/\bar{H}$  is compact,  $\bar{H}$  being the topological closure of  $H$ . It is shown in [3] that if  $G$  is nilpotent, and simply connected, then there is a unique subgroup  $\tilde{H}$  in which the subgroup  $H$  is full, and  $\tilde{H}$  may also be characterized as the unique analytic subgroup of  $G$  in which  $H$  is uniform.

We shall often use without explicit reference the following fact: Let  $G$  be a Lie group,  $H$  a closed subgroup, and  $F$  a closed subgroup invariant under inner automorphism by elements of  $H$ , and such that  $F \cap H$  is uniform in  $F$ . Then  $HF = \{h \cdot f \mid h \in H, f \in F\}$  is a closed subgroup of  $G$ . Clearly  $HF$  is a subgroup, and to see that it is closed we need only to observe that  $HF/H$ , being a continuous image of the compact space  $F/F \cap H$ , is a compact, hence closed, subset of  $G/H$ , and consequently the inverse image  $HF$  of  $HF/H$  in  $G$  is closed.

If  $\rho$  is a representation of the Lie group  $G$ , we say that a subgroup  $H$  of  $G$  is  $\rho$ -reductive if  $\rho(H)$  is a reductive (i.e., fully reducible) subgroup of  $\rho(G)$ . It is known that a connected solvable matrix group is reductive if and only if it is abelian and consists only of semisimple matrices (a matrix is called semisimple if its minimal polynomial has no repeated roots).

All linear transformations considered will be on vector spaces over the real or complex numbers. A linear transformation  $x$  is said to be unipotent if, for some integer  $n$ ,  $(x - I)^n = 0$ ,  $I$  being the identity. Any nonsingular linear transformation  $x$  may be written uniquely in the form  $x = su = us$ , where  $s$  is semisimple and  $u$  is unipotent;  $s$  and  $u$  are called respectively the semisimple and unipotent parts of  $x$ , or the Jordan components of  $x$ .

By an algebraic matrix group we shall mean a subgroup of  $GL(n, R)$  (resp.  $GL(n, C)$ ) which is the intersection of some algebraic variety in  $E^{n^2}$  with  $GL(n, R)$  (or in  $E^{2n^2}$  with  $GL(n, C)$ ),  $GL(n)$  being embedded

in euclidean space in the usual way. If  $H$  is a matrix group, the intersection of all algebraic groups containing  $H$  is called the algebraic group hull of  $H$ . If  $x$  and  $y$  are in  $GL(n)$ ,  $y$  is called an adherent of  $x$  if  $y$  is in every algebraic group containing  $x$ . In particular, the Jordan components of  $x$  are adherents of  $x$ .

We shall say that the space  $X$  has a euclidean fibering if  $X$  is the total space of a fiber bundle having a compact base space, euclidean fiber, and a group whose action on the fiber is equivalent to the action of a linear group.

The term  $\Gamma$ -invariant exp-set will be used as defined in [5], namely: Let  $\Gamma$  be a group of automorphisms of a Lie group  $G$ ,  $S$  be a subset of  $G$ , and let  $d\Gamma$  denote the group of automorphisms of  $\mathfrak{G}$  induced by  $\Gamma$ .  $S$  is called a  $\Gamma$ -invariant exp-set if there exist linearly independent subspaces  $\mathfrak{S}_1, \dots, \mathfrak{S}_n$  of  $\mathfrak{G}$ , each invariant under  $d\Gamma$ , such that the mapping  $s_1 + s_2 + \dots + s_n \rightarrow \exp s_1 \cdot \exp s_2 \cdots \exp s_n$  is a homeomorphism of  $\mathfrak{S}_1 + \dots + \mathfrak{S}_n = \mathfrak{S}$  onto  $S$  ( $s_i$  in  $\mathfrak{S}_i$ ). Thus the operation of  $\Gamma$  on  $S$  is equivalent to the operation of the linear group  $d\Gamma$  on  $\mathfrak{S}$ . If  $H$  is a subgroup of  $G$  and  $\Gamma_H$  is the group of inner automorphisms of  $G$  by elements in  $H$ , we shall also call a  $\Gamma_H$ -invariant exp-set an  $H$ -invariant exp-set.

Topological terms applied to matrix groups will always refer to the euclidean topology induced on  $GL(n)$ , not the Zariski topology, unless otherwise stated.

**3. Splittability of solvable groups.** In this section, we consider different definitions of splittability for solvable groups. We first define splittability for linear groups.

DEFINITION. Let  $G$  be a solvable linear group, and  $H$  a subgroup.

(1)  $G$  is said to be Jordan-splittable if it contains the Jordan components of each of its elements;

(2)  $G$  is said to be splittable if  $G = A \cdot N$  (semi-direct), where  $A$  is a maximal abelian subgroup of semisimple elements of  $G$ , and  $N$  is the subgroup of unipotent elements in  $G$ ;

(3)  $H$  is said to split in  $G$  if there is a semi-direct decomposition  $G = A \cdot N$  as in (2), with  $H = (H \cap A) \cdot (H \cap N)$  (semi-direct).

It has been shown by Togo in [9] that for connected linear groups, (1) and (2) above are equivalent. We generalize this result in the following theorem.

**THEOREM 3.1.** *Let  $G$  be a connected splittable solvable linear group, and  $H$  a subgroup of  $G$  with  $H \cap N$  connected, where  $N$  is the subgroup of unipotent matrices in  $G$ . Then (1), (2), and (3) above are equivalent conditions on  $H$ . Furthermore, if any of these are*

satisfied, then maximal reductive subgroups of  $H$  are conjugate by inner automorphism from  $H \cap N$ .

*Proof.* It is obvious that (3) implies (2). Conversely, if  $H$  is splittable, so that  $H = A_H(H \cap N)$  (semi-direct), then  $A_H$  is contained in a maximal reductive subgroup  $A$  of  $G$ ; since  $G$  is connected we have, by the result in [9],  $G = A \cdot N$  (semi-direct), and  $A_H = H \cap A$ , so  $H$  satisfies (3).

Proof that (1) implies (2). Let  $\tilde{H}$  be the algebraic group hull of  $H$ , and  $x$  a unipotent element of  $\tilde{H}$ , so  $x$  is an adherent of some element  $y \in H$ . Since  $x$  is a unipotent adherent of  $y$ ,  $x$  is an adherent of the unipotent part of  $y$  [1]; but  $H \cap N$ , being a connected group of unipotent elements, is algebraic, so  $x \in H \cap N$ . Thus  $H \cap N = \tilde{H} \cap N$ . Let  $M_i$  be a maximal reductive subgroup of  $\tilde{H}$  containing the maximal reductive subgroup  $A_i$  of  $H$ ,  $i = 1, 2$ . By Theorem 7.1 in [7], there is  $n \in \tilde{H} \cap N$  with  $nM_1n^{-1} = M_2$ ; and  $\tilde{H} = M_i \cdot (\tilde{H} \cap N)$  (semi-direct). Since  $A_i = M_i \cap H$ , and  $n \in H$ , it follows that  $nA_1n^{-1} = A_2$ , and  $H = (H \cap M_i) \cdot (\tilde{H} \cap N)$ . We have thus shown that (1) implies (2), and that (1) implies that maximal reductive subgroups of  $H$  are conjugate by inner automorphism from  $H \cap N$ .

Proof that (2) implies (1). Suppose  $H = A_H \cdot (N \cap H)$  (semi-direct). If  $\tilde{A}_H$  is the algebraic group hull of  $A_H$ , then by Proposition 5.5 in [1],  $\tilde{H} = \tilde{A}_H \cdot (N \cap H)$  is algebraic, since  $N \cap H$ , being a connected group of unipotent elements, is algebraic. If  $x \in H$ , and  $x = s \cdot u$  is the Jordan decomposition for  $x$ , then  $u \in \tilde{H}$ , so  $u$ , being unipotent, is in  $N \cap \tilde{H} = N \cap H$ . Thus  $s = xu^{-1}$  is in  $H$ , and it follows that  $H$  is Jordan-splittable.

Definitions (1) and (2) are not equivalent unless  $H \cap N$  is connected, as counterexamples of Togo show ([10], [11]). It is obvious, however, that (3) always implies (2), and the first part of the proof did not use the assumption that  $H \cap N$  is connected, so (2) and (3) are equivalent for subgroups of connected splittable linear groups.

We now make analogous definitions for simply connected abstract groups.

DEFINITION. Let  $G$  be a connected, simply connected solvable Lie group, and  $H$  a subgroup.

(1')  $H$  is said to be Jordan  $Ad$ -splittable if  $AdH$ , the image of  $H$  in the adjoint group of  $G$ , is Jordan-splittable;

(2')  $H$  is said to be splittable if  $H = A_H \cdot (H \cap N)$  (semi-direct) where  $A_H$  is an  $Ad$ -reductive subgroup of  $H$ , and  $N$  is the maximum

analytic nilpotent normal subgroup of  $G$ .

(3')  $H$  is said to split in  $G$  if  $G = A \cdot N'$ ,  $H = (A \cap H) \cdot (H \cap N')$  (both semi-direct), where  $A$  is  $Ad$ -reductive and  $N'$  is an analytic subgroup of  $N$  normal in  $G$ .

For a suitable faithful representation of  $G$ , Def. 3 is equivalent to Def. 3'. We deal in this paper with spaces  $G/H$  where  $G$  is a solvable linear group and  $H$  is a closed subgroup. When we refer to  $H$  splitting in  $G$  it will always be in the sense of Def. 3.

Taking  $H = G$  in definitions (1'), (2'), we obtain equivalent conditions on  $G$ . To see this, observe that if  $G$  satisfies (1'), then  $AdG$ , being Jordan-splittable, can be written  $AdG = A' \cdot AdN$ , where  $A'$  is a maximal reductive subgroup of  $AdG$ . Letting  $\mathfrak{A}$  be a complementary subspace to  $\mathfrak{N}$  in  $\mathfrak{G}$  invariant under  $A'$ , it follows easily that  $\mathfrak{G} = \mathfrak{A} + \mathfrak{N}$  (semi-direct), and that the analytic group  $A$  determined by  $\mathfrak{A}$  is  $Ad$ -reductive, so  $G$  is splittable. Conversely, if  $G = A \cdot N$  (semi-direct), with  $A$  and  $N$  being as in Definition (2'), then  $AdG = AdA \cdot AdN$  (semi-direct), so by Theorem (3.1),  $AdG$  is Jordan-splittable.

We now prove a partial analogue of Theorem 3.1 for abstract groups.

**THEOREM 3.2.** *Let  $G$  be a connected, simply connected solvable splittable Lie group, and  $H$  a subgroup of  $G$  containing no proper analytic subgroup central in  $G$ . Assume that  $H \cap N$  is connected, and that  $Ad(H \cap N) = AdH \cap AdN$ , where  $N$  is the maximum analytic nilpotent normal subgroup of  $G$ . Then  $H$  satisfies (1') if and only if it satisfies (2'), and in this case maximal  $Ad$ -reductive subgroups of  $H$  are conjugate by inner automorphism from  $H \cap N$ .*

Proof that (1') implies (2'). Let  $h \in H$ . Since  $AdH$  is Jordan-splittable, we can find  $s', u \in H$  with  $Ads'$  semisimple,  $Adu \in AdH \cap AdN$ , and  $Adh = Ads' Adu = Adu Ads'$ . Since  $AdH \cap AdN = Ad(H \cap N)$ , we can assume  $u \in H \cap N$ . Now  $Ad(h^{-1}s'u) = e$ , so  $h^{-1}s'u = z \in H \cap Z$ , where  $Z$  is the center of  $G$ . Letting  $s = s'z^{-1}$ , we have  $h = su$ . Now  $Ad(sus^{-1}u^{-1}) = e$ , so  $sus^{-1}u^{-1} \in Z \cap H \cap N$ , which is an analytic subgroup of  $H$  central in  $G$ , and therefore reduces to the identity. Thus  $su = us$ . A similar argument shows that  $s$  and  $u$  are unique, if we require  $u \in H \cap N$ .

Let  $A_H$  be a maximal  $Ad$ -reductive subgroup of  $H$ . Then  $Ad(A_H)$  is a maximal reductive subgroup of  $AdH$ . For suppose  $T$  is a maximal reductive subgroup of  $AdH$  containing  $Ad(A_H)$ , and let  $t \in T$ . There is  $h \in H$  with  $Adh = t$ ; for any  $a \in A_H$ ,  $Ad(aha^{-1}h^{-1}) = (Ada)t(Ada)^{-1}t^{-1} = e$ , so  $aha^{-1}h^{-1} \in H \cap N \cap Z = (e)$ . Thus  $h$  commutes with all of  $A_H$ , and  $Adh$  is semisimple, so  $h \in A_H$ , and  $t \in Ad(A_H)$ , whence  $T \subset Ad(A_H)$ .

If  $A_H$  and  $A'_H$  are maximal  $Ad$ -reductive subgroups of  $H$ , then

there is  $n = Adu$ , with  $u \in H \cap N$ , such that  $n(AdA_H)n^{-1} = Ad(A'_H)$ , so  $Ad(uA_Hu^{-1}) = Ad(A'_H)$ . Thus for any  $a \in A_H$ , there is  $a' \in A'_H$  with  $Ad(uau^{-1}a') = e$ , so  $(uau^{-1}a') \in Z \cap H$ . Now any element of  $Z \cap H$  is contained in every maximal  $Ad$ -reductive subgroup of  $H$ , so  $uau^{-1}a' \in A'_H$ , and hence  $uau^{-1} \in A'_H$ . It follows that  $uA_Hu^{-1} \subset A'_H$ , and, by maximality of  $A_H$ ,  $uA_Hu^{-1} = A'_H$ . We have shown that maximal  $Ad$ -reductive subgroups of  $H$  are conjugate by inner automorphism from  $H \cap N$ .

To show that  $H$  is a semi-direct product of the form  $A'_H(H \cap N)$ , we observe that if  $h = s \cdot u$  is the unique decomposition given above for  $h$ , then since  $s$  is contained in a maximal  $Ad$ -reductive subgroup of  $H$ , there is  $n \in H \cap N$  with  $nsn^{-1} \in A_H$ , so  $h = (nsn^{-1}) \cdot (ns^{-1}n^{-1}su)$  is a decomposition of the desired form. Since  $A_H \cap (H \cap N)$  is contained in  $H \cap N \cap Z$ , which consists of the identity alone, the decomposition is semi-direct.

Proof that (2') implies (1'). If  $H = A_H \cdot (H \cap N)$ , then  $AdH = AdA_H \cdot Ad(H \cap N)$ , and since  $Ad(H \cap N)$  is connected, it follows from Theorem 3.1 that  $AdH$  is Jordan-splittable.

The definitions obtained by requiring that  $AdH$  satisfy (2) or (3) are not in general equivalent with (2') and (3') as stated, as simple counterexamples show.

That the assumption  $Ad(H \cap N) = AdH \cap AdN$  is necessary for the theorem to hold, is shown by the following example: Let  $G$  be the simply connected Lie group whose Lie algebra is spanned by  $X, Y, Z$ , with bracket relations  $[X, Y] = Z$ ;  $[X, Z] = -Y$ ;  $[Y, Z] = 0$ ; let  $h = \exp(2\pi X) \cdot \exp Y$ , and let  $H$  be the cyclic subgroup of  $G$  generated by  $h$ . Then  $AdH$  is a discrete group of unipotent elements, so  $H$  is Jordan  $Ad$ -splittable, and  $H \cap N = (e)$ ; thus if  $H$  were splittable in the sense of (2'),  $H$  would be an  $Ad$ -reductive subgroup of  $G$ , which is not the case.

**4. A lemma on nilpotent groups.** We prove here a generalization of Case 2a in Theorem 2.1 of [5] which will be used to apply the results of the preceding section to fiberings of solvmanifolds.

**LEMMA 4.1.** *Let  $G$  be a connected, simply connected nilpotent Lie group,  $R$  a fully reducible group of automorphisms of  $G$  (i.e., the group  $dR$  of induced automorphisms of  $\mathfrak{G}$  is fully reducible, considering  $\mathfrak{G}$  as vector space), and  $H$  an analytic subgroup of  $G$  invariant under  $R$ . Then there is a closed subset  $E$  of  $G$ , invariant under  $R$  and homeomorphic to a euclidean space, such that the mapping of  $E \times H$  onto  $G$  given by  $(e, h) \rightarrow e \cdot h$  is a homeomorphism.*

*Proof.* We prove the lemma first in the case that  $H$  is normal

in  $G$ . Since  $\mathfrak{H}$  is invariant under  $dR$ , there is a complementary subspace  $\mathfrak{C}$  in  $\mathfrak{G}$  invariant under  $dR$ . Since  $\mathfrak{G} = \mathfrak{C} + \mathfrak{H}$ ,  $G = \exp(\mathfrak{C} + \mathfrak{H})$ . Let  $g \in G$ . Then there is a unique element  $X \in \mathfrak{G}$  with  $g = \exp X$ , and  $X = Y_1 + Y_2$  with  $Y_1 \in \mathfrak{C}$ ,  $Y_2 \in \mathfrak{H}$ . Let  $h = \exp(-Y_1) \exp(Y_1 + Y_2)$ . By the Campbell-Hausdorff formula,  $h = \exp(Y_2 + Q)$ , where  $Q$  is a finite linear combination of terms of the form  $ad^{p_1}(Y_1 + Y_2) \cdot ad^{q_1}Y_1 \cdots ad^{p_k}(Y_1 + Y_2) \cdot ad^{q_k}Y_1(Y_1 + Y_2)$ , with  $\sum(p_i + q_i) \leq \dim \mathfrak{G}$ . Since  $adY_1(Y_1 + Y_2) = [Y_1, Y_2] \in \mathfrak{H}$ , and  $adZ(\mathfrak{H}) \subset \mathfrak{H}$  for all  $Z \in \mathfrak{G}$ , we see that  $Q \in \mathfrak{H}$ .

Consequently  $Y_2 + Q \in \mathfrak{H}$ , and  $h \in H$ . Thus, we have  $g = (\exp Y_1)h = e \cdot h$ , with  $e \in E = \exp \mathfrak{C}$ . To show uniqueness, we suppose  $e_1 h_1 = e_2 \cdot h_2$  with  $e_1, e_2 \in \exp \mathfrak{C}$ ,  $h_1, h_2 \in H$ . Then  $e_1 = e_2 h_2 h_1^{-1} = e_2 h'$ , so  $\exp(\log e_1) = \exp(\log e_2) \cdot \exp(\log h')$ . Again using the Campbell-Hausdorff formula, we have  $\exp(\log e_2) \cdot \exp(\log h') = \exp(\log e_2 + Y)$ , where  $Y \in \mathfrak{H}$ . Then  $\exp(\log e_1) = \exp(\log e_2 + Y)$ , whence  $\log e_1 = \log e_2 + Y$ , so  $Y \in \mathfrak{H} \cap \mathfrak{C} = (0)$ , and consequently  $e_1 = e_2$ ,  $h_1 = h_2$ .

The map of  $(E \times H) \rightarrow G$  given by  $(e, h) \rightarrow eh$  is continuous, by continuity of group multiplication; let  $\pi$  denote the projection (as vector space) of  $\mathfrak{G}$  onto  $\mathfrak{C}$  given by  $\pi(Y_1 + Y_2) = Y_1$ , where  $Y_1 \in \mathfrak{C}$ ,  $Y_2 \in \mathfrak{H}$ . Since  $\pi$  is linear, it is continuous. It is clear from the construction that, to obtain the representation  $e \cdot h$  for an element  $g \in G$ , we define  $e$  by  $e = \exp(\pi(\log g))$ . Since  $\log$ ,  $\pi$ , and  $\exp$  are continuous,  $e$  depends continuously on  $g$ , and hence  $h = e^{-1}g$  does also. Thus  $\varphi: (e, h) \rightarrow eh$  is a homeomorphism. That  $E$  is homeomorphic to euclidean space follows from the fact that  $\exp: \mathfrak{C} \rightarrow E$  is a homeomorphism.

In the case that  $H$  is not normal in  $G$ , let  $H_1$  be the normalizer of  $H$  in  $G$ , and define inductively  $H_{i+1}$  to be the normalizer of  $H_i$  in  $G$  ( $i = 1, 2, \dots$ ). For some integer  $K$ ,  $H_K = G$ . That this is true is seen by observing that for all  $i$ ,  $H_i$  is connected, so either  $\dim H_i > \dim H_{i-1}$  or  $H_i = H_{i-1}$ . But if  $H_i = H_{i-1}$ , then the  $\mathfrak{G}/\mathfrak{H}_i$  part of  $ad\mathfrak{H}_i$  has no nonzero eigenvector corresponding to the eigenvalue zero, and this contradicts the nilpotency of  $\mathfrak{G}$ . Therefore,  $\dim H_i > \dim H_{i-1}$ , so  $H_K = G$  for some  $K \leq \dim G$ .

Since  $H_{i-1}$  is normal in  $H_i$  (letting  $H_0 = H$ ),  $H_i = E_i \cdot H_{i-1}$ , by the first part of the proof, where  $E_i$  is a euclidean subspace of  $H_i$  invariant under  $R$  (clearly  $H_i$  is invariant under  $R$  for all  $i$ ). Then the map of  $E_K \times E_{K-1} \times \dots \times E_1 \times H \rightarrow G$  given by  $(e_K, \dots, e_1, h) \rightarrow e_K \cdot e_{K-1} \cdots e_1 h$  can be factored into

$$\begin{aligned} (E_K \times E_{K-1} \times \dots \times E_1 \times H) &\rightarrow (E_K \times \dots \times E_2 \times H_1) \\ &\rightarrow (E_K \times \dots \times E_3 \times H_2) \rightarrow \dots \rightarrow (E_K \times H_{K-1}) \rightarrow G, \end{aligned}$$

each of which is a homeomorphism, and, letting  $E = E_K E_{K-1} \cdots E_1$ , the image of  $E_K \times E_{K-1} \times \dots \times E_1$  under this map, we see that  $E$  is



homeomorphic to euclidean space, and is closed in  $G$ , since  $E \times (id.)$  is closed in  $E \times H$ .

It is clear from the construction that  $E$  is an  $R$ -invariant exp-set.

**5. Applications to fiberings of solvmanifolds.** Before considering more general types of solvmanifolds, we wish to single out one special case, in which the fibers are permuted transitively by a reductive subgroup of  $G$ .

**THEOREM 5.1.** *Let  $G$  be a connected splittable solvable linear group,  $H$  a closed subgroup which splits in  $G$ , and assume  $H \cap N$  is connected. Then  $G/H$  has a euclidean fibering such that the fibers are permuted transitively by a reductive subgroup of  $G$ .*

*Proof.* Let  $G = A \cdot N$ ,  $H = (H \cap A) \cdot (H \cap N)$  be the simultaneous semi-direct decompositions of  $G$  and  $H$ .  $A/H \cap A$ , being a connected abelian group, may be written as the direct product of a toroidal group and a vector group. Let  $P$  denote the inverse image in  $A$  of the toroidal group, and let  $V$  denote a complementary subgroup to  $P$  in  $A$ , so  $V$  is a vector group, and an  $A$ -invariant exp-set, and  $A = P \cdot V$  (direct).

By Lemma 4.1, there is an  $A_H$ -invariant exp-set  $E$  such that  $N = E \cdot (H \cap N)$ , since  $H \cap N$  is analytic. Now we may write  $G = P \cdot V \cdot E \cdot (H \cap N)$ , or letting  $W = \{v \cdot e \mid v \in V, e \in E\}$ ,  $G = P \cdot W \cdot (H \cap N)$ ,  $H = (P \cap H) \cdot (H \cap N)$ , where  $P \cap H = A_H$ , and  $W$  and  $H \cap N$  are  $A_H$ -invariant exp-sets. By Lemma 4.1 in [5], it follows that  $G/H$  is a fiber bundle over  $P/H \cap P$ , which is a toroidal group, with fiber  $W$ , which is a euclidean space, and the group  $P$  permutes the fibers transitively.

We now prove a lemma on fiber bundles which we shall need later.

**LEMMA 5.2.** *Let  $B$  be the total space of a fiber bundle with base space  $X$ , fiber  $F_1 \times F_2$ , and group  $G$ , where  $F_1, F_2$  are topological spaces on which the Lie group  $G$  operates, and the operation of  $G$  on  $F_1 \times F_2$  is given by  $g \cdot (f_1, f_2) = (g \cdot f_1, g \cdot f_2)$ . Then  $B$  is the total space of a fiber bundle with fiber  $F_2$  and group  $G$ , whose base space  $Y$  is itself a fiber bundle with base space  $X$ , fiber  $F_1$ , and group  $G$ .*

*Proof.* We can regard the space  $B$  as the quotient space of the space

$$T_1 = \bigcup_{j \in J} \{(x, f_1, f_2, j) \mid x \in U_j, f_1 \in F_1, f_2 \in F_2\}$$

by the relation

$$R_1: (x, f_1, f_2, j)R_1(x', f'_1, f'_2, k) \text{ if } x = x', \\ f'_1 = g_{kj}(x) \cdot f_1, \quad f'_2 = g_{kj}(x) \cdot f_2,$$

since  $G$  operates componentwise on  $F_1 \times F_2$ ; here  $\{U_j\}$  is the family of coordinate neighborhoods on  $X$ , and  $g_{kj}(x): U_k \cap U_j \rightarrow G$  are the coordinate transformations.

Let  $Y$  be the fiber bundle with base space  $X$ , fiber  $F_1$ , group  $G$ , coordinate neighborhoods  $\{U_j\}$  and coordinate transformations  $g_{kj}$ , whose existence and uniqueness up to equivalence is proved in Theorem 3.2 in [8], and let  $\psi_j: U_j \times F_1 \rightarrow p^{-1}(U_j)$  be the coordinate functions, where  $p: Y \rightarrow X$  is the projection.  $Y$  may be regarded as the quotient space of

$$T_2 = \bigcup_{j \in J} \{(x, f_1, j) \mid x \in U_j, f_1 \in F_1\}$$

by the relation

$$R_2: (x, f_1, j)R_2(x', f'_1, k) \text{ if } x = x', f'_1 = g_{kj}(x) \cdot f_1.$$

Denote by  $q_1$  the map of  $T_1$  onto  $B$  given by mapping a point into its  $R_1$ -equivalence class, and by  $q_2$  the map of  $T_2$  onto  $Y$  given by mapping a point into its  $R_2$ -equivalence class, so  $q_1$  and  $q_2$  are open and continuous.

Define  $\pi: B \rightarrow Y$  by  $\pi(q_1(x, f_1, f_2, j)) = q_2(x, f_1, j)$ .  $\pi$  is well-defined, for if  $q_1(x, f_1, f_2, j) = q_1(x', f'_1, f'_2, k)$ , then  $x = x'$ ,  $f'_1 = g_{kj}(x) \cdot f_1$ , so  $q_2(x, f_1, j) = q_2(x', f'_1, k)$ . It is clear that  $\pi$  is open and continuous, since  $q_1$  and  $q_2$  are both open and continuous.

Let  $V_j = p^{-1}(U_j)$ , so  $\{V_j\}$  is an open covering of  $Y$ , and define  $\varphi_j: V_j \times F_2 \rightarrow \pi^{-1}(V_j)$  by  $\varphi_j(y, f_2) = q_1(\psi_j^{-1}(y), f_2, j)$ . Since  $p\pi(q_1(\psi_j^{-1}(y), f_2, j)) = pq_2(\psi_j^{-1}(y), j) = p(y) \in U_j$ ,  $\varphi_j(V_j \times F_2) \subset \pi^{-1}(V_j)$ . To see that  $\varphi_j$  is onto, observe that if  $b = q_1(x, f_1, f_2, k)$ , then  $b = q_1(x, g_{jk}(x) \cdot f_1, g_{jk}(x) \cdot f_2, j)$ , so  $b = \varphi_j(\psi_j^{-1}(x, g_{jk}(x) \cdot f_1), g_{jk}(x) \cdot f_2, j)$ . The openness and continuity of  $\varphi_j$  follow from that of  $q_1$  and  $\psi_j^{-1}$ .

To show that  $\varphi_j$  is one-to-one, suppose that  $\varphi_j(y, f_2) = \varphi_j(y', f'_2)$ . Then  $q_1(\psi_j^{-1}(y), f_2, j) = q_1(\psi_j^{-1}(y'), f'_2, j)$ , hence  $f_2 = f'_2$ , and  $\psi_j^{-1}(y) = \psi_j^{-1}(y')$ . Since  $\psi_j$  is one-to-one,  $y = y'$ , so  $\varphi_j$  is one-to-one.

We have thus shown that  $B$  has a fibering of the desired type, and we need only show that  $G$  is the group of the bundle. Suppose  $y \in V_j \cap V_k$ , and consider  $f'_2 = \varphi_{k,y}^{-1} \cdot \varphi_{j,y}(f_2)$ . Since  $\varphi_j(y, f_2) = \varphi_k(y, f'_2)$  we have  $q_1(\psi_j^{-1}(y), f_2, j) = q_1(\psi_k^{-1}(y), f'_2, k)$ , or, denoting  $\psi_j^{-1}(y)$  by  $(x, f_1)$ ,  $\psi_k^{-1}(y)$  by  $(x, f'_1)$ ,  $q_1(x, f_1, f_2, j) = q_1(x, f'_1, f'_2, k)$ , so  $f'_2 = g_{kj}(x) \cdot f_2$ . Since  $g_{kj}(x) \in G$ , the lemma is proved.

We now apply our previous results to prove two theorems on the existence of euclidean fiberings.

**THEOREM 5.3.** *Let  $G$  be a connected solvable linear Lie group, and  $H$  a closed subgroup which splits in  $G$ . Then  $G/H$  has a euclidean fibering.*

*Proof.* Let  $N$  denote the analytic subgroup of unipotent elements of  $G$ , let  $A_H$  be a maximal reductive subgroup of  $H$  such that  $H = A_H \cdot (H \cap N)$ , and let  $A$  be a maximal reductive subgroup of  $G$  containing  $A_H$ , so  $G = A \cdot N$ ,  $A_H = H \cap A$ . If we denote by  $F$  the minimum analytic subgroup of  $N$  containing  $H \cap N$ , then  $H$  is contained in the normalizer of  $F$ , and the subgroup  $HF$  is closed in  $G$ , since  $HF/H = F/F \cap H$  is compact.  $HF$  splits in  $G$ , since  $HF = A_H \cdot (H \cap N)F = A_H \cdot F$  (semi-direct), and  $HF \cap N = F$ .

Let  $\tilde{A}_H$  be the minimum analytic subgroup of  $A$  containing  $A_H$  and the maximum compact subgroup of  $A$ , and let  $V$  be a subgroup of  $A$  complementary to  $\tilde{A}_H$ . Then  $V$  is a vector group,  $A = \tilde{A}_H \cdot V$ , and  $G = \tilde{A}_H \cdot V \cdot N$ . Letting  $S = V \cdot N$ , we have  $HS = A_H \cdot S$  (semi-direct),  $HS$  is closed,  $A_H S/A_H = S/S \cap A_H = S$  is euclidean, and  $G/HS = \tilde{A}_H \cdot S/A_H \cdot S$  is homeomorphic to  $\tilde{A}_H/A_H$ , which is compact.

Now  $G/H$  is a fiber bundle with base space  $G/HS$ , fiber  $HS/H$ , and group  $HS$ , acting on  $HS/H$  by left translation. Since  $HS/A_H = A_H S/A_H$  is euclidean, and hence solid, we can reduce the group of the bundle to  $A_H$ , acting on  $HS/H$  by left translation. (Theorem 12.5 in [8]).

Letting  $E$  be an  $A_H$ -invariant exp-set such that  $N = E \cdot F$  (the existence of  $E$  is proved in Lemma 4.1), we obtain  $HS = A_H \cdot V \cdot E \cdot F = V \cdot E \cdot F \cdot A_H$ . Let  $A_H$  operate on  $V \times E \times F \times A_H$  by  $a(v, e, f, a') = (ava^{-1}, aea^{-1}, afa^{-1}, aa')$ , so the homeomorphism of  $HS$  onto  $V \times E \times F \times A_H$  is equivariant with respect to the action of  $A_H$  on  $HS$ , by left translation, and on the product space as defined above. The map of  $HS/H$  onto  $V \times E \times (F \cdot A_H/H)$  given by  $sH \rightarrow (v, e, hH)$ , where  $s = v \cdot e \cdot h$ ,  $v \in V$ ,  $e \in E$ ,  $h \in F \cdot A_H$ , is clearly a homeomorphism, also equivariant with respect to the action of  $A_H$ .

Thus the fiber  $HS/H$  of the bundle whose total space is  $G/H$  is homeomorphic to the Cartesian product of a euclidean space  $V \times E$ , and the compact connected nilmanifold  $F \cdot A_H/(F \cap H) \cdot A_H = F/F \cap H$ , and the group  $A_H$  of the bundle acts componentwise. Applying Lemma 5.2, we thus obtain the stated result, that  $G/H$  is a fiber bundle with compact base space and euclidean fiber, and the action of the group is equivalent to the action of a linear group. Proof of the theorem is complete.

In case the eigenvalues of the elements of  $Ad(A_H)$  are all positive,

we can obtain Mostow's result of [4] that the fibering is trivial.

**COROLLARY.** *If the eigenvalues of the elements of  $Ad(A_H)$  are positive, then  $G/H = Y \times E'$ , where  $E'$  is a euclidean space, and  $Y$  is compact.*

*Proof.* The action of the group  $A_H$  on the fiber  $E' (= V \times E)$  is equivalent to the action of  $Ad(A_H)$  on the tangent space  $\mathfrak{G}'$  of  $E'$  at the identity. Taking a basis in  $\mathfrak{G}'$  with respect to which the matrices of  $Ad(A_H)$  are diagonal, and letting  $A$  denote the minimum analytic subgroup of diagonal matrices with positive eigenvalues containing  $Ad(A_H)$ , we observe that  $A$  operates naturally on  $\mathfrak{G}'$ . Defining  $\lambda(a)(\exp e_i) = \exp(a(e_i))$ , for  $a \in A$ ,  $e_i \in \mathfrak{G}_i$  ( $\mathfrak{G}_i$  being the subspaces of  $\mathfrak{G}'$  which make  $E'$  an  $A_H$ -invariant exp-set), we see that  $\lambda$  defines an operation of  $A$  on  $E'$ , and the action of  $A_H$  is equivalent to the action of a subgroup of  $A$ , so we may regard  $A$  as the group of the bundle.  $A$  is euclidean, and hence solid, so, by Theorem 12.5 in [8], we may reduce the group of the bundle to the identity. Thus  $G/H = Y \times E'$ .

The theorem just proved made assumptions about both  $G$  and  $H$ . By making a somewhat stronger assumption on  $G$ , we are able to drop all restrictions on  $H$ , except that it be closed.

**THEOREM 5.4.** *Let  $G$  be a connected solvable matrix group, and assume that  $G$  is of finite index in its algebraic group hull. Then for any closed subgroup  $H$  of  $G$ ,  $G/H$  has a euclidean fibering.*

*Proof.* Let  $H'$  be the intersection of  $G$  with the algebraic group hull of  $H$ , so  $H'$  has a finite number of connected components. Denote by  $H^0$  the identity component of  $H'$ , by  $H^*$  the intersection of  $H$  with  $H^0$ , and let  $F = H \cdot H^0$ , so  $F/H^0$ , being contained in the finite group  $H'/H^0$ , has a finite number of connected components, and  $F$  is closed in  $G$ .  $H^* \cap [H^0, H^0]$  is uniform in  $[H^0, H^0]$ , since  $[H^0, H^0]$  is the minimum analytic group containing  $[H, H] \subset H^*$ . Thus  $H^* \cdot [H^0, H^0]$  is closed, and normal in  $H^0$ . The factor group  $H^0/H^* \cdot [H^0, H^0]$ , being abelian and connected, is the direct product of a toroidal group and a vector group. Let  $T$  be the inverse image in  $H^0$  of the toroidal group, so  $H^0/T$  is a vector group, and  $T/H^*$ , being a fiber bundle over the compact base space  $T/H^* \cdot [H^0, H^0]$  with compact fiber  $H^* \cdot [H^0, H^0]/H^* = [H^0, H^0]/H^* \cap [H^0, H^0]$ , is compact.

We note that, since  $H^0/H^*$  is a fiber bundle with euclidean base space  $H^0/T$  and compact fiber  $T/H^*$ , by Corollary 1.6 in [8],  $H^0/H^* = F/H$  is homeomorphic with the product space  $(H^0/T) \times (T/H^*)$ .

$HT$ , being a finite extension of  $T$  (since  $H/H \cap T = H/H^*$  is finite), is closed in  $G$ , and has a finite number of connected components,

so  $HT = K \cdot E$ , where  $K$  is a maximal compact subgroup of  $HT$  and  $E$ , being a  $K$ -invariant exp-set, is euclidean (cf. [5], [6]). Observe that  $K$  leaves  $T$  invariant under inner automorphism, since  $T = HT \cap H^0$ , so there is a  $K$ -invariant exp-set  $V$  in  $H^0$  with  $F = V \cdot E \cdot K = V \cdot HT$ .

Since  $F$  is Jordan-splittable and  $F \cap N$  is connected,  $N$  being the group of unipotent matrices in  $G$ ,  $F$  splits in  $G$ , so  $FN$  is closed and normal in  $G$ . Then  $G/FN$  is a connected abelian group, so is a direct product of a toroidal group and a vector group. Let  $G_1$  be the inverse image in  $G$  of the toroidal group, so  $G/G_1$  is euclidean and  $G_1/FN$  is compact.

Since  $F = V \cdot HT$ , and  $H \cap N$  is connected, by Lemma 4.1 we can write  $FN = W' \cdot V \cdot HP$ , where  $W'$  is a  $K$ -invariant exp-set complementary to  $F \cap N$  in  $N$ .

Now  $G_1/H$  is a bundle with compact base space  $G_1/FN$ , fiber  $FN/H$ , and group  $FN$  acting by left translation on  $FN/H$ . Noting that  $FN/K = W' \cdot V \cdot E \cdot K/K \approx W' \times V \times E$ , we see that  $FN/K$  is solid, and we can reduce the group of the bundle to  $K$  acting on  $FN/H$  by left translation. Letting  $W = W' \cdot V$ , we see that if  $g \in FN$  and  $g = w \cdot t$  is the unique decomposition with  $w \in W$ ,  $t \in HT$ , then the map  $gH \rightarrow (w, tH)$  is a homeomorphism of  $FN/H$  onto  $W \times HT/H$ , and this map is equivariant with respect to the action of  $K$ , where  $K$  acts componentwise on  $W \otimes HT/H$ , by inner automorphism on  $W$  and left translation on  $HT/H$ . By Lemma 5.2,  $G_1/H$  is a fiber bundle over a compact base space  $X$  with euclidean fiber and compact group  $K$ , whose action on the fiber is equivalent to the action of a linear group.

$G/H$ , being a bundle over the euclidean space  $G/G_1$  with fiber  $G_1/H$ , is homeomorphic to the product bundle  $(G/G_1) \times (G_1/H)$ , by Corollary 1.6 in [8]. It is easily seen that  $G/H$  is thus a bundle over  $X$  with euclidean fiber  $(G/G_1) \times W$ , and group  $K$  acting componentwise, where  $K$  acts trivially on  $G/G_1$  and by inner automorphism on  $W$ . Proof of the theorem is complete.

The proof of Theorem 5.4 included an investigation of the factor space of  $H^0/H^*$ , which can also be applied to  $H'/H$ , where  $H'$  is the minimum subgroup of finite index in the algebraic group hull of  $H$ , and containing  $H$ . If  $\tilde{H}$  is the algebraic group hull of  $H$ , then  $\tilde{H}/H$  is homeomorphic with the cartesian product of  $H'/H$  with a finite set  $S$ , having the discrete topology, and each component is homeomorphic to  $H'/H$ , which is the cartesian product of a compact space  $C$  and a euclidean space  $E$ . Since the space  $C \times S$  is compact, and  $\tilde{H}/H \approx (C \times S) \times E$ , we have thus proved the following special result:

**THEOREM 5.5** *Let  $G$  be a solvable algebraic matrix group, and*

*H any closed subgroup whose algebraic group hull is G. Then  $G/H \approx C \times E$ , where C is a compact space and E is euclidean.*

## REFERENCES

1. Armand Borel, *Groupes lineaires algebriques*, Ann. Math. **64** (1956), 20-82.
2. A. I. Malcev, *Solvable Lie algebras*, Amer. Math. Soc. Translation No. 27, Series 1, 1950.
3. ———, *On a class of homogeneous spaces*, Amer. Math. Soc. Translation No. 39, Series 1, 1951.
4. G. D. Mostow, *Factor spaces of solvable groups*, Ann. Math. **60** (1954), 1-27.
5. ———, *On covariant fiberings of Klein spaces*, Amer. J. Math. **77** (1955), 247-278.
6. ———, *Self-adjoint groups*, Ann. Moth. **62** (1955), 44-55.
7. ———, *Fully reducible subgroups of algebraic groups*, Amer. J. Math. **78** (1956), 200-221.
8. Norman Steenrod, *Topology of fibre bundles*, Princeton University Press, 1951.
9. Shigeaki Togo, *On splittable linear Lie algebras*, J. Sci. Hiroshima Univ., Ser. A, **18** (1955), 289-306.
10. ———, *On splittable linear groups*, Rend. Circ. Mat. Palermo (2) **8** (1959), 49-76.
11. ———, *On linear groups in which all elements can be decomposed in Jordan products*, Math. Zeit. **75** (1961), 305-324.

RESEARCH INSTITUTE FOR ADVANCED STUDY (RIAS), BALTIMORE, MARYLAND  
NOW AT WESTINGHOUSE ELECTRIC CORPORATION, BALTIMORE, MARYLAND



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California

R. M. BLUMENTHAL

University of Washington  
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California  
Los Angeles, California 90007

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.



# Pacific Journal of Mathematics

Vol. 15, No. 2

October, 1965

Patrick Robert Ahern, <i>On the generalized F. and M. Riesz theorem</i> . . . . .	373
A. A. Albert, <i>On exceptional Jordan division algebras</i> . . . . .	377
J. A. Anderson and G. H. Fullerton, <i>On a class of Cauchy exponential series</i> . . . . .	405
Allan Clark, <i>Hopf algebras over Dedekind domains and torsion in H-spaces</i> . . . . .	419
John Dauns and D. V. Widder, <i>Convolution transforms whose inversion functions have complex roots</i> . . . . .	427
Ronald George Douglas, <i>Contractive projections on an <math>L_1</math> space</i> . . . . .	443
Robert E. Edwards, <i>Changing signs of Fourier coefficients</i> . . . . .	463
Ramesh Anand Gangolli, <i>Sample functions of certain differential processes on symmetric spaces</i> . . . . .	477
Robert William Gilmer, Jr., <i>Some containment relations between classes of ideals of a commutative ring</i> . . . . .	497
Basil Gordon, <i>A generalization of the coset decomposition of a finite group</i> . . . . .	503
Teruo Ikebe, <i>On the phase-shift formula for the scattering operator</i> . . . . .	511
Makoto Ishida, <i>On algebraic homogeneous spaces</i> . . . . .	525
Donald William Kahn, <i>Maps which induce the zero map on homotopy</i> . . . . .	537
Frank James Kosier, <i>Certain algebras of degree one</i> . . . . .	541
Betty Kvarda, <i>An inequality for the number of elements in a sum of two sets of lattice points</i> . . . . .	545
Jonah Mann and Donald J. Newman, <i>The generalized Gibbs phenomenon for regular Hausdorff means</i> . . . . .	551
Charles Alan McCarthy, <i>The nilpotent part of a spectral operator. II</i> . . . . .	557
Donald Steven Passman, <i>Isomorphic groups and group rings</i> . . . . .	561
R. N. Pederson, <i>Laplace's method for two parameters</i> . . . . .	585
Tom Stephen Pitcher, <i>A more general property than domination for sets of probability measures</i> . . . . .	597
Arthur Argyle Sagle, <i>Remarks on simple extended Lie algebras</i> . . . . .	613
Arthur Argyle Sagle, <i>On simple extended Lie algebras over fields of characteristic zero</i> . . . . .	621
Tôru Saitô, <i>Proper ordered inverse semigroups</i> . . . . .	649
Oved Shisha, <i>Monotone approximation</i> . . . . .	667
Indranand Sinha, <i>Reduction of sets of matrices to a triangular form</i> . . . . .	673
Raymond Earl Smithson, <i>Some general properties of multi-valued functions</i> . . . . .	681
John Stuepnel, <i>Euclidean fiberings of solvmanifolds</i> . . . . .	705
Richard Steven Varga, <i>Minimal Gerschgorin sets</i> . . . . .	719
James Juei-Chin Yeh, <i>Convolution in Fourier-Wiener transform</i> . . . . .	731