

# Pacific Journal of Mathematics

## **PROPERTIES OF SOLUTIONS OF $n$ -TH ORDER LINEAR DIFFERENTIAL EQUATIONS**

THOMAS LAWRENCE SHERMAN

## PROPERTIES OF SOLUTIONS OF $N^{\text{th}}$ ORDER LINEAR DIFFERENTIAL EQUATIONS

THOMAS L. SHERMAN

**The purpose of this paper is to establish zero properties for solutions of the general  $n^{\text{th}}$  order linear differential equation**

$$(1) \quad L_n y = \sum_{i=0}^n r_i(x) y^{(i)} = 0,$$

**where  $r_n(x) \neq 0$  and all the coefficients are continuous.**

This work was stimulated by the recent investigations of fourth-order equations by Leighton and Nehari [11], Barrett [3, 4, 5] and Howard [8]. For example, if  $\eta_1(a)$  is defined to be the first point  $b > a$  for which there exist a solution of

$$(2) \quad (r(x)y'')'' - p(x)y = 0 \quad (r, p > 0, r \in C_2, p \in C)$$

with four zeros in  $[a, b]$  (counting multiplicities), then it is known [11] that there is a solution which vanishes with double zeros at  $a$  and  $\eta_1(a)$  ( $\eta_1(a)$  is termed the first (right) conjugate point of  $a$ ). If  $\mu_1(a)$  is defined to be the first point  $b > a$  for which there exists a non-trivial solution of (2) satisfying the conditions  $y(a) = y'(a) = (ry'')(b) = (ry'')(b) = 0$ , then it is known [3] that  $a < \mu_1(a) < \eta_1(a)$  ( $a$  is called the first (left) focal point of  $\mu_1(a)$ ). These results were subsequently extended to general even-order self-adjoint equations by Reid [13] and Hunt [9].

In §1 we derive some basic results which are used in later sections.

In §2 we extend the definition of conjugate point found in [11]. Using this definition we obtain generalizations (in a direction different from that of [9] and [13]) of results of [11].

In §3 we define a notion of focal point for (1) and extend the discussions of [3, 4, 5] to obtain results similar to those of §2.

In §4 we develop an eigenvalue relationship from which we can easily obtain a generalization of the following theorem of Leighton and Nehari:

**THEOREM.** *Equation (2) is disconjugate (i.e.,  $\eta_1(a)$  does not exist)*

---

Received June 12, 1963. Supported, in part, by the National Science Foundation under grant NSF G23719. This paper was presented to the Mathematics Department of the University of Utah in partial fulfillment of the requirements for the degree of Doctor of Philosophy.

on  $[a, \infty)$  if and only if the minimum eigenvalue of the associated problem

$$(ry'')'' - py = \lambda y, \quad y(a) = y'(a) = y(b) = y'(b) = 0,$$

is positive for all  $b > a$ .

**1. Preliminaries.** We begin our discussion with a number of theorems concerning properties of real functions.

For reference we state the following basic lemma, which was proven by Leighton and Nehari [11].

**LEMMA 1.** *Let  $u(x)$  and  $v(x)$  be of class  $C_1$  in  $(a, b)$  and let  $v(x)$  be of constant sign ( $\neq 0$ ) in the interval. If  $x = \alpha$  and  $x = \beta$  ( $a < \alpha < \beta < b$ ) are consecutive zeros of  $u(x)$ , then there exists a constant  $\mu$  such that the function  $u(x) - \mu v(x)$  has at least a double zero in  $(\alpha, \beta)$ .*

This lemma has been extended by Azbelev and Chaluk [1] to the case where  $\beta$  is allowed to be a double zero of  $u(x)$  and a simple zero of  $v(x)$ . We shall use the method of the latter together with Lemma 1 to prove

**THEOREM 1.** *Let  $u(x)$  be a function such that  $u(x)$  has a zero of order  $n \geq 1$  at  $x = a$  and a zero of order  $m \geq 1$  at  $x = b$ , and  $u(x)$  is of constant sign ( $\neq 0$ ) in  $(a, b)$ . Let  $v(x)$  be a function such that  $v(x)$  has a zero of order  $n_1 < n$  at  $x = a$  and a zero of order  $m_1 < m$  at  $x = b$ , and  $v(x)$  is of constant sign ( $\neq 0$ ) in  $(a, b)$ . Further suppose  $u(x)$  and  $v(x)$  are both of class  $C_M[a, b]$ , where  $M = \max(n_1, m_1)$ . Then there exists a linear combination  $z(x)$  of  $u(x)$  and  $v(x)$  such that  $z(x)$  has a double zero in  $(a, b)$ .*

*Proof.* We may as well suppose that  $u(x)$  and  $v(x)$  are both positive on  $(a, b)$ . Let  $c \in (a, b)$  and let

$$w(x) = v(x) - \frac{v(c) + 1}{u(c)} u(x).$$

There exists an  $h_1 \in (a, c)$  such that  $w(h_1) > 0$ , since, by the zero properties of  $u(x)$  and  $v(x)$ ,  $w^{(n_1)}(a) = v^{(n_1)}(a) > 0$  and  $w^{(i)}(a) = 0$  ( $i = 0, 1, \dots, n_1 - 1$ ). Similarly there is an  $h_2 \in (c, b)$  such that  $w(h_2) > 0$ , since  $(-1)^{m_1} w^{(m_1)}(b) > 0$  and  $w^{(i)}(b) = 0$  ( $i = 0, 1, \dots, m_1 - 1$ ). Also  $w(c) = -1 < 0$ ; thus  $w(x)$  has a zero in  $(a, c)$  and a zero in  $(c, b)$ . Since  $w(x)$  is continuous, there exists  $\alpha$  and  $\beta$  ( $a < \alpha < c < \beta < b$ ) such that  $w(\alpha) = w(\beta) = 0$  and  $w(x) < 0$  on  $(\alpha, \beta)$ . Applying Lemma 1 to the functions  $w(x)$  and  $v(x)$  on  $(\alpha, \beta)$ , have our result.

**THEOREM 2.** *Suppose  $r(x)f(x) \in C_n$  and  $f(x) \in C_n$  in a neighborhood of the point  $x = a$ . If  $f^{(i)}(a) = 0$  ( $i = 0, 1, \dots, n$ ) and if  $r(x)$  is a bounded function in a neighborhood of  $a$ , then  $(rf)^{(i)}(a) = 0$  ( $i = 0, 1, \dots, n$ ).*

*Proof.* The theorem is trivial for  $n = 0$ . For  $n = 1$ ,

$$\begin{aligned} |(rf)'(a)| &= \left| \lim_{h \rightarrow 0} \frac{(rf)(a+h) - (rf)(a)}{h} \right| \\ &= \left| \lim_{h \rightarrow 0} \frac{r(a+h)f(a+h) - r(a)f(a)}{h} \right| \\ &\leq \overline{\lim}_{h \rightarrow 0} |r(a+h)| \left| \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h} \right| \\ &= M |f'(a)| = 0, \end{aligned}$$

where  $M = \overline{\lim}_{x \rightarrow a} |r(x)|$ . Now using the formula

$$(rf)^{(k)}(a) = \lim_{h \rightarrow 0} \frac{\sum_{i=0}^k (-1)^i C(k, i) (rf)(a + (k-i)h)}{h^k}$$

we shall show that  $(rf)^{(k)}(a) = 0$  ( $1 \leq k \leq n$ ). Applying L'Hospital's rule, and using the zero properties of  $f(x)$  at  $x = a$  and the boundedness of  $r$  at  $x = a$ , we have

$$\begin{aligned} |(rf)^{(k)}(a)| &= \left| \lim_{h \rightarrow 0} \sum_{i=0}^k \frac{(-1)^i C(k, i) r(a + (k-i)h) f(a + (k-i)h)}{h^k} \right| \\ &\leq M \sum_{i=0}^{k-1} \left| \frac{(k-i)^k}{i! (k-i)!} \lim_{h \rightarrow 0} \frac{f^{(k-1)}(a + (k-i)h) - f^{(k-1)}(a)}{(k-i)h} \right| \\ &= M |f^{(k)}(a)| \sum_{i=0}^{k-1} \frac{(k-i)^k}{i! (k-i)!}. \end{aligned}$$

Since  $f^{(k)}(a) = 0$ , the result follows.

**COROLLARY.** *Suppose  $f(x) \in C_n$ ,  $r(x)f(x) \in C_n$ , and  $r(x) \in C$  in a neighborhood of  $x = a$ . If  $f^{(i)}(a) = 0$  ( $i = 0, 1, \dots, n-1$ ),  $(rf)^{(n)}(a) = 0$ , and  $r(a) \neq 0$ , then  $f^{(n)}(a) = 0$ .*

*Proof.* Since  $r(x)$  is continuous at  $x = a$ , we can replace  $M$  by  $r(a)$ , “ $\leq$ ” by “ $=$ ”, and drop the absolute value signs in the inequality in the proof of Theorem 2, and obtain a valid equality; the result is then immediate.

In the next theorem and in the remainder of the paper we shall deal with the equation

$$(1) \quad L_n y = \sum_{i=0}^n r_i(x) y^{(i)}(x) = 0,$$

where  $r_n(x) \neq 0$  and the coefficients  $r_i (i = 0, 1, \dots, n)$  are real valued continuous functions.

The following basic existence theorem is apparently sufficiently well-known so that references to it in the literature are scarce; we include it here for the sake of completeness.

**THEOREM 3.** *If  $a_i \in [a, b]$  ( $i = 1, \dots, k \leq n - 1$ ) and  $a_1 < a_2 < \dots < a_k$ , then there exists a nontrivial solution of (1) which satisfies the boundary conditions*

$$\begin{aligned} y^{(i_1)}(a_1) &= 0 & i_1 &= n_{11}, n_{12}, \dots, n_{1p_1} \\ y^{(i_2)}(a_2) &= 0 & i_2 &= n_{21}, n_{22}, \dots, n_{2p_2} \\ & \vdots & & \\ y^{(i_k)}(a_k) &= 0 & i_k &= n_{k1}, n_{k2}, \dots, n_{kp_k} \end{aligned}$$

where  $0 \leq n_{j1} < n_{j2} < \dots < n_{jp_j} < n$  ( $j = 1, \dots, k$ ) and  $\sum_{j=1}^k p_j \leq n - 1$ .

*Proof.* Let  $y_1(x), y_2(x), \dots, y_n(x)$  be a fundamental set of solutions of (1). We wish to find a nontrivial set of constants  $C_1, \dots, C_n$  such that  $y = C_1 y_1 + \dots + C_n y_n$  satisfies the above boundary conditions.

Applying the boundary conditions to  $y(x)$ , we have

$$\begin{aligned} y^{(n_{11})}(a_1) &= C_1 y_1^{(n_{11})}(a_1) + \dots + C_n y_n^{(n_{11})}(a_1) = 0 \\ & \vdots \\ y^{(n_{1p_1})}(a_1) &= C_1 y_1^{(n_{1p_1})}(a_1) + \dots + C_n y_n^{(n_{1p_1})}(a_1) = 0 \\ & \vdots \\ y^{(n_{21})}(a_2) &= C_1 y_1^{(n_{21})}(a_2) + \dots + C_n y_n^{(n_{21})}(a_2) = 0 \\ & \vdots \\ y^{(n_{2p_2})}(a_2) &= C_1 y_1^{(n_{2p_2})}(a_2) + \dots + C_n y_n^{(n_{2p_2})}(a_2) = 0 \\ & \vdots \\ y^{(n_{k1})}(a_k) &= C_1 y_1^{(n_{k1})}(a_k) + \dots + C_n y_n^{(n_{k1})}(a_k) = 0 \\ & \vdots \\ y^{(n_{kp_k})}(a_k) &= C_1 y_1^{(n_{kp_k})}(a_k) + \dots + C_n y_n^{(n_{kp_k})}(a_k) = 0. \end{aligned}$$

This is a system of  $\sum_{j=1}^k p_j$  (i.e., less than  $n$ ) homogeneous equations in  $n$  unknowns  $C_1, \dots, C_n$ , and so there always exists a nontrivial solution.

We shall make repeated use of this theorem throughout the remainder of this paper.

2. **Zero properties of solutions.** In this section we shall be concerned with zero properties of solutions of (1). The principal stimulation for this investigation is the work of Leighton and Nehari [11]. Their concern was with the equation

$$(2) \quad (r(x)y'')'' - p(x)y = 0,$$

where  $r(x) \in C_2$ ,  $p(x) \in C$ , and  $r(x) > 0$ . They defined the first conjugate point of the point  $a$ , denoted by  $\eta_1(a)$ , to be the minimum point  $b > a$  such that there is a nontrivial solution  $y(x)$  of (2) vanishing at  $x = a$ , with four zeros, counted in their multiplicities, in  $[a, b]$ . They showed that if there is a nontrivial solution with four zeros on  $[a, \infty)$ , then  $\eta_1(a)$  exists. It was then shown that, if  $p(x) > 0$  in (2), a solution which vanishes four times in  $[a, \eta_1(a)]$  has double zeros at  $x = a$  and  $x = \eta_1(a)$  and does not vanish in  $(a, \eta_1(a))$ . Barrett [3, 4, 5] then extended this work to equations of the form

$$[(r(x)y'')' + q(x)y']' + p(x)y = 0,$$

where  $r(x)$ ,  $q(x)$  and  $p(x)$  are all continuous. Hunt [9] and Reid [13] in turn extended this work to self-adjoint differential equations of order  $2n$ , defining conjugate point as the minimum point  $b > a$  such that there exists a nontrivial solution with a zero of order  $n$  at  $a$  and a zero of order  $n$  at  $b$ .

Hanan [7], using a definition of conjugate point similar to that of Leighton and Nehari, investigated the third order equation

$$y''' + p(x)y'' + q(x)y' + r(x)y = 0.$$

Investigations similar to Hanan's were undertaken by Azbelev and Chaluk [1].

We shall here be interested in the extension of this work to the equation (1), using the conjugate point concept of Leighton and Nehari. Some of the results obtained in this section have been reported independently by A. Ju. Levin [12] who uses the methods of Green's functions. It should be noted that self-adjointness and assumptions on the signs of the coefficients have been of fundamental importance in the work cited above (except for that of Levin); in general we make no such assumptions here. Following Barrett [2, 3], we make the following definition:

**DEFINITION 1.** Equation (1) is said to be *disconjugate* in  $[a, \infty)$  if there exists no nontrivial solution of (1) which vanishes at  $a$ , and has  $n$  zeros, counting multiplicities, in  $[a, \infty)$ .

We now prove a basic theorem, special cases of which have been

used by Leighton and Nehari and Barrett, and a special case of which was proved by Hanan.

**THEOREM 4.** *Suppose that for each positive integer  $k$  there is an  $n^{\text{th}}$  order linear boundary operator  $U_k$ , operating at points  $b_{k1}, \dots, b_{kj_k}$ , such that*

- (i) (1) has a nontrivial solution  $y_k(x)$  satisfying  $U_k(y_k(x)) = 0$ ;
- (ii) there exist points  $a_{k1}, \dots, a_{kn}$  such that  $y_k^{(i-1)}(a_{ki}) = 0$  ( $i = 1, \dots, n$ );
- (iii)  $\min_i b_{ki} \leq \min_i a_{ki} < \max_i a_{ki} \leq \max_i b_{ki}$ ;
- (iv)  $\min_i b_{ki} \leq \min_i b_{ji}$  and  $\max_i b_{ji} \leq \max_i b_{ki}$  if  $k < j$ .

Then  $\lim_{k \rightarrow 0} (\max_i b_{ki} - \min_i b_{ki}) > 0$ .

**REMARK.** If  $y_k(x)$  is a nontrivial solution of (1) having  $n$  zeros on an interval, counted in accordance with their multiplicities, then repeated application of Rolle's Theorem shows the existence of the desired set of points  $a_{k1}, a_{k2}, \dots, a_{kn}$ .

*Proof of Theorem 4.* Let  $A_k = \min_i a_{ki}$ ,  $B_k = \max_i a_{ki}$ ,  $C_k = \min_i b_{ki}$  and  $D_k = \max_i b_{ki}$ . Conditions (iv) and (iii) assure us that  $\{D_k\}$  forms a decreasing sequence bounded below; hence there is a  $D$  such that  $D_k \rightarrow D$ . Similarly  $\{C_k\}$  forms an increasing sequence bounded above; hence there is a  $C$  such that  $C_k \rightarrow C$  and  $D \geq C$ . We need only show that equality does not hold.

Let  $\phi_1(x), \phi_2(x), \dots, \phi_n(x)$  be the fundamental set of solutions of (1) satisfying  $\phi_{i+1}^{(j)}(D) = \delta_{ij}$  ( $i, j = 0, 1, \dots, n - 1$ ). Then  $y_k(x)$  can be written

$$y_k(x) = \sum_{i=1}^n c_{ik} \phi_i(x).$$

We normalize  $y_k(x)$  by letting

$$\bar{y}_k(x) = \frac{y_k(x)}{\left(\sum_{i=1}^n c_{ik}^2\right)^{1/2}} = \sum_{i=1}^n d_{ik} \phi_i(x),$$

so that  $\sum_{i=1}^n d_{ik}^2 = 1$ . Then  $\bar{y}_k(x)$  is a solution of (1) possessing the same zero properties as  $y_k(x)$ ; in particular,  $\bar{y}_k(x)$  satisfies condition (ii). Further, on any closed interval about  $D$ ,  $\{\bar{y}_k(x)\}$  is a uniformly bounded and equicontinuous family. Hence there is a uniformly convergent subsequence  $\{\bar{y}_{k_i}(x)\}$  whose limit function  $\bar{y}(x)$  is a solution of (1).

Now suppose  $D = C$ ; then by condition (iii)

$$A = \lim_{k \rightarrow 0} A_k = \lim_{k \rightarrow 0} B_k = B .$$

But now by condition (ii), for each  $j(0 \leq j < n)$ ,  $A$  is a limit point of zeros of  $\{\bar{y}_{k_i}^{(j)}(x)\}$ ; hence  $\bar{y}^{(j)}(A) = 0$  ( $j = 0, 1, \dots, n - 1$ ), and so  $\bar{y}(x) \equiv 0$ . This is impossible, since  $\sum_{i=1}^n d_{i,k}^2 = 1$ , which completes the proof.

We are now in a position to make

**DEFINITION 2.** The first conjugate point  $\eta_1(a)$  of the point  $a$  is the smallest number  $b > a$  such that there exists a nontrivial solution of (1) which vanishes at  $a$  and has  $n$  zeros, counting multiplicities in  $[a, b]$ .

We now establish one of the main results of this section.

**THEOREM 5.** *If (1) is not disconjugate, then  $\eta_1(a)$  exists, and there is a nontrivial solution of (1) which has a total of at least  $n$  zeros at  $a$  and  $\eta_1(a)$  and does not vanish in  $(a, \eta_1(a))$ .*

*Proof.* The existence of  $\eta_1(a)$  is guaranteed by Theorem 4. Let  $Y = \{y(x) \mid y(x) \text{ is a nontrivial solution of (1) which has } n \text{ zeros in } [a, \eta_1(a)]\}$ ,  $R = \{r \mid \text{there is a } y(x) \in Y \text{ with a zero of order } r \text{ at } x = a\}$ ,  $n_0 = \max r \in R$ ,  $M = \{m \mid \text{there is a } y \in Y \text{ with a zero of order } n_0 \text{ at } x = a, \text{ and } m \text{ zeros in } (a, \eta_1(a))\}$ ,  $\bar{m} = \max m \in M$ , and let  $\phi(x) \in Y$  be a solution with  $\bar{m}$  zeros in  $(a, \eta_1(a))$  and a zero of order  $n_0$  at  $x = a$ . Let  $a = a_0 < a_1 < \dots < a_p = \eta_1(a)$  be the zeros of  $\phi(x)$  on  $[a, \eta_1(a)]$ , with respective multiplicities  $n_0, n_1, \dots, n_p$ . Then  $n_1 + n_2 + \dots + n_{p-1} = \bar{m}$ . Let  $\bar{n} = n_0 + \dots + n_{p-1}$ . We claim that  $\phi(x)$  exhibits the property asserted. For suppose the theorem false for  $\phi(x)$ , i.e., suppose  $p > 1$ . We shall show the existence of a  $\psi(x) \in Y$  with a zero of order  $n_0$  at  $x = a$  and a zero of order  $\bar{m} + 1$  in  $(a, \eta_1(a))$ , which will contradict the maximality of  $\bar{m}$ . To this end let us consider a nontrivial solution  $z(x)$  of (1) with zeros at  $a_0, a_1, \dots, a_p$  of multiplicities  $n_0 + 1, n_1, \dots, n_{p-1} - 1, n - \bar{n} - 1$  respectively; such a solution exists (note that  $p > 1$  implies  $n_0 < n - 1$ ) since only  $n - 1$  zeros (counting multiplicities) are specified. Since the zero of  $z(x)$  at  $a_0$  is of order  $n_0 + 1$ ,  $z(x) \notin Y$ , and so  $z(x)$  does not vanish on  $(a_{p-1}, a_p)$ , nor are any of the zeros of multiplicity greater than that specified. Hence we can apply Theorem 1 to  $z(x)$  and  $\phi(x)$  on  $(a_{p-1}, a_p)$  to obtain a linear combination  $\psi(x)$  with a double zero at  $c$ , say, on  $(a_{p-1}, a_p)$ . Now  $\psi(x)$  has zeros at  $a_0, a_1, \dots, a_{p-2}, a_{p-1}, c, a_p$  of multiplicities  $n_0, n_1, \dots, n_{p-2}, n_{p-1} - 1, 2, n - \bar{n} - 1$ ; i.e.,  $\psi(x) \in Y$  with a zero at  $a$  of order  $n_0$ , and with a total multiplicity of  $\bar{m} + 1$  of zeros on  $(a, \eta_1(a))$ , contradicting the maximality of  $\bar{m}$ ; this established the theorem.

**COROLLARY.** *Let  $\phi(x)$  be a nontrivial solution of (1) with zeros*



of orders  $n_0$  and at least  $n - n_0$  at  $a$  and  $\eta_1(a)$  respectively, where  $n_0$  is maximal. Then  $\phi(x)$  is essentially unique i.e., unique except for multiplication by a constant factor). (We note that a similar result holds for a  $\psi(x) \in Y$ , not vanishing on  $(a, \eta_1(a))$ , and with a zero of maximal order at  $\eta_1(a)$ ).

*Proof.* Let  $\psi(x)$  be any such solution. By the maximality of  $n_0$  we have  $\phi^{(n_0)}(a) \neq 0 \neq \psi^{(n_0)}(a)$ . Consider the solution

$$\chi(x) = \psi^{(n_0)}(a)\phi(x) - \phi^{(n_0)}(a)\psi(x) .$$

$\chi(x)$  clearly has a zero of order  $n + 1$  at  $x = a$  and a zero of order  $n - n_0$  at  $x = \eta_1(a)$ ; hence, by the maximality of  $n_0$ ,  $\chi(x) \equiv 0$ .

In order to examine the behavior of  $\eta_1(a)$  as a function of  $a$  we prove the following theorems. The principal theorem, Theorem 7, in this sequence was obtained by Leighton and Nehari [11] for equation (2) and by Hunt [9] for the special definition of  $\eta_1$  mentioned at the beginning of this section, for an equation more general than that in [11].

**THEOREM 6.** *Let  $a$  be any point for which  $\eta_1(a)$  exists; then, for any  $b < a$ ,  $\eta_1(b)$  exists.*

*Proof.* Let  $a_0$  be the largest number  $c \geq a$  such that  $\eta_1(c) = \eta_1(a)$  (the existence of  $a_0$  is guaranteed by Theorem 4). Let  $\phi(x)$  be a solution such that  $\phi(a_0) = 0$  and  $\phi(x)$  has  $n$  zeros in  $[a_0, \eta_1(a)]$ , where the order of the zero of  $\phi(x)$  at  $\eta_1(a)$  is minimal over the set of all such solutions. Let  $m_0, m_1, \dots, m_r$  be the orders of the zeros of  $\phi(x)$  at  $a_0 < a_1 < \dots < a_r = \eta_1(a)$  respectively. We are supposing  $m_r$  is minimal. Let  $M = m_0 + \dots + m_{r-1} \leq n - 1$ . Clearly  $m_r \geq n - M$ .

Now suppose the theorem is not true. Let  $\psi(x)$  be a nontrivial solution of (1) such that  $\psi(b) = 0$  and  $\psi(x)$  has zeros of multiplicities  $m_0, m_1, \dots, m_{r-2}, m_{r-1} - 1, n - M - 1$  at  $a_0, a_1, \dots, a_{r-2}, a_{r-1}, a_r$  respectively. We have specified (counting multiplicities)  $n - 1$  zeros of  $\psi(x)$  in  $[b, \eta_1(a)]$ ; hence, since we are supposing  $\eta_1(b)$  does not exist,  $\psi(x)$  has zeros only at these points and only of the multiplicities specified. By Theorem 1, applied to  $\phi(x)$  and  $\psi(x)$ , there must exist a linear combination  $\chi(x)$  of  $\phi(x)$  and  $\psi(x)$  with a double zero in  $(a_{r-1}, a_r)$ . Further,  $\chi(x)$  has zeros of orders  $m_0, m_1, \dots, m_{r-2}, m_{r-1} - 1, n - M - 1$  at  $a_0, a_1, \dots, a_{r-1}, a_r$  respectively. Hence  $\chi(x)$  has  $n$  zeros on  $[a_0, \eta_1(a)]$ . Now if  $r = 1 = m_0 = m_r$  then  $\chi(x)$  has a double zero and is also a solution of a second order equation; hence  $\chi(x) \equiv 0$ . If  $r = 1 = m_0$  and  $m_r > 1$  then  $\chi(\eta_1(a)) = 0$  and  $\chi(x)$  has  $n$  zeros in  $(a_0, \eta_1(a)]$  which contradicts the maximality of  $a_0$ . If  $r = 1 = m_r$  and  $m_0 > 1$  then  $\chi(x)$  has  $n$  zeros

in  $[a_0, \eta_1(a))$  and vanishes at  $a_0$ , which contradicts the fact that  $\eta_1(a^0) = \eta_1(a)$ . In all other cases  $\chi(x)$  has  $n$  zeros in  $[a_0, \eta_1(a)]$  with a zero of order  $n - M - 1 < m_r$  at  $\eta_1(a)$ , which contradicts the minimality of  $m_r$ .

**COROLLARY 1.** *Let  $a$  be any point for which  $\eta_1(a)$  exists; then, for any  $b < a$ , there exists a solution which vanishes at  $b$  and which vanishes  $n - 1$  times, counting multiplicity, in  $[a, \eta_1(a)]$ .*

*Proof.* The function  $\psi(x)$  constructed in the proof of Theorem 5 is such a function.

**COROLLARY 2.** *Let  $a$  be any point for point for which  $\eta_1(a)$  exists; then no solution has more than  $n - 1$  zeros in  $[a, \eta_1(a))$ .*

*Proof.* Suppose this is not true. Then there exists a solution with  $n$  zeros in  $[a, \eta_1(a))$ . Hence there is a  $b \in [a, \eta_1(a))$  such that  $\eta_1(b) < \eta_1(a)$ ; clearly such a  $b$  cannot equal  $a$ . Then, by Corollary 1, there exists a solution  $\phi(x)$  which vanishes at  $a$  and which vanishes  $n - 1$  times, counting multiplicity, in  $[b, \eta_1(b)]$ . Hence  $\phi(x)$  vanishes at  $a$  and has  $n$  zeros in  $[a, \eta_1(b)]$ , which contradicts the definition of  $\eta_1(a)$ .

It should be noted that, for  $n = 2$ , Corollaries 1 and 2 combine to give the classical Sturm separation theorem.

**COROLLARY 3.**  $\eta_1(a)$  is a nondecreasing function of  $a$ .

**THEOREM 7.**  $\eta_1(a)$  is an increasing function of  $a$ .

*Proof.* Suppose the theorem is not true. We know by Corollary 3 that  $\eta_1(a)$  is a nondecreasing function of  $a$ . Thus there exist  $a, \eta_1(a), b$  and  $\eta_1(b)$  such that  $a < b$  and  $\eta_1(a) = \eta_1(b)$ . Hence, by Theorem 5 and Corollary 3, for every  $x \in [a, b]$ ,  $\eta_1(x)$  exists and equals  $\eta_1(a)$ . For each  $\alpha \in [a, b]$  let  $\phi_\alpha(x)$  be the essentially unique solution which vanishes at least  $n$  times at  $\alpha$  and  $\eta_1(a)$ , where the order of the zero at  $\eta_1(a)$  is maximal over all such solutions, and  $\phi_\alpha(x) \neq 0$  for  $x \in (\alpha, \eta_1(a))$ , as in the corollary to Theorem 5.

Let  $S_i = \{\alpha \mid \phi_\alpha(x) \text{ has a zero of exactly order } i \text{ at } x = \eta_1(a)\}$ , ( $i = 1, \dots, n - 1$ ). Clearly  $\bigcup_{i=1}^{n-1} S_i = [a, b]$ . Hence there is a subinterval  $[c, d] \subseteq [a, b]$  and an  $i$  ( $1 \leq i < n$ ) such that  $S_i$  is dense in  $[c, d]$  (see Simmons [14], page 74). Pick the maximum such  $i$  and denote it by  $m$ ; denote the subinterval  $[c, d]$  corresponding to this value of  $i$  by  $[a_m, b_m]$ .

Let  $\{y_1, y_2, \dots, y_n\}$  be the fundamental set of solution satisfying  $y_{i+1}^{(j)}(\eta_i(a)) = \delta_{ij}$ . Now the condition that  $S_m$  is dense in  $[a_m, b_m]$  is equivalent to the condition that the determinant

$$W_{m+1} = \begin{vmatrix} y_{m+1} & y_{m+2} & \cdots & y_n \\ y'_{m+1} & y'_{m+2} & \cdots & y'_n \\ \vdots & \vdots & \ddots & \vdots \\ y_{m+1}^{(n-m-1)} & y_{m+2}^{(n-m-1)} & \cdots & y_n^{(n-m-1)} \end{vmatrix}$$

vanish at a dense set of points in  $[a_m, b_m]$ . This follows since  $W_{m+1}(\alpha) = 0$  is the condition for a solution with a zero of order  $m$  at  $\eta_i(a)$  to have a zero of order  $n - m$  at a point  $\alpha \neq \eta_i(a)$ . Further, since  $W_{m+1}$  is continuous, we have  $W_{m+1} \equiv 0$  in  $[a_m, b_m]$ . Moreover from the maximality of  $m$  we know that

$$W_{m+2} = \begin{vmatrix} y_{m+2} & y_{m+3} & \cdots & y_n \\ y'_{m+2} & y'_{m+3} & \cdots & y'_n \\ \vdots & \vdots & \ddots & \vdots \\ y_{m+2}^{(n-m-2)} & y_{m+3}^{(n-m-2)} & \cdots & y_n^{(n-m-2)} \end{vmatrix} \neq 0$$

on some subinterval  $[a'_m, b'_m] \subseteq [a_m, b_m]$ .

Now consider the differential equation

$$\begin{vmatrix} y & y_{m+2} & \cdots & y_n \\ y' & y'_{m+2} & \cdots & y'_n \\ \vdots & \vdots & \ddots & \vdots \\ y^{(n-m-1)} & y_{m+2}^{(n-m-1)} & \cdots & y_n^{(n-m-1)} \end{vmatrix} = 0$$

formed by substituting  $y$  for  $y_{m+1}$  in the equation  $W_{m+1} \equiv 0$ . This is an equation of order  $n - m - 1$  with non-vanishing leading coefficient  $W_{m+2}$  on the interval  $[a'_m, b'_m]$ . The solutions are  $y_{m+2}, y_{m+3}, \dots, y_m$ , a total of  $n - m - 1$  solutions. These are linearly independent, since  $W_{m+2} \neq 0$  on  $[a'_m, b'_m]$ ; hence they form a solution basis. But  $y_{m+1}$  is also a solution, since  $W_{m+1} \equiv 0$  on  $[a'_m, b'_m]$ ; therefore  $y_{m+1}, y_{m+2}, \dots, y_n$  are linearly dependent on  $[a'_m, b'_m]$ , which contradicts the fact that  $y_1, \dots, y_n$  are solutions of (1) which are independent on  $[a, b]$ . This completes the proof.

3. Zero properties of derivatives of solutions. We now turn our attention to the behavior of the derivatives of solutions. The principal motivation for this type of investigation is the work of Barrett [2, 3, 4, 5], in which are references to earlier work in this field. Barrett [2] discussed the behavior of solutions of the equation

$$(p(x)y')' + f(x)y = 0$$

in relation to the boundary conditions  $y(a) = y'(b) = 0$ ; the minimum such  $b > a$  for which these conditions are satisfied was denoted by  $\mu_1(a)$ , and  $a$  was called a (left) focal point of the point  $\mu_1(a)$ . This discussion was extended to the equation (2), with a focal point being defined in terms of the first point  $b > a$ , for which conditions of the type  $y(a) = y'(a) = y''(b) = y'''(b) = 0$  are satisfied. This was in turn extended by Hunt [9] and Reid [13] to more general even order self-adjoint differential equations of the form

$$[r_n(x)y^{(n)}]^{(n)} + \sum_{i=0}^{n-1} [(-1)^{n+i+1}r_i(x)y^{(i)}]^{(i)} = 0 .$$

Their definition of focal point involved the conditions

$$\begin{aligned} y(a) &= y'(a) = \dots = y^{(n-1)}(a) = 0 \\ &= (r_n y^{(n)})(b) = (r_n y^{(n)})'(b) = \dots = (r_n y^{(n)})^{(n-1)}(b) . \end{aligned}$$

It should be noted that the use of the terms ‘‘focal point’’ and ‘‘conjugate point’’ stems from considerations in the calculus of variations, in which field the majority of the investigations have taken place (see for example Reid [13] and the references in that paper).

In the present paper, we propose to extend the definition of focal point in a different direction, to which the techniques of differential equations seem more applicable. To this end we make the following definition.

**DEFINITION 3.** Suppose that there exists a point  $c > a$ , an integer  $k(0 < k < n)$ , and a nontrivial solution  $\phi(x)$  of (1) which has a zero of order  $k$  at  $x = a$  and such that  $\phi^{(k)}(x)$  vanishes  $n - k$  times, counting multiplicities, in  $(a, c]$ . Then the minimum such  $c$ , over all such solutions for all possible  $k$ , which exists by Theorem 4, will be denoted by  $\tilde{\mu}_1(a)$ .

**DEFINITION 4.** Suppose that there exists a point  $c > a$ , an integer  $k(0 < k < n)$ , and a nontrivial solution  $\phi(x)$  of (1) such that  $\phi(x)$  has a zero at  $x = a$  of order  $k$  and such that  $\phi^{(k)}(x)$  has a zero at  $x = c$  of order  $n - k$ . By Theorem 4 there is a minimum such  $c$ , over all such solutions for all possible  $k$ , to be denoted by  $\mu_1(a)$ ;  $a$  will then be called the first (left) focal point of  $\mu_1(a)$ .

**THEOREM 8.** *If  $\tilde{\mu}_1(a)$  exists, then  $\mu_1(a)$  exists and  $\mu_1(a) = \tilde{\mu}_1(a)$ .*

*Proof.* Let  $U = \{u(x) \mid u(x)$  is a nontrivial solution of (1) and there exists a  $k(0 < k < n)$  such that  $u(x)$  has a zero of order  $k$  at  $x = a$  and such that  $u^{(k)}(x)$  has  $n - k$  zeros (counting multiplicities) in  $(a, \tilde{\mu}_1(a)]\}$ ,

$K = \{k \mid \text{there is a } u(x) \in U \text{ with a zero of order } k \text{ at } x = a\}$ ,  $m_0 = \max k \in K$ ,  $P = \{p \mid \text{there is a } u(x) \in U \text{ with a zero of order } m_0 \text{ at } x = a, \text{ and such that } u^{(m_0)}(x) \text{ has } p \text{ zeros in } (a, \tilde{\mu}_1(a))\}$ ,  $p_0 = \max p \in P$ , and let  $\psi(x) \in U$  be a solution with a zero of order  $m_0$  at  $x = a$  and such that  $\psi^{(m_0)}(x)$  has  $p_0$  zeros in  $(a, \tilde{\mu}_1(a))$ . Let the zeros of  $\psi^{(m_0)}(x)$  be of orders  $m_1, m_2, \dots, m_j$  at  $a_1, a_2, \dots, a_j$  respectively, where  $a < a_1 < \dots < a_j = \tilde{\mu}_1(a)$ ; by definition of  $p_0$ ,  $m_1 + m_2 + \dots + m_{j-1} = p_0$ .

Now suppose the conclusion does not hold; i.e., suppose  $j > 1$ . Let  $z(x)$  be a nontrivial solution of (1) with a zero of order  $m_0 + 1$  at  $x = 0$  (note that  $z(x) \notin U$ ) and with  $z^{(m_0)}(x)$  having zeros of multiplicities  $m_1, m_2, \dots, m_{j-1}, m_{j-1} - 1, n - m_0 - 1 - p_0$  at  $a_1, a_2, \dots, a_{j-2}, a_{j-1}, a_j$  respectively, except that if  $m_{j-1} = 1$  we make no specification at  $a_{j-1}$ . We have specified  $n - 1$  zeros of  $z(x)$  and  $z^{(m_0)}$  on  $[a, \tilde{\mu}_1(a)]$ ,  $n - m_0 - 1$  zeros of  $z^{(m_0)}(x)$  on  $[a, \tilde{\mu}_1(a)]$ ,  $p_0$  zeros of  $z^{(m_0)}(x)$  on  $[a, \tilde{\mu}_1(a))$ , and  $p_0 - 1$  zeros of  $z^{(m_0)}(x)$  on  $(a, \tilde{\mu}_1(a))$ . Hence by Rolle's Theorem  $z^{(m_0+1)}(x)$  has at least  $n - m_0 - 2$  zeros on  $[a, \tilde{\mu}_1(a)]$ , and, in fact, since  $z(x) \notin U$ ,  $z^{(m_0)}(x)$  has exactly  $n - m_0 - 2$  zeros on  $[a, \tilde{\mu}_1(a)]$ . Therefore,  $z^{(m_0)}(x)$  does not vanish at any point on  $(a_{j-1}, a_j)$ , nor are any of the specified zeros of multiplicities greater than that specified. We can now apply Theorem 1 to the functions  $z^{(m_0)}(x)$  and  $\psi^{(m_0)}(x)$  on the interval  $[a_{j-1}, a_j]$ , concluding that there is a linear combination  $\chi(x)$  of  $\psi(x)$  and  $z(x)$  such that  $\chi^{(m_0)}(x)$  has a double zero in  $(a_{j-1}, a_j)$ . We now have a nontrivial solution  $\chi(x)$  of (1) with a zero of order  $m_0$  at  $x = a$ , such that  $\chi^{(m_0)}(x)$  has at least  $n - m_0$  zeros on  $[a, \tilde{\mu}_1(a)]$  (hence  $\chi(x) \in U$ ), and  $\chi^{(m_0)}(x)$  has at least  $p_0 + 1$  zeros on  $(a, \tilde{\mu}_1(x))$  which contradicts the maximal property of  $p_0$ . Hence  $j = 1$  and the conclusion holds.

**THEOREM 9.** *If  $\eta_1(a)$  exists, then  $\mu_1(a)$  exists and  $\mu_1(a) < \eta_1(a)$ .*

*Proof.* Let  $\phi(x)$  be the essentially unique solution of Theorem 5, with its zero at  $a$  of order  $n_0$  (where  $n_0$  is maximal over the orders of the zero of all nontrivial solutions vanishing  $n$  times at  $a$  and  $\eta_1(a)$ ); then its zero at  $\eta_1(a)$  is of order at least  $n - n_0$ . By Rolle's Theorem,  $\phi'(x)$  vanishes at least once on  $(a, \eta_1(a))$  and at least  $n - n_0$  times on  $(a, \eta_1(a)]$ . After repeated application of Rolle's Theorem we find that  $\phi^{(n_0)}(x)$  vanishes at least  $n_0$  times on  $(a, \eta_1(a))$  and at least  $n - n_0$  times on  $(a, \eta_1(a)]$ . Hence  $\tilde{\mu}_1(a)$  exists, by Theorem 4; clearly  $\tilde{\mu}_1(a) \leq \eta_1(a)$ , and by Theorem 8  $\mu_1(a) = \tilde{\mu}_1(a)$ .

To show that  $\mu_1(a) < \eta_1(a)$ , suppose  $\mu_1(a) \geq \eta_1(a)$ . Then, by the above,  $\mu_1(a) = \tilde{\mu}_1(a) = \eta_1(a)$ . Let  $R = \{r \mid \text{there exists a } u(x) \in U \text{ with a zero of order } r \text{ at } x = a \text{ and such that } u^{(r)}(x) \text{ vanishes in } (a, \tilde{\mu}_1(a))\}$ . By assumption,  $\tilde{\mu}_1(a) = \eta_1(a)$ ; hence, by the first paragraph of this proof,  $R$  is nonvoid. Let  $r_0 = \max r \in R$ . Let  $S = \{s \mid \text{there exists a}$

$u(x) \in U$  with a zero of order  $r_0$  at  $x = a$  and such that  $u^{(r_0)}(x)$  has  $s$  zeros (counting multiplicities) in  $(a, \tilde{\mu}_1(a))$ . Let  $s_0 = \max s \in S$ . We now apply the argument used in the proof of Theorem 8 to obtain a contradiction.

**COROLLARY 1.** *If  $\phi(x)$  is a nontrivial solution of (1) with a zero of order  $k$  at  $x = a$  and such that  $\phi^{(k)}(x)$  has a zero of order  $n - k$  at  $x = \mu_1(a)$ , where  $k$  is maximal over all such solutions, then  $\phi^{(i)}(x) \neq 0$  in  $(a, \mu_1(a)]$  ( $i = 0, 1, \dots, k - 1$ ) and  $\phi^{(k)}(x) \neq 0$  in  $(a, \mu_1(a))$ .*

*Proof.* Theorem 8 tells us that  $\phi^{(k)}(x) \neq 0$  in  $(a, \mu_1(a))$ ; and if  $\phi^{(i)}(x) = 0$  at some point  $x \in (a, \mu_1(a)]$ , ( $0 \leq i \leq k - 1$ ), Rolle's Theorem allows us to conclude that there is a point  $x_0 \in (a, \mu_1(a))$  where  $\phi^{(k)}(x_0)$  is zero.

**COROLLARY 2.** *The conclusions of Theorems 8 and 9 are valid under the weaker conditions that there exists a nontrivial solution  $\phi(x)$  of (1) satisfying the following:  $\phi(x)$  has a zero of order  $q$  at  $x = a$ , and either*

- (i)  $\phi(x)$  has  $n - 1$  zeros (counting multiplicities) on  $[a, \infty)$  and there exists a  $p$  ( $1 \leq p \leq q$ ) such that  $\phi^{(p)}(x)$  vanishes at least once past the  $(n - 1)^{\text{st}}$  zero; or
- (ii) there exists a  $p$  ( $0 \leq p \leq q$ ) such that  $\phi^{(p)}(x)$  has  $n - q$  zeros (counting multiplicities) on  $(a, \infty)$ .

*Proof.* As in the proof of Theorem 9, Rolle's Theorem guarantees the existence of  $\tilde{\mu}_1(a)$ . The remainder of the proof is identical with that of Theorem 8.

**4. An eigenfunction result.** The basic relationships between conjugate points of equation (2) and eigenvalues of the associated boundary value problem were investigated in detail by Leighton and Nehari [11]. Subsequently Howard [8], using variational methods, discussed the relationships between focal points of (2) and eigenvalues of an eigenvalue problem, with focal point boundary conditions, associated with (2), making various assumptions on the coefficients. Barrett [3, 5] extended these results to fourth order equations of more general form. These results were extended by Reid [13] to more general even order self-adjoint differential equations.

After obtaining a general relationship between the solutions of (1) and those of

$$(3) \quad L_n^+ y = \sum_{i=0}^n (-1)^i (r_i y)^{(i)} = 0$$

(the Lagrangian adjoint equation), we shall derive an eigenvalue relationship, from which we will develop a disconjugacy criterion. It should be noted that Theorem 10 is but a part of the more general theory of boundary operators ([10, Chapter 9], [6, Chapter 11]) included here to demonstrate the intimate relationship between boundary value problems associated with equation (1) and those associated with equation (3).

**THEOREM 10.** *If  $u(x)$  is a solution of (1) with a zero of order exactly  $k$  at  $x = a$  and a zero of order  $n - k$  at  $x = b$ , then there exists a nontrivial solution  $v(x)$  of (3) having a zero of order  $k$  at  $x = b$  and a zero of order  $n - k$  at  $x = a$ . Further, if  $k$  is maximal in the sense of Theorem 5, then  $v(x)$  is essentially unique (i.e., unique except for multiplication by a constant factor).*

*Proof.* The operators  $L$  and  $L^+$  in the equations (1) and (3) are connected by Green's identity:

$$\begin{aligned}
 (4) \quad (vL_n, u) - (u, L_n^+v) &= \int_a^b (vL_n u - uL_n^+v) dx \\
 &= \sum_{i=1}^n \left( \sum_{j=0}^{i-1} (-1)^j (vr_i)^{(j)}(x) u^{(i-j-1)}(x) \right) \Big|_a^b.
 \end{aligned}$$

Let  $u$  be the solution of  $L_n y = 0$  whose existence we have assumed; let  $v$  be a nontrivial solution of  $L_n^+ y = 0$  with a zero of order  $n - k - 1$  at  $x = a$  and a zero of order  $k$  at  $x = b$ . Then, upon applying Theorem 2, (4) becomes

$$0 = (-1)^{n-k} (vr_n)^{(n-k-1)}(a) u^{(k)}(a).$$

Since  $u^{(k)}(a) \neq 0$  by hypothesis, this implies that  $(vr_n)^{n-k-1}(a) = 0$ ; hence, by the corollary to Theorem 2,  $v^{(n-k-1)}(a) = 0$ , and  $v(x)$  has a zero of order  $n - k$  at  $x = a$ .

To prove the second part of the theorem suppose  $k$  is maximal and  $v(x)$  is not unique; i.e., suppose that there are two nontrivial solutions  $v_1(x)$  and  $v_2(x)$  of (3) with zeros of order  $n - k$  at  $x = a$ , and zeros of order  $k$  at  $x = b$ . Either  $v_1(x), v_2(x)$  or a linear combination of  $v_1(x)$  and  $v_2(x)$  has a zero of order  $k + 1$  at  $x = b$  and a zero of order  $n - k$  at  $a$ . Let  $v(x)$  be the nontrivial solution of (3) with a zero of order at least  $n - k$  at  $x = a$  and a maximal order zero, at  $x = b$ . Interchanging the roles of  $u$  and  $v$  in the previous argument, we find that there is a solution  $u(x)$  with a zero of order  $k + 1$  at  $x = a$  and a zero of order  $n - k - p$  at  $x = b$ , where  $p > 0$ ; this contradicts the maximality of  $k$ .

We now derive an eigenfunction relationship between equations (1) and (3).

**THEOREM 11.** *Consider the adjoint  $n^{\text{th}}$  order linear differential operators  $L_n$  and  $L_n^+$  and the adjoint problems*

$$(5) \quad Ly = \lambda y, \quad Uy = 0,$$

$$(6) \quad L^+y = \lambda y, \quad U^+y = 0,$$

where  $U$  and  $U^+$  are self-adjoint or regular adjoint linear homogeneous boundary operators and  $\lambda$  is a parameter. Suppose further that Green's function for (5) has only simple poles. Let the eigenfunctions of (5) and (6) be  $\phi_i(x)$  and  $\phi_i^+(x)$  respectively, normalized so that  $(\phi_i, \phi_i^+) = 1$ . Let  $f(x)$  and  $g(x)$  be any two functions such that  $L_n g$  exists,  $Ug = 0$ ,  $L_n^+ f$  exists, and  $U^+ f = 0$ . Further let  $a_i = (f, \phi_i)$  and  $b_i = (g, \phi_i^+)$  and let  $\lambda_i$  be the  $i$ th eigenvalue of (5). Then

$$(f, L_n g) = \sum_{i=0}^{\infty} a_i b_i \lambda_i.$$

*Proof.* We freely use the well-known properties of problems (5), (6) (see for instance Ince [5] Chapter 11, and Coddington and Levinson [6] Chapter 12).

$$\begin{aligned} (f, L_n g) &= \left( \sum_{i=0}^{\infty} a_i \phi_i^+, L_n g \right) = \sum_{i=0}^{\infty} a_i (\phi_i^+, L_n g) \\ &= \sum_{i=0}^{\infty} a_i (L_n^+ \phi_i^+, g) = \sum_{i=0}^{\infty} a_i \left( \lambda_i \phi_i^+, \sum_{j=0}^{\infty} b_j \phi_j \right) \\ &= \sum_{i=0}^{\infty} a_i \lambda_i \sum_{j=0}^{\infty} b_j (\phi_i^+, \phi_j) \\ &= \sum_{i=0}^{\infty} a_i \lambda_i \sum_{j=0}^{\infty} b_j \delta_{ij} = \sum_{i=0}^{\infty} a_i b_i \lambda_i. \end{aligned}$$

If we specialize this result to the self-adjoint case, letting  $f = g$  in (7), we obtain

$$(8) \quad (f, L_n f) = \sum_{i=0}^{\infty} a_i^2 \lambda_i.$$

Further suppose  $L_n y = 0$  assumes the special form

$$(9) \quad L_n y = [r_n(x)y^{(n)}]^{(n)} + \sum_{i=0}^{n-1} [(-1)^{n+i+1} r_i(x)y^{(i)}]^{(i)} = 0$$

where  $r_i(x) > 0$  ( $i = 0, 1, \dots, n$ ).

We now state a disconjugacy criterion due to Reid [13].

**THEOREM (Reid [13]).** *Equation (9) is disconjugate in the sense of Reid (i.e., no solution possesses two  $n$ th order zeros) on the interval  $[a, \infty)$  if and only if for all values  $b$  on  $(a, \infty)$  and all functions*



$y(x)$  such that  $y(x) \in C_{n-1}[a, b]$ ,  $y(x)$  is absolutely continuous on  $[a, b]$ ,  $(y^{(n)})^2$  is integrable on  $[a, b]$  and  $y(x)$  has  $n^{\text{th}}$  order zeros at  $x = a$  and  $x = b$ ,  $(y, L_n y) > 0$  (where  $L_n y$  is as in equation (9)).

Using this result we prove the following theorem. (Our impression is that Theorem 12 is in the literature, but the author has been unable to find it).

**THEOREM 12.** *Equation (9) is disconjugate in the sense of Reid if and only if the minimum eigenvalue  $\lambda_0$  is positive (all functions are assumed to be of the class described in the above theorem of Reid).*

*Proof.* Suppose (9) is disconjugate. Letting  $f(x) = \phi_0(x)$  in (8) and applying the theorem of Reid, we have  $0 < (\phi_0, L_n \phi_0) = \lambda_0$ , since  $(\phi_i, \phi_j) = \delta_{ij}$ . The sufficiency is immediate from equation (8).

#### BIBLIOGRAPHY

1. N. V. Azbelev, and Z. B. Chalyuk, *On the question of the distribution of the zeros of solution of a third order linear differential equation*, Mat. Sbornik, **51** (1960), 475-486 (Russian).
2. J. H. Barrett, *Disconjugacy of second-order linear differential equations with non-negative coefficients*, Proc. Amer. Math. Soc. **10** (1959), 552-561.
3. ———, *Disconjugacy of a self-adjoint differential equation of the fourth order*, Pacific J. Math. **11** (1961), 25-37.
4. ———, *Fourth order boundary value problems and comparison theorems*, Canadian J. Math. **13** (1961), 625-638.
5. ———, *Two-point boundary problems for linear self-adjoint differential equations of the fourth order with middle term*, Duke Math. J. **29** (1962), 543-554.
6. E. A. Coddington, and N. Levinson, *Theory of Ordinary Differential Equations*, McGraw-Hill, New York, 1955.
7. M. Hanan, *Oscillation criteria for third-order linear differential equations*, Pacific J. Math. **11** (1961), 919-944.
8. H. Howard, *Oscillation criteria for fourth-order linear differential equations*, Trans. Amer. Math. Soc. **96** (1960), 296-311.
9. R. W. Hunt, *The behavior of solutions of ordinary, self-adjoint differential equations of arbitrary even order*, Pacific J. Math. **12** (1962), 945-961.
10. E. L. Ince, *Ordinary differential equations*, Dover, New York, 1956.
11. W. Leighton, and Z. Nehari, *On the oscillation of solutions of self-adjoint differential equations of the fourth order*, Trans. Amer. Math. Soc. **89** (1958), 325-377.
12. A. Ju. Levin, *Some questions on the oscillations of solutions of linear differential equations*, Doklady Akad. Nauk. **148** (1963), 512-515 (Russian).
13. W. T. Reid, *Oscillation criteria for self-adjoint differential systems*, Trans. Amer. Math. Soc. **101** (1961), 91-106.
14. G. F. Simmons, *Introduction to topology and modern analysis*, McGraw-Hill, New York, 1963.

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California

R. M. BLUMENTHAL

University of Washington  
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California  
Los Angeles, California 90007

\*RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. No separate author's resumé is required. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

\* Basil Gordon, Acting Managing Editor until February 1, 1966.

David R. Arterburn and Robert James Whitley, <i>Projections in the space of bounded linear operators</i> .....	739
Robert McCallum Blumenthal, Joram Lindenstrauss and Robert Ralph Phelps, <i>Extreme operators into <math>C(K)</math></i> .....	747
L. Carlitz, <i>A note on multiple exponential sums</i> .....	757
Joseph A. Cima, <i>A nonnormal Blaschke-quotient</i> .....	767
Paul Civin and Bertram Yood, <i>Lie and Jordan structures in Banach algebras</i> .....	775
Luther Elic Claborn, <i>Dedekind domains: Overrings and semi-prime elements</i> .....	799
Luther Elic Claborn, <i>Note generalizing a result of Samuel's</i> .....	805
George Bernard Dantzig, E. Eisenberg and Richard Warren Cottle, <i>Symmetric dual nonlinear programs</i> .....	809
Philip J. Davis, <i>Simple quadratures in the complex plane</i> .....	813
Edward Richard Fadell, <i>On a coincidence theorem of F. B. Fuller</i> .....	825
Delbert Ray Fulkerson and Oliver Gross, <i>Incidence matrices and interval graphs</i> .....	835
Larry Charles Grove, <i>Tensor products over <math>H^*</math>-algebras</i> .....	857
Deborah Tepper Haimo, <i><math>L^2</math> expansions in terms of generalized heat polynomials and of their Appell transforms</i> .....	865
I. Martin (Irving) Isaacs and Donald Steven Passman, <i>A characterization of groups in terms of the degrees of their characters</i> .....	877
Donald Gordon James, <i>Integral invariants for vectors over local fields</i> .....	905
Fred Krakowski, <i>A remark on the lemma of Gauss</i> .....	917
Marvin David Marcus and H. Minc, <i>A subdeterminant inequality</i> .....	921
Kevin Mor McCrimmon, <i>Norms and noncommutative Jordan algebras</i> .....	925
Donald Earl Myers, <i>Topologies for Laplace transform spaces</i> .....	957
Olav Njstad, <i>On some classes of nearly open sets</i> .....	961
Milton Philip Olson, <i>A characterization of conditional probability</i> .....	971
Barbara Osofsky, <i>A counter-example to a lemma of Skornjakov</i> .....	985
Sidney Charles Port, <i>Ratio limit theorems for Markov chains</i> .....	989
George A. Reid, <i>A generalisation of <math>W^*</math>-algebras</i> .....	1019
Robert Wells Ritchie, <i>Classes of recursive functions based on Ackermann's function</i> .....	1027
Thomas Lawrence Sherman, <i>Properties of solutions of <math>n</math>th order linear differential equations</i> .....	1045
Ernst Snapper, <i>Inflation and deflation for all dimensions</i> .....	1061
Kondagunta Sundaresan, <i>On the strict and uniform convexity of certain Banach spaces</i> .....	1083
Frank J. Wagner, <i>Maximal convex filters in a locally convex space</i> .....	1087
Joseph Albert Wolf, <i>Translation-invariant function algebras on compact groups</i> .....	1093