

# Pacific Journal of Mathematics

**BOUNDARY MEASURES OF ANALYTIC DIFFERENTIALS AND  
UNIFORM APPROXIMATION ON A RIEMANN SURFACE**

Laura Ketchum Kodama

# BOUNDARY MEASURES OF ANALYTIC DIFFERENTIALS AND UNIFORM APPROXIMATION ON A RIEMANN SURFACE

LAURA KETCHUM KODAMA

**A classical theorem of F. and M. Riesz establishes a one-to-one correspondence between analytic differentials of class  $H_1$  on the interior of the unit disc and finite complex-valued Borel measures on the boundary of the disc which are orthogonal to polynomials. The main result of this paper gives a similar correspondence when the unit disc is replaced by a compact subset, satisfying a finite connectivity condition, of any noncompact Riemann surface. The analytic differentials on the interior of the set satisfy a boundedness condition analogous to the classical  $H_1$  differentials and the measures on the boundary of the set are those orthogonal to all meromorphic functions with a finite number of poles in the complement of the set. This result is then used to obtain theorems on uniform approximation on the set by such meromorphic functions.**

This paper extends results of Bishop in [2] and [5] where he considers compact subsets of the plane satisfying a simple connectivity condition.<sup>1</sup> He obtained such a one-to-one correspondence between boundary measures and analytic differentials and used his result together with an approximation theorem for nowhere dense sets to give a proof of Mergelyan's approximation theorem [6]. We are able to extend Mergelyan's theorem to our more general sets and also show that "local" approximation implies approximation on the whole set.

## I. Boundary measures of analytic differentials.

### A. DEFINITIONS AND PRELIMINARIES.

In this section  $S$  will denote an open Riemann surface. If  $K$  is a compact subset of  $S$ , we denote by  $C(K)$  the algebra of all continuous complex-valued functions on  $K$  with norm  $\|f\| = \sup_{x \in K} |f(x)|$ , and by  $A(K)$  the closed subalgebra of  $C(K)$  consisting of those functions which are limits of meromorphic functions on  $S$  with finitely many poles in

---

Received January 10, 1964. This paper is a portion of a doctoral dissertation written at the University of California at Berkeley under the direction of Professor Errett Bishop. The author wishes to thank Professor Bishop for proposing the problem and for his helpful suggestions and encouragement. This work was partially supported by contracts NSFG-9103 and NSFG-18974.

<sup>1</sup> The case with smooth boundary is discussed by Royden in [7].

$S \sim K$ . By Runge's Theorem when  $S$  is the plane, or by the extension of Runge's Theorem due to Behnke and Stein [1, p. 445 and p. 456] in the general case,  $A(K)$  can also be characterized as all functions of  $C(K)$  which are uniform limits on  $K$  of functions analytic in a neighborhood of  $K$ .

The sets for which our results are obtained are defined as follows.

**DEFINITION 1.** A compact subset  $K$  of  $S$  will be called *n-balanced* if there exists a finite family  $\{U_i\}_{i=1}^n$  of components of  $S \sim K$  such that any point of the boundary of  $K$  lies on the boundary of one of the  $U_i$ . An open subset of  $S$  will be called *n-balanced* if it is the interior of its closure and its closure is a compact *n-balanced* set.

The following properties are clear.

**LEMMA 1.** *The interior of a compact n-balanced set is an open m-balanced set for some  $m \leq n$ . The boundary of a compact n-balanced set is a nowhere dense compact n-balanced set.*

The measures on the boundary of  $K$  to be considered are now defined.

**DEFINITION 2.** If  $K$  is a compact subset of  $S$ , we denote by  $M(K)$  all finite complex-valued Borel measures  $\mu$  on the boundary of  $K$  such that  $\int f d\mu = 0$  for all  $f \in A(K)$ .

Several preliminary definitions will be necessary to describe the boundedness condition on the analytic differentials to be studied.

By an arc we will mean a continuous map  $f: [a, b] \rightarrow S$  of a closed interval  $a \leq t \leq b$  into  $S$ . We will identify arcs  $f: [a, b] \rightarrow S$  and  $g: [c, d] \rightarrow S$  whenever  $b - a = d - c$  and  $g(t) = f(t + a - c)$ . The image of  $[a, b]$  under  $f$  will be denoted by  $|f|$ . By a subarc of  $f$  we mean the restriction of  $f$  to a subinterval  $[c, d]$ ,  $a \leq c < d \leq b$ . If  $g: [a_1, b_1] \rightarrow S$  is such that  $f(b) = g(a_1)$  then by the product of  $f$  and  $g$ , written  $fg$ , we mean the arc  $h: [a, b + b_1 - a_1] \rightarrow S$  defined by

$$h(t) = \begin{cases} f(t) & \text{if } a \leq t \leq b \\ g(t + a_1 - b) & \text{if } b \leq t \leq b + b_1 - a_1. \end{cases}$$

An arc  $f: [a, b] \rightarrow S$  is an analytic arc if  $f$  can be extended to be analytic with nonzero derivative in a neighborhood of  $[a, b]$ . A piecewise analytic arc is a product of a finite number of analytic arcs. A simple closed curve is an arc  $f: [a, b] \rightarrow S$  such that  $f(a) = f(b)$ , and if  $x \neq a$  and  $x \neq b$  then  $f(x) \neq f(a)$  and  $f$  is one-to-one on the open interval  $(a, b)$ .

DEFINITION 3. If  $U$  is an open subset of  $S$  we say that a sequence  $\{\gamma_i\}$  delimits  $U$  if

(i) each  $\gamma_i$  is a finite family of disjoint piecewise analytic simple closed curves  $\alpha_{ij}$  such that  $|\alpha_{ij}| \subset U$  and  $\bigcup_j |\alpha_{ij}|$  is the boundary of an open set  $V_i \subset U$  and each  $\alpha_{ij}$  is positively oriented with respect to  $V_i$ .

(ii) if  $T$  is any compact subset of  $U$ , then for all sufficiently large  $i$ ,  $T \subset V_i$ .

DEFINITION 4. If  $U$  is an open subset of  $S$  with compact closure  $K$  and  $\gamma$  is a finite family of piecewise analytic curves  $\alpha_j$  such that  $|\alpha_j| \subset U$  and  $\omega$  is an analytic differential on  $U$ , we denote by  $\|\omega\|_\gamma$  the norm of the linear functional  $F$  on  $C(K)$  defined by  $F(h) = \int_\gamma h\omega$ .

DEFINITION 5. Let  $U$  be an open subset of  $S$  with compact closure. The class  $H(U)$  consists of all analytic differentials  $\omega$  on  $U$  such that there exists a sequence  $\{\gamma_i\}$  which delimits  $U$  and an  $M > 0$  such that  $\|\omega\|_{\gamma_i} < M$  for all  $i$ .

Our aim is to establish, in case  $K$  is an  $n$ -balanced set, a one-to-one correspondence between  $M(K)$  and  $H(U)$ , where  $U$  is the interior of  $K$ . The correspondence will be between a differential and its boundary measure, in the following sense.

DEFINITION 6. Let  $U$  be an open subset of  $S$  with compact closure and let  $B$  be its boundary. A finite complex-valued Borel measure  $\mu$  on  $B$  is said to be a boundary measure of  $\omega \in H(U)$  if the sequence of Definition 5 can be chosen so that

$$\int_{\gamma_i} h\omega \rightarrow \int h d\mu \quad \text{as } i \rightarrow \infty$$

for all  $h \in C(U \cup B)$ .

We do not need any restrictions on  $K$  other than compactness in order to show the existence of a boundary measure for every differential  $\omega \in H(U)$ . The following theorem has the same proof as Theorem 1 in [5].

**THEOREM 1.** *Let  $U$  be an open subset of  $S$  with compact closure  $K$ . Then any  $\omega \in H(U)$  has a boundary measure  $\mu \in M(K)$ .*

In order to "fit together" sequences which delimit two different open sets to obtain a sequence which delimits the union, we will need the following lemma.

**LEMMA 2.** *Let  $\gamma$  and  $\delta$  each be a finite family of disjoint piece-*

wise analytic simple closed curves,  $\alpha_j$  and  $\beta_j$  respectively, such that  $\bigcup_j |\alpha_j|$  is the boundary of an open set  $\Gamma$  with each  $\alpha_j$  positively oriented with respect to  $\Gamma$  and similarly  $\bigcup_j |\beta_j|$  is the boundary of an open set  $\Delta$  with each  $\beta_j$  positively oriented with respect to  $\Delta$ . Then there exists a finite collection of analytic coordinate functions  $h_i$  with domain  $V_i$ ,  $V_i$  a neighborhood of a point  $p_i \in S$  (the  $p_i$  need not be distinct), so that given any neighborhood  $U_i$  of  $h_i(p_i)$  such that  $U_i \subset h_i(V_i)$  and any  $\varepsilon_i > 0$ , there exists  $\varphi$ , a finite family of disjoint piecewise analytic simple closed curves  $\psi_j$ , such that  $\bigcup_j |\psi_j|$  is the boundary of an open set  $\Phi$  and

- (i) each  $\psi_j$  is positively oriented with respect to  $\Phi$
- (ii) each  $\psi_j$  is the product of a finite number of subarcs, each of which is either a subarc of some  $\alpha_j$  or  $\beta_j$  or is an arc  $f$  such that for some  $i$ , the arc  $h_i \circ f$  has length less than  $\varepsilon_i$  and  $|h_i \circ f| \subset U_i$ .
- (iii)  $\Gamma \cup \Delta \subset \Phi \subset \Gamma \cup \Delta \cup \bigcup_{i=1}^n h_i^{-1}(U_i)$

The proof is left as an exercise for the reader.<sup>2</sup>

## B. PLANE SETS.

In this section we consider the special case where  $S$  is the plane.

The proofs of the following lemma and theorem are the same as Lemma 4 and Theorem 1 in [5].

**LEMMA 3.** *If  $K$  is a compact  $n$ -balanced subset of the plane and if  $\mu$  and  $\nu$  are both in  $M(K)$  and  $\int (t-z)^{-1} d\mu(t) = \int (t-z)^{-1} d\nu(t)$  for all  $z$  in the interior of  $K$ , then  $\mu = \nu$ .*

**THEOREM 2.** *Let  $U$  be an  $n$ -balanced open subset of the plane and  $K$  be its closure. Then given  $\omega \in H(U)$ , its boundary measure, which exists by Theorem 1, is unique and if  $\omega = f(z)dz$  then*

$$f(z) = (2\pi i)^{-1} \int (t-z)^{-1} d\mu(t)$$

for all  $z \in U$ .

The next lemma is a modification of Lemma 6 in [3]. The assumption that  $\nu$  is orthogonal to all functions analytic in a neighborhood of  $K$  rather than just all polynomials enables us to obtain the measure  $\beta_{x_0}$  with support in  $K$ . The proof is not given, as the same proof applies with only obvious minor modifications and we prove a general version for any open Riemann surface as Lemma 5 below.

<sup>2</sup> A proof may be found in the author's thesis.

LEMMA 4. *Let  $K$  be a compact subset of the complex plane. Let  $\nu$  be a measure on  $K$  orthogonal to  $A(K)$ . Then for almost all real numbers  $x_0$ , there exists a measure  $\beta_{x_0}$  on the set  $K \cap \{z : \operatorname{Re} z = x_0\}$  such that*

$$\int_{R_{x_0}} h d\nu = - \int_{L_{x_0}} h d\nu = \int h d\beta_{x_0}$$

for all  $h \in A(K)$ , where

$$R_{x_0} = K \cap \{z : \operatorname{Re} z \geq x_0\} \quad \text{and} \quad L_{x_0} = K \cap \{z : \operatorname{Re} z \leq x_0\} .$$

THEOREM 3. *Let  $K$  be a compact  $n$ -balanced subset of the complex plane with interior  $U$ . Then if  $\mu \in M(K)$ , there exists an analytic differential  $\omega \in H(U)$  such that  $\mu$  is the boundary measure of  $\omega$ .*

*Proof.* The proof is by induction on  $n$ . If  $n = 1$ ,  $K$  is balanced in the sense of [5] and Theorem 3 of [5] is the required result.

Suppose for  $n > 1$  the theorem is true for  $m$ -balanced sets for all  $m < n$ . For  $z \in U$ , define

$$f(z) = (2\pi i)^{-1} \int (t - z)^{-1} d\mu(t) .$$

Now suppose  $x_0$  is as in Lemma 4 and furthermore that  $\{z : \operatorname{Re} z = x_0\}$  intersects the interior of at least one of the bounded components  $U_i$  of Definition 1. Then  $L_{x_0}$  and  $R_{x_0}$  are both  $m$ -balanced for some  $m < n$ . Thus since  $\mu|_{L_{x_0}} + \beta_{x_0} \in M(L_{x_0})$  and  $\mu|_{R_{x_0}} - \beta_{x_0} \in M(R_{x_0})$  by Lemma 4 and Runge's theorem, the induction hypothesis applies and they are boundary measures of analytic differentials  $f_1(z)dz$  and  $f_2(z)dz$  respectively.

For  $z$  in the interior of  $L_{x_0}$ ,

$$\begin{aligned} f_1(z) &= (2\pi i)^{-1} \int (t - z)^{-1} d(\mu|_{L_{x_0}} + \beta_{x_0})(t) \\ &= (2\pi i)^{-1} \int (t - z)^{-1} d(\mu|_{L_{x_0}} + \beta_{x_0})(t) \\ &\quad + (2\pi i)^{-1} \int (t - z)^{-1} d(\mu|_{R_{x_0}} - \beta_{x_0})(t) \\ &= (2\pi i)^{-1} \int (t - z)^{-1} d\mu(t) = f(z) \end{aligned}$$

and for  $z$  in the interior of  $R_{x_0}$  we have similiary

$$f_2(z) = (2\pi i)^{-1} \int (t - z)^{-1} d\mu(t) = f(z) .$$

Now let  $x_0 < x_1$  both restricted as above. Then  $\mu | R_{x_0} - \beta_{x_0}$  is a boundary measure for  $f(z)dz$  on the set  $R_{x_0}$ . Denote the delimiting sequence by  $\{\gamma_j\}$ . Also  $\mu | L_{x_1} + \beta_{x_1}$  is a boundary measure for  $f(z)dz$  on the set  $L_{x_1}$ . Denote the delimiting sequence by  $\{\delta_j\}$ . Suppose  $\Gamma_j$  is the open set bounded by  $\gamma_j$  and  $\Delta_j$  the open set bounded by  $\delta_j$ , as required in Definition 3. We apply Lemma 2 to  $\gamma_j, \delta_j, \Gamma_j, \Delta_j$  where  $U_i$  are chosen so that  $h_i^{-1}(U_i) \subset U$  and  $\varepsilon_i$  chosen so that the length of the arc in  $U_i$  which is not from  $\delta_j$  or  $\gamma_j$  is less than  $\eta_i$  and  $\sum \eta_i \sup_{z \in U_i} |f(z)| < 1$ .

The lemma yields  $\varphi_j$  a finite union of disjoint piecewise analytic simple closed curves in  $U$  which form the boundary of the open set  $\Phi_j$ , and  $\Gamma_j \cup \Delta_j \subset \Phi_j \subset U$ . If  $S$  is a compact subset of  $U$ , let  $x_0 < x_2 < x_1$ . Then  $S_1 = S \cap \{z : \operatorname{Re} z \leq x_2\}$  is a compact subset of the interior of  $L_{x_1}$  and  $S_2 = S \cap \{z : \operatorname{Re} z \geq x_2\}$  is a compact subset of the interior of  $R_{x_0}$ . Thus for all  $j$  sufficiently large,

$$S_1 \subset \Delta_j \quad \text{and} \quad S_2 \subset \Gamma_j \quad \text{and} \quad S = S_1 \cup S_2 \subset \Delta_j \cup \Gamma_j \subset \Phi_j .$$

Therefore  $\{\varphi_j\}$  is a delimiting sequence for  $U$ . Furthermore,

$$\begin{aligned} \|\omega\|_{\varphi_j} &= \int_{\varphi_j} |f(z)| |dz| \leq \int_{\gamma_j} |f(z)| |dz| + \int_{\delta_j} |f(z)| |dz| \\ &\quad + \sum \eta_i \sup_{z \in U_i} |f(z)| \leq \|\omega\|_{\gamma_j} + \|\omega\|_{\delta_j} + 1 . \end{aligned}$$

Thus  $\omega \in H(U)$ .

By Theorems 1 and 2 there exists a boundary measure  $\nu$  on the closure of  $U$  such that for  $z \in U$ ,

$$f(z) = (2\pi i)^{-1} \int (t - z)^{-1} d\nu(t) \quad \text{and} \quad \nu \in M(\operatorname{clsr} U) \subset M(K) .$$

Applying Lemma 3 to  $\mu$  and  $\nu$  we see that  $\mu = \nu$  and thus  $\mu$  is the boundary measure of  $\omega$ .

### C. SUBSETS OF AN OPEN RIEMANN SURFACE.

In this section we consider the general case where  $S$  is any open Riemann surface. The function  $(t - z)^{-1}$  used in the plane case must be replaced by the elementary differential of Behnke and Stein [1]. The result needed is the following: there exists an elementary differential  $\alpha(p)$  which for fixed  $p$  is a meromorphic differential on  $S$  with exactly one pole, a simple pole at  $p$  with residue 1. Furthermore, if  $h$  is an analytic coordinate function on an open set  $V \subset S$  and  $\alpha(p) = A(z, p)dz$  on  $h(V)$ , then  $A(z, p)$  is meromorphic in  $p$  on  $S$  with exactly one pole, a simple pole at  $h^{-1}(z)$ . Thus if  $h^{-1}(z_0) \notin K$ ,  $A(z_0, p) \in A(K)$ .

We prove the following generalization of Lemma 4.

LEMMA 5. *Let  $K$  be a compact subset of  $S$ . Let  $\nu$  be a measure on  $K$  orthogonal to  $A(K)$ . Then if  $f$  is a nonconstant function analytic on  $S$ , for almost all real numbers  $x_0$ , there exists a measure  $\beta_{x_0}$  on the set  $K \cap \{p : \operatorname{Re} f(p) = x_0\}$  such that*

$$\int_{R_{x_0}} h d\nu = - \int_{L_{x_0}} h d\nu = \int h d\beta_{x_0}$$

for all  $h \in A(K)$  where

$$L_{x_0} = \{p : \operatorname{Re} f(p) \leq x_0\} \cap K \text{ and } R_{x_0} = \{p : \operatorname{Re} f(p) \geq x_0\} \cap K .$$

*Proof.* Since  $f$  is nonconstant, for all but finitely many real numbers  $x$ , the differential of  $f$  does not vanish on  $K \cap \{p : \operatorname{Re} f(p) = x\}$ . Let  $x_1$  have this property and let  $x_2 > x_1$  be such that the differential of  $f$  does not vanish on  $K \cap \{p : x_1 \leq \operatorname{Re} f(p) \leq x_2\}$ . Since the differential of  $f$  does not vanish, there exists a neighborhood of any point of  $K \cap \{p : x_1 \leq \operatorname{Re} f(p) \leq x_2\}$  on which  $f$  is a coordinate function. Cover  $K \cap \{p : x_1 \leq \operatorname{Re} f(p) \leq x_2\}$  by finitely many neighborhoods  $\{U_i\}_{i=1}^n$  such that the closure of  $U_i$  is compact and contained in  $V_i$  and  $f$  is a coordinate function on  $V_i$ . Denote by  $f_i^{-1}$  the inverse of  $f$  as a coordinate function on  $V_i$ .

There exists a nonnegative measure  $\mu$  on  $K$  such that  $|\nu(B)| \leq \mu(B)$  for all Borel sets  $B$ . Let  $\phi$  be the nonnegative, nondecreasing function defined by  $\phi(x) = \mu(\{p : \operatorname{Re} f(p) \leq x\})$ . Then  $\phi'(x)$  will exist for almost all  $x$ . Let  $x_0$  be such that  $\phi'(x_0)$  exists and  $x_1 \leq x_0 \leq x_2$ . Thus  $\nu$  vanishes on all subsets of  $L_{x_0} \cap R_{x_0}$  and since  $h \in A(K)$  implies  $\int h d\nu = 0$  we have

$$\int_{R_{x_0}} h d\nu = - \int_{L_{x_0}} h d\nu \text{ for all } h \in A(K) .$$

Suppose now that  $h$  is a meromorphic function with finitely many poles outside  $K$ . Let  $W$  be an open neighborhood of  $K$  on which  $h$  is analytic. Let  $W_i = W \cap U_i$ . Choose  $\varepsilon, 0 < \varepsilon < 1$  and let

$$T = \bigcup_{i=1}^n \{p \in W_i : \operatorname{Re} f(p) = x_0 \text{ and } \operatorname{dist}(f(p), f(K \cap W_i)) \leq \varepsilon\} .$$

Let  $\|h\| = \sup_{p \in T} |h(p)|$ .

If  $\operatorname{Re} f(p) > x_0$ , define  $h_1(p) = (2\pi i)^{-1} \int_T h \alpha(p)$  and if  $\operatorname{Re} f(p) < x_0$ , define  $h_2(p) = (2\pi i)^{-1} \int_T h \alpha(p)$  where in each case integration is in a positive direction with respect to  $\{p : \operatorname{Re} f(p) \leq x_0\}$ . Suppose  $p_0$  is interior to  $T$  relative to  $\{z : \operatorname{Re} z = x_0\}$ . Then for some  $i_0$ ,  $f(p_0)$  is



interior to  $f(W_{i_0} \cap T)$  relative to  $\{z : \operatorname{Re} z = x_0\}$ . Let  $\tau_{i_0} = f(T \cap W_{i_0})$ . Since the  $W_i$  cover  $T$ , we can choose, for  $i \neq i_0$ , measurable sets  $\tau_i \subset \{z : \operatorname{Re} z = x_0\} \cap f(W_i)$  so that  $f_i^{-1}(\tau_i)$  are pairwise disjoint and each is disjoint from  $f_{i_0}^{-1}(\tau_{i_0})$  and so that  $T = \bigcup_{i=1}^n f_i^{-1}(\tau_i)$ . Then if  $p \in U_{i_0}$ ,  $(2\pi i)^{-1} \int_T h\alpha(p)$  becomes

$$(2\pi i)^{-1} \int_{\tau_{i_0}} h(f_{i_0}^{-1}(\zeta))(\zeta - f(p))^{-1} d\zeta + (2\pi i)^{-1} \sum_{i=1}^n \int_{\tau_i} h(f_i^{-1}(\zeta))g_i(f(p), \zeta) d\zeta$$

where  $g_i$  is analytic in  $f(\operatorname{clsr} U_{i_0})$  in the first variable and in  $f(\operatorname{clsr} U_i)$  in the second variable. The first term has continuous boundary values both from the right and the left at  $p_0$  with difference  $h(p_0)$  and the integrals in the summation are all continuous in  $p$  at  $p_0$ . Thus  $h_1$  and  $h_2$  have continuous boundary values  $h_1(p_0)$  and  $h_2(p_0)$  and

$$h_1(p_0) - h_2(p_0) = h(p_0) .$$

If we define  $h_1(p) = h(p) + h_2(p)$  in  $\operatorname{Re} f(p) < x_0$  and  $h_2(p) = h(p) + h_1(p)$  in  $\operatorname{Re} f(p) > x_0$ , then  $h_1$  and  $h_2$  are analytic in a neighborhood of  $K$  and  $h = h_1 - h_2$ . Thus

$$\int_{R_{x_0}} h d\nu = \int_{R_{x_0}} h_1 d\nu - \int_{R_{x_0}} h_2 d\nu = \int_{R_{x_0}} h_1 d\nu + \int_{L_{x_0}} h_2 d\nu$$

and

$$\begin{aligned} \left| \int_{R_{x_0}} h_1 d\nu \right| &= \left| \int_{R_{x_0}} \left[ (2\pi i)^{-1} \int_T h\alpha(p) \right] d\nu(p) \right| \\ &\leq \int_{\{p : \operatorname{Re} f(p) > x_0\}} \left| \int_T h\alpha(p) \right| d\mu(p) \\ &\quad + \sum_{i=1}^n \int_{R_{x_0} \cap U_i} \left| \int_T h\alpha(p) \right| d\mu(p) . \end{aligned}$$

Cover  $K \cap \{p : \operatorname{Re} f(p) \geq x_0\}$  by a finite number of open analytic neighborhoods, which are the domains of analytic coordinate functions  $\psi_k$ , each with range the unit circle  $D$ . Continuing the inequalities we have

$$\begin{aligned} &\leq \|h\| \sum_{i=1}^n \int_{R_{x_0} \cap U_i} \left( \int_{\tau_i} |\zeta - f(p)|^{-1} d\zeta \right) d\mu(p) \\ &\quad + \|h\| \sum_{i=1}^n \int_{R_{x_0} \cap U_i} \left( \sum_{j=1}^n \int_{\tau_j} |g_{ij}(f(p), \zeta)| d\zeta \right) d\mu(p) \\ &\quad + \|h\| \sum_k \int_{\psi^{-1}(D)} \left( \sum_{i=1}^n \int_{\tau_i} |\gamma_{ki}(\psi_k(p), \zeta)| d\zeta \right) d\mu(p) \end{aligned}$$

where  $g_{ij}$  is analytic in the first variable in  $f(\operatorname{clsr} U_i)$  and in the

second variable in  $f(\text{clsr } U_j)$  and  $\gamma_{ki}$  is analytic in the first variable in  $D$  and the second variable in  $f(\text{clsr } U_i)$ . The  $g_{ij}$  and  $\gamma_{ki}$  are therefore bounded and we have a constant  $L$ , independent of  $\varepsilon$ , so that the above expression is less than or equal to

$$\|h\| \left\{ n \int_{x_0}^{x_2} \int_{-M}^M [(x_0 - x)^2 + t^2]^{-1/2} dt d\phi(x) + L \right\}$$

where  $M$  is chosen, independent of  $\varepsilon$ , so that if  $y = \text{Im } f(p)$  and  $v = \text{Im } \zeta$  where  $p \in R_{x_0}$  and  $\zeta$  is in some  $\tau_i$ , then  $|y - v| < M$ .

A bound  $N$ , independent of  $\varepsilon$ , is found for

$$\int_{x_0}^{x_2} \int_{-M}^M [(x_0 - x)^2 + t^2]^{-1/2} dt d\phi(x)$$

as in [3, p. 42]. Thus

$$\left| \int_{R_{x_0}} h_1 d\nu \right| \leq \|h\| (nN + L)$$

and a similar estimate can be made for  $\left| \int_{L_{x_0}} h_2 d\nu \right|$ . Combining these we have

$$\left| \int_{R_{x_0}} h d\nu \right| \leq Q \|h\| .$$

where  $Q$  is independent of  $\varepsilon$ , and thus

$$\left| \int_{R_{x_0}} h d\nu \right| \leq Q \sup \{ |h(p)| : p \in K \cap \{p : \text{Re } f(p) = x_0\} \} .$$

Therefore  $h \rightarrow \int_{R_{x_0}} h d\nu$  is a bounded linear functional on a dense subset of  $A(K) | K \cap \{p : \text{Re } f(p) = x_0\}$  and therefore on  $A(K) | K \cap \{p : \text{Re } f(p) = x_0\}$ . By the Hahn-Banach theorem we can extend this bounded linear functional to  $C(K \cap \{p : \text{Re } f(p) = x_0\})$  and then apply the Riesz representation theorem to obtain the desired measure  $\beta_{x_0}$ .

**LEMMA 6.** *Suppose  $K$  is an  $n$ -balanced compact subset of  $S$ . Suppose  $f$  is a nonconstant analytic function on  $S$  and  $K_1 = K \cap \{p : \text{Re } f(p) \geq x_0\}$ . Then  $K_1$  is a compact  $m$ -balanced set for some  $m \leq n$ .*

*Proof.*  $K_1$  is clearly compact.

Let  $\{U_i\}_{i=1}^n$  be the finite set of components of  $S \sim K$  from Definition 1. A point  $q$  on the boundary of  $K_1$  is either on the boundary

of  $K$  or in the intersection of the interior of  $K$  with the boundary of  $\{p: \operatorname{Re} f(p) \geq x_0\}$ . In the former case,  $q$  is on the boundary of some  $U_i$  and therefore on the boundary of the component of  $S \sim K$  which contains  $U_i$ , which we call  $V_i$ . There are  $n$   $U_i$  and therefore  $m$   $V_i$  with  $m \leq n$ . In the later case,  $q$  is on the boundary of some component  $Q$  of  $\{p: \operatorname{Re} f(p) < x_0\}$ . Suppose  $Q \subset K$ , then  $\operatorname{clsr} U \subset K$  and  $\operatorname{clsr} Q$  is compact.  $Q$  is open, so  $\operatorname{Re} f(p) = x_0$  on the boundary of  $Q$ . Since  $\operatorname{clsr} Q$  is compact,  $\operatorname{Re} f(p)$  must assume its minimum on  $\operatorname{clsr} Q$  is compact,  $\operatorname{Re} f(p)$  must assume its minimum on  $\operatorname{clsr} Q$  and by the minimum modulus theorem for real parts of analytic functions, the minimum must be assumed on the boundary, but there  $\operatorname{Re} f(p) = x_0$ . Thus  $\operatorname{Re} f(p) \geq x_0$  on  $Q$  which is a contradiction. Since  $Q$  is not contained in  $K$ , it must intersect some  $U_i$ . Therefore  $Q \subset V_i$  and  $q$  is on the boundary of  $V_i$ . This shows  $K_1$  is  $m$ -balanced.

LEMMA 7. *Under the hypotheses of Lemma 5, the measure  $\nu | R_{x_0} - \beta_{x_0}$  is orthogonal to  $A(R_{x_0})$  and the measure  $\nu | L_{x_0} + \beta_{x_0}$  is orthogonal to  $A(L_{x_0})$ .*

*Proof.* Let  $h$  be a rational function on  $S$  with poles at  $p_1, p_2, \dots, p_n$  in  $S \sim R_{x_0} = S \sim K \cap \{p: \operatorname{Re} f(p) < x_0\}$ . Let  $p_1, \dots, p_k$  be those poles not in  $S \sim K$ . Each  $p_i, i = 1, \dots, k$  is in some component  $Q_i$  of  $\{p: \operatorname{Re} f(p) < x_0\}$ . By the proof of Lemma 6, such a component cannot be contained in  $K$ . Thus we may choose  $q_i, i = 1, \dots, k, q_i \in Q_i \sim K$  and let  $J_i$  be a curve in  $Q_i$  joining  $p_i$  and  $q_i$ . Let

$$B = S \sim \bigcup_{i=1}^k J_i \sim \bigcup_{i=1}^k \{p_i\} \quad \text{and} \quad \tilde{B} = S \sim \bigcup_{i=1}^k \{q_i\} \sim \bigcup_{i=k+1}^n \{p_i\}.$$

Then by Theorem 6 in [4],  $h$ , which is analytic on  $B$ , can be uniformly approximated on  $R_{x_0}$ , a compact subset of  $B$ , by functions  $f_j$  analytic on  $\tilde{B}$ . Now letting  $B_1 = \tilde{B}$  and  $\tilde{B}_1 = S$  we apply Theorem 13 in [1, p. 456] and approximate  $f_j$  on  $R_{x_0}$  by meromorphic functions  $g_j$  with poles on the boundary of  $B_1$ , i.e., at the points  $q_1, \dots, q_k, p_{k+1}, \dots, p_n$ . But these are all in  $S \sim K$ . Thus  $g_j \in A(K)$ . By Lemma 5,  $\int g_j d(\nu | R_{x_0} - \beta_{x_0}) = 0$ . Thus  $\int h d(\nu | R_{x_0} - \beta_{x_0}) = 0$  and  $\nu | R_{x_0} + \beta_{x_0}$  is orthogonal to  $A(R_{x_0})$ . The same argument shows  $\nu | L_{x_0} + \beta_{x_0}$  is orthogonal to  $A(L_{x_0})$ .

Given any finite collection of functions  $\{g_k\}_{k=1}^l$  on  $S$ , and a real number  $x_0$ , we define an equivalence relation on the points of  $S$  as follows. The points  $p$  and  $q$  are equivalent, if, for all  $k$ ,  $\operatorname{Re} f_k(p) \leq x_0$  if and only if  $\operatorname{Re} f_k(q) \leq x_0$ .

LEMMA 8. *Let  $K$  be a compact subset of  $S$  and  $\{U_i\}$  an open*

covering of  $K$ . Then there exists a finite collection of nonconstant functions, each analytic on  $S$ , such that given any  $x_0$ ,  $1/4 < x_0 < 3/4$ , each equivalence class of the relation defined with these functions lies in a single member of the covering.

*Proof.* Fix a metric on  $S$ . By the Lebesgue covering lemma, there exists  $\rho > 0$  such that any set of diameter less than or equal to  $\rho$ , containing a point of  $K$ , lies in a single member of the covering  $\{U_i\}$ . Cover  $K$  by a finite number of sets of diameter less than  $\rho/3$  which are homeomorphic to a closed disc. Call these sets  $\{D_i\}_{i=1}^m$ . For  $i, j$  such that  $D_i \cap D_j$  is empty, let  $f_{ij}$  be a function analytic on  $S$  such that  $Re f_{ij} < 1/4$  on  $D_i$  and  $Re f_{ij} > 3/4$  on  $D_j$ . This is possible since by the Behnke-Stein extension of Runge's theorem [1, p. 445 and p. 456] we can approximate a function which is identically zero on a neighborhood of  $D_1$  and identically one on a neighborhood of  $D_j$  by functions analytic on  $S$ .

Now if  $A$  is an equivalence class of the equivalence relation defined by these functions, we will show  $diam A \leq \rho$ .

Let  $p_0 \in A$ . Then for some  $i_0, p_0 \in D_{i_0}$ . Let  $i_0, i_1, \dots, i_k$  be all  $i$  such that  $D_{i_0} \cap D_i$  is not empty. Let  $p \in K \cap \{p : Re f_{i_0 j}(p) \leq 3/4, \text{ all } j \neq i_0, i_1, \dots, i_k\}$ . Suppose  $p \notin \bigcup_{i=i_0}^{i_k} D_i$ . Then since  $p \in K \subset \bigcup_{i=1}^m D_i$ ,  $p \in D_{j_0}$ , some  $j_0 \neq i_0, \dots, i_k$ . Thus  $f_{i_0 j_0}(p) > 3/4$  which contradicts the choice of  $p$ . We have shown

$$K \cap \left\{ p : Re f_{i_0 j}(p) \leq \frac{3}{4} \text{ all } j \neq i_0, \dots, i_k \right\} \subset \bigcup_{i=i_0}^{i_k} D_i .$$

Now since  $p_0 \in D_{i_0}$ ,  $Re f_{i_0 j}(p) < 1/4$ , all  $j \neq i_0, \dots, i_k$ , but  $p_0 \in A$ , so for all  $p \in A$ , we have  $Re f_{i_0 j}(p) < 1/4$  for  $i \neq i_0, \dots, i_k$ . Therefore

$$A \subset K \cap \left\{ p : Re f_{i_0 j}(p) \leq \frac{3}{4} \text{ all } j \neq i_0, \dots, i_k \right\} \subset \bigcup_{i=i_0}^{i_k} D_i .$$

Each  $D_{i_0}, \dots, D_{i_k}$  intersects  $D_{i_0}$  and  $diam D_i < \rho/3$ . Therefore  $diam \bigcup_{i=i_0}^{i_k} D_i < \rho$  and the proof is complete.

**THEOREM 4.** *If  $K$  is a compact subset of  $S$  and  $\{U_i\}_{i=1}^n$  is an open covering of  $K$ , and if  $\mu$  is a measure on  $K$  which is orthogonal to  $A(K)$ , then there exist measures  $\nu_i$  with support contained in a compact set  $T_i \subset K \cap U_i$  such that  $\nu_i$  is orthogonal to  $A(T_i)$  and  $\mu = \nu_1 + \nu_2 + \dots + \nu_n$ .*

*Proof.* Let  $f_k$  be the functions of Lemma 8,  $k = 1, \dots, l$ . The proof will be by induction on  $l$ . If  $l = 0$ , let  $T = K \subset U_{i_0}$  and  $\nu_{i_0} = \mu$  is orthogonal to  $A(K) = A(T)$ .

Suppose the theorem is true for  $l - 1$ . Let  $1/4 < x_0 < 3/4$  and  $R_{x_0} = K \cap \{p : \operatorname{Re} f_l(p) \geq x_0\}$  and  $L_{x_0} = K \cap \{p : \operatorname{Re} f_l(p) \leq x_0\}$ .  $R_{x_0}$  and  $L_{x_0}$  are both compact, and  $\{U_i\}_{i=1}^n$  is a covering for each. An equivalence class of points of  $R_{x_0}$  of the relation defined by  $f_1, \dots, f_{l-1}$  lies in a single member of  $\{U_i\}_{i=1}^n$ . Similarly for an equivalence class of points of  $L_{x_0}$ . Thus we may apply the induction hypothesis to the measures  $\mu_1 = \mu | R_{x_0} - \beta_{x_0}$  which is orthogonal to  $A(R_{x_0})$  by Lemma 7, and  $\mu_2 = \mu | L_{x_0} + \beta_{x_0}$  which is orthogonal to  $A(L_{x_0})$  by Lemma 7. Thus we have measures  $\nu_{ji}$  with support contained in a compact set  $T_{ji} \subset U_i \cap K$  which is orthogonal to  $A(T_{ji})$   $j = 1, 2, i = 1, \dots, n$  and

$$\mu_1 = \nu_{11} + \nu_{12} + \dots + \nu_{1n}, \quad \mu_2 = \nu_{21} + \nu_{22} + \dots + \nu_{2n}.$$

Thus  $\mu = \mu_1 + \mu_2 = (\nu_{11} + \nu_{21}) + (\nu_{12} + \nu_{22}) + \dots + (\nu_{1n} + \nu_{2n})$  and  $\nu_{1i} + \nu_{2i}$  has support contained in  $T_{1i} \cup T_{2i} \subset U_i \cap K$ . If  $f \in A_{1i}(T_{1i} \cup T_{2i})$ , then  $f | T_{1i} \in A(T_{1i})$  and  $f | T_{2i} \in A(T_{2i})$  and

$$\int f d(\nu_{1i} + \nu_{2i}) = \int f d\nu_{1i} + \int f d\nu_{2i} = 0.$$

Thus  $\nu_{1i} + \nu_{2i}$  is orthogonal to  $A(T_{1i} \cup T_{2i})$  and the theorem is proved.

**THEOREM 5.** *If  $K$  is a compact subset of  $S$  and if for every  $p \in K$ , there is a closed neighborhood  $W$  of  $p$  such that  $f | W \in A(K \cap W)$ , then  $f \in A(K)$ .*

*Proof.* Suppose  $f \notin A(K)$ . Then there exists a measure  $\mu$  on  $K$  such that  $\mu$  is orthogonal to  $A(K)$  and  $\int f d\mu \neq 0$ . Let  $V$  be the interior of  $W$ . Then  $\{V\}$  is an open covering of  $K$ . Let  $\{V_i\}_{i=1}^n$  be a finite subcovering. Apply the last theorem with this covering to get measures  $\nu_i$  with support contained in a compact set  $T_i \subset V_i \cap K \subset W_i \cap K$  and  $\nu_i$  is orthogonal to  $A(T_i)$  and  $\mu = \nu_1 + \nu_2 + \dots + \nu_n$ .  $f | W \in A(K \cap W_i)$  implies  $f | T_i \in A(T_i)$ . Thus  $\int f d\nu_i = 0$  and  $\int f d\mu = 0$  which is contradiction.

**COROLLARY 1.** *If  $K$  is a compact subset of  $S$  and for every  $p \in K$  there exists an analytic coordinate function  $h$  with  $h(p) = 0$  and the range of  $h$  is  $\{z : |z| < 1\}$  and an  $r, 0 < r < 1$ , such that  $A(h(K) \cap \{z : |z| \leq r\}) = C(h(K) \cap \{z : |z| \leq r\})$ , then  $A(K) = C(K)$ .*

*Proof.* Let  $f \in C(K)$ . For every  $p$ ,

$$\begin{aligned} f \circ h^{-1} | h\{z : |z| \leq r\} &\in C(h(K) \cap \{z : |z| \leq r\}) \\ &= A(h(K) \cap \{z : |z| \leq r\}). \end{aligned}$$

Thus  $f|_{K \cap h^{-1}\{z: |z| \leq r\}} \in A(K \cap h^{-1}\{z: |z| \leq r\})$ . Applying the theorem,  $f \in A(K)$ .

Thus a local condition on a compact set in the plane which implies that any continuous function can be uniformly approximated by rational functions, such as Theorems 2.4 and 3.4 in [6], can be applied in coordinate neighborhoods of every point of  $K$  to show  $A(K) = C(K)$ . As a special case, using Theorem 2.4 of [6] we have the next corollary which we will need to prove uniqueness of boundary measures.

**COROLLARY 2.** *If  $K$  is a nowhere dense compact  $n$ -balanced subset of  $S$ , then  $A(K) = C(K)$ .*

We also obtain a generalization from the plane to Riemann surface of the approximation theorem of Bishop [4].

**COROLLARY 3.** *If  $K$  is a compact nowhere dense subset of an open Riemann surface and  $M$  is the minimal boundary of  $A(K)$ , then  $M = K$  implies  $A(K) = C(K)$ .*

*Proof.* Let  $h$  be an analytic coordinate function at  $p \in K$  such that  $h(p) = 0$  and the range of  $h$  is  $\{z: |z| < 1\}$ . Let  $r$  be  $0 < r < 1$ . Let  $M'$  be the minimal boundary of  $A(h(K) \cap \{z: |z| \leq r\})$ . Let  $z \in h(K)$  and  $|z| \leq r$ , then  $h^{-1}(z) \in K = M$ . There exists  $f \in A(K)$  such that  $f(h^{-1}(z)) = 1$  and  $|f(q)| < 1$  if  $q \in K$  and  $q \neq h^{-1}(z)$ .

$$f \circ h^{-1} \in A(h(K) \cap \{z: |z| \leq r\}), \quad f \circ h^{-1}(z) = 1, \quad |f \circ h^{-1}(z)| < 1$$

if  $\zeta \in h(K)$  and  $|\zeta| \leq r$ ,  $\zeta \neq z$ . Thus  $z \in M'$ . Since

$$M' = h(K) \cap \{z: |z| \leq r\},$$

by Theorem 4 in [4], we have

$$A(h(K) \cap \{z: |z| \leq r\}) = C(h(K) \cap \{z: |z| \leq r\}).$$

Now the theorem applies and we have  $A(K) = C(K)$ .

**LEMMA 9.** *Suppose  $K$  is an  $n$ -balanced compact subset of  $S$ . If  $\mu$  is a measure on the boundary  $B$  of  $K$  which is orthogonal to all rational functions on  $S$  with poles in the interior of  $K$  or in  $S \sim K$ , then  $\mu = 0$ .*

*Proof.* The hypothesis implies  $\mu$  is orthogonal to  $A(B)$ . By Lemma 1,  $B$  is an  $n$ -balanced nowhere dense compact subset of  $S$ . Thus by Corollary 2,  $A(B) = C(B)$  and  $\mu = 0$ .

**THEOREM 6.** *If  $K$  is a compact  $n$ -balanced subset of  $S$  and  $\omega$  is an analytic differential on the interior of  $K$ , then the boundary measure  $\mu$  of  $\omega$  which exists by Theorem 1 is unique, and if  $h$  is an analytic coordinate function on an open set  $V \subset S$  and  $\omega = f(z)dz$  on  $h(V)$ , then*

$$f(z) = (2\pi i)^{-1} \int -A(z, q) d\mu(q)$$

where  $\alpha(p) = A(z, p)dz$  on  $h(V)$ .

*Proof.* Suppose  $\mu$  and  $\nu$  are both boundary measures of  $\omega$ . Let  $g$  be a rational function on  $S$  with poles in the interior or the complement of  $K$ . Then

$$\int gd(\mu - \nu) = \int gd\mu - \int gd\nu = \lim_n \int_{\delta_n} g\omega - \lim_n \int_{\gamma_n} g\omega .$$

If  $n$  is large enough so both  $\delta_n$  and  $\gamma_n$  surround all the poles of  $g$  which lie in the interior of  $K$ , then

$$\int_{\delta_n} g\omega = \sum_{p \in \text{int}K} \text{Res}_p(g\omega) = \int_{\gamma_n} g\omega .$$

Therefore  $\int gd(\mu - \nu) = 0$  and by Lemma 9,  $\mu = \nu$ .

$A(z, q)$  is meromorphic in  $q$  with a simple pole of residue  $-1$  at  $h^{-1}(z)$ . Thus

$$\begin{aligned} (2\pi i)^{-1} \int -A(z, q) d\mu(q) &= (2\pi i)^{-1} \lim_n \int_{\gamma_n} -A(z, q)\omega = \\ &= -\text{Res}_{h^{-1}(z)}(A(z, q)\omega) = f(z) . \end{aligned}$$

**THEOREM 7.** *Let  $K$  be a compact  $n$ -balanced subset of  $S$  with interior  $U$  and let  $\mu \in M(K)$ . Then there exists a differential  $\omega \in H(U)$  such that  $\mu$  is the boundary measure of  $\omega$ .*

*Proof.* Let  $f_1, \dots, f_l$  be the finite set of functions analytic on  $S$  and satisfying the conditions of Lemma 8 using coordinate neighborhoods for the covering. The proof will be by induction on  $l$ . If  $l = 0$ ,  $K$  lies in a single coordinate neighborhood and we may consider  $K$  as a subset of the plane. In this case we have the result in Theorem 3.

Suppose the theorem is true for  $l - 1$ . Let  $1/4 < x_0 < 3/4$  satisfy the conditions of Lemma 5 for  $f_l, \mu, K$ . Let  $L_{x_0}, R_{x_0}$ , and  $\beta_{x_0}$  be as in Lemma 5. By Lemma 6,  $R_{x_0}$  and  $L_{x_0}$  are compact  $m$ -balanced sets for some  $m$  and by Lemma 7,  $\mu|_{R_{x_0}} - \beta_{x_0} \in M(R_{x_0})$  and  $\mu|_{L_{x_0}} + \beta_{x_0} \in M(L_{x_0})$ . Since  $f_1, \dots, f_{l-1}$  partition  $R_{x_0}$  and  $L_{x_0}$  in the sense of Lemma 8, the

induction hypothesis applies. Thus we have analytic differentials  $\omega_1$  and  $\omega_2$  on the interiors of  $R_{x_0}$  and  $L_{x_0}$  for which  $\mu | R_{x_0} - \beta_{x_0}$  and  $\mu | L_{x_0} + \beta_{x_0}$  are the boundary measures respectively.

If  $h_1$  is an analytic coordinate function on  $V_1 \subset \text{int } R_{x_0}$  and  $\omega_1 = f_1(z)dz$  on  $h_1(V_1)$  and  $\alpha(q) = A_1(z, q)dz$  on  $h_1(V_1)$  then

$$\begin{aligned} f_1(z) &= (2\pi i)^{-1} \int - A_1(z, q)d(\mu | R_{x_0} - \beta_{x_0})(q) \\ &= (2\pi i)^{-1} \int - A_1(z, q)d(\mu | R_{x_0} - \beta_{x_0})(q) \\ &\quad + (2\pi i)^{-1} \int - A_1(z, q)d(\mu | L_{x_0} + \beta_{x_0})(q) \\ &= (2\pi i)^{-1} \int - A_1(z, q)d\mu(q) . \end{aligned}$$

Similarly, if  $h_2$  is an analytic coordinate function on  $V_2 \subset \text{int } L_{x_0}$  and  $\omega_2 = f_2(z)dz$  on  $h_2(V_2)$  and  $\alpha(q) = A_2(z, q)dz$  on  $h_2(V_2)$  then

$$f_2(z) = (2\pi i)^{-1} \int - A_2(z, q)d\mu(q) .$$

Since we have this for almost all  $x_0$  between  $1/4$  and  $3/4$  we can define, for any coordinate function  $h$  on  $V \subset U$ , a differential  $\omega = f(z)dz$  on  $h(V)$  with

$$f(z) = (2\pi i)^{-1} \int - A(z, q)d\mu(q)$$

where  $\alpha(q) = A(z, q)dz$  on  $h(V)$ ,  $\omega = \omega_1$  on  $\text{int } R_{x_0}$ , and  $\omega = \omega_2$  on  $\text{int } L_{x_0}$ .

Let  $1/4 < x_1 < x_2 < 3/4$  and both  $x_1$  and  $x_2$  satisfy the conditions of Lemma 5. Let the delimiting sequence of Definition 6 for the boundary measures  $\mu | L_{x_2} + \beta_{x_2}$  and  $\mu | R_{x_1} - \beta_{x_1}$  be  $\{\delta_i\}$  and  $\{\gamma_i\}$  respectively. Let  $\Delta_i$  and  $\Gamma_i$  be the open sets of which  $\delta_i$  and  $\gamma_i$  are the boundaries. Let  $p_j, V_j$  be the finite collection of points and coordinate neighborhoods obtained in Lemma 2 with  $h_j$  the analytic coordinate function on  $V_j$ . Let  $U_j$  be a closed neighborhood of  $p_j$  so that  $U_j \subset V_j \cap U$ . Let  $k_j$  be the maximum of  $|f(h_j(p))|$  for  $p \in U$  where  $\omega = f(z)dz$  on  $V_j$ . Let  $\epsilon_j = (k_j^{-1}2^{-j})$ . Using these  $U_j$  and  $\epsilon_j$  we apply Lemma 2 to get  $\varphi_i$  a finite union of disjoint piecewise analytic simple closed curves forming the boundary of  $\Phi_i$  and  $|\varphi_i| \cup \Phi_i \subset U$ . Furthermore, since  $\{\delta_i\}$  and  $\{\gamma_i\}$  delimit the interiors of  $L_{x_2}$  and  $R_{x_1}$ , respectively, and  $\Gamma_i \cup \Delta_i \subset \Phi_i$ ,  $\{\varphi_i\}$  delimits  $U$ .

Finally we see that

$$\|\omega\|_{\varphi_i} \leq \|\omega\|_{\gamma_i} + \|\omega\|_{\delta_i} + \sum_j k_j \epsilon_j \leq \|\omega\|_{\gamma_i} + \|\omega\|_{\delta_i} + 1 .$$



Therefore  $\omega \in H(U)$  and by Theorem 1,  $\omega$  has a boundary measure  $\nu$  on the boundary of clsr  $U$ .

Now let  $g$  be a rational function on  $S$  with poles in  $U$  or  $S \sim K$ . Choose  $x$ ,  $1/4 < x_0 < 3/4$ , as in Lemma 5 and so that no pole of  $g$  lies on  $\{p: \operatorname{Re} f_i(p) = x_0\}$ . Let  $\{\sigma_i\}$  and  $\{\tau_i\}$  be the delimiting sequence of Definition 6 for the boundary measures  $\mu|L_{x_0} + \beta_{x_0}$  and  $\mu|R_{x_0} - \beta_{x_0}$  respectively. Then

$$\begin{aligned} \int g d(\mu - \nu) &= \int g d(\mu|R_{x_0} - \beta_{x_0}) + \int g d(\mu|L_{x_0} + \beta_{x_0}) - \int g d\nu \\ &= \lim_i \int_{\tau_i} g \omega + \lim_i \int_{\sigma_i} g \omega - \lim_i \int_{\varphi_{n_i}} g \omega. \end{aligned}$$

Letting  $i$  be large enough so that all the poles of  $g$  in  $U$  are surrounded by  $\varphi_{n_i}$  and by either  $\tau_i$  or  $\sigma_i$  and using the residue theorem we have

$$\int g d(\mu - \nu) = \int_{\tau_i} g \omega + \int_{\sigma_i} g \omega - \int_{\varphi_{n_i}} g \omega = 0.$$

Thus by Lemma 9,  $\mu - \nu = 0$  and  $\mu$  is the boundary measure of  $\omega$ .

**COROLLARY 4.** *If  $K$  is a compact  $n$ -balanced subset of  $S$  with interior  $U$ , then  $A(K)$  consists of all functions in  $C(K)$  which are analytic on  $U$ .*

*Proof.* Clearly every function in  $A(K)$  is analytic on  $U$ . Suppose  $A(K)$  does not contain all such functions in  $C(K)$ . Then there exists a continuous linear functional  $L$  orthogonal to  $A(K)$  with  $L(f) \neq 0$  for some  $f \in C(K)$ ,  $f$  analytic on  $U$ . The boundary of  $K$  is the Silov boundary of the algebra of functions in  $C(K)$  analytic on  $U$ , so there exists a measure  $\mu$  on the boundary of  $K$  so that  $\int g d\mu = L(g)$ , all  $g \in C(K)$ , analytic on  $U$ . Thus  $\mu \in M(K)$  and there exists  $\omega \in H(U)$ , so that

$$0 \neq L(f) = \int f d\mu = \lim_i \int_{\gamma_j} f \omega = 0$$

since  $f$  is analytic on  $U$ .

#### BIBLIOGRAPHY

1. H. Behnke and K. Stein, *Entwicklungen analytischer Funktionen auf Riemannschen Flächen*, Math. Annalen, **120** (1949) 430-461.
2. E. Bishop, *The structure of certain measures*, Duke Math. J. **25** (1958), 283-290.
3. ———, *Subalgebras of functions on a Riemann surface*, Pacific J. Math. **8** (1958), 29-50.

4. E. Bishop, *A minimal boundary for function algebras*, Pacific J. Math., **9** (1959), 629-642.
5. ———, *Boundary measures of analytic differentials*, Duke Math. J. **27** (1960), 331-340.
6. S. N. Mergelyan, *Uniform approximations to functions of a complex variable*, Amer. Math. Soc. Translation No. 101, 1954.
7. H. L. Royden, *The boundary values of analytic and harmonic functions*, Math. Zeitschrift, **78** (1962), 1-24.
8. A. Zygmund, *Trigonometrical Series*, New York, 1955.



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California

R. M. BLUMENTHAL

University of Washington  
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California  
Los Angeles, California 90007

\*RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CALIFORNIA RESEARCH CORPORATION  
SPACE TECHNOLOGY LABORATORIES  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. No separate author's resumé is required. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published quarterly, in March, June, September, and December. Effective with Volume 13 the price per volume (4 numbers) is \$18.00; single issues, \$5.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$8.00 per volume; single issues \$2.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

\* Basil Gordon, Acting Managing Editor until February 1, 1966.

# Pacific Journal of Mathematics

Vol. 15, No. 4

December, 1965

Robert James Blattner, <i>Group extension representations and the structure space</i> .....	1101
Glen Eugene Bredon, <i>On the continuous image of a singular chain complex</i> .....	1115
David Hilding Carlson, <i>On real eigenvalues of complex matrices</i> .....	1119
Hsin Chu, <i>Fixed points in a transformation group</i> .....	1131
Howard Benton Curtis, Jr., <i>The uniformizing function for certain simply connected Riemann surfaces</i> .....	1137
George Wesley Day, <i>Free complete extensions of Boolean algebras</i> .....	1145
Edward George Effros, <i>The Borel space of von Neumann algebras on a separable Hilbert space</i> .....	1153
Michel Mendès France, <i>A set of nonnormal numbers</i> .....	1165
Jack L. Goldberg, <i>Polynomials orthogonal over a denumerable set</i> .....	1171
Frederick Paul Greenleaf, <i>Norm decreasing homomorphisms of group algebras</i> .....	1187
Fletcher Gross, <i>The 2-length of a finite solvable group</i> .....	1221
Kenneth Myron Hoffman and Arlan Bruce Ramsay, <i>Algebras of bounded sequences</i> .....	1239
James Patrick Jans, <i>Some aspects of torsion</i> .....	1249
Laura Ketchum Kodama, <i>Boundary measures of analytic differentials and uniform approximation on a Riemann surface</i> .....	1261
Alan G. Konheim and Benjamin Weiss, <i>Functions which operate on characteristic functions</i> .....	1279
Ronald John Larsen, <i>Almost invariant measures</i> .....	1295
You-Feng Lin, <i>Generalized character semigroups: The Schwarz decomposition</i> .....	1307
Justin Thomas Lloyd, <i>Representations of lattice-ordered groups having a basis</i> .....	1313
Thomas Graham McLaughlin, <i>On relative coimmunity</i> .....	1319
Mitsuru Nakai, <i><math>\Phi</math>-bounded harmonic functions and classification of Riemann surfaces</i> .....	1329
L. G. Novoa, <i>On <math>n</math>-ordered sets and order completeness</i> .....	1337
Fredos Papangelou, <i>Some considerations on convergence in abelian lattice-groups</i> .....	1347
Frank Albert Raymond, <i>Some remarks on the coefficients used in the theory of homology manifolds</i> .....	1365
John R. Ringrose, <i>On sub-algebras of a <math>C^*</math>-algebra</i> .....	1377
Jack Max Robertson, <i>Some topological properties of certain spaces of differentiable homeomorphisms of disks and spheres</i> .....	1383
Zalman Rubinstein, <i>Some results in the location of zeros of polynomials</i> .....	1391
Arthur Argyle Sagle, <i>On simple algebras obtained from homogeneous general Lie triple systems</i> .....	1397
Hans Samelson, <i>On small maps of manifolds</i> .....	1401
Annette Sinclair, <i><math> \varepsilon(z) </math>-closeness of approximation</i> .....	1405
Edsel Ford Stiel, <i>Isometric immersions of manifolds of nonnegative constant sectional curvature</i> .....	1415
Earl J. Taft, <i>Invariant splitting in Jordan and alternative algebras</i> .....	1421
L. E. Ward, <i>On a conjecture of R. J. Koch</i> .....	1429
Neil Marchand Wigley, <i>Development of the mapping function at a corner</i> .....	1435
Horace C. Wisner, <i>Embedding a circle of trees in the plane</i> .....	1463
Adil Mohamed Yaqub, <i>Ring-logics and residue class rings</i> .....	1465
John W. Lamperti and Patrick Colonel Suppes, <i>Correction to: Chains of infinite order and their application to learning theory</i> .....	1471
Charles Vernon Coffman, <i>Correction to: Non-linear differential equations on cones in Banach spaces</i> .....	1472
P. H. Doyle, III, <i>Correction to: A sufficient condition that an arc in <math>S^n</math> be cellular</i> .....	1474
P. P. Saworotnow, <i>Correction to: On continuity of multiplication in a complemented algebra</i> .....	1474
Basil Gordon, <i>Correction to: A generalization of the coset decomposition of a finite group</i> .....	1474