

Pacific Journal of Mathematics

**HARNACK'S INEQUALITIES ON THE CLASSICAL CARTAN
DOMAINS**

SHIH-HSIUNG TUNG

HARNACK'S INEQUALITIES ON THE CLASSICAL CARTAN DOMAINS

SHIH-HSIUNG TUNG

Recently an extensive work by L. K. Hua on harmonic analysis in Cartan domains, which are called the classical domains, has been translated into English. Here we give Harnack's inequalities for the four main types of Cartan domains treated by Hua.

Harnack's inequality on a type of Cartan domain was obtained [6] for the case of square matrix spaces. Some of these inequalities are application and extension of the results of [6]. I am grateful to Professor J. Mitchell for her encouragement and comments on writing this paper.

Let z be a matrix of complex entries, $z^* = \bar{z}'$ the complex conjugate of the transposed matrix z' and I the identity matrix. Also $H > 0$ means that a hermitian matrix H is positive definite. The first three types of Cartan domains are defined by $D_k = \{z : I - zz^* > 0\}$, $k = 1, 2, 3$, where for $D_1 \equiv D_1(m, n)$, z is an (m, n) matrix (Since the conditions $I - zz^* > 0$ and $I - z^*z > 0$ are equivalent we assume for definiteness that $m \leq n$), for $D_2 \equiv D_2(n)$, z is a symmetric matrix of order n and for $D_3 \equiv D_3(n)$, z is a skew-symmetric matrix of order n . The fourth type, $D_4 \equiv D_4(1, n)$, is the set of all $(1, n)$ matrices, or n -dimensional vectors ($n > 2$), of complex numbers satisfying the conditions

$$(1) \quad 1 + |zz'|^2 - 2zz^* > 0, \quad |zz'| < 1.$$

It is known that each of the domain D_k possesses a distinguished boundary [1] or characteristic manifold [2, p. 6] $C_k : C_1 \equiv C_1(m, n)$ consists of the (m, n) matrices u satisfying the condition $uu^* = I$. $C_2 \equiv C_2(n)$ consists of all symmetric unitary matrices of order n . $C_3 \equiv C_3(n)$ [2, p. 71] consists of all matrices u of the form $u = w's_1w$, where w is an n -rowed unitary matrix and

$$(2) \quad s_1 = \begin{cases} \left(\begin{matrix} 0 & 1 \\ -1 & 0 \end{matrix} \right) \dot{+} \cdots \dot{+} \left(\begin{matrix} 0 & 1 \\ -1 & 0 \end{matrix} \right) & \text{for even } n \\ \left(\begin{matrix} 0 & 1 \\ -1 & 0 \end{matrix} \right) \dot{+} \cdots \dot{+} \left(\begin{matrix} 0 & 1 \\ -1 & 0 \end{matrix} \right) \dot{+} 0 & \text{for odd } n. \end{cases}$$

$C_4 \equiv C_4(1, n)$ consists of all $(1, n)$ matrices u of the form

$$(3) \quad u = e^{i\theta}x, \quad xx' = 1, \quad 0 \leq \theta \leq \pi$$

where x is a real vector.

We denote the Poisson kernel on D_k by $P_k(z, u)$, its explicit forms [3, 4] being given in following sections. The following Dirichlet problem is solved on D_k [2, 3]: Given a real-valued continuous function $f(u)$ on C_k , the Poisson integral

$$(4) \quad \phi(z) = \int_{\sigma_k} f(u)P_k(z, u)\dot{u}$$

where \dot{u} is Euclidean volume element on C_k , gives the unique function which is harmonic, in the sense given in [3], on the closure of D_k and takes the given boundary values $f(u)$ on C_k . We obtain Harnack's inequality on each D_k as a consequence of evaluating upper and lower bounds for $P_k(z, u)$.

2. Harnack's inequalities on D_1 and D_2 . The Poisson kernel on D_1 [3, 4] is

$$(5) \quad P_1(z, u) = \frac{1}{V_1} \frac{[\det(I - zz^*)]^n}{|\det(I - zu^*)|^{2n}}$$

where $z \in D_1$, $u \in C_1$ and V_1 is the Euclidean volume of C_1 . It is known [4, p. 411] that $P_1 > 0$ and $\int_{\sigma_1} P_1 \dot{u} = 1$. In [6], we obtained bounds of the Poisson kernel (5) for the case $m = n$:

$$(6) \quad \frac{1}{V_1} \prod_{k=1}^n \left(\frac{1 - r_k}{1 + r_k} \right)^n \leq P_1(z, u) \leq \frac{1}{V_1} \prod_{k=1}^n \left(\frac{1 + r_k}{1 - r_k} \right)^n$$

where $z = u_0 R v_0 \in D_1(n, n)$, $u \in C_1(n, n)$, u_0 and v_0 are unitary matrices and $R = (\delta_{jk} r_k)$ is a diagonal matrix with $0 \leq r_k < 1$ for $k = 1, 2, \dots, n$. In order to obtain (6), we proved the inequality

$$(7) \quad \prod_{k=1}^n (1 - r_k)^2 \leq |\det(I - zu^*)|^2 = |\det((v - R)(v^* - R))| \leq \prod_{k=1}^n (1 + r_k)^2$$

where $u, v \in C_1(n, n)$, is any unitary matrix of order n , $z \in D_1(n, n)$ and $v = u_0^* u v_0^*$.

For $z \in D_1(m, n)$ there are unitary matrices u_0 of order m and v_0 of order n such that $z = u_0(R, 0)v_0$, where R is a diagonal submatrix $(\delta_{jk} r_k)$ of order m and 0 is the $(m, n-m)$ zero submatrix [3, p. 1049]. Hence

$$(8) \quad \det(I^{(m)} - zz^*) = \det(I^{(m)} - RR^*) = \prod_{k=1}^m (1 - r_k^2).$$

For the denominator of (5) we have

$$(9) \quad \det(I - zu^*) = \det \left(I - \begin{pmatrix} z \\ 0 \end{pmatrix} \begin{pmatrix} u \\ u_1 \end{pmatrix}^* \right)$$

where 0 is the $(n-m, n)$ zero submatrix and u_1 is chosen so that $U \equiv \begin{pmatrix} u \\ u_1 \end{pmatrix}$ becomes a unitary matrix of order n [5, p.190]. If we also denote $Z \equiv \begin{pmatrix} z \\ 0 \end{pmatrix}$, then $Z \in D_1(n, n)$ and $U \in C_1(n, n)$ for the case $m = n$. Therefore from (7)

$$(10) \quad \max_{U \in O_1} \{ |\det(I - ZU^*)|^2 \} \leq \prod_{k=1}^n (1 + r_k)^2 = \prod_{k=1}^m (1 + r_k)^2$$

and

$$(11) \quad \min_{U \in O_1} \{ |\det(I - ZU)|^2 \} \geq \prod_{k=1}^n (1 - r_k)^2 = \prod_{k=1}^m (1 - r_k)^2$$

since $r_{m+1} = \dots = r_n = 0$ in Z . Finally from (9), (10) and (11) we obtain

$$(12) \quad \prod_{k=1}^m (1 - r_k)^2 \leq |\det(I - zu^*)|^2 \leq \prod_{k=1}^m (1 + r_k)^2$$

where $z = u_0(R, 0)v_0 \in D_1(m, n)$ and $u \in C_1(m, n)$. This and (8) lead to

$$\frac{1}{V_1} \prod_{k=1}^m \left(\frac{1 - r_k}{1 + r_k} \right)^n \leq P_1(z, u) \leq \frac{1}{V_1} \prod_{k=1}^m \left(\frac{1 + r_k}{1 - r_k} \right)^n$$

for $z \in D_1$ and $u \in C_1$. Furthermore from (4) we obtain Harnack's inequality

$$\prod_{k=1}^m \left(\frac{1 - r_k}{1 + r_k} \right)^n \phi(0) \leq \phi(z) \leq \prod_{k=1}^m \left(\frac{1 + r_k}{1 - r_k} \right)^n \phi(0).$$

For $z \in D_2(n)$ it is known that there is a unitary matrix u_0 such that $u_0 z u_0' = (\delta_{jk} r_k)$ where r_1, \dots, r_n are the positive square roots of the characteristic roots of $z\bar{z}$. Since $z \in D_2(n)$ implies $z \in D_1(n, n)$ and the characteristic manifold $C_2(n)$ is a subset of $C_1(n, n)$ we know that (7) and (8) hold for $z \in D_2(n)$ and $u \in C_1(n, n)$, and it can be seen from the Poisson kernel

$$P_2(z, u) = \frac{1}{V_2} \frac{|\det(I - z\bar{z})|^{(n+1)/2}}{|\det(I - zu^*)|^{n+1}}$$

that Harnack's inequality on $D_2(n)$ is

$$\prod_{k=1}^n \left(\frac{1 - r_k}{1 + r_k} \right)^{(n+1)/2} \phi(0) \leq \phi(z) \leq \prod_{k=1}^n \left(\frac{1 + r_k}{1 - r_k} \right)^{(n+1)/2} \phi(0).$$

3. Harnack's inequality on D_3 . The Poisson kernel on D_3 is

$$P_3(z, u) = \frac{1}{V_3} \frac{|\det(I - zz^*)|^a}{|\det(I - zu^*)|^{2a}}$$

where $a = (n - 1)/2$ for even n and $a = n/2$ for odd n . For $z \in D_3(n)$ it is known [2, p. 67] that there is a unitary matrix u_0 such that

$$(13) \quad u_0 z u_0' = s = \begin{cases} \left(\begin{pmatrix} 0 & r_1 \\ -r_1 & 0 \end{pmatrix} \dagger \begin{pmatrix} 0 & r_2 \\ -r_2 & 0 \end{pmatrix} \dagger \cdots \dagger \begin{pmatrix} 0 & r_b \\ -r_b & 0 \end{pmatrix} \right) & \text{for even } n \\ \left(\begin{pmatrix} 0 & r_1 \\ -r_1 & 0 \end{pmatrix} \dagger \begin{pmatrix} 0 & r_2 \\ -r_2 & 0 \end{pmatrix} \dagger \cdots \dagger \begin{pmatrix} 0 & r_{[b]} \\ -r_{[b]} & 0 \end{pmatrix} \right) \dagger 0 & \text{for odd } n \end{cases}$$

where $b = n/2$ and r_1^2, r_2^2, \dots are characteristic roots of zz^* . Thus for even n

$$u_0 z z^* u_0^* = [r_1^2, r_1^2, \dots, r_b^2, r_b^2].$$

And for all n

$$\det(I - zz^*) = \det(I - u_0 z z^* u_0^*) = \prod_{k=1}^{[b]} (1 - r_k^2)^2.$$

We denote the right hand side of (2) by u_1 with the change that the last term 0 be replaced by 1 in the case of odd n . Hence u_1 is a unitary matrix of order n and from (13) su_1^* is a diagonal matrix $R \equiv [r_1, r_1, \dots, r_{[b]}, r_{[b]}]$ for even n and $R_0 \equiv [R, 0]$ for odd n . First we consider the case of even n . We notice from (2) that $u \in C_3$ is skew-symmetric unitary. Therefore for $z \in D_3$ and $u \in C_3$

$$|\det(I - zu^*)| = |\det(I - u_0^* s u_1^* u_1 \bar{u}_0 u^*)| = |\det(v - R)|$$

where $v = u_0 u u_0' u_1^*$ is a unitary matrix of order n . Hence from (7) we have

$$(14) \quad \prod_{k=1}^b (1 - r_k)^2 \leq |\det(I - zu^*)| \leq \prod_{k=1}^b (1 + r_k)^2.$$

For the case of odd n , we notice from (2) that $\det(u) = 0$. It is known [3, p. 1073] that any $v \in C_3(n + 1)$ can be written in the form

$$v = \begin{pmatrix} u & w'h' \\ -hw & 0 \end{pmatrix}, \quad h = (0, \dots, 0, e^{i\theta})$$

where $u = w's_1 w \in C_3(n)$. Hence for odd n

$$\det(I - zu^*) = \det \left(I^{(n+1)} - \begin{pmatrix} z & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} u & w'h' \\ -hw & 0 \end{pmatrix}^* \right)$$

and from (13)

$$\begin{pmatrix} u_0 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} z & 0 \\ 0 & 0 \end{pmatrix} \begin{pmatrix} u_0 & 0 \\ 0 & 1 \end{pmatrix}' = R \dagger \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}.$$

Since $n + 1$ is even we can apply (14) and obtain

$$\prod_{k=1}^{[b]} (1 - r_k)^2 \leq |\det (I - zu^*)| \leq \prod_{k=1}^{[b]} (1 + r_k)^2 .$$

As the result, this inequality is good for both odd and even n . Thus, as in § 2 we can obtain both bounds for the Poisson kernel $P_3(z, u)$ and finally Harnack's inequality on $D_3(n)$ as

$$\prod_{k=1}^{[b]} \left(\frac{1 - r_k}{1 + r_k} \right)^{2a} \phi(0) \leq \phi(z) \leq \prod_{k=1}^{[b]} \left(\frac{1 + r_k}{1 - r_k} \right)^{2a} \phi(0)$$

where $b = n/2$ and $a = (n - 1)/2$ for even n and $a = n/2$ for odd n .

4. Harnack's inequality on D_4 . The Poisson kernel on D_4 is

$$P_4(z, u) = \frac{1}{V_4} \frac{(1 + |zz'|^2 - 2zz^*)^{n/2}}{|1 + zz'uu' - 2zu^*|^n}$$

where $z \in D_4$ and $u \in C_4$. For every fixed $z \in D_4$ there is a real orthogonal matrix t such that [3, p. 1037]

$$(15) \quad z = (z_1, z_2, 0, \dots, 0)t .$$

Thus we have

$$1 + |zz'|^2 - 2zz^* = 1 + |z_1^2 + z_2^2|^2 - 2(|z_1|^2 + |z_2|^2) .$$

Here by denoting $z_1 - iz_2 \equiv w_1 = r_1 e^{i\theta_1}$ and $z_1 + iz_2 \equiv w_2 = r_2 e^{i\theta_2}$, from (1)

$$(16) \quad 0 < 1 + |zz'|^2 - 2zz^* = (1 - |w_1|^2)(1 - |w_2|^2) = (1 - r_1^2)(1 - r_2^2)$$

and

$$0 < 1 - |zz'| = 1 - |z_1^2 + z_2^2| = 1 - |w_1 w_2| = 1 - r_1 r_2 ,$$

hence

$$0 \leq r_k < 1 , \quad k = 1, 2 .$$

Next, from $z \in D_4$ in (15) and $u \in C_4$ in (3)

$$zz' \overline{uu'} = (z_1^2 + z_2^2) e^{-2i\theta} = w_1 w_2 e^{-2i\theta}$$

and

$$2zu^* = 2(z_1, z_2, 0, \dots, 0) t x' e^{-i\theta} .$$

By denoting $t = (t_{jk})$ we have $tx' = (a_1, a_2, \dots, a_n)'$ where $a_j = \sum_{k=1}^n t_{jk} w_k$ and since

$$\sum_{j=1}^n a_j^2 = (tx')'(tx') = xt'tx' = xx' = 1$$

we have $a_1^2 + a_2^2 \leq 1$. Now with $z_1 = (w_1 + w_2)/2$ and $z_2 = i(w_1 - w_2)/2$, we have $2zu^* = (aw_1 + \bar{a}w_2)e^{-i\theta}$ where $a = a_1 + ia_2$ and $|a|^2 \leq 1$. Thus

$$(17) \quad 1 + zz'\overline{uu'} - 2zu^* = 1 + w_1w_2e^{-2i\theta} - (aw_1 + \bar{a}w_2)e^{-i\theta}.$$

We wish to find upper and lower bounds for the absolute values of expression (17). We consider the image of the closed unit disk $|a| \leq 1$ under the mapping $f(a) = aw_1 + \bar{a}w_2$ for $a = re^{i\alpha}$. Here $f(a)$ can be written in the form

$$f(a) = r[r_1(e^{i\beta} + e^{-i\beta}) + (r_2 - r_1)e^{-i\beta}]e^{i(\theta_1 + \theta_2)/2}$$

where $\beta = \alpha + (\theta_1 - \theta_2)/2$. For the case $r_1 = r_2$, $f(a)$ maps the closed unit disk onto a line segment of length $4r_1$. When $r_1 \neq r_2$, the image is a simple closed connected region. Furthermore the image of the unit circle is the line segment when $r_1 = r_2$ and is the boundary of the region when $r_1 \neq r_2$. Hence from the fact that $1 + zz'\overline{uu'} - 2zu^* \neq 0$ [3, p. 1079] we know that the maximum and the minimum of the absolute values of (17) can be found in the case $|a| = 1$. Thus

$$1 + w_1w_2e^{-2i\theta} - (aw_1 + \bar{a}w_2)e^{-i\theta} = (1 - aw_1e^{-i\theta})(1 - \bar{a}w_2e^{-i\theta})$$

and therefore

$$(1 - r_1)(1 - r_2) \leq |1 + zz'\overline{uu'} - 2zu^*| \leq (1 + r_1)(1 + r_2).$$

This and (16) give us

$$\frac{1}{V_4} \prod_{k=1}^2 \left(\frac{1 - r_k}{1 + r_k} \right)^{n/2} \leq P_4(z, u) \leq \frac{1}{V_4} \prod_{k=1}^2 \left(\frac{1 + r_k}{1 - r_k} \right)^{n/2}$$

and the corresponding Harnack's inequality

$$\prod_{k=1}^2 \left(\frac{1 - r_k}{1 + r_k} \right)^{n/2} \phi(0) \leq \phi(z) \leq \prod_{k=1}^2 \left(\frac{1 + r_k}{1 - r_k} \right)^{n/2} \phi(0).$$

REFERENCES

1. S. Bergman, *The kernel function and conformal mapping*, Amer. Math. Soc. New York, 1950.
2. L. K. Hua, *Harmonic analysis of functions of several complex variables in the classical domains*, Amer. Math. Soc. Rhode Island, 1963.
3. L. K. Hua and K. H. Look, *Theory of harmonic functions in classical domains*, Sci. Sinica **8** (1959), 1031-1094.
4. J. Mitchell, *Potential theory in the geometry of matrices*, Trans. Amer. Math. Soc. **79** (1955), 401-422.
5. S. Perlis, *Theory of matrices*, Addison-Wesley, Cambridge, 1952.
6. S. H. Tung, *Harnack's inequality and theorems on matrix spaces*, Proc. Amer. Math. Soc. **15** (1964), 375-381.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

R. M. BLUMENTHAL

University of Washington
Seattle, Washington 98105

*J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$ 8.00; single issues, \$ 3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$ 4.00 per volume; single issues \$ 1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

* Paul A. White, Acting Editor until J. Dugundji returns.

Loren N. Argabright, <i>Invariant means on topological semigroups</i>	193
William Arveson, <i>A theorem on the action of abelian unitary groups</i>	205
John Spurgeon Bradley, <i>Adjoint quasi-differential operators of Euler type</i>	213
Don Deekard and Lincoln Kearney Durst, <i>Unique factorization in power series rings and semigroups</i>	239
Allen Devinatz, <i>The deficiency index of ordinary self-adjoint differential operators</i>	243
Robert E. Edwards, <i>Operators commuting with translations</i>	259
Avner Friedman, <i>Differentiability of solutions of ordinary differential equations in Hilbert space</i>	267
Boris Garfinkel and Gregory Thomas McAllister, Jr., <i>Singularities in a variational problem with an inequality</i>	273
Seymour Ginsburg and Edwin Spanier, <i>Semigroups, Presburger formulas, and languages</i>	285
Burrell Washington Helton, <i>Integral equations and product integrals</i>	297
Edgar J. Howard, <i>First and second category Abelian groups with the n-adic topology</i>	323
Arthur H. Kruse and Paul William Liebnitz, Jr., <i>An application of a family homotopy extension theorem to ANR spaces</i>	331
Albert Marden, I. Richards and Burton Rodin, <i>On the regions bounded by homotopic curves</i>	337
Willard Miller, Jr., <i>A branching law for the symplectic groups</i>	341
Marc Aristide Rieffel, <i>A characterization of the group algebras of the finite groups</i>	347
P. P. Saworotnow, <i>On two-sided H^*-algebras</i>	365
John Griggs Thompson, <i>Factorizations of p-solvable groups</i>	371
Shih-hsiung Tung, <i>Harnack's inequalities on the classical Cartan domains</i>	373
Adil Mohamed Yaqub, <i>Primal clusters</i>	379