

Pacific Journal of Mathematics

HERMITIAN AND ANTI-HERMITIAN PROPERTIES OF GREEN'S MATRICES

EARL A. CODDINGTON AND ANTON ZETTL

HERMITIAN AND ANTI-HERMITIAN PROPERTIES OF GREEN'S MATRICES

E. A. CODDINGTON AND A. ZETTL

In this paper hermitian and anti-hermitian properties of the components of Green's matrices of related boundary value problems are studied. Necessary and sufficient conditions, depending only on the matrices defining the boundary conditions, for the components of the Green's matrix of one problem to be hermitian or anti-hermitian with respect to certain components of the kernel matrix of a related problem, are found. It is also shown-for a wide class of problems-that some components of these Green's matrices cannot be hermitian (anti-hermitian).

The techniques used depend on

- (i) A construction of Green's matrices due to J. W. Neuberger [2, Theorem B],
- (ii) A new vector-matrix formulation of ordinary linear differential equations developed in [3], and
- (iii) Chapter 11 of [1].

Denote by $[a, b]$ a finite interval, each of A, B, P, Q a $k \times k$ complex matrix, and $F = (f_{ij})$ a $k \times k$ matrix of continuous complex valued functions on $[a, b]$ such that

$$(1) \quad f_{ij} = 0 \text{ if } i + j \text{ is even.}$$

Let

$$H = (\bar{f}_{k+1-j, k+1-i}), \quad T = ((-1)^i \delta_{i, k+1-j}).$$

Consider the following vector matrix equations with boundary conditions:

$$(2) \quad Y' = FY, AY(a) + BY(b) = 0,$$

$$(3) \quad X' = HX, PX(a) + QX(b) = 0.$$

Assume that each problem has only the trivial solution. Let M and N be the unique functions such that

$$M(t, u) = I + \int_u^t F(s)M(s, u)ds,$$

and

$$N(t, u) = I + \int_u^t H(s)N(s, u)ds,$$

for all t, u in $[a, b]$.

Define K and L by:

$$(4) \quad K(t, y) = \begin{cases} M(t, a)UM(a, y), & a \leq t < y \leq b, \\ M(t, a)UM(a, y) + M(t, y), & a \leq y < t \leq b, \\ M(t, a)UM(a, y) + \frac{1}{2}M(t, y), & a \leq y = t \leq b, \end{cases}$$

$$(5) \quad L(t, y) = \begin{cases} N(t, a)DN(a, y), & a \leq t < y \leq b, \\ N(t, a)DN(a, y) + N(t, y), & a \leq y < t \leq b, \\ N(t, a)DN(a, y) + \frac{1}{2}N(t, y), & a \leq y = t \leq b. \end{cases}$$

where $U = -[A + BM(b, a)]^{-1}BM(b, a)$ and $D = -[P + QN(b, a)]^{-1}QN(b, a)$. The existence of $[A + BM(b, a)]^{-1}$ and $[P + QN(b, a)]^{-1}$ follows from the assumption that problems (2) and (3), respectively, have only the trivial solution-(see [2]-Theorem A). The Green's functions K and L are modified versions of kernels as constructed by Neuburger in [2]. This modification is obtained with the help of the following property of M (or N):

$$(6) \quad M(t, u)M(u, v) = M(t, v), \quad t, u, v \in [a, b].$$

This property is an immediate consequence of the fact that for any fundamental matrix Φ of $Y' = FY$, $M(t, u) = \Phi(t)\Phi^{-1}(u)$.

The main theorems of this paper are:

THEOREM 1. *Suppose-using the notation of [1, p. 286]-the rank of $(A: B)$ and $(P: Q)$ are both k . Then $L(t, y) = -T^{-1}K^*(y, t)T$ for all $t, y \in [a, b]$ if and only if $PT^{-1}A^* = QT^{-1}B^*$.*

THEOREM 2. *If $U = D$, there exist $t, y \in [a, b]$ such that $L(t, y) \neq T^{-1}K^*(y, t)T$.*

We first establish some lemmas.

LEMMA 1. $N(t, u) = T^{-1}M^*(u, t)T, t, u \in [a, b]$.

For a proof see [4, Theorem 3.3]; it is a consequence of the fact that $H = -T^{-1}F^*T$.

LEMMA 2. $L(t, y) = -T^{-1}K^*(y, t)T$ for all $t, y \in [a, b]$ if and only if $U = -[T^{-1}D^*T + I]$.

Lemma 2 follows from Lemma 1, (4), (5) and (6).

Proof of Theorem 2. Just as in Lemma 2 one can show that $L(t, y) = T^{-1}K^*(y, t)T$ for all $t, y \in [a, b]$ if and only if

$$(7) \quad U = T^{-1}D^*T + I.$$

But for $U = D$ equation (7) implies $u_{11} = \bar{u}_{kk} + 1$ and $u_{kk} = \bar{u}_{11} + 1$. Hence $1 = u_{11} - \bar{u}_{kk} = \bar{u}_{11} - u_{kk} = -1$. This contradiction completes the proof.

Proof of Theorem 1. K is the Green's matrix for

$$(8) \quad \mathfrak{L}Y = Y' - FY = G, AY(a) + BY(b) = 0$$

i.e., for each continuous G the unique solution Y of (8) is given by

$$Y(t) = \int_a^b K(t, s)G(s)ds, t \in [a, b];$$

whereas L is the Green's function of

$$(9) \quad X' - HX = G, PX(a) + QX(b) = 0.$$

Now X satisfies (9) if and only if $Z = TX$ satisfies

$$(10) \quad \mathfrak{L}^+Z = -Z' - F^*Z = \tilde{G}, \tilde{P}Z(a) + \tilde{Q}Z(b) = 0,$$

where $\tilde{G} = -TG, \tilde{P} = PT^{-1}, \tilde{Q} = QT^{-1}$.

Problem (10) is adjoint to problem (8) if and only if

$$(11) \quad \tilde{P}A^* = \tilde{Q}B^*.$$

The argument for this is entirely similar to that for Theorem 3.1, p. 289 in [1]. By problem 6, p. 297 of [1], $\tilde{L}(t, y) = K^*(y, t)$ for $t, y \in [a, b]$, if and only if (11) holds, where \tilde{L} is the Green's matrix for (10). From the definition of \tilde{L} , we have $\tilde{L} = -TLT^{-1}$. Hence $L(t, y) = -T^{-1}K^*(y, t)T$ if and only if $PT^{-1}A^* = QT^{-1}B^*$.

Note. Some special cases of Theorem 1 were obtained in [4]-Theorems 6.13 and 7.1.

THEOREM 3. *If, in addition, F satisfies $f_{ij} = 0$ if $j > i + 1$ and $f_{i, i+1}(t) \neq 0$ for t in $[a, b], i = 1, \dots, k - 1$, then $L_{ik}(t, y) = (-1)^k \bar{K}_{1k}(y, t)$ for all $t, y \in [a, b]$ implies $L(t, y) = -T^{-1}K^*(y, t)T$ for all $t, y \in [a, b]$.*

Proof. Observe that for any y in $[a, b]$ the functions $m_{1i}(t, y), i = 1, \dots, k$, are linearly independent functions of t . Assume $L_{ik}(t, y) = (-1)^k \bar{K}_{1k}(y, t)$ for all t, y in $[a, b]$. Using (4), (5) and (6) we can show that the conditions of Lemma 2 are satisfied.

Comments.

1. In [3] it is shown that

$$Ly = y^{(k)} + \sum_{i=1}^{k-1} p_i y^{(i)} = 0,$$

with p_i continuous, is equivalent to a vector-matrix equation $Y' = FY$ with F satisfying the hypothesis of Theorem 3.

2. The $(1, k)$ component of K, K_{1k} , corresponds to the Green's function of an ordinary boundary value problem. An important special case occurs when F satisfies the hypothesis of Theorem 3 and $F = H$, $A = P, B = Q$. For this case Theorems 1 and 3 yield necessary and sufficient conditions, expressed solely in terms of the matrices defining the boundary conditions, for the ordinary Green's function K_{1k} of even and odd order problems to be hermitian or anti-hermitian, respectively. Furthermore, by Theorem 2, K_{1k} cannot be anti-hermitian for k even and hermitian for k odd.

3. An interesting way of getting examples of matrices A, B, P, Q satisfying the conditions of Theorem 1 is as follows: for any continuous $k \times k$ matrix F satisfying (1) define M and N as above, then for arbitrary B, Q, t , and u , let $A = BM(t, u)$ and $P = QN(t, u)$.

REFERENCES

1. E. Coddington and N. Levinson, *Theory of Ordinary Differential Equations*, McGraw-Hill, New York, 1955.
2. J. W. Neuberger, *Concerning boundary value problems*, Pacific J. Math. **10** (1960), 1385-1392.
3. Anton Zettl, *Adjoint linear differential operators* (to appear in the Proceedings).
4. ———, *Related Two Point Boundary Value Problems and Associated Harmonic Matrices*, Dissertation, University of Tennessee, 1964. Published as O.R.N.L. Report No. 3646.

Received April 21, 1965.

UNIVERSITY OF CALIFORNIA, LOS ANGELES
LOUISIANA STATE UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

R. M. BLUMENTHAL

University of Washington
Seattle, Washington 98105

*J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

* Paul A. White, Acting Editor until J. Dugundji returns.

William George Bade and Philip C. Curtis, Jr., <i>Embedding theorems for commutative Banach algebras</i>	391
Wilfred Eaton Barnes, <i>On the Γ-rings of Nobusawa</i>	411
J. D. Brooks, <i>Second order dissipative operators</i>	423
Selwyn Ross Caradus, <i>Operators with finite ascent and descent</i>	437
Earl A. Coddington and Anton Zettl, <i>Hermitian and anti-hermitian properties of Green's matrices</i>	451
Robert Arnold Di Paola, <i>On sets represented by the same formula in distinct consistent axiomatizable Rosser theories</i>	455
Mary Rodriguez Embry, <i>Conditions implying normality in Hilbert space</i>	457
Garth Ian Gaudry, <i>Quasimeasures and operators commuting with convolution</i>	461
Garth Ian Gaudry, <i>Multipliers of type (p, q)</i>	477
Ernest Lyle Griffin, Jr., <i>Everywhere defined linear transformations affiliated with rings of operators</i>	489
Philip Hartman, <i>On the bounded slope condition</i>	495
David Wilson Henderson, <i>Relative general position</i>	513
William Branham Jones, <i>Duality and types of completeness in locally convex spaces</i>	525
G. K. Kalisch, <i>Characterizations of direct sums and commuting sets of Volterra operators</i>	545
Ottmar Loos, <i>Über eine Beziehung zwischen Malcev-Algebren und Lietripelsystemen</i>	553
Ronson Joseph Warne, <i>A class of bisimple inverse semigroups</i>	563