

Pacific Journal of Mathematics

RELATIVE GENERAL POSITION

DAVID WILSON HENDERSON

RELATIVE GENERAL POSITION

DAVID W. HENDERSON

In this paper we will say that a piecewise linear map $f: K \rightarrow M$ from a finite complex into an n -manifold is a general position (gp) map, if for every pair of simplexes, A, B , contained in K ,

$$\begin{aligned} &(\text{dimension of the singularities of } f|A + B) \\ &\leq (\text{dimension of } A) + (\text{dimension of } B) - n. \end{aligned}$$

By letting $B = \emptyset$, we see that a gp map into an n -manifold is an embedding on each simplex of dimension less than or equal to n . Also note that the restriction of a gp map to a subcomplex is again a gp map. It is well known that every map f of a complex into a combinatorial manifold can be homotopically approximated by a gp map, g , on some subdivision of the complex. One might suppose that, if L is a subcomplex on which f is already a gp map, then $g|L$ could be made equal to $f|L$. However, this cannot be done, in general, even if the manifold is a Euclidean space and the complex is a subdivision of a cell. (See the Remark at the end of § 3.)

In § 3 are two general position theorems which fix the map on a subcomplex on which it is already a gp map, but not without some severe restrictions. These theorems are stated in terms of relative general position (rgp) which applied to maps from a pair into a pair. Section 4 considers maps $f: (D, \text{Bd } D) \rightarrow (M, N)$ of a 2-manifold, D , into a 3-manifold, M , with 2-submanifold, N , with the added restriction that $f(\text{Bd } D) \cdot f(D - \text{Bd } D) = \emptyset$. It is, in general, impossible in this setting to make f into an rgp map while keeping $f| \text{Bd } D$ fixed. However, two "relative normal position theorems" are proved which make the singularities "nice" while not considering a particular subdivision.

The proofs are contained in § 5 through 8.

It should be pointed out that E. C. Zeeman's definition of general position (see [9], p. 59, for general description and [10], Chapter 6, for detailed discussion and proofs) differs from the one used here and avoids most, if not all, of the difficulties encountered in this paper. Thus in a round-about fashion this paper points up several advantages in Zeeman's definition. However, Zeeman's definition may be undesirable for certain purposes. The main difference between the definitions is that Zeeman cannot require that a general position map from a complex K into a manifold to be *both* in general position on each subcomplex of K and a homeomorphism on each simplex of K .

2. Definitions.

2.1. A complex K will be considered to have a given fixed triangulation. L is called a *subcomplex of K* if L is a complex each of whose simplexes is a simplex of K . L is called a *polyhedron in K* if L is a subcomplex of some subdivision of K .

2.2. If K is a complex and L a subset of $[K]$, then *the closed star of L in K* , $\text{st}(L, K)$, is the union of all (closed) simplexes of K which intersect L . *The open star of L in K* , $\text{ost}(L, K)$, is the union of all open simplexes whose closures intersect L .

2.3. A mapping, f , from a complex K to a complex L is called *piecewise linear* (pwl) if the graph of f is a polyhedron in the product complex $K \times L$. If K is finite, then there are subdivisions α, β such that $f: \alpha K \rightarrow \beta L$ is simplicial.

2.4. A *n -manifold M^n* is a separable metric space each of whose points has a closed neighborhood homeomorphic to I^n , the standard n -cell.

The *boundary of M^n* , $\text{Bd } M^n$, is the set of points of M^n which do not have arbitrarily small neighborhoods homeomorphic to E^n , n -dimensional Euclidean space.

A *combinatorial n -manifold* is a complex such that the closed star of each vertex has a rectilinear subdivision which is isomorphic to a rectilinear subdivision of an n -simplex.

It follows easily from [1; 5] that all 2- and 3-manifolds may be given combinatorial triangulations, and henceforth in this paper we shall assume that this has been done.

If N^p and M^n are triangulated manifolds of dimension p and n , respectively, then N^p is a *p -submanifold of M^n* if N^p is a subcomplex of M^n .

2.5. If $f: K \rightarrow L$ is a pwl map of one complex into another, then by the *singularities of f* , $S(f)$, we shall mean the closure of the set of all points in L which are the images under f of more than one point of K . The cardinality of $f^{-1}(x)$ is called the *order of x* .

2.6. Let L be a subcomplex of K and let N^p be a polyhedral p -submanifold of the n -manifold M^n . A pwl map $f: (K, L) \rightarrow (M^n, N^p)$ is said to be a *relative general position (rgp) map* if

(a) for every pair of simplexes (A, B) in K such that B is not in L , $\dim S(f|A + B) \leq \dim A + \dim B - n$

(b) for every pair of simplexes (A, B) in L , $\dim S(f|A + B) \leq \dim A + \dim B - p$.

2.7. Let $f: J \rightarrow K$ be a pwl map of a 1-manifold into a 2-manifold. We shall call f *normal* if

- (a) f is at most 2-to-1 and $S(f)$ is a finite set of points, and
- (b) $f(J)$ crosses itself at each point of $S(f)$.

2.8. Let C^3 denote the solid cube in E^3 whose vertices are the eight points in E^3 which have as each coordinate either 1 or -1 .

2.9. Let $f: (D, \text{Bd } D) \rightarrow (M^3, N^2)$ be a pwl map of a 2-manifold D with boundary into a 3-manifold M^3 with 2-submanifold N^2 . We shall call f a *relative normal position (rnp) map* if

- (a) $f(\text{Bd } D) \cdot f(D - \text{Bd } D) = \emptyset$
- (b) f is at most 3-to-1
- (c) $f| \text{Bd } D: \text{Bd } D \rightarrow N$ is normal

(d) is a 1-dim polyhedron in M^3 consisting of (i) *double curves* (curves along which two sheets of $f(D)$ cross), (ii) *triple points* (points with arbitrarily small neighborhoods N such that $(N, N \cdot f(D))$ is homeomorphic to $(E^3, \text{coordinate planes})$), (iii) *branch points* (points with arbitrarily small closed neighborhoods N such that $(N, N \cdot f(D))$ is homeomorphic to $(C^3, \text{cone from the origin over a singular curve on } \text{Bd } C^3)$), and (iv) *pinched branch points* (points with arbitrarily small closed neighborhoods V such that $(V, V \cdot f(D))$ is homeomorphic to $(C^3, \text{cone from origin over two paths on } \text{Bd } C^3)$).

A *pinched branch point is simple* if the two paths on $\text{Bd } C^3$ are arcs. Note that all pinched branch points must be crossing points of $\text{Bd } D$ on N^2 because of (c).

2.10. If $f: K \rightarrow M$ and $g: L \rightarrow M$ are maps, then the *disjoint sum of f and g* is a map $f \oplus g: K \oplus L \rightarrow M$ defined by

$$(f \oplus g)(x) = \begin{cases} f(x), & \text{if } x \in K \\ g(x), & \text{if } x \in L, \end{cases}$$

where $K \oplus L$ denotes the abstract disjoint sum of K and L .

3. Relative general position theorems.

THEOREM I. *Let $f: (K, L) \rightarrow (M, N)$ be a pwl map of a pair into a pair, where L is a $(\leq p)$ -dimensional subcomplex of the finite and $(\leq n)$ -dimensional complex K and N is a p -submanifold of the combinatorial n -manifold M . Let ε be a positive number.*

If there exists a subdivision α of K and a pwl map $g: \alpha K \rightarrow M$ such that

(3.1) $g|L = f|L: \alpha L \rightarrow N$ is a gp map,

(3.2) g takes each closed star of a simplex in αK into the open star of a vertex in M ,

(3.3) g is a pwl embedding on each simplex of αK , and

(3.4) g is obtained from f by a homotopy of (K, L) into (M, N) which leaves L pointwise fixed and moves each point less than ε ,

then there exists an rgp map $g': (\alpha K, \alpha L) \rightarrow (M, N)$ that satisfies (3.1)–(3.4).

THEOREM II. Let $f: (K, L) \rightarrow (M, N)$ be a pwl map of a pair into a pair, where L is a $(\leq p)$ -dimensional subcomplex of the finite and $(\leq n)$ -dimensional complex K and N is a p -submanifold of the combinatorial n -manifold M . Let ε be a positive number.

If there exists a subdivision β of K such that

(3.5) $f|L: \beta L \rightarrow N$ is a gp map,

(3.6) $S(f|L)$ is a finite set of points, and

(3.7) if A is a simplex in βL such that $A \cdot f^{-1}(S(f|L))$ is not empty, then $f|A$ can be extended to a pwl map $F[A]: \text{st}(A, \beta K) \rightarrow M$ which is a homeomorphism on each simplex;

then there exists a subdivision, α , of K and an rgp map $g: (\alpha K, \alpha L) \rightarrow (M, N)$ that satisfies (3.1)–(3.4).

COROLLARY IIa. If $f: K \rightarrow M$ is a map of a finite $(\leq n)$ -dimensional complex into the combinatorial n -manifold M ,

then there exists a subdivision, α , of K and a pwl gp map $g: \alpha K \rightarrow M$ such that g is arbitrarily close, homotopically, to f .

Moreover, if L is a subcomplex of K and $f|L$ is a pwl homeomorphism, then we may require that $f|L = g|L$.

(To prove the Corollary, apply the relative simplicial approximation theorem [11] and Theorem II to $f: (K, L) \rightarrow (M, M)$.)

REMARK. Theorem II is false with (3.7) deleted. To see this, let $M = N = E^4$, K be some subdivision of 4-simplex, and $L = D_1 + D_2$ be the union of two disjoint 2-simplexes on $\text{Bd } K$. Let D'_1 be a polyhedral disk in E^4 which fails to be locally flat at a point p in the interior of D'_1 , and let D'_2 be any polyhedral disk in E^4 such that

$D'_1 \cdot D'_2 = p$. (For example, consider E^3 to be a 3-hyperplane of E^4 with $p \in E^4 - E^3$, and let D'_i be the cone from p over a sec J_i in E^3 , where J_1 is knotted in E^3 and J_2 links J_1 .) For $i = 1, 2$, let $f|D_i$ be a pwl homeomorphism onto D'_i ; and then extend to the rest of K in any pwl fashion. f satisfies (3.5) and (3.6) but there is no extension of $f|D_1 + D_2$ which is a gp map. This is because if $g: K \rightarrow M = E^4$ is an extension of $f|D_1 + D_2$ and a homeomorphism on each simplex of some subdivision λ of K , then, in order that $g|\lambda(D_1 + D_2)$ be a gp map, a neighborhood of p in D_1 lies on the boundary of the pwl homeomorphic image of a 4-simplex and thus D_1 is locally flat at p .

QUESTION. Can (3.6) be weakened?

4. Relative normal position theorems.

THEOREM III. *If $f: (D, J) \rightarrow (M, N)$ is a pwl map of a pair into a pair, where D is a 2-manifold with boundary J and N is a 2-submanifold of the 3-manifold M , and, in addition, $f(D - J) \cdot f(J) = \emptyset$, $f|J$ is normal, and ε is a positive number,*

then there exists a pwl map $g: (D, J) \rightarrow (M, N)$ such that

(4.1) *g is obtained from f by a homotopy of (D, J) into (M, N) which moves each point less than ε and only moves points at all in an ε -neighborhood of the set of points in M at which f fails to be in rnp (see 2.9),*

(4.2) *g is a rnp map, and*

(4.3) *$g|J = f|J$.*

COROLLARY IIIa. *If $f: D \rightarrow M$ is a Dehn surface in the 3-manifold, M , (i.e. $S(f) \cdot f(\text{Bd } D) = \emptyset$ and f is pwl),*

then by a "slight adjustment" of f we can get a pwl map $g: D \rightarrow M$ and a neighborhood N of $\text{Bd } D$ in D such that $g|N = f|N$ and $g: D \rightarrow M$ is a normal Dehn surface (i.e. $S(g)$ consists of double curves, branch points, and triple points (see 2.9)).

THEOREM IV. *Let N be a 2-submanifold of the 3-manifold M and h be a fixed-point-free pwl homeomorphism of (M, N) onto (M, N) such that hh equals the identity map. Let ε be a positive number.*

If $f: (D, J) \rightarrow (M, N)$ is a pwl map of a 2-manifold D with boundary J into (M, N) such that $(f(D - J) + hf(D - J)) \cdot (f(J) + hf(J)) = \emptyset$, and $f \oplus hf|J \oplus J$ is normal [$D \oplus D$ is the abstract disjoint union of two copies of D and $f \oplus hf$ is the mapping which is equal to f

on one copy of D and hf on the other (see 2.10)];

then there exists a pwl map $g: (D, J) \rightarrow (M, N)$ such that

(4.4) g is obtained from p by a homotopy of (D, J) into (M, N) which moves each point less than ε and only moves points at all in an ε -neighborhood of the set of points in M at which $f \oplus hf$ fails to be in rnp.

(4.5) $g \oplus hg: (D_1, J_1) \oplus (D_2, J_2) \rightarrow (M, N)$ is a rnp map, and

(4.6) $g|J = f|J$.

COROLLARY IVa. Let M be a 3-manifold and h a pwl homeomorphism of M onto M such that hh equals the identity map.

If $f: D \rightarrow M$ is an embedding of a surface into M such that $hf(\text{Bd } D) \cdot f(D - \text{Bd } D) = \emptyset$.

then by a "slight adjustment" of f we can get a pwl embedding $g: D \rightarrow M$ and a neighborhood N of $\text{Bd } D$ in D such that $g|N = f|N$ and $g(D) \cdot hg(D)$ is a finite collection of disjoint simple closed curves.

To prove the Corollary apply the Theorem and note that the only singularities possible are double lines.

The Corollaries are used in the various proofs of Dehn's Lemma and the Loop Theorem. (Corollary IIIa is used in [6] and its proof is indicated in [2]. Corollary IVa is used without proof in [7] and [8].) The relative versions in the Theorems are used in the proofs of the author's extensions of Dehn's Lemma and the Loop Theorem, [3] and [4].

It should be noted that if $g: D \rightarrow M$ is a normal Dehn surface, then there are arbitrarily fine subdivisions of D with respect to which g is a gp map. However, there is not necessarily any triangulation with respect to which a given rnp map is a rgp map, because a non-simple pinched branch point must be the image of a vertex if the map is a homeomorphism on each simplex.

5. Proof of Theorem I. Let $\alpha L = L_0 \subset L_1 \subset L_2 \subset \dots \subset L_k = \alpha K$, where for $i = 0, 1, 2, \dots, k-1$, $L_{i+1} = L_i + A_i$, A_i is a simplex of αK , and $A_i \cdot L_i = \text{Bd } A_i$. (This can be accomplished by adding the simplexes of αK_0 not in αL in some order of increasing dimension.) Assume inductively that there is a map $g_m: (K, L) \rightarrow (M, N)$ such that (let $g_0 = g$)

(5.1)_m g_m satisfies (3.1)–(3.4) with "g" replaced by "g_m",

(5.2)_m $g_m|L_m: (L_m, \alpha L) \rightarrow (M, N)$ is a rgp map, and

(5.3)_m $g_m|K - \text{ost}(A_{m-1}, \alpha K) = g_{m-1}|K - \text{ost}(A_{m-1}, \alpha K)$. (Note that $L_{m-1} \subset K - \text{ost}(A_{m-1}, \alpha K)$.)

Let $P = \text{st}(A_m, \alpha K)$ and $OP = \text{ost}(A_m, \alpha K)$. We shall now alter g_m in P , keeping it fixed on $P - OP$. Let Q be closed star of a vertex of M such that $g_m P \subset \text{int } Q$ and let h be a pwl embedding of Q onto the standard n -simplex Δ , which we consider as a convex subcomplex of n -dimensional Euclidean space, E^n . Let β be a subdivision of αK such that $\beta\alpha(g_m^{-1}(Q))$ is a subcomplex of $\beta\alpha K$, and

$$h \circ [g_m | g_m^{-1}(Q)] : \beta\alpha(g_m^{-1}(Q)) \rightarrow \Delta \subset E^n$$

takes simplexes linearly into simplexes, and such that each simplex of βA has at least one vertex in OP . Call $h \circ [g_m | g_m^{-1}(Q)] = H$ and $\beta\alpha(g_m^{-1}(Q)) = R$.

If C is a collection of simplexes in E^n , let $T(C)$ denote the union of all hyperplanes in E^n which contain n vertices of C . For a finite collection of simplexes, C , $E^n - T(C)$ is open and dense and each of its components is convex.

Let v_1, v_2, \dots, v_r be the images of those vertices of βA_m which are not $\beta(\text{Bd } A_m)$. (Remember that g_m , and therefore H , are embeddings on each simplex of αK and therefore are embeddings on each simplex of $\beta\alpha K$.) Let v_1^* be a point "very close" to v_1 in $\Delta \cdot [E^n - T(H(R))]$ such that the straight line segment from v_1 to v_1^* intersects $T(H(R))$ only at v_1 . Define $H_1: R \rightarrow \Delta \subset E^n$ to be equal to H on $R - \text{ost}(H^{-1}v_1, \beta P)$ and $H_1(H^{-1}v_1) = v_1^*$ and extend linearly to the rest of $\text{st}(H^{-1}v_1, \beta P)$. We leave to the reader the easy verification that H_1 is an embedding on each simplex of P . We now repeat the process with " v_2 " and " H_1 " replacing " v_1 " and " H ", and so forth until we get a map $H' = H_r$ that is a linear embedding on each simplex of R and a pwl embedding on each simplex of P , and such that, for each i , $H'(H^{-1}v_i) = v_i^*$ belongs to $[E^n - T(H(R - OP) + v_1^* + \dots + v_{i-1}^*)] \cdot \Delta$. (Note that $H | R - OP = H_1 | R - OP = H_2 | R - OP = \dots = H' | R - OP$.) Define $g_{m+1} | \alpha K - OP = g_m | \alpha K - OP$ and $g_{m+1} | P = h^{-1}H | P$.

We must now show that $(5.1)_{m+1} - (5.3)_{m+1}$ are satisfied. $(5.1)_{m+1}$ is satisfied if we make v_i close enough to v_i^* . $(5.3)_{m+1}$ is inherent in the construction. If $(5.2)_{m+1}$ is not satisfied then there is a b -dimensional simplex B in αK such that (let $a = \text{dimension of } A_m$)

$$\text{dimension}(S(g_{m+1} | A_m + B)) > b + a - n .$$

Then there would be a $(\leq b)$ -dimensional simplex B' in $\beta B \cdot R$ and a $(\leq a)$ -dimensional simplex A' in βA_m such that

$$\text{dimension}((g_{m+1} A' \cdot g_{m+1} B') - g_{m+1}(A' \cdot B')) > b + a - n .$$

However, if v_i^* is the one member of $\{v_1^*, \dots, v_r^*\} \cdot g_{m+1} A'$ with highest subscript, then it is a straightforward exercise in linear algebra to show that v_i^* belongs to

$$T[g_{m+1}(A' + B') - \text{ost}(v_i^*, g_{m+1}A')] \subset T(H(R - OP) + v_i^* + \cdots + v_{i-1}^*)$$

which contradicts our restrictions on the choice of v_i^* .

Thus we may conclude that there exists a map $g' = g_k$ that satisfies (5.1)_k – (5.3)_k. This is the map required in Theorem I.

6. Proof of Theorem II. First, we prove a necessary

LEMMA. *Let C be pwl homeomorphic (under h) to the standard n -simplex (Δ) in E^n , and let A be a ($\leq n$)-dimensional simplex.*

If $f: \alpha A \rightarrow C$ is a pwl map which is a homeomorphism on each simplex of $\alpha(\text{Bd } A)$, for some subdivision α of A ,

then f is arbitrarily close, homotopically, to a pwl map $g: \pi A \rightarrow C$ such that $g|_{\text{Bd } A} = f|_{\text{Bd } A}$ and g is a homeomorphism on each simplex of πA , where π is some subdivision of A and refines α .

Proof. Let π be a subdivision of A such that πA is a subdivision of αA and $hf: \pi A \rightarrow C \rightarrow \Delta \subset E^n$ is linear on each simplex of πA . Further, assume that, for each simplex B in πA , $B \cdot \text{Bd } A$ is a single face (possibly empty) of B . Let v_1, \dots, v_r be the vertices of πA not in $\pi(\text{Bd } A)$. Let v'_1, \dots, v'_r be a collection of points in Δ such that, for each i , v'_i is “close” to $hf(v_i)$ and v'_i does not lie on any linear subspace of E^n that is determined by some collection of $k(\leq n)$ points from $\{v'_1, \dots, v'_{i-1}, v'_{i+1}, \dots, v'_r\} + hf$ (the vertices of $\pi(\text{Bd } A)$). Define $g|_{\text{Bd } A} = f|_{\text{Bd } A}$, $g(v_i) = v'_i$ for each i , and then extend linearly to the rest of A . By making v'_i close enough to $hf(v_i)$, for each i , we can make g as close, homotopically, to f as we please. It is an easy exercise to show that g is a homeomorphism on each simplex of πA .

We now return to the proof of Theorem II. By the *carrier* of a point, p , in βK we shall mean the unique simplex of βK which contains p in its interior. Note that any subdivision μ of βK satisfies (3.5)–(3.7) if, for every point p of $f^{-1}(S(f|L))$, the carrier of p in $\mu\beta K$ has the same dimension as the carrier of p in βK . Therefore we may assume without loss of generality that β is so fine that, for every simplex A of βK , $f(\text{st}(A, \beta K))$ is contained in some open star of a vertex of M . Let A_1, \dots, A_s be the collection of carriers of points of $f^{-1}(S(f|L))$. Since A_i is compact and there is a pwl homeomorphism, keeping A_i fixed, of $\text{st}(A_i, \beta K)$ into any neighborhood of A_i in $\text{st}(A_i, \beta K)$, we may assume that, for each i , $F[A_i](\text{st}(A_i, \beta K))$ is contained in the same open star of a vertex as $f(\text{st}(A_i, \beta K))$. We also suppose that $f[A_i] = f|_{\text{st}(A_i, \beta K)}$, if the latter is a homeomorphism on each simplex. We can now easily obtain a pwl map $f^*: (\beta K, \beta L) \rightarrow (M, N)$ such that $f^*|L = f|L$ and $f^*|_{\text{st}(A_i, \beta K)} = F[A_i]$ and such that (3.4) is satisfied with “ g ” replaced by “ f^* ”. Let B_1, \dots, B_t be the simplexes of βK

not in $\beta L + \sum [\text{st}(A_i, \beta K)]$ and suppose that they are arranged in some order of increasing dimension. Let

$$L_j = \beta L + \sum_{i=1}^{j-1} [\text{st}(A_i, \beta K)] + \sum_{k=1}^{j-1} [B_k],$$

for $j = 0, 1, \dots, t$. Note that $\text{Bd } B_{j+1} \subset L_j$. We assume inductively that we have a subdivision μ_j of K and a pwl map $f_j: (\mu_j K, \mu_j L) \rightarrow (M, N)$ such that

- (6.1)_{*j*} f_j is a homeomorphism on each simplex of $\mu_j L_j$,
- (6.2)_{*j*} $f_j|L_0 = f^*|L_0$,
- (6.3)_{*j*} $\mu_j K$ is a subdivision of βK ,
- (6.4)_{*j*} each A_i is a simplex of $\mu_j K$,
- (6.5)_{*j*} f_j maps the star of each simplex of βK into the open star of a vertex of M , and
- (6.6)_{*j*} f_j is obtained from f by a homotopy of (K, L) into (M, N) which leaves L pointwise fixed and moves each point less than ε .

Note that $f^* = f_0$ satisfies (6.1)₀—(6.6)₀. We will now use the Lemma and μ_j and f_j to construct μ_{j+1} and f_{j+1} . Let C be the open star of some vertex of M such that $f_j(\text{st}(B_{j+1}, \beta K)) \subset C$. Thus we can apply the Lemma and obtain a pwl map $g: \pi B_{j+1} \rightarrow C$ arbitrarily close, homotopically, to $f_j|B_{j+1}$ such that $g|B_{j+1} = f_j|B_{j+1}$, g is a homeomorphism on each simplex of πB_{j+1} , and $\pi(\text{Bd } B_{j+1})$ is a subdivision of $\mu_j(\text{Bd } B_{j+1})$. Then define $\mu_{j+1}(B_{j+1}) = \pi(B_{j+1})$ and extend to a subdivision of $\mu_j K$. Since $\text{st}(A_i, \beta K)$ is contained in L_0 and therefore in L_j , we may do the extending in such a way that (6.4)_{*j+1*} is satisfied. We define $f_{j+1}|B_{j+1} = g$ and

$$f_{j+1}|K - \text{ost}(B_{j+1}, \beta K) = f_j|K - \text{ost}(B_{j+1}, \beta K)$$

and then extend (in C) to the rest of $\text{st}(B_{j+1}, \beta K)$. It is clear that if g is close enough to $f_j|B_{j+1}$ then (6.5)_{*j+1*} and (6.6)_{*j+1*} will be satisfied. (6.1)_{*j+1*} is satisfied because $\mu_{j+1} K$ is a subdivision of $\mu_j K$.

Thus by induction we may assume that there is a pwl map $f_t: (\mu_t K, \mu_t L) \rightarrow (M, N)$ that satisfies (6.1)_{*t*}—(6.6)_{*t*}. We now show that f_t and μ_t satisfy (3.1)—(3.4) with “ g ” replaced by “ f_t ” and “ α ” replaced by “ μ_t ”. (3.1) follows from (6.2)_{*t*}—(6.4)_{*t*}, (3.2) follows from (6.5)_{*t*}, (3.3) follows from (6.1)_{*t*}, and (3.4) follows from (6.6)_{*t*}.

Thus Theorem II now follows from Theorem I.

7. Proof of Theorem III. We first obtain a subdivision λ of D and a pwl map $g': \lambda D \rightarrow M$ that satisfies (4.1) and (4.2) with “ g ”

replaced by “ g' ”, and so that g' is a homeomorphism on each simplex of λD . The proof that such a λ and g' exist follows closely the arguments in § 6, with the arguments dealing with the A_i 's omitted; and therefore we will not give the details here. Suffice it to say that one should find a subdivision β of M so that ε is larger than the diameter of each vertex star of βM and then require that the image under f and g' of each vertex star of λD must be contained in the open star of a vertex of M .

Let α and δ be subdivision of M and D , respectively, so that g' is simplicial from δD to αM . Then $g'(D)$ is a subcomplex of αM and $S(g')$ is a subcomplex of $g'(D)$. Let α^2 denote the second barycentric subdivision of α . Note that $S(g') \cdot g'(J)$ is just the crossing points of $g' | J$.

If ω is a 2-simplex of $S(g')$, then $g'^{-1}(\omega)$ is the union of two or more 2-simplexes of D and $\text{st}(\omega, \alpha^2 M)$ is a 3-cell of which ω is a spanning disk. Being careful not to move things too far we can adjust the interiors of the images of each 2-simplex of $g'^{-1}(\omega)$ so that they still lie in $\text{st}(\omega, \alpha^2 M)$ but are disjoint except for their boundaries. By applying this procedure to each 2-simplex in $S(g')$ we can get a pwl map $g'': D \rightarrow M$ that satisfies (4.1) and (4.3) and $S(g'')$ is a 1-subcomplex of αM .

If τ is a 1-simplex of $S(g'')$, then $\text{st}(\tau, \alpha^2 M)$ is a 3-cell of which τ is a spanning arc and $g''^{-1}(\text{st}(\tau, \alpha^2 M))$ is the union of two or more subdisks of D , the image of each being a spanning disk of $\text{st}(\tau, \alpha^2 M)$. We may then alter these spanning disks slightly in the interior of $\text{st}(\tau, \alpha^2 M)$ so that the only singularities in the interior of $\text{st}(\tau, \alpha^2 M)$ are double lines whose endpoints are the endpoints of τ . In this fashion we can obtain a pwl map g''' that satisfies (4.1) and (4.3) and g''' fails to be a rnp map only at the vertices of $S(g'')$. Analogously, we adjust slightly the images in $\text{st}(v, \alpha^2 M)$ for each vertex, v , of $S(g'') - g''(J)$ so that the only singularities in $\text{st}(v, \alpha^2 M)$ are triple points, branch points, and the ends of double lines. The pwl map then obtained is the desired map g . The vertices of $S(g'') \cdot g''(J)$ are just the crossing points of $g''' | J = g'' | J$ and thus the pinched branch points of $g''' | D$.

8. Proof of Theorem IV. First we wish to pick subdivisions of D and M so that both f and h are simplicial with respect to these subdivisions. Let ν and β be subdivisions so that $f: \nu D \rightarrow \beta M$ is simplicial. Since h is pwl, there exist refinements β_1 and β_2 of β such that $h: \beta_1 M \rightarrow \beta_2 M$ is simplicial. Let $(\beta_1 \cdot \beta_2)M$ denote the convex linear cell complex composed of all cells of the form $\tau_1 \cdot \tau_2$ where τ_i is a simplex of $\beta_i M$. (See [10], Chapter 1, page 5, for a description of

convex linear cell complex.) Now the image under h of a cell of $(\beta_1 \cdot \beta_2)M$ is again a cell of $(\beta_1 \cdot \beta_2)M$. Then, order the cells of $(\beta_1 \cdot \beta_2)M$ in some order of increasing dimension, $\mu_1, \mu_2, \dots, \mu_p$, so that for each t , $\mu_{2t} = h(\mu_{2t-1})$. We now subdivide, one at a time in order, each cell into a simplex, while leaving the subdivision fixed on the boundary of the cell; and in doing this we let $h(\mu_{2t-1})$ determine the subdivision on μ_{2t} , for each t . (See [10], Chapter 1, Lemma 1.) In this way we obtain a subdivision α of M that refines β , and such that $h: \alpha M \rightarrow \alpha M$ is simplicial. By Lemma 5 of Chapter 1 of [10], there is a subdivision δ of D such that $f: \delta D \rightarrow \alpha M$ is simplicial.

Theorem IV can now be proved using essentially the same arguments used in the proof of Theorem III, with the exception that all subdivisions of M should be such that h is simplicial with respect to them and in altering the maps the changes made in the star of a simplex should "agree" with the changes of the maps in the image of that star under h .

REFERENCES

1. R. H. Bing, *Locally tame sets are tame*, Ann. of Math., **59** (1954), 145-158.
2. M. Dehn, *Über die topologie des dreidimensionalen raumes*, Math. Ann. **69** (1910), 137-168.
3. D. W. Henderson, *Extensions of Dehn's lemma and the loop theorem*, Trans. Amer. Math. Soc. **120** (1965), 448-469.
4. ———, *Self-unlinked simple closed curves*, Trans. Amer. Math. Soc. **120** (1965), 470-480.
5. E. E. Moise, *Affine structures in 3-manifolds, V. The triangulation theorem and hauptvermutung*, Ann. of Math. **56** (1952), 96-114.
6. C. D. Papakyriakopoulos, *Dehn's lemma and the asphericity of knots*, Ann. of Math. **66** (1957), 1-26.
7. A. Shapiro and J. H. C. Whitehead, *A proof and extension of Dehn's lemma*, Bull. Amer. Math. Soc. **64** (1958), 174-178.
8. J. Stallings, *On the loop theorem*, Ann. of Math. **72** (1960), 12-19.
9. E. C. Zeeman, *Polyhedral Manifolds*, Topology of 3-manifolds, ed. by M. K. Fort, 1961.
10. ———, *Seminar on Combinatorial Topology*, Institut des Hautes Etudes Scientifiques, 1963, (mimeographed).
11. ———, *Relative simplicial approximation*, Proc. Camb. Phil. Soc. **60** (1964), 39-43.

Received January 30, 1965. This research was supported by a grant from The Institute for Advanced Study.

THE INSTITUTE FOR ADVANCED STUDY
PRINCETON, N.J.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON
Stanford University
Stanford, California

R. M. BLUMENTHAL
University of Washington
Seattle, Washington 98105

*J. DUGUNDJI
University of Southern California
Los Angeles, California 90007

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

* Paul A. White, Acting Editor until J. Dugundji returns.

Pacific Journal of Mathematics

Vol. 18, No. 3

May, 1966

William George Bade and Philip C. Curtis, Jr., <i>Embedding theorems for commutative Banach algebras</i>	391
Wilfred Eaton Barnes, <i>On the Γ-rings of Nobusawa</i>	411
J. D. Brooks, <i>Second order dissipative operators</i>	423
Selwyn Ross Caradus, <i>Operators with finite ascent and descent</i>	437
Earl A. Coddington and Anton Zettl, <i>Hermitian and anti-hermitian properties of Green's matrices</i>	451
Robert Arnold Di Paola, <i>On sets represented by the same formula in distinct consistent axiomatizable Rosser theories</i>	455
Mary Rodriguez Embry, <i>Conditions implying normality in Hilbert space</i>	457
Garth Ian Gaudry, <i>Quasimeasures and operators commuting with convolution</i>	461
Garth Ian Gaudry, <i>Multipliers of type (p, q)</i>	477
Ernest Lyle Griffin, Jr., <i>Everywhere defined linear transformations affiliated with rings of operators</i>	489
Philip Hartman, <i>On the bounded slope condition</i>	495
David Wilson Henderson, <i>Relative general position</i>	513
William Branham Jones, <i>Duality and types of completeness in locally convex spaces</i>	525
G. K. Kalisch, <i>Characterizations of direct sums and commuting sets of Volterra operators</i>	545
Ottmar Loos, <i>Über eine Beziehung zwischen Malcev-Algebren und Lie tripelsystemen</i>	553
Ronson Joseph Warne, <i>A class of bisimple inverse semigroups</i>	563