

Pacific Journal of Mathematics

***K*- AND *L*-KERNELS ON AN ARBITRARY RIEMANN SURFACE**

MYRON GOLDSTEIN

K - AND L -KERNELS ON AN ARBITRARY RIEMANN SURFACE

MYRON GOLDSTEIN

The l -kernel which was first considered by Schiffer for plane regions is extended to arbitrary open Riemann surfaces for a number of significant subspaces of the space of square integrable harmonic differentials Γ_h . The l -kernel for each of the subspaces considered is expressed in terms of the principal functions. Thus if W is an open Riemann surface and p and q the L_1 principal functions of W with singularities $\operatorname{Re} 1/z$ and $\operatorname{Im} 1/z$ respectively, then the following result is proved.

THEOREM. The differential $dp - dq^*$ is an l -kernel for the space Γ_h .

The l -kernel and another kernel function called the k -kernel by Schiffer are applied to the solution of some well known extremal problems on open Riemann surfaces.

It should be noted here that these problems have also been considered by G. Weill [9]. Finally, the properties of the kernel functions are used to obtain a test for when a surface is of class 0_{AD} .

M. Schiffer in [7] defined the k - and l -kernels for plane regions G . The k -kernel reproduces the value of every square integrable analytic function on G at a prescribed point while the l -kernel is orthogonal to the space of square integrable analytic functions on G with Dirichlet metric. Schiffer showed that these kernel functions can be expressed in terms of the Green's function thus enabling one to actually construct them for a given region.

An important question is whether the k - and l -kernels can be generalized to arbitrary open Riemann surfaces and, if so, whether they can be expressed in terms of functions depending only on the surface as in the case of plane regions. We shall answer this question in the affirmative for a number of significant subspaces of the space of square integrable harmonic differentials. In addition, we shall see that these generalized k - and l -kernels have important extremal properties.

1. Principal functions

2. In this section we shall briefly review certain results on principal functions (see cf. [1] pp. 148–186) that will be needed later on.

Let W be the interior of a compact bordered Riemann surface \bar{W}

with border β . Consider a regularly imbedded boundary neighborhood $W' \subset W$ with compact complement and with relative boundary α , and a continuous function f on α . Then $L_0 f$ is defined to be the harmonic function on W' with boundary values f on α and with vanishing normal derivative on β . $L_0 f$ is unique.

Let P be a partition of the contours of W . That is, $\beta = \cup \beta_i$, where the β_i are disjoint unions of contours. We then define $(P)L_1 f$ to be the harmonic function on W' with boundary values f on α and constant values on each β_i such that the flux across β_i vanishes. L_0 and $(P)L_1$ are called the principal operators.

Let W now be an arbitrary open Riemann surface, and W' a fixed regularly imbedded subregion $\subset W$ with compact complement and relative boundary α , negatively oriented with respect to W' . We can extend the definitions of the principal operators to W' as follows. Let $\{\Omega_n\}$, with borders $\{\alpha \cup \beta_n\}$, be a regular exhaustion of W' . $L_0(\Omega_n)$ and $(P)L_1(\Omega_n)$, the principal operators on Ω_n , are well defined and by taking the limit as Ω_n tends to W' , we obtain the principal operators on W' . For brevity, we shall use L_1 for $(P)L_1$.

3. The principal functions p_0 and p_1 of an arbitrary open Riemann surface W are defined as follows.

DEFINITION. Suppose at a finite number of points $\zeta_j \in W$, there are given singularities of the form

$$(3.1) \quad \operatorname{Re} \sum_{n=1}^{\infty} b_n^{(j)} (z - \zeta_j)^{n-1} + c^{(j)}$$

where the $c^{(j)}$ are real and subject to the condition $\sum_j c^{(j)} = 0$. The principal functions p_0 and p_1 are by definition harmonic on W , except for the singularities (3.1), such that $L_0 p_0 = p_0$ and $L_1 p_1 = p_1$ in W' .

These functions are unique and independent of W' , save for an additive constant (cf. [1] p. 169).

2. Harmonic differentials on Riemann surfaces

4. We shall consider here the space of square integrable harmonic differentials Γ_h and some significant subspaces of Γ_h . The main tool to be used is the general formula for partial integration which states that if f is of class C^1 , and if x is a 2-chain, then

$$(4.1) \quad \int_x (df)\omega = \int_{\partial x} f\omega - \int_x f d\omega$$

holds for all differentials $\omega \in C^1$ and thus in particular for all $\omega \in \Gamma_h$.

The k -kernel of any closed subspace $\Gamma_{h'}$ of Γ_h is defined as follows.

DEFINITION. A differential ψ is said to a k -kernel of $\Gamma_{h'}$ provided

$$(4.2) \quad \psi \in \Gamma_{h'}$$

$$(4.3) \quad \text{for all } \omega \in \Gamma_{h'}, (\omega, \psi) = \frac{\partial}{\partial x} u(\zeta)$$

where $\omega = du$ near ζ .

The l -kernel of Γ_h is defined as follows.

DEFINITION. A harmonic differential θ on $W - \{\zeta\}$ with a harmonic pole at ζ is said to be an l -kernel of Γ_h provided

$$(4.4) \quad (\omega, \theta) = 0 \text{ for all } \omega \in \Gamma_h \text{ where the inner product is taken in the Cauchy sense.}$$

The k -kernel is easily seen to be unique while the l -kernel depends on the singularity. However the following result on the l -kernel is valid.

THEOREM 1. *If θ_1 and θ_2 are two l -kernels of Γ_h with the same singularity, then $\theta_1 = \theta_2$.*

Proof. Since θ_1 and θ_2 have the same singularity, $\theta_1 - \theta_2 \in \Gamma_h$. By property 4.4, $(\theta_1 - \theta_2, \theta_1 - \theta_2) = 0$ and thus $\theta_1 = \theta_2$.

Consequently, we can say that the l -kernel of Γ_h is unique up to a singularity.

We shall now relate the k - and l -kernels of Γ_h to the principal functions and thus obtain constructive proofs of the existence of the k - and l -kernels.

Let W be an arbitrary open Riemann surface with ideal boundary β , and denote by r , the L_1 principal function of W with respect to the identity partition of β and for any singularity. The following auxiliary result is valid.

LEMMA 1 (Rodin [6]). If $\omega \in \Gamma_h$, then

$$(5.1) \quad \int_{\beta} r \omega = 0.$$

Proof. Let Ω denote a canonical subregion of W with border β_{Ω} , and r_{Ω} the L_1 principal function of Ω with respect to the identity partition of β_{Ω} . Since r_{Ω} is constant on β_{Ω} , it follows that

$$(5.2) \quad \int_{\beta_{\Omega}} r_{\Omega} \omega = 0.$$

By 4.1, the right hand side of the above equation is dominated by $|(dr - dr_\Omega, \bar{\omega}^*)_\Omega|$. Consequently, using Schwarz's inequality, we find that

$$(5.3) \quad \left| \int_{\beta_\Omega} r\omega \right| \leq \|dr - dr_\Omega\|_\Omega \|\omega\|_\Omega$$

In Ahlfors and Sario [1], it is shown that $\lim_{\Omega \rightarrow W} \|dr - dr_\Omega\| = 0$. Hence the right hand side of 5.3 tends to 0 as Ω tends to W which implies the result of the lemma.

Let ζ denote an arbitrary point on W and p and q the L_1 principal functions of W with respect to the identity partition of β and singularities $\operatorname{Re} 1/z$ and $\operatorname{Im} 1/z$ respectively at ζ . We can now state the following result.

THEOREM 2. *The differential $dp - dq^*$ is an l -kernel of the Hilbert space Γ_h . Explicitly*

$$(5.4) \quad (\omega, dp - dq^*) = 0$$

for all $\omega \in \Gamma_h$.

Proof. Let Δ denote a parametric disc with center at ζ and corresponding to $\{z: |z| \leq r < 1\}$. In Δ , $p = \operatorname{Re}(1/z) + h(z)$ and $q = \operatorname{Im}(1/z) + v(z)$ where $h(z)$ and $v(z)$ are harmonic. From the definition of the inner product, it follows that

$$(5.5) \quad (\omega, dp - dq^*)_{W-\Delta} = \iint_{W-\Delta} (dp)\omega^* - \iint_{W-\Delta} (dq)\omega.$$

By 4.1 and Lemma 1, we may rewrite 5.5 as follows.

$$(5.6) \quad (\omega, dp - dq^*)_{W-\Delta} = \int_{|z|=r} q\omega - p\omega^*.$$

By the linearity of the scalar product, we may assume without loss of generality that ω is real. Therefore in Δ , we can write $\omega = du$ where u is harmonic. Since $\operatorname{Re}(1/z) = \operatorname{Im}(i/\bar{z})$ and $\operatorname{Im}(1/z) = \operatorname{Im}(1/\bar{z})$, we obtain from 5.6, the following result.

$$(5.7) \quad (\omega, dp - dq^*)_{W-\Delta} = \operatorname{Im} \int_{|z|=r} \frac{du + i du^*}{\bar{z}} + \int_{|z|=r} h du^* + v du.$$

On the circle $|z| = r$, $\bar{z} = r^2/z$ and therefore

$$(5.8) \quad \operatorname{Im} \int_{|z|=r} \frac{du + i du^*}{\bar{z}} = 0.$$

Since u and u^* are of bounded variation on $|z| = r$, it follows that

$$(5.9) \quad \int_{|z|=r} h du^* + v du = o(1).$$

Consequently,

$$(5.10) \quad (\omega, dp - dq^*)_{W-A} = o(1).$$

Letting r tend to 0, we obtain the desired conclusion.

It was shown by Rodin [6] in his doctoral dissertation that the corresponding k -kernel of Γ_h is $-(1/2\pi)(dp + dq^*)$.

6. We shall now consider the space of square integrable harmonic exact differentials Γ_{he} . The l -kernel of Γ_{he} is defined as follows.

DEFINITION. A harmonic exact differential θ on $W - \{\zeta\}$ with a harmonic pole at ζ is said to be an l -kernel of Γ_{he} provided

(6.1) $(\omega, \theta) = 0$ for all $\omega \in \Gamma_{he}$ where the inner product is again taken in the Cauchy sense.

We shall now prove the following lemma.

LEMMA 2. *Let W be an arbitrary open Riemann surface with border β , and let p denote the L_0 principal function of W for any singularity. Then*

$$(6.2) \quad \int_{\beta} u dp^* = 0$$

for all $du \in \Gamma_{he}$.

Proof. As before Ω will denote a canonical subregion of W with border β_{Ω} . Since p_{Ω} , the L_0 principal function of Ω with the same singularity as p , has vanishing normal derivative on β_{Ω} , it follows that

$$(6.3) \quad \left| \int_{\beta_{\Omega}} u dp^* \right| = \left| \int_{\beta_{\Omega}} u (dp^* - dp_{\Omega}^*) \right|$$

The rest of the proof now follows in the same manner as that of Lemma 1.

Let p_{is} denote the L_1 principal function of W with respect to the identity partition and singularity $s = \operatorname{Re}(1/z)$, and p_s denote the L_0 principal function of W with singularity s . Applying Lemma 2, we obtain in a manner similar to the proof of Theorem 2 the following result.

THEOREM 3. *The differential $dp_{1Is} + dp_{os}$ is an l -kernel of Γ_{he} .*

It should be noted that the l -kernel of Γ_{he} is also unique up to a singularity. This remark follows in the same manner as that of Theorem 1.

The corresponding k -kernel [6] of Γ_{he} is $-(1/2\pi)(dp_{1Is} - dp_{os})$.

Let us now turn to the space of square integrable harmonic semiexact differentials Γ_{hse} . We recall that a differential (not necessarily square integrable) is said to be harmonic semiexact on an arbitrary open Riemann surface W if it is harmonic with vanishing periods along all dividing cycles of W . This leads us to the following definition.

DEFINITION. A harmonic semiexact differential θ on $W - \{\zeta\}$ with a harmonic pole at ζ is said to be an l -kernel of Γ_{hse} provided

$$(6.4) \quad (\omega, \theta) = 0 \text{ for all } \omega \in \Gamma_{hse}.$$

By a method of proof similar to that of Lemma 1, we obtain the following result.

LEMMA 3. *If p denotes the L_1 principal function of W with respect to the canonical partition for any singularity, then*

$$(6.3) \quad \int_{\beta} p\omega = 0$$

for all $\omega \in \Gamma_{hse}$.

Denote by p_{1ot} the L_1 principal function of W with respect to the canonical partition and with singularity $t = \text{Im}(1/z)$. Applying the result of Lemma 3, we can establish in a manner similar to the proof of Theorem 2 the following.

THEOREM 4. *The differential $(dp_{1Is} - dp_{1ot}^*)$ is an l -kernel of Γ_{hse} .*

We again note that the l -kernel of Γ_{hse} is unique up to a singularity. The corresponding k -kernel [6] is $-(1/2\pi)(dp_{1Is} + dp_{1ot}^*)$.

7. The k -kernel of Γ_{he} can also be characterized in terms of a complete orthonormal set of square integrable harmonic exact differentials on any surface $W \notin 0_{HD}$. In fact, if $\{du_\nu\}$, $\nu = 1, 2, \dots$, is such a complete orthonormal set, then as an immediate consequence of the Riesz Fischer theorem we obtain the following result.

THEOREM 5.

$$-\frac{1}{2\pi} (dp_{1s} - dp_{os}) = \sum_{\nu=1}^{\infty} \frac{\partial u(\zeta)}{dx} du_{\nu}.$$

3. Analytic differentials on Riemann surfaces

8. We shall now consider some important subspaces of the space of square integrable analytic differentials Γ_a . The l -kernel of Γ_a is defined as follows.

DEFINITION. An analytic differential θ on $W - \{\zeta\}$ with a pole at ζ is said to be an l -kernel of Γ_a provided

(8.1) $(\omega, \theta) = 0$ for all $\omega \in \Gamma_a$ where the inner product is taken in the Cauchy sense.

With the same notation as before, we obtain the following result.

THEOREM 6. *The differential $dp - dq^* + i(dp^* + dq)$ is an l -kernel for Γ_a .*

The proof is similar to that of Theorem 2. Of course, the l -kernel is again unique up to a singularity.

Rodin [6] has shown that the k -kernel of Γ_a is

$$-\frac{1}{4\pi} (dp + dq^* + i(dp^* - dq)).$$

Let us now consider the space of square integrable analytic semiexact differentials Γ_{ase} . If in the definition of the l -kernel of Γ_a we replace the word analytic by analytic semiexact and the space Γ_a by the space Γ_{ase} , we obtain the definition of the l -kernel of Γ_{ase} .

Denote by p_{1os} the L_1 principal function of W with singularity $s = \text{Re}(1/z)$ at ζ with respect to the canonical partition. As a consequence of Lemma 3, we have the following result.

THEOREM 7. *The differential $dp_{1os} - dp_{1ot}^* + i(dp_{1os}^* + dp_{1ot})$ is an l -kernel of Γ_{ase} .*

The corresponding k -kernel [6] of Γ_{ase} is

$$-\frac{1}{4\pi} (dp_{1os} + dp_{1ot}^* + i(dp_{1os}^* - dp_{1ot})).$$

We shall now consider the space of square integrable analytic exact differentials. If in the definition of the l -kernel of Γ_a we

replace the word analytic by analytic exact and Γ_a by Γ_{ae} , we obtain the definition of the l -kernel of the space of square integrable exact differentials Γ_{ae} .

Let us now consider a planar Riemann surface W . On a planar surface every cycle is dividing and consequently $P_1 = p_{1\sigma_s} + ip_{1\sigma_s}^*$ and $P_0 = p_{0\sigma_s} + ip_{0\sigma_s}^*$ become single valued meromorphic functions. Thus we obtain the following result.

COROLLARY 1. *On a planar surface, the differential $dP_1 + dP_0$ is an l -kernel of Γ_{ae} .*

The corresponding k -kernel [6] is $-(1/4\pi)(dP_1 - dP_0)$.

9. The k -kernel of Γ_a can also be characterized in terms of a complete orthonormal set of square integrable analytic differentials. To be precise, if $W \notin 0_{AD}$ and if $a_\nu(z)dz, \nu = 1, 2, \dots$, is such a complete orthonormal set, then the following result is valid.

THEOREM 8.

$$-\frac{1}{4\pi} (dp + dq^* + i(dp^* - dq)) = \sum_{\nu=1}^{\infty} a_\nu(\zeta) a_\nu(z) dz.$$

The proof is similar to that of Theorem 5.

4. Extremal problems on Riemann surfaces

The properties of the k - and l -kernels of the space Γ_{ae} can be used to solve certain extremal problems for harmonic functions on an arbitrary open Riemann surface. Thus if we let $B(p) = \int_{\beta} dp^*$, we obtain the following result.

THEOREM 9. *The function $(1/2)(p_{1Is} + p_{0s})$ minimizes the expression $B(p)$ in the class of all harmonic functions p with singularity s .*

Proof. Since $dp_{1Is} + dp_{0s}$ is an l -kernel of Γ_{ae} , it follows that

$$(9.1) \quad 0 \leq \left\| d\left(p - \frac{1}{2}(p_{1Is} + p_{0s})\right) \right\|_{W-A}^2 = \|dp\|_{W-A}^2 \\ - (dp, dp_{1Is} + dp_{0s})_{W-A} + \frac{1}{4} \|dp_{1Is} + dp_{0s}\|_{W-A}^2$$

and that

$$(9.2) \quad \left\| d\left(p - \frac{1}{2}(p_{1Is} + p_{os})\right) \right\|_{W-A}^2 = \|dp\|_{W-A}^2 - \frac{1}{2}(dp, dp_{1Is} + dp_{os})_{W-A} + o(1).$$

Consequently

$$(9.3) \quad (dp, dp_{1Is} + dp_{os})_{W-A} = \frac{1}{2} \|dp_{1Is} + dp_{os}\|_{W-A}^2 + o(1),$$

and therefore

$$(9.4) \quad \frac{1}{4} \|dp_{1Is} + dp_{os}\|_{W-A}^2 \leq \|dp\|_{W-A}^2 + o(1).$$

Applying 4.1 to 9.4, we obtain the following.

$$(9.5) \quad B\left(\frac{1}{2}(p_{1Is} + p_{os})\right) - \int_{|z|=r} \frac{1}{2}(p_{1Is} + p_{os})d\left(\frac{1}{2}(p_{1Is} + p_{os})\right)^* \\ = B(p) - \int_{|z|=r} p dp^* + o(1).$$

Letting r tend to 0, we obtain the desired result.

Theorem 9 was first proved by Sario [9] by another method. The proof presented here is considerably shorter thus showing the power of the l -kernel concept.

We also have the following extremal property of the k -kernel of the space Γ_{he} .

THEOREM 10. *In the class of all harmonic functions u , the expression*

$$(9.6) \quad \|du\|^2 - 4\pi \frac{\partial u(\zeta)}{\partial x}$$

is minimized by the function $p_{os} - p_{1Is}$.

Proof. By the reproducing property of the k -kernel, it follows that

$$(9.7) \quad \|du\|^2 - 4\pi \frac{\partial u(\zeta)}{\partial x} \\ = \|du - (dp_{os} - dp_{1Is})\|^2 - \|dp_{os} - dp_{1Is}\|^2.$$

However

$$(9.8) \quad \|du - (dp_{os} - dp_{1Is})\|^2 - \|dp_{os} - dp_{1Is}\|^2 \\ \leq -\|dp_{os} - dp_{1Is}\|^2 = \|dp_{os} - dp_{1Is}\|^2 - 4 \frac{\partial(p_{os} - p_{1Is})(\zeta)}{\partial x}.$$

Actually in the course of the proof we have shown more, namely that the value of the minimum is $-\|dp_{os} - dp_{1Is}\|^2$. Again we see that the proof is very easy once the reproducing property of

$$-\frac{1}{2\pi} (dp_{1Is} - dp_{os})$$

is established. For another proof of this theorem the reader is referred to [9].

10. The k -kernel of the space Γ_{ae} also possesses an extremal property with respect to the class of analytic functions. Let $dk(z, \zeta)$ denote the k -kernel of Γ_{ae} . Then the following result is valid.

THEOREM 11. *The function $4\pi k(z, \zeta)$ minimizes the expression $\|df\|^2 - 4\pi \operatorname{Re} a(f)$ in the class of all analytic functions f on W where $a(f)$ is the coefficient of $z - \zeta$ in the Taylor expansion of f about ζ . Moreover $a(4\pi k(z, \zeta))$ is nonnegative and $2\pi a(4\pi k(z, \zeta)) = D(4\pi k(z, \zeta))$ where in general $D(f)$ denotes the Dirichlet integral of f .*

The proof follows from the reproducing property of the k -kernel and is similar to that of Theorem 10. The existences of the k -kernel of Γ_{ae} follows from the existence of the k -kernel of Γ_a and the orthogonal decomposition $\Gamma_a = \Gamma_{ae} + \Gamma_{as}$ where Γ_{as} denotes the space of all square integrable analytic Schottky differentials.

If we call $a(4\pi k(z, \zeta))$ the span, then by Theorem 11, we have the following result.

THEOREM 12. *An arbitrary open Riemann surface is of class 0_{AD} if and only if the span vanishes for all choices of ζ .*

The span $a(4\pi k(z, \zeta))$ is a generalization of the well known Schiffer span which was defined by him for planar surfaces.

BIBLIOGRAPHY

1. L. Ahlfors and L. Sario, *Riemann Surfaces*, Princeton University Press, 1960.
2. N. Aronszajn, *Theory of reproducing kernels*, Trans. Amer. Math. Soc. **68** (1950), 337-404.
3. S. Bergman, *The kernel function and conformal mapping*, Math. Surveys No. V, Amer. Math. Soc., 1950.

4. ———, *Ueber die Kernfunktion eines Bereiches und ihr Verhalten am Rande*, J. Reine Angew. Math. **169** (1930), 1-42.
5. R. Courant, *Dirichlet's Principle*, Academic Press Inc., 1950.
6. B. Rodin, *Reproducing formulas on Riemann Surfaces*, Technical Report No. 26, Contract No. DA-04-495-ORD-1959, OOR Project No. 1517.
7. M. Schiffer, *The kernel function of an orthonormal system*, Duke Math. J. **13** (1946), 529-540.
8. M. Schiffer and D. Spencer, *Functionals of Finite Riemann Surfaces*, Princeton University Press, 1954.
9. G. Weill, *Some extremal properties of linear combinations of kernels on Riemann surfaces*, Pacific J. Math. **12** (1962), 1459-1465.

Received April 10, 1965. This paper is part of the author's doctoral thesis, and the author is indebted to Professor Leo Sario for his guidance in its preparation.

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

J. P. JANS

University of Washington
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$ 8.00; single issues, \$ 3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$ 4.00 per volume; single issues \$ 1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

S. J. Bernau, <i>The spectral theorem for unbounded normal operators</i>	391
Lu-san Chen, <i>Asymptotic behavior of solutions of parabolic equations of higher order</i>	407
Lawrence William Conlon, <i>An application of the Bott suspension map to the topology of EIV</i>	411
Neal Eugene Foland and John M. Marr, <i>Sets with zero-dimensional kernels</i>	429
Stanley Phillip Franklin and R. H. Sorgenfrey, <i>Closed and image-closed relations</i>	433
William Jesse Gray, <i>A note on topological transformation groups with a fixed end point</i>	441
Myron Goldstein, <i>K- and L-kernels on an arbitrary Riemann surface</i>	449
George Joseph Kertz and Francis Regan, <i>The exponential analogue of a generalized Weierstrass series</i>	461
Walter Leighton, <i>On Liapunov functions with a single critical point</i>	467
Bernard Werner Levinger and Richard Steven Varga, <i>On a problem of O. Taussky</i>	473
Lowell Duane Loveland, <i>Tame subsets of spheres in E^3</i>	489
Erik Andrew Schreiner, <i>Modular pairs in orthomodular lattices</i>	519
K. N. Srivastava, <i>On dual series relations involving Laguerre polynomials</i>	529
Arthur Steger, <i>Diagonability of idempotent matrices</i>	535
Walter Strauss, <i>On continuity of functions with values in various Banach spaces</i>	543
Robert Vermes, <i>On the zeros of a linear combination of polynomials</i>	553
Elliot Carl Weinberg, <i>On the scarcity of lattice-ordered matrix rings</i>	561
Harold Widom, <i>Toeplitz operators on H_p</i>	573
Neal Zierler, <i>On the lattice of closed subspaces of Hilbert space</i>	583
Irving Leonard Glicksberg, <i>Correction to: "Maximal algebras and a theorem of Radó"</i>	587
John Spurgeon Bradley, <i>Correction to: "Adjoint quasi-differential operators of Euler type"</i>	587
William Branham Jones, <i>Erratum: "Duality and types of completeness in locally convex spaces"</i>	588
Stanley P. Gudder, <i>Erratum: "Uniqueness and existence properties of bounded observables"</i>	588