

# Pacific Journal of Mathematics

## **A TRANSPLANTATION THEOREM FOR JACOBI COEFFICIENTS**

RICHARD ALLEN ASKEY

## A TRANSPLANTATION THEOREM FOR JACOBI COEFFICIENTS

RICHARD ASKEY

Let  $f(\theta)$  be integrable on  $(0, \pi)$  and define

$$\alpha_n^{\alpha, \beta} = t_n^{\alpha, \beta} \int_0^\pi f(\theta) P_n^{(\alpha, \beta)}(\cos \theta) \left(\sin \frac{\theta}{2}\right)^{\alpha+(1/2)} \left(\cos \frac{\theta}{2}\right)^{\beta+(1/2)} d\theta$$

where  $P_n^{(\alpha, \beta)}(x)$  is the Jacobi polynomial of degree  $n$ , order  $(\alpha, \beta)$  and

$$[t_n^{\alpha, \beta}]^2 = \frac{(2n + \alpha + \beta + 1)\Gamma(n + 1)\Gamma(n + \alpha + \beta + 1)}{\Gamma(n + \alpha + 1)\Gamma(n + \beta + 1)}.$$

Then if  $\alpha, \beta, \gamma, \delta \geq -1/2$  we have

$$\sum_{n=0}^{\infty} |\alpha_n^{(\gamma, \delta)}|^p (n + 1)^\sigma \leq A \sum_{n=0}^{\infty} |\alpha_n^{(\alpha, \beta)}|^p (n + 1)^\sigma$$

for  $1 < p < \infty$ ,  $-1 < \sigma < p - 1$  whenever the right hand side is finite.

From this result any norm inequality for Fourier coefficients can be transplanted to give a corresponding norm inequality for Fourier-Jacobi coefficients.

Let  $P_n^{(\alpha, \beta)}(x)$  be defined by  $(-1)^n 2^n n! (1-x)^\alpha (1+x)^\beta P_n^{(\alpha, \beta)}(x) = (d/dx)^n \{(1-x)^{n+\alpha} (1+x)^{n+\beta}\}$ ,  $\alpha, \beta > -1$ . The functions  $P_n^{(\alpha, \beta)}(\cos \theta)$  are orthogonal on  $(0, \pi)$  with respect to the measure

$$\left(\sin \frac{\theta}{2}\right)^{2\alpha+1} \left(\cos \frac{\theta}{2}\right)^{2\beta+1} d\theta$$

and

$$(1) \quad \int_0^\pi [P_n^{(\alpha, \beta)}(\cos \theta)]^2 \left(\sin \frac{\theta}{2}\right)^{2\alpha+1} \left(\cos \frac{\theta}{2}\right)^{2\beta+1} d\theta = \frac{\Gamma(n + \alpha + 1)\Gamma(n + \beta + 1)}{(2n + \alpha + \beta + 1)\Gamma(n + \alpha + \beta + 1)\Gamma(n + 1)} = [t_n^{\alpha, \beta}]^{-2}.$$

Observe that  $t_n^{\alpha, \beta} = An^{1/2} + O(n^{-1/2})$  where  $A$  is a constant whose numerical value is of no interest to us. For simplicity we set  $R_n^{\alpha, \beta}(\theta) = t_n^{\alpha, \beta} P_n^{(\alpha, \beta)}(\cos \theta) [\sin(\theta/2)]^{\alpha+(1/2)} [\cos(\theta/2)]^{\beta+(1/2)}$ . The functions  $\{R_n^{\alpha, \beta}(\theta)\}_{n=0}^{\infty}$  form a complete orthonormal sequence of functions on  $(0, \pi)$ . Also  $R_n^{-1/2, -1/2}(\theta) = A \cos n\theta$  and  $R_n^{1/2, 1/2}(\theta) = A \sin(n+1)\theta$ .

If  $f(\theta) \in L^1(0, \pi)$  we define its Fourier-Jacobi coefficients by

$$(2) \quad \alpha_n^{\alpha, \beta} = \int_0^\pi f(\theta) R_n^{\alpha, \beta}(\theta) d\theta.$$

We define  $l^{p, \sigma}$  to be the space of sequences  $\{\alpha_n\}$  such that  $\|\alpha_n\|_{p, \sigma} =$

$\left[ \sum_{n=0}^{\infty} |a_n|^p (n+1)^\sigma \right]^{1/p}$  is finite. Our main theorem follows.

**THEOREM 1.** *Let  $\alpha, \beta, \gamma, \delta \geq -1/2$  and  $f(\theta) \in L^1(0, \pi)$ . Let  $a_n^{\alpha, \beta}$  and  $a_n^{\gamma, \delta}$  be defined by (2). Then if  $1 < p < \infty$ ,  $-1 < \sigma < p - 1$  and if either  $\|a_n^{\alpha, \beta}\|_{p, \sigma}$  or  $\|a_n^{\gamma, \delta}\|_{p, \sigma}$  is finite so is the other and*

$$(3) \quad A \leq \|a_n^{\alpha, \beta}\|_{p, \sigma} / \|a_n^{\gamma, \delta}\|_{p, \sigma} \leq A$$

where  $A$  is independent of  $f$  and thus of  $a_n^{\alpha, \beta}$  and  $a_n^{\gamma, \delta}$ .

For  $\alpha = \beta$ ,  $\gamma = \delta$  this theorem was proved in [1]. The last section of [1] gives two applications of this theorem. They can be carried over word for word to Jacobi coefficients. If all of the formulas for ultraspherical polynomials that were used in [1] were known for Jacobi polynomials, the proof of Theorem 1 could be exactly the same as the proof of the special case of it in [1]. While it is undoubtedly true that the relevant facts stated in [1] do generalize they are at present unknown. An example of such a fact is the following. Consider  $P_n^{(\alpha, \beta)}(x)P_m^{(\alpha, \beta)}(x)$ . This is a polynomial of degree  $n + m$  and so

$$P_n^{(\alpha, \beta)}(x)P_m^{(\alpha, \beta)}(x) = \sum_{k=0}^{n+m} \alpha_k P_k^{(\alpha, \beta)}(x).$$

If  $\alpha \geq \beta$  the conjecture is that  $\alpha_k \geq 0$ . This is true for  $\alpha = \beta$  and was used in [1]. The limiting result  $\alpha \rightarrow \infty$  is also true and is stated in [4] as a result for Laguerre polynomials. For  $\alpha = \beta + 1$  it was proven in [6].

2. In this section we give various results that we need to prove Theorem 1.

For  $0 < \theta < \pi/2$ ,  $\alpha \geq -1/2$ , we have the following two inequalities

$$(4) \quad |P_n^{\alpha, \beta}(\cos \theta)| = O(n^\alpha),$$

$$(5) \quad |R_n^{\alpha, \beta}(\theta)| = \left| t_n^{\alpha, \beta} \left( \sin \frac{\theta}{2} \right)^{\alpha + (1/2)} \left( \cos \frac{\theta}{2} \right)^{\beta + (1/2)} P_n^{\alpha, \beta}(\cos \theta) \right| \leq A.$$

See [7, (7.32.6)]. In (5) the power of  $\cos \theta/2$  can be changed at will since  $\cos \theta/2$  is bounded away from zero for  $0 \leq \theta \leq \pi/2$ .

$$(6) \quad \frac{d}{dx} P_n^{\alpha, \beta}(x) = \frac{1}{2} (n + \alpha + \beta + 1) P_{n-1}^{(\alpha+1, \beta+1)}(x).$$

See [7, (4.21.7)].

The asymptotic formula we need is an easy consequence of two known results which we now state.

If  $\alpha > -1$ ,  $\beta$  real and  $0 < \theta \leq \pi - \varepsilon$ ,  $\varepsilon > 0$ , then

$$\begin{aligned}
 (7) \quad & \left(\sin \frac{\theta}{2}\right)^\alpha \left(\cos \frac{\theta}{2}\right)^\beta P_n^{(\alpha, \beta)}(\cos \theta) \\
 & = N^{-\alpha} \frac{\Gamma(n + \alpha + 1)}{\Gamma(n + 1)} (\theta/\sin \theta)^{1/2} J_\alpha(N\theta) + R_n(\theta)
 \end{aligned}$$

where  $N = n + (\alpha + \beta + 1)/2$  and

$$R_n(\theta) = \begin{cases} \theta^{1/2} O(n^{-3/2}) & n^{-1} \leq \theta \leq \pi - \varepsilon \\ \theta^{\alpha+2} O(n^\alpha) & 0 < \theta < n^{-1}. \end{cases}$$

$J_\alpha(x)$  is the Bessel function of the first kind of order  $\alpha$ . See [7, (8. 21. 17)]. We also need a known asymptotic formula for  $J_\alpha(x)$ .

$$\begin{aligned}
 (8) \quad x^{1/2} J_\alpha(x) & = A \cos(x - \alpha\pi/2 - \pi/4)[1 + O(x^{-2})] \\
 & + A \sin(x - \alpha\pi/2 - \pi/4)[Ax^{-1} + O(x^{-3})], \quad x \rightarrow \infty.
 \end{aligned}$$

See [7, (1. 71. 8)]. Combining (7), (8) and the asymptotic formula for  $t_n^{\alpha, \beta}$  we get

$$\begin{aligned}
 (9) \quad R_n^{\alpha, \beta}(\theta) & = A \cos(N\theta - \alpha\pi/2 - \pi/4) + A \sin(N\theta - \alpha\pi/2 - \pi/4)/(N\theta) \\
 & + O(N^{-1}) + O(N^{-2}\theta^{-2}), \quad 0 < c/n \leq \theta \leq \pi/2.
 \end{aligned}$$

Finally we need a simple estimate for an integral.

$$(10) \quad \int_N^\infty \frac{\cos y}{y} dy = O(N^{-1}), \quad N \rightarrow \infty.$$

This follows on integrating by parts.

3. We assume that  $f(\theta)$  is smooth enough, say  $C^2$  and vanishing near 0 and  $\pi$ , so that the series  $\sum a_n R_n^{\alpha, \beta}(\theta)$  converges uniformly on  $[0, \pi]$ . These conditions are sufficient for  $a_n = O(n^{-2})$ , integrate by parts twice, and  $|R_n^{\alpha, \beta}(\theta)| \leq A$ . We remove this condition after the following argument.

$$\alpha_n^{\gamma, \delta} = \int_0^\pi f(\theta) R_n^{\gamma, \delta}(\theta) d\theta = \sum_{k=0}^\infty a_k^{\alpha, \beta} \int_0^\pi R_k^{\alpha, \beta}(\theta) R_n^{\gamma, \delta}(\theta) d\theta = \sum_{k=0}^\infty a_k^{\alpha, \beta} R(k, n).$$

Since  $P_n^{(\alpha, \beta)}(-x) = (-1)^n P_n^{(\beta, \alpha)}(x)$ , [7, (4. 1. 3)] it is sufficient to estimate  $S(k, n) = \int_0^{\pi/2} R_k^{\alpha, \beta}(\theta) R_n^{\gamma, \delta}(\theta) d\theta$ . Also because we have made no assumptions about the relationships among  $\alpha, \beta, \gamma, \delta$  it is sufficient to consider the case  $k \geq n$ . We do this in two stages,  $n \leq k \leq 2n$  and  $k \geq 2n$ . For  $n \leq k \leq 2n$  the method is the same as in [1]. We repeat it here for convenience and because the other estimate is handled by a refinement of this argument.

$$S(k, n) = \int_0^{\pi/2} = \int_0^{1/k} + \int_{1/k}^{\pi/2} .$$

The first integral is  $O(k^{-1})$  since  $R_k^{\alpha, \beta}(\theta) = O(1)$ , see (5). In the second integral we use (9),

$$R_n^{\alpha, \beta}(\theta) = A \cos\left(N\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) + A \sin\left(N\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) / N\theta + O(N^{-1}) + O(N^{-2}\theta^{-2})$$

to get

$$S(k, n) = A \int_{1/k}^{\pi/2} \cos\left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) \cos\left(N\theta - \frac{\gamma\pi}{2} - \frac{\pi}{4}\right) d\theta + \frac{A}{K} \int_{1/k}^{\pi/2} \sin\left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) \cos\left(N\theta - \frac{\gamma\pi}{2} - \frac{\pi}{4}\right) \frac{d\theta}{\theta} + \frac{A}{N} \int_{1/k}^{\pi/2} \cos\left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) \sin\left(N\theta - \frac{\gamma\pi}{2} - \frac{\pi}{4}\right) \frac{d\theta}{\theta} + O(K^{-1}) .$$

The first integral is  $A/(K - N) + O(K^{-1})$ , the second is

$$\frac{A}{K} \log \frac{K}{K - N} + O(K^{-1}) ,$$

the third is  $(A/N) \log N/(K - N) + O(K^{-1})$  by a simple computation. The details are in [1]. The one time this argument breaks down is when  $K = N$ . In this case  $S(k, n) = O(1)$  by (5).

Now we consider the case  $k > 2n$ . This time we need not be so careful, i.e., all our estimates may be  $O$  estimates, but the details turn out to be harder than in the above case. This probably isn't necessary but we have not found a simple proof of the following estimates. There is one case,  $\gamma = \alpha + 2$ ,  $\beta = \delta$ , in which it is possible to give easy estimates as we will show later. But this is a very singular case.

As before  $S(k, n) = \int_0^{1/k} + \int_{1/k}^{\pi/2}$ . The first integral is  $O(K^{-1})$  by (5). Next we show that in the second integral we may replace  $R_K(\theta)$  by  $\cos\{K\theta - (\alpha\pi/2) - (\pi/4)\}$ . Using (9) we see it is sufficient to show that

$$\frac{n^{1/2}}{K} \int_{1/k}^{\pi/2} \sin\left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) P_n^{(\gamma, \delta)}(\cos \theta) \left(\sin \frac{\theta}{2}\right)^{\gamma+(1/2)} \left(\cos \frac{\theta}{2}\right)^{\delta+(1/2)} \theta^{-1} d\theta = O(K^{-1}) .$$

Integrating by parts and estimating we have

$$\begin{aligned} & \frac{n^{1/2}}{K^2} \int_{1/k}^{\pi/2} P_n^{(\gamma, \delta)}(\cos \theta) \left(\sin \frac{\theta}{2}\right)^{\gamma+(1/2)} \left(\cos \frac{\theta}{2}\right)^{\delta+(1/2)} \theta^{-1} d \cos \left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) \\ &= O\left(\frac{n^{1/2}}{K^2} P_n^{(\gamma, \delta)}(\cos \theta) \theta^{\gamma-(1/2)}\right) \Big|_{1/k}^{\pi/2} + O\left[\frac{n^{3/2}}{K^2} \int_{1/k}^{\pi/2} \cos \left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) \right. \\ &\quad \cdot P_{n-1}^{(\gamma+1, \delta+1)}(\cos \theta) \left(\sin \frac{\theta}{2}\right)^{\gamma+(3/2)} \left(\cos \frac{\theta}{2}\right)^{\delta+(3/2)} \theta^{-1} d\theta \Big] \\ &\quad + O\left(\frac{n^{1/2}}{K^2} \int_{1/k}^{\pi/2} \cos \left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) P_n^{(\gamma, \delta)}(\cos \theta) \left(\sin \frac{\theta}{2}\right)^{\gamma-(1/2)} \right. \\ &\quad \cdot \left.\left(\cos \frac{\theta}{2}\right)^{\delta+(3/2)} \theta^{-1} d\theta\right) \\ &\quad + \text{similar terms.} \end{aligned}$$

The integrated term is  $O(K^{-1})$  by (5). The second integral is

$$O\left(K^{-2} \int_{1/k}^{\pi/2} \theta^{-2} d\theta\right) = O(K^{-1}).$$

by (5). The first integral we write as  $\int_{1/k}^{1/n} + \int_{1/n}^{\pi/2}$ . Using (4) in  $\int_{1/k}^{1/n}$  we have the bound  $(n^{3/2}/K^2) \int_{1/k}^{1/n} n^{\gamma+1} \theta^{\gamma+(1/2)} d\theta = O(K^{-1})$ . In  $\int_{1/n}^{\pi/2}$  we use (9) to get

$$\begin{aligned} & \frac{An}{K^2} \int_{1/n}^{\pi/2} \cos \left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{2}\right) \left[ \cos \left(N\theta - \frac{\gamma\pi}{2} - \frac{\pi}{4}\right) \right. \\ &\quad \left. + \frac{A \sin \left(N\theta - \frac{\gamma\pi}{2} - \frac{\pi}{4}\right)}{N\theta} + O(N^{-1}) + O(N^{-2}\theta^{-2}) \right] \theta^{-1} d\theta \\ &= \frac{An}{K^2} \int_{1/n}^{\pi/2} \frac{\cos(K-N)\theta}{\theta} d\theta + O(K^{-1}) \\ &\quad + \text{terms similar to the first.} \end{aligned}$$

Changing variables we get  $\int_{(k-n)/n}^{(\pi/2)(k-n)} (\cos y/y) dy$ . Since  $k > 2n$  we have  $(k-n)/n \geq 1$ . Using (10) we get an estimate for the first term of the form  $(An/K^2)\{n/(K-n)\} = O(K^{-1})$ . Thus it is sufficient to consider

$$t_n^{\gamma, \delta} \int_{1/k}^{\pi/2} \cos \left(K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4}\right) P_n^{(\gamma, \delta)}(\cos \theta) \left(\sin \frac{\theta}{2}\right)^{\gamma+(1/2)} \left(\cos \frac{\theta}{2}\right)^{\delta+(1/2)} d\theta.$$

As above we break this integral into two parts  $\int_{1/k}^{1/n} + \int_{1/n}^{\pi/2}$ . We treat the first of these first. Integrating by parts and estimating we have

$$\int_{1/k}^{1/n} = O(K^{-1}) + \frac{t_n^{\gamma, \delta}}{K} \int_{1/k}^{1/n} \sin \left( K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4} \right) \frac{d}{d\theta} \left[ P_n^{(\gamma, \delta)}(\cos \theta) \cdot \left( \sin \frac{\theta}{2} \right)^{\gamma+(1/2)} \left( \cos \frac{\theta}{2} \right)^{\delta+(1/2)} \right] d\theta .$$

Using (6) and (4) we see that this integral is  $O(K^{-1})$ .

In our one remaining integral we may use (9). However to get an estimate of the form  $O(K^{-1})$  we must first integrate by parts. Then we get

$$\begin{aligned} & \frac{t_n^{\gamma, \delta}}{2K} \left( \gamma + \frac{1}{2} \right) \int_{1/n}^{\pi/2} \sin \left( K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4} \right) P_n^{(\gamma, \delta)}(\cos \theta) \\ & \quad \cdot \left( \sin \frac{\theta}{2} \right)^{\gamma-(1/2)} \left( \cos \frac{\theta}{2} \right)^{\delta+(3/2)} d\theta \\ & + \frac{t_n^{\gamma, \delta}}{K} [n + \gamma + \delta + 1] \int_{1/n}^{\pi/2} \sin \left( K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4} \right) P_{n-1}^{(\gamma+1, \delta+1)}(\cos \theta) \\ & \quad \cdot \left( \sin \frac{\theta}{2} \right)^{\gamma+(3/2)} \left( \cos \frac{\theta}{2} \right)^{\delta+(3/2)} d\theta + O(K^{-1}) \text{ (by (5)).} \end{aligned}$$

For the first integral we have the estimate

$$\begin{aligned} & O \left[ \frac{1}{K} \int_{1/n}^{\pi/2} \sin \left( K\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4} \right) \left[ \cos \left( N\theta - \frac{\gamma\pi}{2} - \frac{\pi}{4} \right) \right. \right. \\ & \quad \left. \left. + A(N\theta)^{-1} \sin \left( N\theta - \frac{\gamma\pi}{2} - \frac{\pi}{4} \right) + O(N^{-1}) + O(\theta^{-2}N^{-2}) \right] \left( \sin \frac{\theta}{2} \right)^{-1} d\theta \right] \\ & = O \left[ \frac{1}{K} \int_{1/n}^{\pi/2} \frac{\sin(K-N)\theta}{\theta} d\theta \right] + O(K^{-1}) \end{aligned}$$

by (9) and the fact that  $(1/\sin \theta) - (1/\theta)$  is bounded. As above this leads to the estimate  $O(n/K^2) + O(K^{-1}) = O(K^{-1})$ .

A simple computation shows that  $t_n^{\gamma, \delta} = At_{n-1}^{\gamma+1, \delta+1} [1 + O(n^{-1})]$  so the second integral may also be estimated by using (9). The estimate is

$$\begin{aligned} & O \left( \frac{n}{K} \int_{1/n}^{\pi/2} \sin \left( K\theta - \frac{\gamma\pi}{2} - \frac{\pi}{4} \right) \left[ \cos \left( N\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4} \right) \right. \right. \\ & \quad \left. \left. + \frac{A \sin \left( N\theta - \frac{\alpha\pi}{2} - \frac{\pi}{4} \right)}{N\theta} + O(N^{-1}) + O(N^{-2}\theta^{-2}) \right] d\theta \right) \\ & = O \left( \frac{n}{K} \int_{1/n}^{\pi/2} \sin(K-N)\theta d\theta \right) + O \left( \frac{n}{K} \int_{1/n}^{\pi/2} \frac{\cos(K-N)\theta}{N\theta} d\theta \right) \\ & \quad + \text{similar terms} + O(K^{-1}) \\ & = O \left( \frac{n}{K(K-N)} \right) + O(K^{-1}) = O(K^{-1}) \end{aligned}$$

by the same type of arguments that have been used often above. Combining all of the above estimates we see that

$$\begin{aligned}
 \alpha_n^{\gamma, \delta} &= O\left[\frac{1}{n} \sum_{k=0}^{[n/2]} |a_k^{\alpha, \beta}|\right] + A \sum_{k=[n/2]}^{2n} \frac{\alpha_k^{\alpha, \beta}}{K - N} \\
 (11) \quad &+ \frac{A}{n} \sum_{k=[n/2]}^{2n} \alpha_k^{\alpha, \beta} \log \left| \frac{N}{K - N} \right| + A \sum_{k=[n/2]}^{2n} \frac{\alpha_k^{\alpha, \beta}}{K} \log \left| \frac{K}{K - N} \right| \\
 &+ \left[ \sum_{k=2n}^{\infty} \frac{|a_k^{\alpha, \beta}|}{K} \right].
 \end{aligned}$$

As in [1] all of the terms on the right are bounded operators in  $l^{p, \sigma}$ ,  $1 < p < \infty$ ,  $-1 < \sigma < p - 1$ . Thus  $\| \alpha_n^{\gamma, \delta} \|_{p, \sigma} \leq A \| \alpha_n^{\alpha, \beta} \|_{p, \sigma}$  which is (3).

Let  $g(\theta) \in C^2$  and vanish near 0 and  $\pi$  and let  $f(\theta) \in L^1(0, \pi)$ . Define their Fourier-Jacobi coefficients,  $b_n^{\alpha, \beta}$  and  $a_n^{\alpha, \beta}$  respectively, by (2). Then

$$\sum_{n=0}^{\infty} b_n^{\gamma, \delta} \alpha_n^{\gamma, \delta} = \int_0^{\pi} f(\theta) g(\theta) d\theta = \sum_{n=0}^{\infty} b_n^{\alpha, \beta} \alpha_n^{\alpha, \beta}$$

and thus {with  $(1/p) + (1/q) = 1$ }

$$\begin{aligned}
 \| \alpha_n^{\gamma, \delta} \|_{p, \sigma} &= \left[ \sum_{n=0}^{\infty} | \alpha_n^{\gamma, \delta} |^p (n + 1)^{\sigma} \right]^{1/p} = \sup \sum \alpha_n^{\gamma, \delta} b_n^{\delta} = \sup \sum \alpha_n^{\alpha, \beta} b_n^{\alpha, \beta} \\
 &\leq \sup \| \alpha_n^{\alpha, \beta} \|_{p, \sigma} \| b_n^{\alpha, \beta} \|_{q, -q\sigma/p} \leq A \| \alpha_n^{\alpha, \beta} \|_{p, \sigma}
 \end{aligned}$$

by (3). Here the sup is taken over the sequences  $b_n^{\gamma, \delta}$  with

$$\sum | b_n^{\gamma, \delta} |^q (n + 1)^{\sigma(1-q)} \leq 1.$$

This completes the proof of Theorem 1.

There is a simple substitute in  $l^1$  which follows easily from (11).

**THEOREM 2.** *Let  $\alpha, \beta, \gamma, \delta$  be as in Theorem 1 and assume  $\sum | \alpha_n^{\alpha, \beta} | \log(n + 2) < \infty$ . Then  $\sum | \alpha_n^{\gamma, \delta} | < \infty$  where*

$$\alpha_n^{\gamma, \delta} = \int_0^{\pi} f(\theta) R_n^{\gamma, \delta}(\theta) d\theta$$

$$\text{and } f(\theta) = \sum \alpha_n^{\alpha, \beta} R_n^{\alpha, \beta}(\theta).$$

The inequalities that are needed to prove Theorem 2 from (11) are in [3], where this result was proven for  $\alpha = \beta = -1/2$ ,  $\gamma = \delta = 1/2$ . To be pedantic here we must be careful for unless  $\alpha = -1/2$ ,  $R_n^{(\alpha, \beta)}(0) = 0$  and so  $f(\theta) = \sum \alpha_n^{\alpha, \beta} R_n^{\alpha, \beta}(\theta)$  must vanish at  $\theta = 0$ . Thus if  $\alpha = -1/2$  we must assume  $f(0) = 0$  and similarly for  $\beta = -1/2$ ,  $\theta = \pi$ . Theorem 2 is the one place where the above proof is an improvement over the proof in [1] (even in the case  $\alpha = \beta$ ,  $\gamma = \delta$ ) for using the proof in [1] we



must add higher powers to the logarithm if  $\alpha$  and  $\gamma$  are far apart. Even this can be done away with if we use Theorem 4 which follows in the next section. However this problem would again arise if one tried to prove Theorem 2 for Jacobi (not ultraspherical) coefficients by the method in [1] by say holding  $\beta$  fixed first and then varying it with fixed  $\alpha$ .

4. We conclude this paper with two simple theorems that hold in  $l^1$ . Since the details are easier we first give a theorem for Laguerre coefficients and then finally we give the corresponding theorem for Jacobi coefficients.

The Laguerre polynomial  $L_n^\alpha(x)$  is defined by

$$\sum_{n=0}^\infty L_n^\alpha(x)r^n = (1 - r)^{-\alpha-1} \exp\left(\frac{-xr}{1-r}\right).$$

These functions satisfy

$$(12) \quad \int_0^\infty L_n^\alpha(x)L_m^\alpha(x)x^\alpha e^{-x}dx = \frac{\Gamma(n + \alpha + 1)}{\Gamma(n + 1)} \delta_{n,m}, \quad \alpha > -1$$

Let  $f(x) \in L^1(0, \infty)$  and define its Fourier-Laguerre coefficient by

$$(13) \quad a_n^\alpha = t_n^\alpha \int_0^\infty f(x)L_n^\alpha(x)x^{\alpha/2}e^{-x/2}dx.$$

where

$$t_n^\alpha = \left[ \frac{\Gamma(n + 1)}{\Gamma(n + \alpha + 1)} \right]^{1/2}.$$

We need one more fact about these functions.

$$(14) \quad L_n^{\alpha+1}(x) = \sum_{j=0}^n L_j^\alpha(x).$$

From this we see that

$$(15) \quad L_n^\alpha(x) = L_n^{\alpha+1}(x) - L_{n-1}^{\alpha+1}(x).$$

**THEOREM 3.** *Let  $f(x) \in L^1(0, \infty)$  and define  $a_n^\alpha, a_n^{\alpha+2}$  by (13). Then if  $\alpha > 0$ ,*

$$(16) \quad A \leq [\sum |a_n^\alpha|^p]^{1/p} / [\sum |a_n^{\alpha+2}|^p]^{1/p} \leq A$$

for  $1 \leq p < \infty$ . If  $-1 < \alpha < 0$  then (16) holds for  $2/(2 + \alpha) < p < -2/\alpha$ .

Using (12) and (13) we see that

$$\begin{aligned} a_n^{\alpha+2} &= t_n^{\alpha+2} \int_0^\infty f(x) L_n^{\alpha+2}(x) x^{(\alpha/2)+1} e^{-x/2} dx \\ &= t_n^{\alpha+2} \sum_{k=0}^\infty t_k^\alpha a_k^\alpha \int_0^\infty L_k^\alpha(x) L_n^{\alpha+2}(x) x^{\alpha+1} e^{-x} dx \\ &= t_n^{\alpha+2} \sum_{k=0}^{n+1} t_k^\alpha a_k^\alpha \int_0^\infty L_k^\alpha(x) L_n^{\alpha+2}(x) x^{\alpha+1} e^{-x} dx . \end{aligned}$$

Then using (14) and (15) we have

$$\begin{aligned} a_n^{\alpha+2} &= t_n^{\alpha+2} t_0^\alpha a_0^\alpha \int_0^\infty L_0^\alpha(x) \sum_{j=0}^n L_j^{\alpha+1}(x) x^{\alpha+1} e^{-x} dx \\ &\quad + t_n^{\alpha+2} \sum_{k=1}^n t_k^\alpha a_k^\alpha \int_0^\infty [L_k^{\alpha+1}(x) - L_{k-1}^{\alpha+1}(x)] \left[ \sum_{j=0}^n L_j^{\alpha+1}(x) \right] x^{\alpha+1} e^{-x} dx \\ &\quad + t_n^{\alpha+2} t_{n+1}^\alpha a_{n+1}^\alpha \int_0^\infty -L_n^{\alpha+1}(x) L_n^{\alpha+1}(x) x^{\alpha+1} e^{-x} dx \\ &= t_n^{\alpha+2} t_0^\alpha [t_0^{\alpha+1}]^{-2} a_0^\alpha + t_n^{\alpha+2} \sum_{k=1}^n t_k^\alpha a_k^\alpha [(t_k^{\alpha+1})^{-2} - (t_{k-1}^{\alpha+1})^{-2}] \\ &\quad - t_n^{\alpha+2} t_{n+1}^\alpha [t_n^{\alpha+1}]^{-2} a_{n+1}^\alpha . \end{aligned}$$

Thus

$$\begin{aligned} |a_n^{\alpha+2}| &\leq A |a_0^\alpha| n^{-(\alpha/2)-1} + A \sum_{k=1}^n n^{-(\alpha/2)-1} k^{-\alpha/2} k^\alpha |a_k^\alpha| + A |a_{n+1}^\alpha| \\ &\leq A |a_0^\alpha| + A \sum_{k=1}^n |a_k^\alpha| (k/n)^{\alpha/2} n^{-1} + A |a_{n+1}^\alpha| . \end{aligned}$$

Similarly one can show that

$$|a_k^\alpha| \leq A |a_{k-1}^{\alpha+2}| + A \sum_{n=k}^\infty |a_n^{\alpha+2}| (k/n)^{\alpha/2} n^{-1} .$$

Theorem 3 then follows from problem 346 in [5]. Actually there is one application of Theorem 3 and surprisingly it is for  $\alpha$  negative. In a paper which will appear, Wainger and I prove the following theorem.

**THEOREM A.** *Let  $\alpha \geq 0$ ,  $f \in L^1(0, \infty)$  and define*

$$a_n^\alpha = \int_0^\infty f(x) t_n^\alpha L_n^\alpha(x) x^{\alpha/2} e^{-x/2} dx .$$

*Let  $t(x)$  be a bounded function which is of bounded variation on  $(0, \infty)$ , with  $\int_0^\infty |dt(x)| \leq C$ . Define*

$$T a_n^\alpha = \int_0^\infty t(x) f(x) t_n^\alpha L_n^\alpha(x) x^{\alpha/2} e^{-x/2} dx .$$

*Then this operator is bounded in  $l^p$ ,  $4/3 < p < 4$ , i.e.*

$$[\sum |T a_n^\alpha|^p]^{1/p} \leq AC[\sum |a_n^\alpha|^p]^{1/p}$$

where  $A$  is independent of  $f(x)$  and of  $t(x)$ .

We used asymptotic estimates of Erdélyi which have only been proven for  $\alpha \geq 0$ . See [2] where the dual result is proven. We can now extend this result to  $\alpha \geq -1/2$  by using Theorem 3. Similar applications are given in [1] and we will not repeat the details here.

It would be interesting to extend Theorem 3 to get a theorem which corresponds to Theorem 1. The estimates of Erdélyi are probably not sufficient to allow one to prove this but they can probably be extended to give two terms plus an error and this might suffice.

The proof of the following theorem for Jacobi coefficients is exactly the same as the proof of Theorem 3 but the coefficients are not as simple so it looks more complicated.

**THEOREM 4.** *Let  $f(\theta) \in L^1(0, \pi)$  and define  $a_n^{\alpha, \beta}$ ,  $\alpha_n^{\alpha+2, \beta}$  by (2). Then if  $\alpha > -1/2$ ,  $\beta > -1$ , and if either  $\sum |a_n^{\alpha, \beta}|$  or  $\sum |\alpha_n^{\alpha+2, \beta}|$  converges so does the other and*

$$0 < A \leq \sum |a_n^{\alpha, \beta}| / \sum |\alpha_n^{\alpha+2, \beta}| \leq A < \infty$$

with  $A$  independent of  $f$ .

$$\begin{aligned} \alpha_n^{\alpha+2, \beta} &= \int_0^\pi f(\theta) R_n^{\alpha+2, \beta}(\theta) d\theta = \sum_{k=0}^\infty a_k^{\alpha, \beta} \int_0^\pi R_n^{\alpha+2, \beta}(\theta) R_k^{\alpha, \beta}(\theta) d\theta \\ &= t_n^{\alpha+2, \beta} \sum_{k=0}^\infty a_k^{\alpha, \beta} t_k^{\alpha, \beta} \int_0^\pi P_n^{(\alpha+2, \beta)}(\cos \theta) P_k^{(\alpha, \beta)}(\cos \theta) \\ &\quad \cdot \left(\sin \frac{\theta}{2}\right)^{2\alpha+3} \left(\cos \frac{\theta}{2}\right)^{2\beta+1} d\theta \\ &= 2^{-\alpha-\beta-2} t_n^{\alpha+2, \beta} \sum_{k=0}^\infty a_k^{\alpha, \beta} t_k^{\alpha, \beta} \int_{-1}^1 P_n^{(\alpha+2, \beta)}(x) P_k^{(\alpha, \beta)}(x) (1-x)^{\alpha+1} (1+x)^\beta dx \\ &= 2^{-\alpha-\beta-2} t_n^{\alpha+2, \beta} \sum_{k=0}^{n+1} a_k^{\alpha, \beta} t_k^{\alpha, \beta} R(k, n). \end{aligned}$$

To estimate  $R(k, n)$  we use the following.

$$(17) \quad P_n^{(\alpha+1, \beta)}(x) = \frac{\Gamma(n + \beta + 1)}{\Gamma(n + \alpha + \beta + 2)} \cdot \sum_{j=0}^n \frac{(2j + \alpha + \beta + 1) \Gamma(j + \alpha + \beta + 1)}{\Gamma(j + \beta + 1)} P_j^{(\alpha, \beta)}(x),$$

$$(18) \quad \begin{aligned} P_n^{(\alpha, \beta)}(x) &= \frac{n + \alpha + \beta + 1}{2n + \alpha + \beta + 1} P_n^{(\alpha+1, \beta)}(x) \\ &\quad - \frac{n + \beta}{2n + \alpha + \beta + 1} P_{n-1}^{(\alpha+1, \beta)}(x). \end{aligned}$$

(18) follows from (17) and (17) is (4.5.3) in [7]. Using (17) and (18) we see that

$$R(k, n) = \sum_{j=0}^n \frac{\Gamma(n + \beta + 1)(2j + \alpha + \beta + 2)\Gamma(j + \alpha + \beta + 2)}{\Gamma(n + \alpha + \beta + 3)\Gamma(j + \beta + 1)(2k + \alpha + \beta + 1)} \cdot \int_{-1}^1 P_j^{(\alpha+1, \beta)}(x) [(k + \alpha + \beta + 1)P_k^{(\alpha+1, \beta)}(x) - (k + \beta)P_{k-1}^{(\alpha+1, \beta)}(x)](1 - x)^{\alpha+1}(1 + x)^\beta dx .$$

For  $1 \leq k \leq n$  we have

$$\begin{aligned} R(k, n) &= \frac{\Gamma(n + \beta + 1)}{\Gamma(n + \alpha + \beta + 3)} \left[ \frac{(2k + \alpha + \beta + 2)\Gamma(k + \alpha + \beta + 2)}{\Gamma(k + \beta + 1)(2k + \alpha + \beta + 1)} \right. \\ &\quad \cdot \frac{(k + \alpha + \beta + 1)}{[t_k^{\alpha+1, \beta}]^2} - \left. \frac{(2k + \alpha + \beta)\Gamma(k + \alpha + \beta + 1)(k + \beta)}{\Gamma(k + \beta)(2k + \alpha + \beta + 1)[t_{k-1}^{\alpha+1, \beta}]^2} \right] \\ &= \frac{\Gamma(n + \beta + 1)}{\Gamma(n + \alpha + \beta + 3)(2k + \alpha + \beta + 1)} \\ &\quad \cdot \left[ \frac{\Gamma(k + \alpha + 2)(k + \alpha + \beta + 1)}{\Gamma(k + 1)} - \frac{\Gamma(k + \alpha + 1)(k + \beta)}{\Gamma(k)} \right] \\ &= \frac{\Gamma(n + \beta + 1)\Gamma(k + \alpha + 1)}{\Gamma(n + \alpha + \beta + 3)(2k + \alpha + \beta + 1)\Gamma(k + 1)} \\ &\quad \cdot [(k + \alpha + 1)(k + \alpha + \beta + 1) - k(k + \beta)] \\ &= O(n^{-\alpha-2}k^\alpha) . \end{aligned}$$

For  $k = 0$ ,  $R(k, n) = O(n^{-\alpha-2})$  follows easily from (17), (18) and (1). For  $k = n + 1$ ,  $R(k, n) = O(n^{-1})$  also follows easily from these same formulas. Thus we have

$$|a_n^{\alpha+2, \beta}| \leq An^{-1} \sum_{k=0}^{n+1} |a_k^{\alpha, \beta}| (k/n)^{\alpha+1/2}$$

and

$$\sum_{n=0}^{\infty} |a_n^{\alpha+2, \beta}| \leq A \sum_{n=0}^{\infty} |a_n^{\alpha, \beta}|$$

follows easily by interchanging the order of summation. The other inequality follows by the same argument.

### REFERENCES

1. R. Askey and S. Wainger, *A transplantation theorem for ultraspherical coefficients*, Pacific J. Math. **16** (1966), 393-405.
2. ———, *Mean convergence of expansions in Laguerre and Hermite series*, Amer. J. Math. **87** (1965), 695-708.
3. R. P. Boas, *Absolute convergence and integrability of trigonometric series*, J. Rat. Mech. Anal. **5** (1956), 621-632.

4. A. Erdélyi, *On some expansions in Laguerre polynomials*, J. Lond. Math. Soc. **13** (1938), 154-156.
5. G. H. Hardy, J. E. Littlewood, and G. Polya, *Inequalities*, Cambridge, 1952.
6. E. Hylleraas, *Linearization of products of Jacobi polynomials*, Math. Scand. **10** (1962), 189-200.
7. G. Szegő, *Orthogonal Polynomials*, Amer. Math. Soc. Coll. Pub. Vol. 23, revised edition (1959), New York.

Received April 1966. Supported in part by NSF grant GP-3483.

THE UNIVERSITY OF WISCONSIN  
MADISON, WISCONSIN

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON  
Stanford University  
Stanford, California

J. P. JANS  
University of Washington  
Seattle, Washington 98105

J. DUGUNDJI  
University of Southern California  
Los Angeles, California 90007

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*

AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$ 8.00; single issues, \$ 3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$ 4.00 per volume; single issues \$ 1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Richard Allen Askey, <i>A transplantation theorem for Jacobi coefficients</i> . . . .	393
Raymond Balbes, <i>Projective and injective distributive lattices</i> . . . . .	405
Raymond Balbes and Alfred Horn, <i>Order sums of distributive lattices</i> . . . . .	421
Donald Charles Benson, <i>Nonconstant locally recurrent functions</i> . . . . .	437
Allen Richard Bernstein, <i>Invariant subspaces of polynomially compact operators on Banach space</i> . . . . .	445
Robert F. Brown, <i>Fixed points and fibre</i> . . . . .	465
David Geoffrey Cantor, <i>On the Stone-Weierstrass approximation theorem for valued fields</i> . . . . .	473
James Walton England, <i>Stability in topological dynamics</i> . . . . .	479
Alessandro Figà-Talamanca and Daniel Rider, <i>A theorem on random Fourier series on noncommutative groups</i> . . . . .	487
Sav Roman Harasymiv, <i>A note of dilations in <math>L^p</math></i> . . . . .	493
J. G. Kalbfleisch, <i>A uniqueness theorem for edge-chromatic graphs</i> . . . . .	503
Richard Paul Kelisky and Theodore Joseph Rivlin, <i>Iterates of Bernstein polynomials</i> . . . . .	511
D. G. Larman, <i>On the union of two starshaped sets</i> . . . . .	521
Henry B. Mann, Josephine Mitchell and Lowell Schoenfeld, <i>Properties of differential forms in <math>n</math> real variables</i> . . . . .	525
John W. Moon and Leo Moser, <i>Generating oriented graphs by means of team comparisons</i> . . . . .	531
Veikko Nevanlinna, <i>A refinement of Selberg's asymptotic equation</i> . . . . .	537
Ulrich Oberst, <i>Relative satellites and derived functors of functors with additive domain</i> . . . . .	541
John Vincent Ryff, <i>On Muirhead's theorem</i> . . . . .	567
Carroll O. Wilde and Klaus G. Witz, <i>Invariant means and the Stone-Čech compactification</i> . . . . .	577