

# Pacific Journal of Mathematics

**NONCONSTANT LOCALLY RECURRENT FUNCTIONS**

DONALD CHARLES BENSON

# NONCONSTANT LOCALLY RECURRENT FUNCTIONS

DONALD C. BENSON

The purpose of this paper is to develop a new method of using the Baire Category Theorem to obtain counterexamples in analysis. The method is used to show that a certain class of nonconstant locally recurrent functions is of second category in a suitable metric space of continuous functions. In § 1 an explicit example is given of a nonconstant locally recurrent function. This example is included because it clarifies the category argument in § 4.

1. A simple example of a nonconstant locally recurrent function.

DEFINITION 1. A real-valued continuous function  $f$  of a real variable is said to be locally recurrent if for any  $x$  in its domain of definition and any neighborhood  $N$  of  $x$ , there exists  $y \neq x$  in  $N$  such that  $f(x) = f(y)$ .

K. A. Bush [2] has given an example of a nonconstant locally recurrent function. The author believes that the example given below is simpler.

A sequence  $\{f_n\}_{n=0}^{\infty}$  of functions on  $[0, 1]$  will be defined. These functions are continuous and piecewise linear. Further,  $f_m$  is linear in any interval of the form  $[n/9^m, (n+1)/9^m]$  where  $n$  and  $m$  are non-negative integers such that  $0 \leq n < 9^m$ . Thus the function  $f_m$  is described completely if we give the values of  $f_m(n/9^m)$ ,  $(0 \leq n \leq 9^m)$ . These functions will be defined inductively. We start with  $f_0(x) = x$ . Now suppose  $f_k$  is defined for some  $k$ . We define  $f_{k+1}$  as follows:


$$(a) \quad f_{k+1}(3m/9^{k+1}) = f_k(3m/9^{k+1}), \quad 0 \leq 3m \leq 9^{k+1}.$$

$$(b) \quad f_{k+1}((3m+1)/9^{k+1}) = f_k((3m+3)/9^{k+1}), \quad 0 \leq 3m+1, 3m+3 \leq 9^{k+1}.$$

$$(c) \quad f_{k+1}((3m+2)/9^{k+1}) = f_k(3m/9^{k+1}), \quad 0 \leq 3m, 3m+2 \leq 9^{k+1}.$$

The figure shows a portion of the graphs of  $f_k$  and  $f_{k+1}$ .

An important feature of these functions is the relation  $f_l(n/9^k) = f_l(n/9^k)$  for  $l \geq k$  and  $0 \leq n \leq 9^k$ . Also notice that on any interval of the form  $[n/9^k, (n+1)/9^k]$ , the values of  $f_m$ ,  $m \geq k$  must lie between  $f_k(n/9^k)$  and  $f_k((n+1)/9^k)$ . It is not hard to see that the  $f_n$  converge uniformly and thus the limit function  $f(x) = \lim_{n \rightarrow \infty} f_n(x)$  is continuous. It is obviously locally recurrent at points of the form  $n/9^k$ . That  $f$  is locally recurrent at any point  $x$  in  $[0, 1]$  follows from an application of the intermediate value theorem for continuous functions. In fact, for any  $k \geq 0$ ,  $x$  must lie in an interval of the form  $I = [3m/9^k,$



FIGURE

$(3m + 3)/9^k]$ , an interval of length  $3/9^k$ . We have  $f((3m + 1)/9^k) = f((3m + 3)/9^k) = m_1$  and  $f((3m + 2)/9^k) = f(3m/9^k) = m_2$ . Also, either  $\sup_{x \in I} f(x) = m_1$  and  $\inf_{x \in I} f(x) = m_2$ , or  $\sup_{x \in I} f(x) = m_2$  and  $\inf_{x \in I} f(x) = m_1$ . The intermediate value theorem shows that every value of the function in  $I$ , except  $m_1$  and  $m_2$ , must occur at least three times in  $I$ . Of course,  $m_1$  and  $m_2$  occur at least twice.

2. A class of complete metric spaces. In this section certain abstract tools will be developed for the purpose of showing the existence of various functions with pathological properties. Using these tools one may show that the set of functions with a certain pathology, for example, the pathology of the function of the previous section, is of the second category in a suitable complete metric space. An early result of this type was obtained by Banach [1] and Mazurkiewicz [3] who showed that the nowhere differentiable functions are of second category in the space of continuous functions with the uniform metric. The space and the metric considered here will be different.

Let  $\mathcal{F}$  be the set of all functions which map a set  $T$  into a set  $S$ . We require that  $T$  is the union of nonempty, disjoint subsets  $T_i$ ,  $i = 1, 2, \dots$ . We provide  $\mathcal{F}$  with a metric  $d$  as follows. If  $f$  and  $g$  belong to  $\mathcal{F}$ , we define

$$d(f, g) = 1/k$$

where  $k$  is the smallest integer such that  $f(x) \neq g(x)$  for some  $x$  in  $T_k$ . If  $f(x) = g(x)$  for all  $x \in T$ , we put  $d(f, g) = 0$ . It is easy to see that  $d$  is a metric.

Let  $\mathcal{G}$  be the set of functions which, for some positive integer  $k$ , maps  $\bigcup_{j=1}^k T_j$  into  $S$ . Further, let us put  $\mathcal{F}^* = \mathcal{F} \cup \mathcal{G}$ .

DEFINITION 2. A subset  $\mathcal{H}$  of  $\mathcal{F}^*$  is said to be *hereditary* if  $f$  belongs to  $\mathcal{H}$  if and only if every restriction of  $f$  which is in  $\mathcal{F}^*$  also belongs to  $\mathcal{H}$ . (A function  $g$  is a restriction of  $f$  if the domain

of definition of  $g$  is a *proper* subset of the domain of definition of  $f$  and if  $f(x)=g(x)$  for any  $x$  in the domain of definition of  $g$ .)

DEFINITION 3. A subset  $\mathcal{A}$  of  $\mathcal{H}(\mathcal{H} \subset \mathcal{F}^*)$  is said to be *absorbing* with respect to  $\mathcal{H}$  if the following two conditions hold.

1. If  $g \in \mathcal{A}$  and  $g$  is a restriction of  $f \in \mathcal{H}$ , then  $f \in \mathcal{A}$ .
2. For any  $g \in \mathcal{G} \cap \mathcal{H}$ , there exists  $f$  in  $\mathcal{G} \cap \mathcal{A}$  such that  $g$  is a restriction of  $f$ .

THEOREM 1. Let  $\mathcal{H}$  be a hereditary subset of  $\mathcal{F}^*$ . Then  $\mathcal{H} \cap \mathcal{F}$  is complete with respect to the metric  $d$ .

*Proof.* Let  $\{f_n\}_{n=1}^\infty$  be a Cauchy sequence in  $\mathcal{H} \cap \mathcal{F}$ . Then  $f_n(x)$  is constant for all sufficiently large values of  $n$ . This constant value will be denoted  $f(x)$ . Thus we have  $\lim_{n \rightarrow \infty} f_n(x) = f(x)$  for all  $x$  in  $T$ . It remains to show  $f \in \mathcal{H}$ . In fact, any restriction  $g$  of  $f$  ( $g \in \mathcal{F}^*$ ) is also the restriction of some  $f_n$ ; and since each  $f_n$  belongs to  $\mathcal{H}$  and since  $\mathcal{H}$  is hereditary, we have that  $g$  belongs to  $\mathcal{H}$ . Again, since  $\mathcal{H}$  is hereditary, and since an arbitrary restriction in  $\mathcal{F}^*$  of  $f$  is in  $\mathcal{H}$ , it follows that  $f$  belongs to  $\mathcal{H}$ .

THEOREM 2. Let  $\mathcal{A}$  be absorbing with respect to a hereditary set  $\mathcal{H}$ . Then  $(\mathcal{H} - \mathcal{A}) \cap \mathcal{F}$  is nowhere dense in  $\mathcal{H} \cap \mathcal{F}$  with respect to the metric  $d$ .

*Proof.* Let  $f$  be an arbitrary element of  $\mathcal{H} \cap \mathcal{F}$ . We shall show that, given any positive integer  $k$ , there exists  $g$  in  $\mathcal{A} \cap \mathcal{F}$  and a positive integer  $l$  such that

$$(1) \quad \{h \in \mathcal{H} \cap \mathcal{F} : d(h, f) < 1/k\} \supset \{h \in \mathcal{H} \cap \mathcal{F} : d(h, g) < 1/l\}$$

and

$$(2) \quad \{h \in \mathcal{H} \cap \mathcal{F} : d(h, g) < 1/l\} \subset \mathcal{A} \cap \mathcal{F}.$$

This will show that  $(\mathcal{H} - \mathcal{A}) \cap \mathcal{F}$  is nowhere dense in  $\mathcal{H} \cap \mathcal{F}$ .

In fact, let  $f_k$  be the restriction of  $f$  to the set  $\bigcup_{j=1}^{k-1} T_j$ . Because  $\mathcal{A}$  is absorbing, there exists  $h_l$  in  $\mathcal{G} \cap \mathcal{A}$  such that  $f_k$  is a restriction of  $h_l$ . Let the domain of definition of  $h_l$  be  $\bigcup_{j=1}^l T_j$ . Because  $\mathcal{A}$  is absorbing,  $h_l$  can be extended to  $h_{l+1}$  on  $\bigcap_{j=1}^{l+1} T_j$  ( $h_{l+1} \in \mathcal{A}$ ), and we proceed inductively to define the successive extensions  $h_k \in \mathcal{A}$   $k = l, l+1, \dots$ . The totality of these extensions defines a function  $g$  defined on all of  $T$ . Since  $h_k \in \mathcal{A} \subset \mathcal{H}$  and  $\mathcal{H}$  is hereditary,  $g$  belongs to  $\mathcal{H}$ , and therefore also to  $\mathcal{A}$  since  $\mathcal{A}$  is absorbing. The function  $g$  and the integer  $l$  have the properties required in (1) and (2).

3. A lemma uniformly continuous functions. The following well-known result is needed later. The proof is routine and will be omitted.

LEMMA 1. Suppose that the real function  $f$  is defined on a set  $S$  of real numbers and is uniformly continuous there. Then  $f$  can be extended in exactly one way to a uniformly continuous function on the closure of  $S$ .

4. Application of methods of §2 to nonconstant locally recurrent functions. Let  $T_i$  consist of all rational numbers of the form  $n/9^{i-1}$  where  $n$  and  $i$  are nonnegative integers such that  $0 \leq n \leq 9^{i-1}$ ,  $i > 0$ , and  $n$  is not divisible by 9. Let  $T = \bigcup_{i=0}^{\infty} T_i$ . Let  $S$  be the set of all real numbers, and  $\mathcal{F}$  the set of real valued functions on  $T$ . As in §2,  $\mathcal{G}$  consists of all real functions defined on sets  $\bigcup_{i=0}^N T_i$ , and  $\mathcal{F}^* = \mathcal{F} \cup \mathcal{G}$ .

Let  $\mathcal{H}$  consist of elements  $f$  of  $\mathcal{F}^*$  which satisfy the following conditions:

Condition 1. Suppose the domain of definition of  $f$  contains  $r/3^k$  and  $(r+1)/3^k$ . Let

$$(3) \quad m = \min (f(r/3^k), f((r+1)/3^k))$$

and

$$(4) \quad M = \max (f(r/3^k), f((r+1)/3^k)).$$

Then  $M \neq m$  and for all  $x$  in the domain of definition of  $f$  such that  $r/3^k < x < (r+1)/3^k$  we have  $m \leq f(x) \leq M$ .

Condition 2. If the domain of definition of  $f$  contains  $r/3^k$  and  $(r+1)/3^k$ , then

$$|f(r/3^k) - f((r+1)/3^k)| \leq 3^{k/2}.$$

Let  $\mathcal{A}_n$  ( $n = 0, 1, \dots$ ) be the subset of  $\mathcal{H}$  consisting of functions  $f$  which satisfy, in addition to the foregoing two conditions, the following:

Condition 3. There exists  $i \geq n$  such that  $T_i$  belongs to the domain of definition of  $f$ , and for any  $r$  such that  $0 \leq 3r < 9^{i-1}$  we have

$$(6) \quad \text{a. } f(3r/9^{i-1}) = f((3r+2)/9^{i-1})$$

$$(7) \quad \text{b. } f((3r+1)/9^{i-1}) = f((3r+3)/9^{i-1}).$$

**THEOREM 3.**  $\mathcal{H}$  is hereditary, and for each  $n$  ( $n = 0, 1, \dots$ )  $\mathcal{A}_n$  is absorbing.

*Proof.* It is easy to verify that  $\mathcal{H}$  is hereditary. The essential thing is that Conditions 1 and 2 involve only universal quantifiers, and no existential quantifiers. If these Conditions hold for all points in the domain of  $f$ , and if  $g$  is a restriction of  $f$ , then they also hold, *a fortiori*, for all points in the domain of  $g$ , because the latter set is a subset of the former.

Now we shall show that  $\mathcal{A}_n$  is absorbing. It is clear that part 1 of Definition 3 is satisfied. To show part 2, suppose  $g \in \mathcal{G} \cap \mathcal{H}$ . Let the domain of definition of  $g$  be  $\bigcup_{i=1}^N T_i = R_N$ . Let us define  $f$  on  $\bigcup_{i=1}^{N+1} T_i$  so that if  $x \in R_N$ , then  $f(x) = g(x)$ .

We define  $f$  at points of the form  $(9r + 3)/9^N$  and  $(9r + 6)/9^N$  by linear interpolation, i.e.,

$$\begin{aligned} f((9r + 3)/9^N) &= \frac{2}{3} f(r/9^{N-1}) + \frac{1}{3} f((r + 1)/9^{N-1}) \\ f((9r + 6)/9^N) &= \frac{1}{3} f(r/9^{N-1}) + \frac{2}{3} f((r + 1)/9^{N-1}). \end{aligned}$$

On the remaining points of  $T_{N+1}$ ,  $f$  is now uniquely determined by imposing requirements  $a$  and  $b$  above for  $\mathcal{A}_n$  with  $n$  replaced by  $N + 1$ :

$$\begin{aligned} f(3r/9^N) &= f((3r + 2)/9^N) \\ f((3r + 1)/9^N) &= f((3r + 3)/9^N). \end{aligned}$$

The author hopes that the geometry of this construction is made clear by the figure. The example of §1 is based on this construction. (However, the functions illustrated in the figure are defined for all  $x$  in  $[0, 1]$  whereas the function  $f$  above is defined only at finitely many points.)

It is rather clear from the construction that  $f$  satisfies the condition on maxima and minima in Condition 1 above. We now show that Condition 2 holds. We have given  $g \in \mathcal{H}$  and hence the condition holds for pairs of points  $r, (r + 1)/3^k$  in case  $k$  is less than  $2N - 1$ . We must show that the condition holds for pairs of points of the type  $3r/9^N, 3r + 3/9^N$ , and then for pairs of points of the type  $s/9^N, (s + 1)/9^N$ . For a pair of the former type we have

$$\begin{aligned} &|f(3r/9^N) - f((3r + 3)/9^N)| \\ &= \frac{1}{3} |f([3r/9]/9^{N-1}) - f([3r/9]/9^{N-1})| \\ &\leq 3^{-1} 9^{-(n-k)/2} < (3/9^N)^{1/2}. \end{aligned}$$

For a pair of the latter type we have

$$\begin{aligned}
& |f(s/9^N) - f((s+1)/9^N)| \\
&= |f(3[s/3]/9^N) - f(3([s/3]+1)/9^N)| \\
&= \frac{1}{3} |f([s/9]/9^{N-1}) - f([s/9]+1)/9^{N-1})| \\
&\leq 3^{-1} 9^{-(N-1)/2} = 1/9^{N/2}.
\end{aligned}$$

Now, from Theorem 2,  $(\mathcal{H} - \mathcal{A}_n) \cap \mathcal{F}$  is nowhere dense in  $\mathcal{H} \cap \mathcal{F}$ , for each  $n$ , with respect to the metric  $d$ . It follows that  $(\mathcal{H} - \bigcup_{n=0}^{\infty} \mathcal{A}_n) \cap \mathcal{F}$  is of the first category, and thus by the Category Theorem we have the following.

**THEOREM 4.** *The set of functions  $f$  on  $T$  such that for any  $n$  there exist  $i \geq n$  such that Condition 3 is satisfied is of the second category with respect to the metric  $d$  in the space of functions of  $T$  which satisfy Conditions 1 and 2.*

Now we wish to extend our functions, using Lemma 1, to functions defined on the whole interval  $[0, 1]$ .

**THEOREM 5.** *The functions in  $\mathcal{H} \cap \mathcal{F}$  are uniformly continuous. In fact, they satisfy a Hölder condition with the exponent  $1/2$ .*

*Proof.* Let  $x$  and  $y$  ( $x < y$ ) be arbitrary points in  $T$ . The interval  $[x, y]$  can be expressed uniquely as a countable union of intervals.  $I_n^{\#} = [a_n, b_n]$  disjoint except possibly for end points, such that the following conditions are satisfied.

1. Each  $I_n$  is of the form  $[r/3^k, (r+1)/3^k]$ .
2. No  $I_n$  is a subset of an interval  $J$  of the above form such that  $J$  is a subset of  $[x, y]$ .

It is clear that for any  $k$ , there can be among the  $I_n$  at most four intervals of length  $1/3^k$ . Let  $L_n$  denote the length of  $I_n$ . We have for any  $f$  in  $\mathcal{H} \cap \mathcal{F}$ ,

$$\begin{aligned}
|f(y) - f(x)| &\leq \sum |f(b_n) - f(a_n)| \\
&\leq \sum L_n^{1/2} < 4 \sum_{k=0}^{\infty} \left( \frac{y-x}{3^k+1} \right)^{1/2} \\
&< \frac{4\sqrt{3}}{\sqrt{3}-1} (y-x)^{1/2}.
\end{aligned}$$

Now by Lemma 1 the functions in  $\mathcal{H} \cap \mathcal{F}$  can be extended to all of  $[0, 1]$  in a unique way. Further, it is clear that these functions satisfy a Hölder condition with the exponent  $1/2$ .

**THEOREM 6.** *The continuous extensions of the functions in  $(\bigcap_{n=0}^{\infty} \mathcal{A}_n) \cap \mathcal{F}$  are nonconstant and everywhere locally recurrent.*

*Proof.* Let  $f$  be in  $(\bigcap_{n=0}^{\infty} \mathcal{A}_n) \cap \mathcal{F}$  and let  $x$  be any point in  $[0, 1]$ . For any  $\varepsilon > 0$ ,  $x$  belongs to an interval of the form  $I = [3r/9^k, (3r + 1)/9^k]$  of length less than  $\varepsilon$ , such that

$$f(3r/9^k) = f((3r + 2)/9^k)$$

and

$$f((3r + 1)/9^k) = f((3r + 3)/9^k).$$

By the intermediate value theorem for continuous functions, there exists  $y \in I$ ,  $x \neq y$ , such that  $f(x) = f(y)$ . This proves local recurrence. The functions are nonconstant because of Condition 1.

5. **Other applications.** It is hoped that the methods of § 2 will be useful as a tool for the construction of various counterexamples.

The methods of § 2 can be used in a routine way to give the following results.

**PROPOSITION 1.** The set of real functions which are not convex in any interval is of second category with respect to the metric  $d$  in the space of monotone functions.

**PROPOSITION 2.** The set of real functions  $f$  such that for any interval  $I \subset [0, 1]$  one has

$$\limsup_{x, y \in I} \frac{f(y) - f(x)}{y - x} = 1$$

and

$$\liminf_{x, y \in I} \frac{f(y) - f(x)}{y - x} = -1$$

is of second category with respect to the metric  $d$  in the space of functions which satisfy the Lipschitz condition  $|f(y) - f(x)| \leq |x - y|$  for all  $x$  and  $y$  in  $[0, 1]$ .

## References

1. S. Banach, *Über die Baire'sche Kategorie gewisser Funktionenmengen*, *Studia Math.* **3** (1931), 174-180.
2. K. A. Bush, *Locally recurrent functions*, *American Math. Monthly* **69** (1962), 199-206.
3. S. Mazurkiewicz, *Sur les fonctions non dérivables*, *Studia Math.* **3** (1931), 92-94.

Received May 27, 1966.

UNIVERSITY OF CALIFORNIA, DAVIS





# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. SAMELSON

Stanford University  
Stanford, California

J. P. JANS

University of Washington  
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California  
Los Angeles, California 90007

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*

AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$ 8.00; single issues, \$ 3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$ 4.00 per volume; single issues \$ 1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), No. 6, 2-chome, Fujimi-cho, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Richard Allen Askey, <i>A transplantation theorem for Jacobi coefficients</i> . . . .	393
Raymond Balbes, <i>Projective and injective distributive lattices</i> . . . . .	405
Raymond Balbes and Alfred Horn, <i>Order sums of distributive lattices</i> . . . . .	421
Donald Charles Benson, <i>Nonconstant locally recurrent functions</i> . . . . .	437
Allen Richard Bernstein, <i>Invariant subspaces of polynomially compact operators on Banach space</i> . . . . .	445
Robert F. Brown, <i>Fixed points and fibre</i> . . . . .	465
David Geoffrey Cantor, <i>On the Stone-Weierstrass approximation theorem for valued fields</i> . . . . .	473
James Walton England, <i>Stability in topological dynamics</i> . . . . .	479
Alessandro Figà-Talamanca and Daniel Rider, <i>A theorem on random Fourier series on noncommutative groups</i> . . . . .	487
Sav Roman Harasymiv, <i>A note of dilations in <math>L^p</math></i> . . . . .	493
J. G. Kalbfleisch, <i>A uniqueness theorem for edge-chromatic graphs</i> . . . . .	503
Richard Paul Kelisky and Theodore Joseph Rivlin, <i>Iterates of Bernstein polynomials</i> . . . . .	511
D. G. Larman, <i>On the union of two starshaped sets</i> . . . . .	521
Henry B. Mann, Josephine Mitchell and Lowell Schoenfeld, <i>Properties of differential forms in <math>n</math> real variables</i> . . . . .	525
John W. Moon and Leo Moser, <i>Generating oriented graphs by means of team comparisons</i> . . . . .	531
Veikko Nevanlinna, <i>A refinement of Selberg's asymptotic equation</i> . . . . .	537
Ulrich Oberst, <i>Relative satellites and derived functors of functors with additive domain</i> . . . . .	541
John Vincent Ryff, <i>On Muirhead's theorem</i> . . . . .	567
Carroll O. Wilde and Klaus G. Witz, <i>Invariant means and the Stone-Čech compactification</i> . . . . .	577