

Pacific Journal of Mathematics

**A SEMIGROUP UNION OF DISJOINT LOCALLY FINITE
SUBSEMIGROUPS WHICH IS NOT LOCALLY FINITE**

THOMAS CRAIG BROWN

A SEMIGROUP UNION OF DISJOINT LOCALLY FINITE SUBSEMIGROUPS WHICH IS NOT LOCALLY FINITE

T. C. BROWN

The semigroup S of the title is the free semigroup F on four generators factored by the congruence generated by the set of relations $\{w^2 = w^3 \mid w \in F\}$. The following lemma is proved by examining the elements of a given congruence class of F :

LEMMA. **If $x, y \in S$ and $x^2 = y^2$, then either $xy = x^2$ or $yx = x^2$.**

From the Lemma it then easily follows that the (disjoint) subsemigroups $\{y \in S \mid y^2 = x^2\}$ of S are locally finite.

This note answers in the negative a question raised by Shevrin in [2].

THEOREM. *There exists a semigroup S with disjoint locally finite subsemigroups S_e such that $S = \cup S_e$ and S is not locally finite.*

Let F be the free semigroup with identity on four generators. Let \sim denote the smallest congruence on F containing the set $\{(x^2, x^3) \mid x \in F\}$. That is, for $w, w' \in F$, $w \sim w'$ if and only if a finite sequence of "transitions", of either of the types $ab^2c \rightarrow ab^3c$ or $ab^3c \rightarrow ab^2c$, transforms w into w' .

The equivalence classes of F with respect to \sim are taken as the elements of S , and multiplication in S is defined in the natural way.

There is given in [1] a sequence on four symbols in which no block of length k is immediately repeated, for any k . Thus the left initial segments of this sequence give elements of F containing no squares. Since no transition of the form $ab^2c \rightarrow ab^3c$ or $ab^3c \rightarrow ab^2c$ can be applied to an element of F containing no squares, the equivalence classes containing these elements consist of precisely one element each; thus the semigroup S is infinite, and hence not locally finite.

In what follows, the symbols $\alpha, \alpha_1, \alpha_2, \dots$ refer to transformations (on elements of F) of the form

$$ab \rightarrow ayb, \quad \text{where } a \sim ay, \quad \text{and } a, b, y \in F.$$

The symbols $\beta, \beta_1, \beta_2, \dots$ refer to transformations of the type

$$axb \rightarrow ab, \quad \text{where } a \sim ax, \quad \text{and } a, b, x \in F.$$

Note that $ab^2c \rightarrow ab^3c$ is an α , and $ab^3c \rightarrow ab^2c$ is a β .

LEMMA 1. *If $w, w' \in F$, and $w\beta\alpha = w'$, then there are α_1, β_1 such that $w\alpha_1\beta_1 = w'$.*

Proof. Let

$$w = axb, \quad w\beta = ab, \quad \text{where } a \sim ax.$$

Let

$$w\beta = AB, \quad w\beta\alpha = AyB, \quad \text{where } A \sim Ay.$$

There are two cases:

(i) A is contained in a . That is,

$$a = Aa' \quad \text{and} \quad w' = wba = a\beta\alpha = Aa'ba = Aya'b.$$

Here let

$$w\alpha_1 = axb\alpha_1 = Aa'xb\alpha_1 = Aya'xb.$$

Now since

$$Aya' \sim Aa' = a \sim ax = Aa'x \sim Aya'x,$$

we may let

$$w\alpha_1\beta_1 = Aya'xb\beta_1 = Aya'b = w'.$$

(ii) A is not contained in a . That is,

$$b = b_1b_2, \quad A = ab_1, \quad A \sim Ay,$$

and

$$w' = w\beta\alpha = ab\alpha = ab_1b_2\alpha = Ab_2\alpha = Ayb_2 = ab_1yb_2.$$

Since

$$axb_1 \sim ab_1 = A \sim Ay = ab_1y \sim axb_1y,$$

we may let

$$w\alpha_1 = axb\alpha_1 = axb_1b_2\alpha_1 = axb_1yb_2,$$

and

$$w\alpha_1\beta_1 = axb_1yb_2\beta_1 = ab_1yb_2 = w'.$$

LEMMA 2. *If $w, w' \in F$, $w\gamma_1\gamma_2 \cdots \gamma_m = w'$, where each γ_i is either an α or a β , then there are $\alpha_1, \dots, \alpha_n, \beta_1, \dots, \beta_k$ such that $w\alpha_1 \cdots \alpha_n\beta_1 \cdots \beta_k = w'$.*

This follows immediately from Lemma 1 by induction.

LEMMA 3. *The word aba contains a left initial segment which is equivalent to a .*

Proof. Let $ab = AB$, $aba = AyB$, where $A \sim Ay$. Again there are two cases:

(i) A is contained in a . That is, $a = Aa'$, $aba = Aa'ba = Aya'b$.

Since $A \sim Ay$, the left initial segment Aya' of aba is equivalent to a .

(ii) A is not contained in a . That is, $b = b_1b_2$, $A = ab_1$, $aba = ab_1b_2a = ab_1yb_2$.

Here, a itself is a left initial segment of aba , and is certainly equivalent to a .

LEMMA 4. *If $x, y \in F$ and $x^2 \sim y^2$, then either $y \sim xa$ for some $a \in F$, or $x \sim yb$ for some $b \in F$.*

Proof. By Lemma 2, there are α_i and β_j such that $xx\alpha_1 \cdots \alpha_m\beta_1 \cdots \beta_n = yy$. Let $w = xx\alpha_1 \cdots \alpha_m = yy\beta_n^{-1} \cdots \beta_1^{-1}$. By Lemma 3, w contains a left initial segment A equivalent to x . Similarly, since each β_i^{-1} is an α , w also contains a left initial segment B equivalent to y . Depending on which segment contains the other, either $B = Aa$ for some a , or $A = Bb$ for some b . In the first case, $y \sim B = Aa \sim xa$; in the second, $x \sim A = Bb \sim yb$.

LEMMA 5. *In this lemma, “=” will denote equality in S . Let e be an idempotent element of S : $e = e^2$. Let $S_e = \{x \in S \mid x^2 = e\}$. Then S_e is a locally finite subsemigroup of S .*

Proof. First, we note that $z \in S_e$ implies $ez = ze = e$. For $ez = z^2z = z^2 = e$, and similarly $ze = e$. Now let $x, y \in S_e$, that is, $x^2 = y^2 = e$. By Lemma 4, either $y = xa$ or $x = yb$. In the first case, we obtain $xy = x^2a = x^3a = x^2y = ey = e$. In the second case, we obtain similarly that $yx = e$. Thus $x, y \in S_e$ implies $xy = e$ or $yx = e$. In either case, $(xy)^2 = e$, that is, $xy \in S_e$. Thus S_e is a semigroup.

Now let $x_1, \dots, x_n \in S_e$, and let $\langle x_1, \dots, x_n \rangle$ denote the subsemigroup of S_e generated by x_1, \dots, x_n . If $n = 1$, then $\langle x_1 \rangle$ is clearly finite; so suppose $n > 1$. Then every element of $\langle x_1, \dots, x_n \rangle$ may be expressed as a product of not more than n of the x_i 's. For any product z of more than n x_i 's must contain some x_i twice: $z = ax_i b x_i c$, where $a, b, c \in S_e$. Since either $x_i b = e$ or $b x_i = e$, it follows that $x_i b x_i = e$ and $z = aec = ec = e = x_i x_i$. This shows that $\langle x_1, \dots, x_n \rangle$ is finite, and hence that S_e is locally finite.

The theorem follows immediately from Lemma 5, since clearly $e \neq e'$ implies that S_e and $S_{e'}$ are disjoint, and

$$S = \cup S_e.$$

REFERENCES

1. R. A. Dean, *A sequence without repeats on x, x^{-1}, y, y^{-1}* , Amer. Math. Monthly **72** (1965), 383-385 (Math. Rev. **31**, No. 3500).
2. L. N. Shevrin, *On locally finite semigroups*, Doklady Akad. Nauk SSSR **162** (1965), 770-773 = Soviet Math. Dokl. **6** (1965), 769.

Received December 19, 1966. Partially supported by Canadian NRC grant no. A-3983.

SIMON FRASER UNIVERSITY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. SAMELSON

Stanford University
Stanford, California

J. P. JANS

University of Washington
Seattle, Washington 98105

J. DUGUNDJI

University of Southern California
Los Angeles, California 90007

RICHARD ARENS

University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Charles A. Akemann, <i>Some mapping properties of the group algebras of a compact group</i>	1
A. V. Boyd, <i>Note on a paper by Uppuluri</i>	9
Thomas Craig Brown, <i>A semigroup union of disjoint locally finite subsemigroups which is not locally finite</i>	11
Richard Thomas Bumby and Everett C. Dade, <i>Remark on a problem of Niven and Zuckerman</i>	15
James Calvert, <i>An integral inequality with applications to the Dirichlet problem</i>	19
Jack Gary Ceder and Terrance Laverne Pearson, <i>On products of maximally resolvable spaces</i>	31
William Guignard Faris, <i>The product formula for semigroups defined by Friedrichs extensions</i>	47
Robert S. Freeman, <i>Closed operators and their adjoints associated with elliptic differential operators</i>	71
Thomas Lee Hayden, <i>The extension of bilinear functionals</i>	99
Gloria Conyers Hewitt, <i>Limits in certain classes of abstract algebras</i>	109
Tilla Weinstein, <i>The dilatation of some standard mappings</i>	117
Mitsuru Nakai, <i>On Evans' kernel</i>	125
Ernest Levane Roetman, <i>On the biharmonic wave equation</i>	139
Malcolm Jay Sherman, <i>Operators and inner functions</i>	159
Walter Laws Smith, <i>On the weak law of large numbers and the generalized elementary renewal theorem</i>	171
A. J. Ward, <i>On H-equivalence of uniformities: The Isbell-Smith problem</i>	189