

# Pacific Journal of Mathematics

## **APPLICATION OF INFINITARY LANGUAGES TO METRIC SPACES**

**RALPH DAVID KOPPERMAN**

## APPLICATION OF INFINITARY LANGUAGES TO METRIC SPACES

RALPH KOPPERMAN

We attempt to lay the groundwork for applying the recently-developed theory of models for the infinitary languages  $L_{\pi\epsilon}^t$  to analysis. It will be shown that within one of these languages, axioms may be written whose class of models is precisely the metric spaces. We show that two complete separable metric spaces are elementarily equivalent in this language if and only if they are isomorphic and obtain an elimination of quantifiers for such spaces. A method is developed for transferring results on metric spaces to structures with metrics whose relations are closed under the metric topology. This class includes Banach Spaces.

When not otherwise indicated, definitions, notations, and model-theoretic results used in this paper may be found in [5].

The paper will be divided into three sections, as follows: I. Axiomatization of metric spaces, II. Theorems on metric systems, and III. Metric algebraic systems. Model-theoretic results will be introduced as needed.

I. Axiomatization of metric spaces. Our method of axiomatization of metric spaces is chosen in order to make their definition possible with one sentence in  $L_{\omega_1\omega}^m$  (for the definition of  $L_{\pi\epsilon}$ ,  $\omega_1$ , and  $\omega$ , see [5]). We begin with a definition of the rational numbers:

DEFINITION 1.1.  $Q = \langle Q, +, \cdot, 0, 1, \langle \rangle \rangle$ , a structure of type  $\mathbf{q} = \langle 3, 3, 0, 0, 2 \rangle$  will be called a rational number system if and only if:

- (1)  $Q$  is an ordered field (see [9, pp. 77, 5]).
- (2)  $(\forall v_0)(0 < v_0 \rightarrow \bigvee_{i < \omega} \bigvee_{j < \omega} (v_0 \cdot (1 + \dots + 1) = 1 + \dots + 1))$ .  
 $\qquad\qquad\qquad j \text{ factors} \qquad\qquad i \text{ factors}$

THEOREM 1.2. *Each rational number system is isomorphic to the rational numbers.*

*Proof.* Each ordered field contains an isomorphic copy of the rational numbers, and by axiom 2, each positive (thus each negative) element of our field is the ratio of two "integers".

DEFINITION 1.3. A structure  $\mathfrak{X} = \langle A, M, Q, +, \cdot, 0, 1, \langle, d \rangle$  of type  $\mathbf{m} = \langle 1, 1, 3, 3, 0, 0, 2, 3 \rangle$  is a metric system if and only if

$\langle Q, +, \cdot, 0, 1, \langle \rangle$  is a rational number system, and  $+, \cdot, \langle$  apply only to members of  $Q$ , and:

- (1)  $(\forall v_0)((M(v_0) \vee Q(v_0)) \wedge (\neg M(v_0) \vee \neg Q(v_0)))$
- (2)  $(\forall v/3)(d(v/3) \rightarrow M(v_0) \wedge M(v_1) \wedge 0 < v_2)$
- (3)  $(\forall v/4)(d(v/3) \wedge v_2 < v_3 \rightarrow d(v_0, v_1, v_3))$
- (4)  $(\forall v/3)(\exists v_3)(d(v/3) \rightarrow v_3 < v_2 \wedge d(v_0, v_1, v_3))$
- (5)  $(\forall v/2)(\exists v_2)(M(v_0) \wedge M(v_1) \rightarrow d(v/3))$
- (6)  $(\forall v/3)(d(v/3) \rightarrow d(v_1, v_0, v_2))$
- (7)  $(\forall v/5)(d(v/3) \wedge d(v_1, v_3, v_4) \rightarrow d(v_0, v_3, v_2 + v_4))$
- (8)  $(\forall v/2)(\exists v_2)(M(v_0) \wedge M(v_1) \wedge \neg(v_0 \equiv v_1) \rightarrow 0 < v_2 \wedge \neg d(v/3))$
- (9)  $(\forall v/3)(M(v_0) \wedge v_0 \equiv v_1 \wedge 0 < v_2 \rightarrow d(v/3)).$

The general idea of these axioms is to identify each real number with the set of rationals greater than it. Axioms 1 to 4 simply prepare the groundwork for the above interpretation. Axiom 5 says that to each pair of points there corresponds a distance. Axiom 6 gives the symmetry of distance, 7 gives the triangle inequality, 8 yields the fact that any pair of distinct points differ by a positive distance, and 9 says that the distance from a point to itself is 0. To formalize the above discussion, we have the following representation theorem for metric systems:

**THEOREM 1.4.** (a) *Let  $\mathfrak{A} = \langle A, M, Q, +, \cdot, 0, 1, \langle, d \rangle$  be a metric system. Then  $\mathfrak{A}^* = \langle M, m \rangle$  is a metric space, where  $m(x, y) = \inf \{q / \langle x, y, q \rangle \in d\}$  (for each pair  $x, y \in M$ ). This metric space is called the associated space of the metric system  $\mathfrak{A}$ .*

(b) *Let  $\mathcal{M} = \langle M, m \rangle$  be a metric space. Then the associated system  $\mathcal{M}^* = \langle M \cup Q, M, Q, +, \cdot, 0, 1, \langle, d \rangle$  of  $\mathcal{M}$  is a metric system, where  $Q$  together with its relations and constants  $(+, \cdot, 0, 1, \langle)$  is the rational numbers, our union  $M \cup Q$  is disjoint, and*

$$d = \{ \langle x, y, q \rangle / m(x, y) < q \in Q \} .$$

*Proof.* Consider the completion  $R$  of  $Q$ . If  $x, y \in M$ , by 5  $\{q / \langle x, y, q \rangle \in d\}$  is nonempty, and by 2, all its elements are positive (thus 0 is a lower bound). Thus  $m(x, y)$  exists. It remains to be shown that  $m$  is a metric. To show the triangle inequality we note that axiom 7 gives us the fact that if  $\langle x, y, q \rangle \in d$  and  $\langle y, z, r \rangle \in d$ , then  $\langle x, z, q + r \rangle \in d$ . Thus

$$\inf \{q / \langle x, y, q \rangle \in d\} + \inf \{r / \langle y, z, r \rangle \in d\} \geq \inf \{s / \langle x, z, s \rangle \in d\} ,$$

in other words  $m(x, z) \leq m(x, y) + m(y, z)$ . The other laws are checked with similar case. This shows (a), and under the assumptions of (b), axioms 1,  $\dots$ , 9 for metric systems can easily be shown.

**LEMMA 1.5.** (a) *If  $\mathfrak{A}, \mathfrak{B}$  are metric spaces or metric systems and*

$\mathfrak{A} \cong \mathfrak{B}$ , then  $\mathfrak{A}^* \cong \mathfrak{B}^*$  (where for metric spaces, isomorphism = isometry).

(b) If  $\mathfrak{A}$  is a metric space or a metric system,  $\mathfrak{A} \cong (\mathfrak{A}^*)^*$ .

(c) For metric systems  $\mathfrak{A}, \mathfrak{B}, \mathfrak{A} \in \text{Iso Sub}(\mathfrak{B})$  if and only if  $\mathfrak{A}^* \cong \mathfrak{C}$  for some subspace  $\mathfrak{C}$  of  $\mathfrak{B}^*$ . (We write  $\mathfrak{A} \in \text{Iso Sub}(\mathfrak{B})$  if and only if  $\mathfrak{A} \cong \mathcal{D}$  for some  $\mathcal{D} \subset \mathfrak{B}$ ,  $\mathfrak{A} \in \text{Iso Sub}_i(\mathfrak{B})$  if and only if  $\mathfrak{A} \in \text{Iso Sub}(\mathfrak{B})$  and  $\text{card}(\mathfrak{A}) < \varepsilon$ .)

The proof is routine, though somewhat long, and left to the reader. We will now say that a metric system has property  $P$  if and only if its associated space has that property. We will also say that a metric system and a metric subsystem of it have property  $Q$  if and only if the associated space and subspace have that property (for example, the subspace may be dense). Often we will want to translate these properties into our first-order infinitary language. A proof similar to that of Theorem 17, [6] (with  $Q$  in place of  $\mathfrak{A}$ , and  $\omega_1\omega$  in place of  $\varepsilon\varepsilon$ ) shows that completeness cannot be characterized by a single sentence in  $L_{\omega_1\omega}^m$ . It is clear, however, that complete metric systems are precisely those which satisfy the following sentence in

$$L_{\omega_1\omega_1}^m: Ct = (\forall v/\omega)((\forall v_\omega) 0 < v_\omega \rightarrow \mathbf{V}_{i < \omega} \mathbf{\Lambda}_{i < j < \omega} \mathbf{\Lambda}_{i < k < \omega} d(v_j, v_k, v_\omega)) \\ \rightarrow ((\exists v_{\omega+1})(\forall v_{\omega+2}) 0 < v_{\omega+2} \rightarrow (\mathbf{V}_{i < \omega} \mathbf{\Lambda}_{i < j < \omega} d(v_{\omega+1}, v_j, v_{\omega+2}))) ,$$

since this sentence simply says that each sequence  $(v/\omega)$  which is cauchy (i.e., such that for each  $v$  there is an  $i < \omega$  such that for each finite  $j, k$  such that  $j$  and  $k$  are greater than  $i, v_j$  and  $v_k$  are “closer together” than  $v_\omega$ ) there is a limit  $(v_{\omega+1})$ . It is also clear that separable metric systems are simply those satisfying Sep, where

$$\text{Sep} = (\exists v/\omega)(\forall v_\omega)(\forall v_{\omega+1})(M(v_\omega) \wedge 0 < v_{\omega+1} \rightarrow \mathbf{V}_{i < \omega} d(v_\omega, v_i, v_{\omega+1})) .$$

Thus most of our future dealings with metric systems will take place in  $L_{\omega_1\omega_1}^m$ .

## II. Theorems on metric systems.

**THEOREM 2.1.** *Let  $\mathfrak{A} \prec_{\omega_1, \omega_1} \mathfrak{B}$  and let  $\mathfrak{A}$  or  $\mathfrak{B}$  be separable. Then  $\mathfrak{A} = \mathfrak{B}$ . (For definition of  $\prec_{\omega_1, \omega_1}$  see [5], Definition 2.2).*

*Proof.* Assume  $\mathfrak{A}$  is separable. Let  $\mathfrak{C} = \langle \{n_0, n_1, \dots\}, m' \rangle$  be a dense countable subspace of  $\mathfrak{A}^*$ , Then  $\mathfrak{A} \models D[n_0, \dots]$ , where

$$D = (\forall v_\omega)(\forall v_{\omega+1})(M(v_\omega) \wedge 0 < v_{\omega+1} \rightarrow \mathbf{V}_{i < \omega} d(v_\omega, v_i, v_{\omega+1})) .$$

Thus  $\mathfrak{C}$  is dense in  $\mathfrak{B}^*$ , since by our assumption,  $\mathfrak{B} \models D[n_0, \dots]$ .

Now let  $b \in M'$ , where  $\mathfrak{B} = \langle B, M', \dots \rangle$ . Since  $\mathfrak{C}$  is dense in  $\mathfrak{B}^*$ , there is a sequence  $n_{i_0}, n_{i_1}, \dots$  of the  $n_i$ 's approaching  $b$ . Thus  $\mathfrak{B} \models (\exists v_\omega)L[n_{i_0}, \dots]$ , where  $L = (\forall v_{\omega+1})(0 < v_{\omega+1} \rightarrow \mathbf{V}_{i < \omega} \mathbf{\Lambda}_{i < j < \omega} d(v_\omega, v_j, v_{\omega+1}))$ . Thus  $\mathfrak{A} \models (\exists v_\omega)L[n_{i_0}, \dots]$ , but since  $\mathfrak{A}^*$  is a subspace of  $\mathfrak{B}^*$  and limits are unique in metric spaces  $\mathfrak{A}^* = \mathfrak{B}^*$ . If  $\mathfrak{A} = \langle A, M, \dots \rangle$  this implies that  $M = M'$ , so  $A = B$ , thus  $\mathfrak{A} = \mathfrak{B}$ . This proves the theorem in the first case, but if  $\mathfrak{B}$  is separable, then  $\mathfrak{A}$  must also be, from which the theorem must follow in the second case.

It can be shown similarly that if  $\mathfrak{A} \prec_{\omega_1, \omega_1} \mathfrak{B}$  and  $\mathfrak{A}$  is dense in  $\mathfrak{B}$ , then  $\mathfrak{A} = \mathfrak{B}$ . The following definition and results may be found in Tarski.

**DEFINITION 2.2.** Let  $\mathfrak{A}$  be a structure of type  $t$ ,  $\text{Dom}(t) = \varepsilon < \pi$ , and let  $a$  be a well-ordering of  $A$ . Then

$$\text{Des}_a(\mathfrak{A}) = (\mathbf{\Lambda} F) \wedge (\mathbf{\Lambda} F') \wedge (\mathbf{\Lambda}_{t(p) \geq 1} (\mathbf{\Lambda} F_p)) \wedge (\mathbf{\Lambda}_{t(p) \geq 1} (\mathbf{\Lambda} F'_p)),$$

where

$$\begin{aligned} F &= \{v_i \equiv v_j/a_i = a_j\} \cup \{v_i \equiv c_j/a_i = c_j\} \cup \{c_i \equiv c_j/c_i = c_j\}, \\ F' &= \{\neg v_i \equiv v_j/a_i \neq a_j\} \cup \{\neg v_i \equiv c_j/a_i \neq c_j\} \cup \{\neg c_i \equiv c_j/c_i \neq c_j\}, \\ F_p &= \{R_p(x \cdot j/t(p)) / b \cdot j/t(p) \in R_p, \end{aligned}$$

where  $b_{j_i} = a_k$  if  $x_{j_i} = v_k$  and  $b_{j_i} = c_k$  if  $x_{j_i} = c_k$  and

$$F'_p = \{\neg R_p(x \cdot j/t(p)) / b \cdot j/t(p) \notin R_p, b \text{ defined as for } F_p\}.$$

**COROLLARY 2.3.** Let  $\mathfrak{A}$  be a structure of type  $t$ . Then  $\mathfrak{A} \in \text{Iso Sub } (\mathfrak{B})$  if and only if  $\mathfrak{B} \models (\exists X) \text{Des}(\mathfrak{A})$ , where  $X = FV(\text{Des}(\mathfrak{A}))$ .

Note that if  $c(A) < \pi$ , then  $\text{Des}(\mathfrak{A})$  and  $(\exists X) \text{Des}(\mathfrak{A})$  are both formulas (the latter a sentence) of  $L_{\pi\pi}^t$ .

**DEFINITION 2.4.** Let  $\mathfrak{C}$  be a countable metric system,  $c$  a mapping from  $\omega$  one-one onto  $C$ . Then let

$$\begin{aligned} Dc_c(\mathfrak{C}) &= (\exists v/\omega)(\forall v_\omega)(\forall v_{\omega+1})(\text{Des}_c(\mathfrak{C}) \wedge (M(v_\omega) \wedge 0 < v_{\omega+1} \\ &\rightarrow \mathbf{V}_{i < \omega} d(v_i, v_\omega, v_{\omega+1}))) . \end{aligned}$$

Note that  $\mathfrak{A}$  is dense in  $\mathfrak{B}$  for some  $\mathfrak{A} \cong \mathfrak{C}$  if and only if

$$\mathfrak{B} \models Dc_c(\mathfrak{C}).$$

Thus  $Dc_c$  should be viewed as a method of expressing within  $L_{\omega_1\omega_1}^m$  the fact that a countable space is dense in another space. If it is unnecessary to distinguish our well-ordering  $c$ , we simply write  $Dc(\mathfrak{C})$  in place of  $Dc_c(\mathfrak{C})$ .

LEMMA 2.5. *If  $\mathfrak{A}$  is a metric system,  $\mathfrak{A} \models \text{Sep}$  if and only if for some countable metric system  $\mathfrak{C}$ ,  $\mathfrak{A} = \text{Dc}(\mathfrak{C})$ .*

The proof is left to the reader.

THEOREM 2.6. *Let  $\mathfrak{A}$  be a complete separable metric system,  $\mathfrak{B}$  a metric system. Then  $\mathfrak{A} \equiv_{\omega_1, \omega_1} \mathfrak{B}$  if and only if  $\mathfrak{A} \cong \mathfrak{B}$ . (A definition of  $\equiv_{\pi, \varepsilon}$  is found in 2.1, [5].)*

Proof. Let  $\langle K, k \rangle$  be a dense countable subspace of  $\mathfrak{A}^*$ , where

$$\mathfrak{A} = \langle A, M, \dots \rangle, \mathfrak{B} = \langle B, N, \dots \rangle.$$

Then  $\mathfrak{A} \models \text{Dc}(\langle K, k \rangle^*) \wedge \text{Ct}$ , so  $\mathfrak{B} \models \text{Dc}(\langle K, k \rangle^*) \wedge \text{Ct}$ . By the remark following Definition 2.4,  $\mathfrak{D}$  is dense in  $\mathfrak{B}$  for some  $\mathfrak{D} \cong \langle K, k \rangle$ . Thus up to isomorphism,  $\langle K, k \rangle$  is dense in  $\mathfrak{B}^*$  and  $\mathfrak{A}^*$ , two complete metric spaces. It is well-known in the theory of metric spaces that if two complete spaces have the same dense subspace (up to isometry), then the spaces are isometric. Thus by Lemma 1.5, a,  $\mathfrak{A} \cong \mathfrak{B}$ .

We are going on to eliminate quantifiers for complete separable metric systems. First we make the notation  $B(S)$  for all the boolean combinations of a set of formulas  $S$  (see Definition 5.1, [6]), and restate Theorem 5.2 of that paper.

THEOREM 2.7. *Let  $S$  be a consistent set of sentences (i.e., one with models) in  $L_{\pi_\varepsilon}^t$ ,  $E$  a set of sentences in  $L_{\pi_\varepsilon}^t$  such that there is a set  $B'(E)$  with the property that for each  $\phi \in B(E)$  there is a  $\phi' \in B'(E)$  with  $S \models \phi \leftrightarrow \phi'$ . For any consistent set of sentences  $T \subset L_{\pi_\varepsilon}^t$ ,  $S \subset T$ , assume there is an  $A \subset E$  such that  $T \subset T_{\pi_\varepsilon}(S \cup A)$  (and  $S \cup A$  consistent). Then for any sentence  $\phi \in L_{\pi_\varepsilon}^t$  there is a  $\theta \in B(E)$  such that  $S \models \phi \leftrightarrow \theta$ .*

LEMMA 2.8. *Let  $S \subset L_{\pi_\varepsilon}^t$  be a consistent set of sentences,  $E \subset L_{\pi_\varepsilon}^t$  a set of sentences such that (1) for any models  $\mathfrak{A}, \mathfrak{B}$  of  $S$ , if  $\mathfrak{A} \models \theta$ ,  $\mathfrak{B} \models \theta$ ,  $\theta \in E$ , then  $\mathfrak{A} \equiv_{\pi, \varepsilon} \mathfrak{B}$ , and (2) for any model  $\mathfrak{A}$  of  $S$  there is a  $\theta \in E$  such that  $\mathfrak{A} \models \theta$ . Then (a) if  $T \subset L_{\pi_\varepsilon}^t$  is a consistent set of sentences and  $S \subset T$ , then there is a  $\theta \in E$  such that  $T \subset T_{\pi_\varepsilon}(S \cup \{\theta\})$  and  $S \cup \{\theta\}$  is consistent, (b) if  $\phi \in B(E)$ ,  $S \models \phi \leftrightarrow \mathbf{V} A$  for some  $A \subset E$ , and (c) for any sentence  $\zeta \in L_{\pi_\varepsilon}^t$  there is a  $\phi \in B(E)$  such that  $S \models \zeta \leftrightarrow \phi$ .*

Proof. (a) If  $\mathfrak{A}$  is a model of  $T$ , then  $\mathfrak{A}$  is a model of  $S$ , so there is a  $\theta \in E$  such that  $\mathfrak{A} \models \theta$ . Now suppose  $\mathfrak{B}$  is a model of  $S$  such that  $\mathfrak{B} \models \theta$ . Then  $\mathfrak{A} \equiv_{\pi, \varepsilon} \mathfrak{B}$ , so  $\mathfrak{B} \models T$ . Thus  $T \subset T_{\pi_\varepsilon}(S \cup \{\theta\})$ . (b)  $\phi = \mathbf{V}_{i \in I} \bigwedge_{j \in J_i} \theta_{ij}$ ,  $\theta_{ij} \in E$  or  $\neg \theta_{ij} \in E$  for each  $i, j$ . Let  $\phi_i = \bigwedge_{j \in J_i} \theta_{ij}$

be a disjunct. If for some  $i, j, \theta_{ij} \in E$ , we assert that either

$$S \models \phi_i \leftrightarrow \theta_{ij}$$

or  $S \models \neg \phi_i$ . To show this, assume not  $S \models \phi_i \leftrightarrow \theta_{ij}$ . Clearly,  $S \models \phi_i \rightarrow \theta_{ij}$ , thus not  $S \models \theta_{ij} \rightarrow \phi_i$ . But then by (1),  $S \models \theta_{ij} \rightarrow \neg \phi_i$ , therefore  $S \models \neg \phi_i$ . If for each  $j$ ,  $\neg \theta_{ij} \in E$ ,  $\phi_i = \bigwedge_{j \in J_i} \neg (\neg \theta_{ij})$ . Thus by (2)  $S \models \phi_i \leftrightarrow \bigvee E - \{\theta_{ij}/j \in J_i\}$ . Thus in any of the above cases  $S \models \phi \leftrightarrow \bigvee_{k \in K_j} \theta_k$ , with each  $\theta_i \in E$ . Thus

$$S \models \phi \leftrightarrow \bigvee_{i \in I} \bigvee_{k \in K_i} \theta_k,$$

and eliminating repetitions,  $S \models \phi \leftrightarrow \bigvee A$ , which is condition (b) (note that while  $\bigvee A$  may not be in  $L_{\pi, \varepsilon}^t$ , it will be in  $L_{\pi', \varepsilon}^t$  for some  $\pi' \geq \pi$ , and all our statements have meaning in that language). (c) This is immediate for (a) and (b) and Theorem 2.7, using

$$B'(E) = \{\bigvee A/A \subset E\}.$$

**THEOREM 2.9.** *Let  $\phi \in L_{\omega_1 \omega_1}^m$  be a sentence. Then if  $\Sigma$  is the conjunction of the axioms for a metric system together with Sep and Ct, (i.e., is an axiom for complete separable metric systems) then  $\Sigma \models \phi \leftrightarrow \theta$  for some  $\theta \in B(E)$ , where  $E = \{Dc(\mathbb{C})/\mathbb{C} \text{ a countable metric system}\}$ .*

*Proof.* Let  $S = \{\Sigma\}$ . Then  $S$  is a consistent set of sentences in  $L_{\omega_1 \omega_1}^m$ .  $E$  is also a set of sentences in  $L_{\omega_1 \omega_1}^m$ , and  $S, E$  satisfy conditions (1) (by the proof of Theorem 2.6) and (2) (by Lemma 2.5) of Lemma 2.8. Thus our conclusion follows from (c) of that lemma.

To eliminate quantifiers for all formulas in  $L_{\omega_1 \omega_1}^m$  for the theory of complete separable metric spaces, we shall need the following:

**DEFINITION 2.10.** Let  $t$  be a type,  $i$  an ordinal. Then  $t * i$  is the type defined by  $t = t * i / \text{Dom}(t)$  and if  $j < i$ , then  $t * i(\text{Dom}(t) + j) = 0$ .

**DEFINITION 2.11.** Let  $\mathbb{C} = \langle C, K, \dots \rangle$  be a countable metric system,

$$\mathbb{C}' = \langle C, K, \dots, e_j \rangle_{j < \omega}, \quad \{e_j/j < \omega\} = D \subset K.$$

Then if  $c$  is a mapping of  $\omega$  one-one onto  $C$ ,  $DC_c(\mathbb{C}, D)$  is the sentence of  $L_{\omega_1 \omega_1}^{m, \omega}$  defined as follows:

$$DC_c(\mathbb{C}, D) = (\exists v/\omega)(\forall v_\omega)(\forall v_{\omega+1})(\text{Des}_c(\mathbb{C}') \\ \wedge (M(v_\omega) \wedge 0 < v_{\omega+1} \rightarrow \bigvee_{i < \omega} d(v_i, v_\omega, v_{\omega+1}))) .$$

If distinguishing  $c$  is unnecessary, we again leave it out and

write  $DC(\mathfrak{C}, D)$  in place of  $DC_c(\mathfrak{C}, D)$ .  $DC(\mathfrak{C}, D)$  should be considered as  $Dc(\mathfrak{C})$  with some constants added.

**THEOREM 2.12.** *Let  $\mathfrak{A}, \mathfrak{B}$  be  $m * \omega$ -structures such that  $\mathfrak{A}/m, \mathfrak{B}/m$  are complete separable metric systems. If  $\mathfrak{A} \models DC(\mathfrak{C}, D)$  and  $\mathfrak{B} \models DC(\mathfrak{C}, D)$ , then  $\mathfrak{A} \cong \mathfrak{B}$ .*

*Proof.* Let  $C = \{c_0, c_1, \dots\}$ , where  $\mathfrak{C}$  is dense in  $\mathfrak{A}/m$ . Let

$$\theta = DC_c(\mathfrak{C}, D),$$

where  $c = \langle c_0, c_1, \dots \rangle$ . Suppose  $\mathfrak{B} \models \theta$ . Then there is a sequence  $\{b_n/n < \omega\}$  satisfying  $\theta$  without the existential quantifier. The map  $f: \mathfrak{C} \rightarrow \mathfrak{C}'$  such that  $f(c_n) = b_n$  is an isomorphism because of  $\text{Des}(\mathfrak{C})$ . In particular, it is an isomorphism from  $\mathfrak{C}$  to  $\mathfrak{C}'$  preserving constants. Since  $\mathfrak{C}$  is dense in  $\mathfrak{A}/m$ ,  $\mathfrak{C}'$  is dense in  $\mathfrak{B}/m$ , thus this isomorphism can be extended to an isomorphism from  $\mathfrak{A}$  to  $\mathfrak{B}$ .

**COROLLARY 2.13.** *Let  $\theta \in L_{\omega_1 \omega_1}^{m * \omega}$  be a sentence. Then  $\Sigma \models \theta \leftrightarrow \phi$ , where  $\phi \in B(H)$ ,  $H = \{DC_c(\mathfrak{C}, D) \mid \mathfrak{C} \text{ countable}, D \subset C, \text{ and } c \text{ a map from } \omega \text{ one-one onto } C\}$ .*

*Proof.* Clearly  $S = \{\Sigma\}$  is a consistent set of sentences in  $L_{\omega_1 \omega_1}^{m * \omega}$ , and  $H$  is also a set of sentences in  $L_{\omega_1 \omega_1}^{m * \omega}$ . By Theorem 2.12, condition (1) of Lemma 2.8 is satisfied. Condition (2) follows from the fact that any separable metric space with a countable number of constants has a countable dense subsystem containing those constants. If  $D$  is that set of constants, and  $\mathfrak{C}$  is that system, then  $\mathfrak{C} \models DC_c(\mathfrak{C}, D)$  for any mapping of  $\omega$  one-one onto  $C$ . Thus our Corollary follows from (c) of that lemma.

We now need the concept of substitution for free occurrences of variable and for constants in a formula. Intuitively, by

$$Sb\{y_i/i \in I\}, \{x_i/i \in I\}\theta$$

we mean the formula obtained by substituting  $y_i$  for  $x_i$  at each free occurrence of  $x_i$  or constant  $x_i$ , and leaving unchanged other terms in  $\theta$ . Somewhat more formally, we consider the function  $f: V \cup C \rightarrow V \cup C$  (see Definition 1.1, [5]) such that for each  $i \in I, f(x_i) = y_i$ , and if  $x_i \in V \cup C, i \notin I, f(x_i) = x_i$ , and:

**DEFINITION 2.14.**  $Sb_f \theta$  is the formula defined inductively as follows:

- (1)  $Sb_j(x_i \equiv x_j) = f(x_i) \equiv f(x_j)$ ,
- (2) If  $j \in (V \cup C)^{\xi(p)}$ , then  $Sb_f(P_p(x \circ j)) = P_p(f \circ x \circ j)$ ,



$$(3) \quad Sb_f(\neg \theta) = \neg Sb_f\theta,$$

$$(4) \quad Sb_f(\mathbf{V} X) = \mathbf{V} \{Sb_f\theta/\theta \in X\},$$

$$(5) \quad Sb_f(\mathbf{\wedge} X) = \mathbf{\wedge} \{Sb_f\theta/\theta \in X\},$$

$$(6) \quad Sb_f(\mathbf{\forall} W)\theta = (\mathbf{\forall} W)Sb_f\theta, \text{ where } f'(x) = f(x) \text{ for } x \in V \cup C - W, \\ f'(x) = x \text{ if } x \in W,$$

$$(7) \quad Sb_f(\mathbf{\exists} W)\theta = (\mathbf{\exists} W)Sb_f\theta.$$

We also define  $Sb(\{y_i/i \in I\}, \{x_i/i \in I\})\theta = Sb_f\theta$  with  $f$  as constructed in the last sentence before our definition.

An immediate consequence of the definition is the fact that if  $\mathfrak{A} \models Sb(\{y_i/i \in I\}, \{x_i/i \in I\})(\theta \rightarrow \phi)$  and none of the  $x_i$  occur in  $\theta$ , then  $\mathfrak{A} \models \theta \rightarrow Sb(\{y_i/i \in I\}, \{x_i/i \in I\})\phi$ . Also, if a sentence  $\theta \models \phi$  and none of the  $x_i$  occur in  $\theta$ , then  $\theta \models Sb(\{y_i/i \in I\}, \{x_i/i \in I\})\phi$ .

**THEOREM 2.15.** *If  $F \in L_{\omega_1}^m$  is a formula, then  $\Sigma \models F \leftrightarrow \theta$  for some  $\theta \in B(E')$ , where  $\phi \in E'$  if and only if*

$$\phi = Sb(\{v_i/i \in \omega\}, \{e_i/i \in \omega\})DC(\mathfrak{C}, D),$$

with  $\mathfrak{C}, D$  as in Definition 2.11,  $D = \{e_i/i \in \omega\}$ .

*Proof.* We can always put  $F$  in an equivalent form  $F'$  such that  $\{v_i/i \in \omega\}$  is the set of free variables of  $F'$ . Thus we lose no generality in assuming  $\{v_i/i \in \omega\}$  is the set of free variables of  $F$ . Now consider the sentence  $\phi = Sb(\{e_i/i \in \omega\}, \{v_i/i \in \omega\})F \in L_{\omega_1}^{m*\omega}$ . For some  $\theta \in B(H)$ ,  $\Sigma \models \phi \leftrightarrow \theta$ . Since  $\Sigma$  contains no reference to the  $e_i$ , we also have  $\Sigma \models Sb(\{v_i/i \in \omega\}, \{e_i/i \in \omega\})(\phi \leftrightarrow \theta)$ , thus  $\Sigma \models F \leftrightarrow Sb(\{v_i/i \in \omega\}, \{e_i/i \in \omega\})\theta$ . But by (3), (4), and (5) of Definition 2.14,  $Sb(\{v_i/i \in \omega\}, \{e_i/i \in \omega\})\theta \in B(E')$ .

Thus we have an elimination of quantifiers in the theory of complete separable metric systems for all formulas of  $L_{\omega_1}^m$ . This elimination does not always take place within that language. We shall proceed to universal equivalence for complete metric systems. Using Corollary 2.3, Tarski has shown that two structures are universally equivalent in  $L_{\pi}^t$  (in symbols  $\mathfrak{A} \equiv_{U^t_{\pi}} \mathfrak{B}$ ) if and only if  $\text{Iso Sub}_x(\mathfrak{C}) = \text{Iso Sub}_x(\mathfrak{B})$ . Thus if we allow  $U = U_{\omega_1}^m$ , we have the fact that  $\mathfrak{A} \equiv_U \mathfrak{B}$  if and only if  $\text{Iso Sub}_{\omega_1}(\mathfrak{A}) = \text{Iso Sub}_{\omega_1}(\mathfrak{B})$  for any two metric systems.

**DEFINITION 2.16.** Let  $K$  be a class of complete metric systems. Then  $SC(K) = \{\mathfrak{B}/\text{for some } \mathfrak{A} \in K, \mathfrak{B} \subset \mathfrak{A} \text{ and } \mathfrak{B} \text{ is a separable complete metric system}\}$ .  $ISC(K) = \{\mathfrak{D}/\mathfrak{D} \cong \mathfrak{B} \text{ for some } \mathfrak{B} \in SC(K)\}$ .

$$ISC(\{\mathfrak{A}\}) = ISC(\mathfrak{A}).$$

**THEOREM 2.17.** *Let  $\mathfrak{A}, \mathfrak{B}$  be complete metric systems. Then  $\mathfrak{A} \equiv_U \mathfrak{B}$*

if and only if  $ISC(\mathfrak{A}) = ISC(\mathfrak{B})$ .

*Proof.* Assume first that  $\text{Iso Sub}_{\omega_1}(\mathfrak{A}) = \text{Iso Sub}_{\omega_1}(\mathfrak{B})$  (i.e., that  $\mathfrak{A} \equiv_v \mathfrak{B}$ ). Let  $\mathfrak{C} \in ISC(\mathfrak{A})$ . Since  $\mathfrak{C}$  is separable and complete there is a countable  $\mathfrak{D} \subset \mathfrak{A}$  such that  $\mathfrak{D}$  is dense in an isomorphic copy of  $\mathfrak{C}$ . By our assumption, there is a countable  $\mathfrak{E} \subset \mathfrak{B}$  such that  $\mathfrak{E} \cong \mathfrak{D}$ . Since  $\mathfrak{B}$  is complete, the completion of  $\mathfrak{E}$  is contained in  $\mathfrak{B}$  (by previous remarks, this is isomorphic to  $\mathfrak{C}$ ). Thus  $ISC(\mathfrak{A}) \subset ISC(\mathfrak{B})$ . Similarly,  $ISC(\mathfrak{B}) \subset ISC(\mathfrak{A})$ .

Conversely, assume  $ISC(\mathfrak{A}) = ISC(\mathfrak{B})$ . Let  $\mathfrak{C} \in \text{Iso Sub}_{\omega_1}(\mathfrak{A})$ . Since  $\mathfrak{A}$  is complete, the completion of an isomorphic copy of  $\mathfrak{C}$  is contained in  $\mathfrak{A}$ , thus another isomorphic copy of its completion is contained in  $\mathfrak{B}$ . Thus an isomorphic copy of  $\mathfrak{C}$  is contained in  $\mathfrak{B}$ . This shows that  $\text{Iso Sub}_{\omega_1}(\mathfrak{A}) \subset \text{Iso Sub}_{\omega_1}(\mathfrak{B})$ , and the reverse inclusion is shown similarly.

**COROLLARY 2.18.** *Let  $\mathfrak{A}, \mathfrak{B}$  be complete separable metric systems. Then  $\mathfrak{A} \equiv_v \mathfrak{B}$  if and only if for some  $\mathfrak{C} \subset \mathfrak{B}, \mathfrak{A} \cong \mathfrak{C}$  and for some  $\mathfrak{D} \subset \mathfrak{A}, \mathfrak{B} \cong \mathfrak{D}$ .*

Universal equivalence, even for complete separable spaces, is not as strong as elementary equivalence. The following example, when formalized, provides a counterexample. Let

$$N = \{f: \omega \rightarrow \mathbf{R} / \sum_{i=1}^{\infty} (f(i))^2 < \infty\},$$

where  $\mathbf{R}$  is the real numbers. Define  $m(f, g) = \sqrt{\sum_{i=1}^{\infty} (f(i) - g(i))^2}$ . Then  $m$  can be shown to be a metric on  $N$ . Let

$$S = \{f \in N / 0 \leq f(1) \leq 1\}.$$

Then  $S \subset N$  and the following isometry takes  $N$  into  $S$ . Let  $F: N \rightarrow S$  by  $F(f)(i) = f(i - 1), i > 1$  and  $F(f)(1) = 0$ .

If  $\mathfrak{B}$  is the metric system associated with  $N, \mathfrak{A}$  that associated with  $S$ , then  $\mathfrak{A}, \mathfrak{B}$  are complete and separable,  $\mathfrak{A} \subset \mathfrak{B}$ , and  $\mathfrak{B} \cong \mathfrak{C} \subset \mathfrak{A}$ , but not  $\mathfrak{A} \cong \mathfrak{B}$ . If we take  $S' = \{f \in N / f(0) \in Q\}$  ( $Q$  the set of rationals), we have an example (using  $S', N$ ) of a sepeable complete metric system universally equivalent to a separable incomplete system. There are also examples of separable systems universally equivalent to inseparable systems.

**III. Metric algebraic systems.** The theory of Banach Algebras (see Naimark) has been one of the most important in analysis during the past twenty years. The theories of metric groups (see Montgomery-Zippin), Banach Spaces (see Dunford-Schwartz), normed spaces (see Day), and many others have been essential to functional analysis.

Measure theory can, in many respects, be reduced to the theory of certain metric Boolean algebras, c.f. [4, pp. 165-174]. All these theories contain a common feature: they are theories of algebraic structures with metric topologies. Most of them share a second feature: they are defined in terms of a norm rather than a metric. A metric, however, can easily be defined by allowing  $m(a, b) = \|a - b\|$ . In this section we will extend the results obtained for metric spaces to algebraic structures with metric topologies.

DEFINITION 3.1. A metric algebraic system is a structure

$$\mathfrak{A} = \langle A, M, Q, +, 0, 1, <, d, R_p \rangle_{p < \omega}$$

with  $o \leq \omega$  and:

- (a1)  $\mathfrak{A}/m$  is a metric system.
- (a2) If  $o = t(p)$ , we have:  $M(c_p)$ . If  $0 < t(p)$ , we have:

$$(\forall v/t(p))(R_p(v/t(p)) \rightarrow \bigwedge_{i < t(p)} M(v_i)) .$$

- (a3)  $(\forall v/t(p))(\exists v_{2t(p)})(\forall t(p)/v/2t(p))(R_p(v/t(p)) \vee \neg [0 < v_{2t(p)} \wedge (R_p(t(p)/v/2t(p)) \wedge (\bigwedge_{i < t(p)} d(v_i, v_{t(p)+i}, v_{2t(p)}))]) .$

(For notation used here see [5], discussion following Def. 1.1.)

In the above axioms, (a2) simply restricts the relations (and constants) to the metric space, while (a3) closes all the relations (for the definition of a closed relation extend Dunford-Schwartz, p. 57, Def. 3 in the obvious manner). Since continuous operations are closed as relations, it is clear that metric topological groups, etc. can be considered by use of the above axioms. Banach Spaces can also be considered, despite the fact that they are defined by use of two metric spaces, rather than one. We simply work with the disjoint union of the universes of the metric spaces involved, and define a metric on that set by setting it equal to the appropriate of the two existing metrics for two elements of the same set, and setting it equal to 1 otherwise. Both sets now become closed and we proceed with the axiomatization in the obvious manner. The remaining systems mentioned at the beginning of this section can similarly be axiomatized as metric algebraic systems.

DEFINITION 3.2. Let  $\mathfrak{A}$  be a metric algebraic system, and let  $W$  be a predicate defined in terms of the relations of  $\mathfrak{A}/m$ . Then  $\mathfrak{A}$  is said to have the property  $W$  if and only if  $\mathfrak{A}/m$  satisfies  $W$ .

In the above definition,  $W$  could define the property of being separable, complete, etc. Note here that any substructure of a metric algebraic system is itself a metric algebraic system, provided it remains a metric system.

DEFINITION 3.3. Let  $A' \subset R_p$ . Then  $A'$  is dense in  $R_p$  if and only if for any  $\langle a_0, \dots, a_{t(p)-1} \rangle \in R_p$  and any  $q > 0$ , there is an

$$\langle a'_0, \dots, a'_{t(p)-1} \rangle \in A'$$

such that for each  $j < t(p)$ ,  $\langle a_j, a'_j, q \rangle \in d$ . A metric algebraic subsystem  $\mathbb{C}$  of a metric algebraic system  $\mathfrak{A}$  is called dense in  $\mathfrak{A}$  if and only if  $\mathbb{C}/m$  is dense in  $\mathfrak{A}/m$  and the restriction of each  $R_p$  to  $C$  is dense in  $R_p$ .

Thus a subset of a relation is called dense here if and only if it is dense in the product topology. The above definition is the "correct" one for density of subsystems, as shown by the following fact: we can now make Definition, Theorem, Lemma, or Corollary 3.x ( $x \geq 4$ ) from the corresponding Definition, Theorem, Lemma, or Corollary 2.x by making the following alterations (where necessary) in their texts: change "metric system" to "metric algebraic system of type  $t$ " and change " $m$ " (the type of metric systems) to " $t$ ". The proofs must be altered somewhat, although only one (done below) creates any problem in the new setting. Finally, the meaning of the words has been changed somewhat. Our new  $DC_c(\mathbb{C}, D)$ , for example, now refers to all the relations in our system, rather than only the metric relations. A separable Banach Space, for example, is determined by the relation of vector sum (as well as norm, i.e., our metric) and scalar product on a dense countable subset.

LEMMA 3.5. (Corresponding to Lemma 2.5). If  $\mathfrak{A}$  is a metric algebraic system of type  $t$ ,  $\mathfrak{A} \models \text{Sep}$  if and only if for some countable metric algebraic system of type  $t$ ,  $\mathbb{C}$ ,  $\mathfrak{A} \models Dc(\mathbb{C})$ .

*Proof.* First note that if  $N_p$  is any countable subset of  $A^{t(p)}$ , then there is a countable  $C_p \subset A$  such that  $N_p \subset C_p^{t(p)}$  (for example, let  $C_p$  be the collection of all points which are any coordinate for any point of  $N_p$ ). Since the domain  $o$  of  $t$  is countable, we can take  $C = \bigcup_{p < o} C_p \cup Q$ , and let  $\mathbb{C}$  be  $C$  together with the restrictions of all the relations of  $\mathfrak{A}$ . Then  $\mathbb{C}$  is clearly countable and  $\mathfrak{A} \models Dc(\mathbb{C})$ .

A remark left implicit in § II is that two separable complete systems are isomorphic if and only if they contain isomorphic dense sub-

systems. This remains true here only because our relations are required to be closed.

#### BIBLIOGRAPHY

1. M. Day *Normed Linear Spaces*, Springer Verlag, Berlin, 1958.
2. J. Dieudonne, *Foundations of Mathematical Analysis*, Academic Press, 1960.
3. N. Dunford, and J. Schwartz, *Linear Operators, Part I*, Interscience Publishers, Inc., New York, 1958.
4. P. R. Halmos, *Measure Theory*, D. van Nostrand, Inc., Princeton, 1950.
5. R. Kopperman, *The  $L_{\omega_1\omega_1}$ -theory of Hilbert spaces*, (to appear in the Journal of Symbolic Logic).
6. ———, *On the axiomatizability of uniform spaces*, (To appear in the Journal of Symbolic Logic).
7. D. Montgomery, and L. Zippin, *Topological Transformation Groups*, Interscience Publishers, Inc., New York, 1955.
8. M. Naimark, *Normed Rings*, Translated by Boron, L. P. Noordhoff, NV, 1959.
9. A. Robinson, *Complete Theories*, North Holland Publishing Company, Amsterdam, 1956.
10. A. Tarski, *Remarks on predicate logic with infinitely long expressions*, Colloquium Mathematicum, VI (1958), pp. 171-176.

Received April 20, 1966. This research was supported in part by National Science Foundation Grants GP-4361 and GP-5710.

UNIVERSITY OF RHODE ISLAND  
AND  
CITY COLLEGE OF NEW YORK

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN  
Stanford University  
Stanford, California

J. DUGUNDJI  
Department of Mathematics  
Rice University  
Houston, Texas 77001

J. P. JANS  
University of Washington  
Seattle, Washington 98105

RICHARD ARENS  
University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Herbert Stanley Bear, Jr. and Bertram John Walsh, <i>Integral kernel for one-part function spaces</i> .....	209
Mario Borelli, <i>Some results on ampleness and divisorial schemes</i> .....	217
John A. Erdos, <i>Unitary invariants for nests</i> .....	229
Nathaniel Grossman, <i>The volume of a totally-geodesic hypersurface in a pinched manifold</i> .....	257
D. M. Hyman, <i>A generalization of the Borsuk-Whitehead-Hanner theorem</i> .....	263
I. Martin (Irving) Isaacs, <i>Finite groups with small character degrees and large prime divisors</i> .....	273
I. Martin (Irving) Isaacs, <i>Two solvability theorems</i> .....	281
William Lee Johnson, <i>The characteristic function of a harmonic function in a locally Euclidean space</i> .....	291
Ralph David Kopperman, <i>Application of infinitary languages to metric spaces</i> .....	299
John Lauchlin MacDonald, <i>Relative functor representability</i> .....	311
Mahendra Ganpatrao Nadkarni, <i>A class of measures on the Bohr group</i> .....	321
Keith Lowell Phillips, <i>Hilbert transforms for the <math>p</math>-adic and <math>p</math>-series fields</i> .....	329
Norman R. Reilly and Herman Edward Scheiblich, <i>Congruences on regular semigroups</i> .....	349
Neil William Rickert, <i>Measures whose range is a ball</i> .....	361
Gideon Schwarz, <i>Variations on vector measures</i> .....	373
Ronald Cameron Riddell, <i>Spectral concentration for self-adjoint operators</i> .....	377
Haskell Paul Rosenthal, <i>A characterization of restrictions of Fourier-Stieltjes transforms</i> .....	403