

Pacific Journal of Mathematics

VARIATIONS ON VECTOR MEASURES

GIDEON SCHWARZ

VARIATIONS ON VECTOR MEASURES

GIDEON SCHWARZ

Let μ be a signed measure, and denote the total measure of its positive and negative parts by P and N . Since the total variation of such a measure is $V = P + |N|$, and the maximum of the absolute value of the measure is $M = \max(P, |N|)$, we have the inequality $M \leq V \leq 2M$. We consider the following question.

What should replace the constant 2 in this inequality when we pass to higher-dimensional vector-valued measures?

This question has been answered by Kaufman and Rickert [1]. In addition to their own result, they describe a geometric proof of the two-dimensional case, due to Kakutani, and state that they know of no geometric proof for $n \geq 3$. In this note we extend Kakutani's proof to all dimensions. The crucial step in his proof is the observation that the total variation of a 2-dimensional vector measure is proportional to the circumference of the convex hull of its range. The "obvious" attempt to generalize to higher dimensions by replacing "circumference" by "surface area" must fail, as a simple dimension analysis shows. The generalization succeeds, however, if we first replace "circumference" in Kakutani's observation by "average width".

For easy reference, we recall the definition of total variation:

The total variation of a vector measure μ is the supremum over all partitions $\{E_1, \dots, E_n\}$ of the measure space of $\sum_{i=1}^n \|\mu(E_i)\|$, where $\|\cdot\|$ is Euclidean length.

THEOREM. *The total variation of an n -vector measure equals c_n times the average width of the range of the measure where $c_{2n} = 4^{-n}n(2n)! \pi/(n!)^2$ and $c_{2n+1} = 4^n(n!)^2/(2n)!$.*

Proof. If the measure space consists of a single atom, the range of the measure consists of two vectors: the vector \mathbf{a} assigned to the whole space, and the zero vector. The total variation is in this case just $\|\mathbf{a}\|$, the length of \mathbf{a} , and the average width of the two-point set is seen to be $\|\mathbf{a}\|/c_n$ by a simple calculation; therefore the theorem holds for a trivial measure space. Since direct sum formation of measure spaces gives rise to (group theoretic) addition of the ranges, their support functions undergo addition as well (see [4]), and so do their average widths. Clearly, total variation is additive as well, and

the theorem follows for measure spaces with finitely many elements. By continuity, of widths and variations, it follows for arbitrary n -vector valued measures.

COROLLARY 1. (*Kaufman and Rickert*). *There exists a set in the measure space, the length of whose measure is at least $(2c_n)^{-1}$ times the total variation V .*

Proof. The maximal width M of the range is at most twice the length L of the longest vector in the range, yet it cannot be less than the average width A . Therefore $Vc_n^{-1} = A \leq M \leq 2L$.

For $n \geq 2$ we also have

COROLLARY 2. *If and only if there is no set the length of whose measure exceeds $(2c_n)^{-1}V$, the range of the measure is a ball centered at the origin.*

Proof. The "only if" is trivial. For the "if": When the maximal width does not exceed the average width, the width must be constant, and the closed convex hull of range is a ball. When the longest vector does not exceed half the maximal width, the ball must be centered at zero.

If the measure space is nonatomic, the range is closed and convex by a well known theorem of Liapounoff [2], and is therefore itself a ball centered at zero.

If, on the other hand, there is an atom in the measure space, the convex closure of the range is the group theoretic sum of an interval and a closed convex set. Hence its boundary contains an interval, and in 2 or more dimensions it cannot be a ball.

REMARK 1. After applying a vector form of the Radon-Nikodym theorem (see the preceding paper [5] in this journal), these results can be translated from vector measures to probabilities and yield the following: If U is an n -dimensional unit vector valued random variable on a probability space, there is at least one event B such that

$$E(U | B)P(B) \geq (2c_n)^{-1} .$$

REMARK 2. The numbers c_n occur in a bound, calculated by A. E. Mayer [3], for the diameter of a polyhedron with a one-dimensional skeleton of given total length.

REFERENCES

1. R. P. Kaufman and N. W. Rickert, *An inequality concerning measures*, Bull. Amer. Math. Soc. **72** (1966), 672-676.
2. A. Liapounoff, *Sur les fonctions—vecteurs complètement additives*, Bull. Acad. Sci. URSS Sér. Math. **4** (1940), 465-478. (Russian, French summary).
3. A. E. Mayer, *Grösste Polygone mit gegebenen Seitenvektoren*, Com. Math. Helvetici. **10** (1938), 288-301.
4. H. Minkowski, *Theorie der Konvexen Körper, insbesondere Begründung ihres Oberflächenbegriffs*, Ges. Abh. 2, Leipzig und Berlin, 1911, 131-229.
5. N. W. Rickert, *Measures whose range is a ball*, Pacific J. Math. **23** (1967), 361-371.

Received May 3, 1967.

UNIVERSITY OF CALIFORNIA, BERKELEY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
Rice University
Houston, Texas 77001

J. P. JANS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.

Herbert Stanley Bear, Jr. and Bertram John Walsh, <i>Integral kernel for one-part function spaces</i>	209
Mario Borelli, <i>Some results on ampleness and divisorial schemes</i>	217
John A. Erdos, <i>Unitary invariants for nests</i>	229
Nathaniel Grossman, <i>The volume of a totally-geodesic hypersurface in a pinched manifold</i>	257
D. M. Hyman, <i>A generalization of the Borsuk-Whitehead-Hanner theorem</i>	263
I. Martin (Irving) Isaacs, <i>Finite groups with small character degrees and large prime divisors</i>	273
I. Martin (Irving) Isaacs, <i>Two solvability theorems</i>	281
William Lee Johnson, <i>The characteristic function of a harmonic function in a locally Euclidean space</i>	291
Ralph David Kopperman, <i>Application of infinitary languages to metric spaces</i>	299
John Lauchlin MacDonald, <i>Relative functor representability</i>	311
Mahendra Ganpatrao Nadkarni, <i>A class of measures on the Bohr group</i>	321
Keith Lowell Phillips, <i>Hilbert transforms for the p-adic and p-series fields</i>	329
Norman R. Reilly and Herman Edward Scheiblich, <i>Congruences on regular semigroups</i>	349
Neil William Rickert, <i>Measures whose range is a ball</i>	361
Gideon Schwarz, <i>Variations on vector measures</i>	373
Ronald Cameron Riddell, <i>Spectral concentration for self-adjoint operators</i>	377
Haskell Paul Rosenthal, <i>A characterization of restrictions of Fourier-Stieltjes transforms</i>	403