

# Pacific Journal of Mathematics

**CRITERIA FOR ZERO CAPACITY OF IDEAL BOUNDARY  
COMPONENTS OF RIEMANNIAN SPACES**

WELLINGTON HAM OW

# CRITERIA FOR ZERO CAPACITY OF IDEAL BOUNDARY COMPONENTS OF RIEMANNIAN SPACES

WELLINGTON H. OW

Capacities of ideal boundary components of Riemannian spaces are introduced to measure their magnitude with respect to harmonic functions on the spaces. The main purpose of this paper is to find zero capacity criteria.

The modular criterion, well-known for Riemann surfaces, i.e. for 2-dimensional Riemannian spaces, is shown to be valid for general Riemannian spaces. The so-called metric criterion, however, brings forth entirely new aspects for higher dimensions.

## CAPACITY OF A SUBBOUNDARY

1. **Subboundaries.** Throughout this paper we denote by  $R$  a non-compact orientable connected  $C^\infty$  Riemannian space. A relatively compact region whose relative boundary is smooth will be called a regular region. A sequence  $\{R_n\}_1^\infty$  of regular regions  $R_n \subset R$  such that  $\bar{R}_n \subset R_{n+1}$  and  $R = \bigcup_1^\infty R_n$  is called an exhaustion of  $R$ .

The ideal boundary component of Kerékjártó and Stoilow may be related to  $\{R_n\}$ ; here  $R - \bar{R}_n$  can be assumed to consist of a finite number of relatively noncompact regions  $F_{ni}$  with corders  $\beta_{ni}$ . Choose a sequence  $F_1 = F_{1i_1}, F_2 = F_{2i_2}, \dots$  such that  $\bar{F}_{n+1} \subset F_n$ . Then  $\{F_n\}_1^\infty$  defines a boundary component  $\gamma$ . We denote by  $\gamma_n$  the relative boundary  $\partial F_n$  of  $F_n$ .

A *subboundary*, also to be denoted by  $\gamma$ , is a union of boundary components.

2. **Capacity function.** Let  $B$  be a parametric ball about  $a \in R$  with compact  $\bar{B}$ . Suppose  $\gamma$  is a subboundary of  $R$ , and  $\gamma_n$  the union of all  $\partial F_n$  such that  $\{F_n\}_1^\infty$  defines a boundary component belonging to  $\gamma$ .

Consider the family  $P = \{p\}$  of harmonic functions  $p$  on  $R - a$  such that (a)  $p = -g_a + h$  in  $B$  where  $g_a$  is the Green's function of  $B$  with pole at  $a$ , and  $h$  a harmonic function on  $B$  with  $h(a) = 0$ , (b)  $\int_{\gamma_n} * dp = 1$  and  $\int_{\beta_{ni} \cap \gamma_n} * dp = 0$  for large  $n$ , where the  $\beta_{ni}$  are components of  $\partial R_n$ .

We use the conventional notations

$$\int_\gamma * dp = \lim_{n \rightarrow \infty} \int_{\gamma_n} * dp .$$

$$\int_{\beta} p * dp = \lim_{n \rightarrow \infty} \int_{\beta_n} p * dp ,$$

$\beta$  being the entire boundary of  $R$ .

Amalgamating the method of Sario [1] with the existence theorem of principal functions in Sario-Schiffer-Glasner [2], we can easily see that  $P$  is not empty and that there exists a function  $p_{\gamma} \in P$  such that

$$(1) \quad k_{\gamma} = \min_P \int_{\beta} p * pd = \int_{\beta} p_{\gamma} * dp_{\gamma} .$$

Here  $-\infty < k_{\gamma} \leq \infty$ , and if  $k_{\gamma} < \infty$ , then  $p_{\gamma}$  is unique. This follows from the identity

$$(2) \quad \int_{\beta} p * dp = D(p - p_{\gamma}) + \int_{\beta} p_{\gamma} * dp_{\gamma} ,$$

where  $D$  indicates the Dirichlet integral.

The function  $p_{\gamma}$  shall be referred to as a *capacity function* for  $\gamma$ . The quantity  $c_{\gamma} = e^{-k_{\gamma}}$  for  $\dim R = 2$ , and  $k_{\gamma}^{-(m-2)}$  for  $\dim R = m \geq 3$  will be called the *capacity* of  $\gamma$ .

### MODULAR CRITERION

3. **Moduli.** Let  $\Omega$  be a union of disjoint regular regions  $\Omega_j$ ,  $j = 1, \dots, k$ . Suppose that  $\partial\Omega_j$  consists of two nonempty disjoint sets  $\beta'_j$  and  $\beta''_j$  which are unions of components of  $\partial\Omega_j$ . Set  $\beta' = \bigcup_1^k \beta'_j$  and  $\beta'' = \bigcup_1^k \beta''_j$ . Let  $u_0$  be the continuous function on  $\bar{\Omega}$  which is harmonic on  $\Omega$  with  $u_0|_{\beta'} = 0$ ,  $u_0|_{\beta''} = \log \mu$ , and  $\int_{\beta'} * du_0 = 1$ . The constant  $\mu > 1$  is called the *modulus* of the configuration  $(\Omega, \beta', \beta'')$ ,

$$\mu = \text{mod}(\Omega, \beta', \beta'') .$$

The function  $u_0$  is referred to as the *modulus function*.

Consider the family  $U = U(\Omega, \beta', \beta'')$  of  $C^1$ -functions  $u$  on  $\bar{\Omega}$  which are harmonic on  $\Omega$  with  $\int_{\beta'} * du = 1$ . Then we have

$$(3) \quad \min_U D_{\Omega}(u) = D_{\Omega}(u_0) .$$

This follows from the identity

$$(4) \quad D_{\Omega}(u) = D_{\Omega}(u_0 - u) + D_{\Omega}(u_0)$$

for every  $u \in U$ .

4. **Modular criterion.** Let  $\tilde{F}_n$  be the sum of those  $F_n$  for which  $\{F_n\}_1^{\infty}$  defines a boundary component in the subboundary  $\gamma$ . Consider

$E_n = (R_{n+1} - \bar{R}_n) \cap \tilde{F}_n$  and set  $\gamma_n = \partial F_n$ ,  $\gamma'_n = \partial E_n - \gamma_n$ . In terms of

$$(5) \quad \mu_{n\gamma} = \text{mod}(E_n, \gamma_n, \gamma'_n)$$

we state:

**THEOREM 1.** *If there exists an exhaustion of  $R$  with*

$$(6) \quad \prod_{n=1}^{\infty} \mu_{n\gamma} = \infty,$$

*then the capacity of  $\gamma$  vanishes.*

In fact, let  $p_n$  and  $k_n$  stand for  $p_\gamma$  and  $k_\gamma$  with respect to  $\gamma_n$  and  $R_n$ . By (1) we infer that

$$D_{R_{n+1} - \bar{R}_n}(p_{n+1}) \leq k_{n+1} - k_n,$$

and by (3) that

$$\log \mu_{n\gamma} \leq D_{E_n}(p_{n+1}).$$

Therefore  $\log \mu_{n\gamma} \leq k_{n+1} - k_n$ , and we conclude that (6) implies

$$\lim_{n \rightarrow \infty} k_n = \infty.$$

On the other hand it is not difficult to see that  $k_\gamma = \lim_n k_n$ , whence  $c_\gamma = 0$ .

### METRIC CRITERION

**5. Conformally equivalent metric.** Let  $\lambda$  be a positive  $C^\infty$ -function on  $R$ . The new metric

$$(7) \quad d\sigma = \lambda ds$$

is conformally equivalent to the original metric  $ds$  on  $R$ . We fix a point  $a \in R$  and assume that

$$(8) \quad R(r) = \{x \in R \mid \sigma(x, a) < r\}$$

is relatively compact in  $R$  for  $0 < r < \infty$ , with  $R = \bigcup_{0 < r < \infty} R(r)$ . Consider the minimal union  $\gamma(r)$  of components of  $\beta(r) = \partial R(r)$  which separates  $\gamma$  from  $a$ . Let

$$(9) \quad S_\gamma(r) = \int_{\gamma(r)} dS_\sigma,$$

where  $dS_\sigma$  is the surface element induced by  $d\sigma$ .

**THEOREM 2.** *If there exists an admissible  $\lambda$  such that*

$$(10) \quad \int_{\varepsilon}^{\infty} \frac{dr}{S_r(r)} = \infty \quad (\varepsilon > 0),$$

then  $\gamma$  has vanishing capacity for  $R$  with  $\dim R = 2$ . If, moreover, there exists a constant  $M$  such that

$$(11) \quad 0 < \frac{1}{M} \leq \lambda \leq M,$$

then the same conclusion holds regardless of the dimension of  $R$ .

For the proof we choose a sequence  $\{r_n\}_1^{\infty}$  such that  $\varepsilon < r_n < r_{n+1} < \infty$  and  $\lim_n r_n = \infty$ , with  $R_n = R(r_n)$  regular. As in §4 we define  $E_n = (R_{n+1} - \bar{R}_n) \cap \bar{F}_n$  and  $\mu_{n\gamma}$ . We also denote by  $u_n$  the corresponding modulus function.

The proof in the case  $\dim R = 2$  will be given in §6 and that in the general case under the assumption (11), in §7.

6. The case  $\dim R = 2$ . Observe that

$$(12) \quad \int_{E_n} |\nabla_{\sigma} u_n|^2 dV_{\sigma} = \int_{r_n}^{r_{n+1}} \left[ \int_{\gamma(r)} |\nabla_{\sigma} u_n|^2 dS_{\sigma} \int_{\gamma(r)} dS_{\sigma} \right] \frac{dr}{S_r(r)}.$$

By the Schwarz inequality we have

$$(13) \quad \int_{\gamma(r)} |\nabla_{\sigma} u_n|^2 dS_{\sigma} \int_{\gamma(r)} dS_{\sigma} \geq \left( \int_{\gamma(r)} *_\sigma du_n \right)^2.$$

Since  $*_{\sigma} = *$  and  $|\nabla_{\sigma} u_n|^2 dV_{\sigma} = |\nabla u_n|^2 dV$ , it is seen that (12), (13), and  $D_{E_n}(u_n) = \log \mu_{n\gamma}$  imply

$$(14) \quad \log \prod_1^n \mu_{k\gamma} \geq \int_{r_1}^{r_{n+1}} \frac{dr}{S_r(r)}.$$

We conclude that (10) implies (6), and consequently  $c_{\gamma} = 0$ .

7. The case  $\dim R = m > 2$ . By (11) we see that

$$(15) \quad \int_{E_n} |\nabla_{\sigma} u_n|^2 dV_{\sigma} \leq M^{m-2} D_{E_n}(u_n),$$

$$(16) \quad \int_{\gamma(r)} *_\sigma du_n \geq M^{-(m-2)} \int_{\gamma(r)} * du_n.$$

Therefore (14) must be modified to give

$$\log \prod_1^n \mu_{k\gamma} \geq M^{-3(m-2)} \int_{r_1}^{r_{n+1}} \frac{dr}{S_r(r)}.$$

But this sufficient to conclude that  $c_i = 0$ .

REMARK. Condition (11) cannot be suppressed in the case of higher dimensions.

The author is greatly indebted to Professor Leo Sario, chairman of his doctoral committee, who guided his research, and also to Professor Mitsuru Nakai, with whom the author had many invaluable discussions.

#### REFERENCES

1. L. Sario, *Capacity of the boundary and of a boundary component*, Ann. of Math. **59** (1954), 135-144.
2. L. Sario, M. Schiffer, and M. Glasner, *The span and principal functions in Riemannian spaces*, J. Analyse Math. **15** (1965), 115-134.

Received January 30, 1967. This work was sponsored in part by the U. S. Army Research Office-Durham, Grant DA-AROD-31-124-G742. The results of this paper are part of the author's dissertation at the University of California, Los Angeles.

UNIVERSITY OF CALIFORNIA, LOS ANGELES



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN

Stanford University  
Stanford, California

J. DUGUNDJI

Department of Mathematics  
Rice University  
Houston, Texas 77001

J. P. JANS

University of Washington  
Seattle, Washington 98105

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL ORDNANCE TEST STATION

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be typewritten (double spaced). The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens at the University of California, Los Angeles, California 90024.

50 reprints per author of each article are furnished free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$ 8.00; single issues, \$ 3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$ 4.00 per volume; single issues \$ 1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.



|  |     |
|--|-----|
| A. A. Aucoin, <i>Diophantine systems</i> .....   | 419 |
| Charles Ballantine, <i>Products of positive definite matrices. I</i> .....   | 427 |
| David Wilmot Barnette, <i>A necessary condition for d-polyhedrality</i> .....  | 435 |
| James Clark Beidleman and Tae Kun Seo, <i>Generalized Frattini subgroups of finite groups</i> .....  | 441 |
| Carlos Jorge Do Rego Borges, <i>A study of multivalued functions</i> .....   | 451 |
| William Edwin Clark, <i>Algebras of global dimension one with a finite ideal lattice</i> .....   | 463 |
| Richard Brian Darst, <i>On a theorem of Nikodym with applications to weak convergence and von Neumann algebras</i> .....   | 473 |
| George Wesley Day, <i>Superatomic Boolean algebras</i> .....   | 479 |
| Lawrence Fearnley, <i>Characterization of the continuous images of all pseudocircles</i> .....   | 491 |
| Neil Robert Gray, <i>Unstable points in the hyperspace of connected subsets</i> .....  | 515 |
| Franklin Haimo, <i>Polynomials in central endomorphisms</i> .....  | 521 |
| John Sollion Hsia, <i>Integral equivalence of vectors over local modular lattices</i> .....  | 527 |
| Jim Humphreys, <i>Existence of Levi factors in certain algebraic groups</i> .....  | 543 |
| E. Christopher Lance, <i>Automorphisms of postliminal C*-algebras</i> .....  | 547 |
| Sibe Mardesic, <i>Images of ordered compacta are locally peripherally metric</i> .....   | 557 |
| Albert W. Marshall, David William Walkup and Roger Jean-Baptiste Robert Wets, <i>Order-preserving functions: Applications to majorization and order statistics</i> ..... | 569 |
| Wellington Ham Ow, <i>An extremal length criterion for the parabolicity of Riemannian spaces</i> .....   | 585 |
| Wellington Ham Ow, <i>Criteria for zero capacity of ideal boundary components of Riemannian spaces</i> .....   | 591 |
| J. H. Reed, <i>Inverse limits of indecomposable continua</i> .....   | 597 |
| Joseph Gail Stampfli, <i>Minimal range theorems for operators with thin spectra</i> .....  | 601 |
| Roy Westwick, <i>Transformations on tensor spaces</i> .....  | 613 |
| Howard Henry Wicke, <i>The regular open continuous images of complete metric spaces</i> .....  | 621 |
| Abraham Zaks, <i>A note on semi-primary hereditary rings</i> .....   | 627 |
| Thomas William Hungerford, <i>Correction to: "A description of <math>\text{Mult}_r(A^1, \dots, A^n)</math> by generators and relations"</i> .....                        | 629 |
| Uppuluri V. Ramamohana Rao, <i>Correction to: "On a stronger version of Wallis' formula"</i> .....   | 629 |
| Takesi Isiwata, <i>Correction: "Mappings and spaces"</i> .....   | 630 |
| Henry B. Mann, Josephine Mitchell and Lowell Schoenfeld, <i>Correction to: "Properties of differential forms in n real variables"</i> .....                              | 631 |
| James Calvert, <i>Correction to: "An integral inequality with applications to the Dirichlet problem"</i> .....   | 631 |
| K. Srinivasacharyulu, <i>Correction to: "Topology of some Kähler manifolds"</i> .....  | 632 |