

Pacific Journal of Mathematics

POINT-LIKE 0-DIMENSIONAL DECOMPOSITIONS OF S^3

HOWARD WILSON LAMBERT AND RICHARD BENJAMIN SHER

POINT-LIKE 0-DIMENSIONAL DECOMPOSITIONS OF S^3

H. W. LAMBERT AND R. B. SHER

This paper is concerned with upper semicontinuous decompositions of the 3-sphere which have the property that the closure of the sum of the nondegenerate elements projects onto a set which is 0-dimensional in the decomposition space. It is shown that such a decomposition is definable by cubes with handles if it is point-like. This fact is then used to obtain some properties of point-like decompositions of the 3-sphere which imply that the decomposition space is a topological 3-sphere. It is also shown that decompositions of the 3-sphere which are definable by cubes with one hole must be point-like if the decomposition space is a 3-sphere.

In this paper we consider upper semicontinuous decompositions of S^3 , the Euclidean 3-sphere. In particular, we shall restrict ourselves to those decompositions G of S^3 which have the property that the union of the nondegenerate elements of G projects onto a set whose closure is 0-dimensional in the decomposition space of G . We shall refer to such decompositions as 0-dimensional decompositions of S^3 . Numerous examples of such decompositions appear in the literature. (One should note that some of the examples and results to which we refer are in E^3 , Euclidean 3-space, but the corresponding examples and results for S^3 will be obvious in each case.)

In § 3, a technique of McMillan [10] is used to show that point-like 0-dimensional decompositions of S^3 are definable by cubes with handles. Armentrout [2] has shown this in the case where the decomposition space is homeomorphic with S^3 . The proof of this theorem shows that compact proper subsets of S^3 with point-like components are definable by cubes with handles.

In § 4 we give some properties of point-like 0-dimensional decompositions of S^3 which imply that the decomposition space is homeomorphic with S^3 . These properties were suggested by Bing in § 7 of [6].

It is not known whether monotone 0-dimensional decompositions of S^3 which yield S^3 must have point-like elements. Partial results in this direction have been obtained by Armentrout [2], Bean [5], and Martin [9]. Bing, in § 4 of [6], has presented an example of a decomposition of S^3 which yields S^3 even though it is not a point-like decomposition, but this example is not 0-dimensional. In § 5 we show that a 0-dimensional decomposition of S^3 that yields S^3 must have point-like elements if it is definable by cubes with one hole.

2. **Definitions and notation.** Let G be an upper semicontinuous decomposition of S^3 , the 3-sphere. We denote the decomposition space of G by S^3/G , the union of the nondegenerate elements of G by H_G , and the projection map from S^3 onto S^3/G by P .

The decomposition G is said to be *monotone* if each element of G is a continuum. If $\text{cl } P(H_G)$ is 0-dimensional in S^3/G , then G is a *0-dimensional decomposition* of S^3 . If each element of G has a complement in S^3 which is homeomorphic with E^3 , Euclidean 3-space, then G is a *point-like decomposition* of S^3 .

The sequence M_1, M_2, M_3, \dots is a *defining sequence* for G if and only if M_1, M_2, M_3, \dots is a sequence of compact 3-manifolds with boundary in S^3 such that (1) for each positive integer i , $M_{i+1} \subset \text{Int } M_i$, and (2) g is a nondegenerate element of G if and only if g is a nondegenerate component of $\bigcap_{i=1}^{\infty} M_i$. Here, as in the remainder of the paper, subsets of S^3 which are manifolds will be assumed to be polyhedral subsets of S^3 . It is well known that if G is a 0-dimensional decomposition of S^3 , a defining sequence exists for G . If a defining sequence M_1, M_2, M_3, \dots exists for G such that for each positive integer i , each component of M_i is a cube with handles, G is said to be *definable by cubes with handles*. If a defining sequence M_1, M_2, M_3, \dots exists for G such that for each positive integer i , each component of M_i is a cube with one hole, G is said to be *definable by cubes with one hole*.

3. **Some consequences of a result of McMillan.** The following lemma is a special case of Lemma 1 of [11]. Its proof follows from the very useful technique used by McMillan to prove Theorem 1 of [10].

LEMMA 1. (McMillan). *In S^3 , let M' be a compact polyhedral 3-manifold with boundary such that $\text{Bd}M'$ is connected, and let M be a compact polyhedral 3-manifold with boundary such that $M \subset \text{Int } M'$, and each loop in M can be shrunk to a point in $\text{Int } M'$. Then there is a cube with handles C such that $M \subset \text{Int}C \subset C \subset \text{Int } M'$.*

LEMMA 2. *If G is a point-like 0-dimensional decomposition of S^3 , then there is a defining sequence M_1, M_2, M_3, \dots for G such that for each positive integer i , each component of M_i has a connected boundary.*

Proof. Let M'_1, M'_2, M'_3, \dots be a defining sequence for G , let n be a positive integer, and let K be a component of M'_n . Let g be a component of $\bigcap_{i=1}^{\infty} M'_i$ which lies in K and let U be an open subset of K containing g such that $\text{cl } U \cap \text{Bd}K = \emptyset$. Since g is point-like, there is a 3-cell C such that $g \subset \text{Int } C \subset C \subset U$. There is an integer j such that L , the component of M'_j containing g , lies in $\text{Int } C$. Since

C separates no points of BdK in K , L separates no points of BdK in K .

Using compactness of $\bigcap_{i=1}^{\infty} M'_i$, one obtains a finite collection L_1, \dots, L_k of mutually exclusive defining elements whose interiors cover $(\bigcap_{i=1}^{\infty} M'_i) \cap K$ and so that no L_i separates points of BdK in K . It follows easily that $\bigcup_{i=1}^k L_i$ separates no points of BdK in K . By suitable relabeling, we suppose then, that if i is a positive integer and K is a component of M'_i , $K \cap M'_{i+1}$ does not separate points of BdK in K . We construct disjoint arcs in $K - M'_{i+1}$ connecting the boundary components of K and "drill-out" these arcs to replace K by a compact 3-manifold with connected boundary. Doing this for each component of each M'_i , we obtain a defining sequence M_1, M_2, M_3, \dots as required by the conclusion of the lemma.

THEOREM 1. *If G is a point-like 0-dimensional decomposition of S^3 , then G is definable by cubes with handles.*

Proof. Using Lemma 2, there is a defining sequence M'_1, M'_2, M'_3, \dots for G such that each component of each M'_i has a connected boundary. Let n be a positive integer and N a component of M'_n . Since G is point-like, there is no loss of generality in supposing that each loop in $M'_{n+1} \cap N$ can be shrunk to a point in $\text{Int } N$. From Lemma 1, there is a cube with handles, C , such that $(M'_{n+1} \cap N) \subset \text{Int } C \subset C \subset \text{Int } N$. Hence, there is a sequence M_1, M_2, M_3, \dots of compact 3-manifolds with boundary such that (1) for each positive integer i , $M'_{i+1} \subset \text{Int } M_i \subset M_i \subset \text{Int } M'_i$, and (2) each component of M_i is a cube with handles. The sequence M_1, M_2, M_3, \dots is a defining sequence for G and so G is definable by cubes with handles.

The proof of the next theorem follows from the proof of Theorem 1.

THEOREM 2. *If M is a closed subset of S^3 such that each component of M is point-like, then there exists a sequence M_1, M_2, M_3, \dots of compact 3-manifolds with boundary such that (1) for each positive integer i , $M_{i+1} \subset \text{Int } M_i$, (2) each component of M_i is a cube with handles, and (3) $M = \bigcap_{i=1}^{\infty} M_i$.*

The concept of equivalent decompositions of S^3 was introduced in [4] and the following theorem follows immediately from Theorem 1 of this paper and Theorem 8 of [4].

THEOREM 3. *If G is a point-like 0-dimensional decomposition of S^3 , then G is equivalent to a point-like 0-dimensional decomposition of S^3 each of whose nondegenerate elements is a 1-dimensional continuum.*

In the remaining two sections, we shall utilize some of the above results to investigate certain properties of 0-dimensional decompositions of S^3 .

4. **Properties of point-like 0-dimensional decompositions of S^3 .** In this section we give two properties, each of which is both necessary and sufficient to imply S^3/G is homeomorphic to S^3 .

A space X will be said to have the *Dehn's Lemma property* if and only if the following condition holds: If D is a disk and f is a mapping of D into X such that on some neighborhood of $f(\text{Bd}D)$, f^{-1} is a function, and U is neighborhood of the set of singular points of $f(D)$, then there is a disk D' in $f(D) \cup U$ such that $\text{Bd}D' = f(\text{Bd}D)$.

A space X will be said to have the *map separation property* if and only if the following condition holds: If D is a disk and f_1, \dots, f_n are maps of D into X such that (1) for each i , on some neighborhood of $f_i(\text{Bd}D)$, f_i^{-1} is a function, (2) if $i \neq j$, $f_i(\text{Bd}D) \cap f_j(D) = \emptyset$, and (3) U is a neighborhood of $f_1(D) \cup \dots \cup f_n(D)$, then there exist maps f'_1, \dots, f'_n of D into X such that (1) for each i , $f'_i|_{\text{Bd}D} = f_i|_{\text{Bd}D}$, (2) $f'_i(D) \cup \dots \cup f'_n(D) \subset U$, and (3) if $i \neq j$, $f'_i(D) \cap f'_j(D) = \emptyset$.

It is a well known (and useful) fact that S^3 has the Dehn's Lemma property and the map separation property.

THEOREM 4. *If G is a point-like 0-dimensional decomposition of S^3 , then S^3/G is homeomorphic with S^3 if and only if S^3/G has the Dehn's Lemma property.*

Proof. The "if" portion of the theorem is the only part that requires proof. Let U be an open set containing $\text{cl}H_G$ and $\varepsilon > 0$. We shall construct a homeomorphism $h_\varepsilon: S^3 \rightarrow S^3$ such that if $x \in S^3 - U$, $h_\varepsilon(x) = x$ and if $g \in G$, $\text{diam } h_\varepsilon(g) < \varepsilon$. It will follow from Theorem 3 of [2] that S^3/G is homeomorphic with S^3 .

By Theorem 1, G is definable by cubes with handles. Hence, there exist disjoint cubes with handles C_1, \dots, C_n such that $\text{cl}H_G \subset \bigcup_{i=1}^n \text{Int } C_i \subset \bigcup_{i=1}^n C_i \subset U$. Let W_1, \dots, W_n be pairwise disjoint neighborhoods of C_1, \dots, C_n respectively such that $\bigcup_{i=1}^n W_i \subset U$. Since C_1 is a cube with (possibly 0) handles, there is a homeomorphism h_0 of S^3 onto S^3 such that $h_0(x) = x$ for $x \in S^3 - W_1$ and $h_0(C_1)$ can be written as the union of a finite number of cubes such that (1) each cube has diameter less than $\varepsilon/2$, (2) no three cubes have a point in common, and (3) the intersection of any two cubes is empty or a disk on the boundary of each. The homeomorphism h_0 can be thought of as pulling C_1 towards a 1-dimensional spine of C_1 . Let D_1, D_2, \dots, D_k be the inverse images under h_0 of the disks obtained by intersecting the various cubes making up $h_0(C_1)$. We note that if a continuum in

C_1 intersects at most one D_i , then its image under h_0 has diameter less than ϵ . For each $i = 1, \dots, k$, let D'_i be a subdisk of D_i such that $D'_i \subset \text{Int } D_i$ and $D_i \cap \text{cl } H_G = \text{Int } D'_i \cap \text{cl } H_G$. Let D be a disk in S^3 such that $\text{Bd } D \cap (\bigcup_{i=1}^n C_i) = \emptyset$ and $\bigcup_{i=1}^k D_i = D \cap (\bigcup_{i=1}^n C_i) = D \cap C_1$. Denote the punctured disk $\text{cl } (D - \bigcup_{i=1}^k D'_i)$ by D' . Now $P_1 = P|D$ is a map of D into S^3/G and P_1^{-1} is a homeomorphism on a neighborhood of $P_1(\text{Bd } D)$. The singular set of $P_1(D)$ is contained in $P_1(\bigcup_{i=1}^k \text{Int } D'_i)$. Let V be an open set in S^3/G containing the singular set of $P_1(D)$ and such that $P^{-1}(V) \subset (\text{Int } C_1) - D'$. By hypothesis there exists a disk E in $P_1(D) \cup V$ bounded by $P_1(\text{Bd } D)$. Let E_1, \dots, E_k be the subdisks of E bounded by $P_1(\text{Bd } D'_1), \dots, P_1(\text{Bd } D'_k)$ respectively, and let U_1, \dots, U_k be open sets whose closures lie in $P(\text{Int } C_1)$ such that for each $i = 1, \dots, k$, $E_i \subset U_i$, and if $i \neq j$, $\text{cl } U_i \cap \text{cl } U_j = \emptyset$. By the proof of Theorem 2.1 of [12], each $\text{Bd } D'_i$ can be shrunk to a point in $P^{-1}(U_i)$. Each map can be "glued" to the annulus $\text{cl } (D_i - D'_i)$ to obtain a map from D_i into $D_i \cup P^{-1}(U_i)$ with no singularities on $D_i - P^{-1}(\text{cl } U_i)$. We now apply Dehn's Lemma in S^3 to these maps to obtain disjoint disks F_1, \dots, F_k such that (1) for each i , $\text{Bd } D_i = \text{Bd } F_i$, (2) $\text{Int } F_i \subset \text{Int } C_1$, and (3) if $g \in G$, g intersects no more than one of the disks F_1, \dots, F_k . Let h'_1 be a homeomorphism of S^3 onto itself fixed on $S^3 - \text{Int } C_1$ such that for each i , $h'_1(F_i) = D_i$. Let $h_1 = h_0 h'_1$. Note that if $g \in G$ and $g \subset C_1$, $\text{diam } h_1(g) < \epsilon$. Let h_2, \dots, h_n be homeomorphisms such as h_1 for the sets C_2, \dots, C_n . We define $h_\epsilon : S^3 \rightarrow S^3$ by $h_\epsilon(x) = h_1 h_2 \dots h_n(x)$.

REMARK. If G is the upper semicontinuous decomposition of S^3 whose only nondegenerate element is a polyhedral 2-sphere, then S^3/G has the Dehn's Lemma property but S^3/G is not homeomorphic with S^3 .

The essential ideas of the proof of the following theorem are so like those of the proof of Theorem 4 that we shall not include the proof here.

THEOREM 5. *If G is a point-like 0-dimensional decomposition of S^3 , then S^3/G is homeomorphic with S^3 if and only if S^3/G has the map separation property.*

5. Decompositions of S^3 which yield S^3 . Let S, T be polyhedral solid tori such that $S \subset \text{Int } T$ and let J be a polygonal center curve of S . Following a definition of Schubert [13] which was used in [7], we let $N(S, T)$ be the $\min_D \{N(J \cap D)\}$: where D is a polyhedral meridional disk of T and $N(J \cap D)$ is the number of points in $J \cap D$.

THEOREM 6. *If G is definable by cubes with one hole and S^3/G*

is homeomorphic to S^3 , then G is point-like.

Proof. Let M_1, M_2, \dots , be the defining sequence for G and let T_0 be a component of some M_n . By hypothesis, T_0 is a cube with one hole. Let g be a component of $\bigcap_{i=1}^{\infty} M_i$ contained in T_0 . We first show that there is a defining stage M_{n+m} such that each loop in the component of M_{n+m} containing g can be shrunk to a point in T_0 .

For $i = 1, 2, 3, \dots$, let T_i be the component of M_{n+i} that contains g . Then each T_i is a cube with one hole, $T_{i+1} \subset \text{Int } T_i$, and $\bigcap_{i=1}^{\infty} T_i = g$. Suppose that there is a positive integer s such that each $T_j, j \geq s$, is a solid torus. If the center curve of each T_{j+1} cannot be shrunk to a point in T_j , then g has nontrivial Čech cohomology, and it follows from Corollary 2 of [8] that S^3/G is not homeomorphic to S^3 , contradicting our hypothesis. Hence there is an m such that the center curve of T_m can be shrunk to a point in T_0 and hence each loop in T_m can be shrunk to a point in T_0 .

Suppose then that infinitely many of the T_i are not solid tori. We may suppose for convenience that each T_i is not a solid torus. By [1], each $T'_i = S^3 - \text{Int } T_i$ is a solid torus. We now have three cases.

Case I. Suppose there is an m such that $N(T'_{m-1}, T'_m) = 0$. This implies that there is a meridional disk D of T'_m such that $D \cap T'_{m-1} = \emptyset$. Then there is a cube K in T'_m such that $T'_{m-1} \subset \text{Int } K$. It then follows that each loop in $T'_m (= S^3 - \text{Int } T'_m)$ can be shrunk to a point in T_0 .

We now show that the remaining two cases cannot occur.

Case II. Suppose that there is a positive integer s such that $N(T'_j, T'_{j+1}) = 1$ for $j \geq s$. Since $P(\bigcap_{i=1}^{\infty} M_i)$ is 0-dimensional there is a positive integer t and a cube K such that $P(T'_{s+t}) \subset \text{Int } K \subset K \subset P(\text{Int } T'_s)$. Let D'_{s+t} be a meridional disk of T'_{s+t} . Using Dehn's Lemma we may adjust $P(D'_{s+t})$ in $P(\text{Int } T'_{s+t})$ so that it is polyhedral, and it follows that $P(T'_{s+t})$ is a solid torus with the adjusted $P(D'_{s+t})$ as a meridional disk. Let J be a longitudinal simple closed curve of T'_{s+t} such that $J \subset \text{Bd } T'_{s+t}$ and J intersects $\text{Bd } D'_{s+t}$ at just one point. Let A be an annulus with boundary components A_1 and A_2 . By [13], $N(T'_s, T'_{s+t}) = 1$. Hence there is a mapping f of A into T'_{s+t} such that $f|_{A_1}$ is a homeomorphism, $f(A_1) = J$, and $f(A_2) \subset T'_s$. Now $P(f(A_2))$ can be shrunk to a point missing K since it is contained in $S^3 - K$; hence $P(f(A_2))$ can be shrunk to a point in $P(T'_{s+t})$. But this implies that the longitudinal simple closed curve $P(J)$ of $P(T'_{s+t})$ can be shrunk to a point in $P(T'_{s+t})$. Hence Case II cannot occur.

Case III. Now assume there is a positive integer s such that $N(T'_j, T'_{j+1}) > 1$ for $j \geq s$. Since each T'_j is knotted in S^3 , we may use an argument similar to that used in [7] to conclude that Case III cannot occur.

These three cases now imply that there is a defining stage M_{n+m} such that each loop in the component of M_{n+m} containing g can be shrunk to a point in T_0 . Since $T_0 \cap (\bigcap_{i=1}^{\infty} M_i)$ is compact, there is a defining stage M_p ($p \geq n+m$) such that each loop in $T_0 \cap M_p$ can be shrunk to a point in T_0 . By Lemma 1 there is a cube with handles C such that $T_0 \cap M_p \subset \text{Int } C \subset C \subset \text{Int } T_0$. It then follows that G is definable by cubes with handles. By Bean's result [5], G is a point-like decomposition, and the proof of Theorem 6 is complete.

COROLLARY. *Let f be a mapping of S^3 onto S^3 and let $H = \text{cl}(\{x : x \in S^3 \text{ and } f^{-1}(x) \text{ is nondegenerate}\})$. If H is a 0-dimensional set which is definable by cubes with one hole, then for each $x \in S^3$, $S^3 - f^{-1}(x)$ is homeomorphic to E^3 .*

Proof. Let $G = \{f^{-1}(x) : x \in S^3\}$. It is not hard to show that G is an upper semicontinuous decomposition of S^3 and that S^3/G is homeomorphic to S^3 . Since H is definable by cubes with one hole, it follows that G is definable by cubes with one hole. By Theorem 6, G is a point-like decomposition of S^3 ; hence if $x \in S^3$, then $S^3 - f^{-1}(x)$ is homeomorphic to E^3 .

REFERENCES

1. J. W. Alexander, *On the subdivision of 3-space by a polyhedron*, Proc. Nat. Acad. Sci. U.S.A. **10** (1924), 6-8.
2. S. Armentrout, *Decompositions of E^3 with a compact 0-dimensional set of non-degenerate elements*, Trans. Amer. Math. Soc. **123** (1966), 165-177.
3. ———, "Monotone decompositions of E^3 ," *Topology Seminar Wisconsin*, 1965, Princeton University Press, 1966.
4. S. Armentrout, L. L. Lininger and D. V. Meyer, "Equivalent decompositions of E^3 ," *Topology Seminar Wisconsin*, 1965, Princeton University Press, 1966.
5. R. J. Bean, *Decompositions of E^3 which yield E^3* , Pacific J. Math. **20** (1967), 411-413.
6. R. H. Bing, "Decompositions of E^3 ," *Topology of 3-manifolds and related topics*, Prentice-Hall, 1962.
7. J. M. Kister and D.R. McMillan, Jr., *Locally Euclidean factors of E^4 which cannot be imbedded in E^3* , Ann. of Math. **76** (1962), 541-546.
8. K. W. Kwun and F. Raymond, *Almost acyclic maps of manifolds*, Amer. J. Math. **86** (1964), 638-650.
9. J. Martin, "Sewings of crumpled cubes which do not yield S^3 ," *Topology Seminar Wisconsin*, 1965, Princeton University Press, 1966.
10. D. R. McMillan, Jr., *Cartesian products of contractible open manifolds*, Bull. Amer. Math. Soc. **67** (1961), 510-514.

11. ———, *A criterion for cellularity in a manifold*, Ann. of Math. **79** (1964), 327-337.
12. T. M. Price, *A necessary condition that a cellular upper semicontinuous decomposition of E^n yield E^n* , Trans. Amer. Math. Soc. **122** (1966), 427-435.
13. H. Schubert, *Knoten und Vollringe*, Acta Math. **90** (1953), 131-286.

Received July 25, 1966.

THE UNIVERSITY OF IOWA
THE UNIVERSITY OF GEORGIA

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
Rice University
Houston, Texas 77001

J. P. JANS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL ORDNANCE TEST STATION

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 24, No. 3

July, 1968

Duane W. Bailey, <i>On symmetry in certain group algebras</i>	413
Lawrence Peter Belluce and Surender Kumar Jain, <i>Prime rings with a one-sided ideal satisfying a polynomial identity</i>	421
L. Carlitz, <i>A note on certain biorthogonal polynomials</i>	425
Charles O. Christenson and Richard Paul Osborne, <i>Pointlike subsets of a manifold</i>	431
Russell James Egbert, <i>Products and quotients of probabilistic metric spaces</i>	437
Moses Glasner, Richard Emanuel Katz and Mitsuru Nakai, <i>Bisection into small annuli</i>	457
Karl Edwin Gustafson, <i>A note on left multiplication of semigroup generators</i>	463
I. Martin (Irving) Isaacs and Donald Steven Passman, <i>A characterization of groups in terms of the degrees of their characters. II</i>	467
Howard Wilson Lambert and Richard Benjamin Sher, <i>Point-like 0-dimensional decompositions of S^3</i>	511
Oscar Tivis Nelson, <i>Subdirect decompositions of lattices of width two</i>	519
Ralph Tyrrell Rockafellar, <i>Integrals which are convex functionals</i>	525
James McLean Sloss, <i>Reflection laws of systems of second order elliptic differential equations in two independent variables with constant coefficients</i>	541
Bui An Ton, <i>Nonlinear elliptic convolution equations of Wiener-Hopf type in a bounded region</i>	577
Daniel Eliot Wulbert, <i>Some complemented function spaces in $C(X)$</i>	589
Zvi Ziegler, <i>On the characterization of measures of the cone dual to a generalized convexity cone</i>	603