

Pacific Journal of Mathematics

**GROWTH TRANSFORMATIONS FOR FUNCTIONS ON
MANIFOLDS**

LEONARD E. BAUM AND GEORGE ROGER SELL

GROWTH TRANSFORMATIONS FOR FUNCTIONS ON MANIFOLDS

LEONARD E. BAUM AND GEORGE R. SELL

In this paper we look at the problem of maximizing a function P defined on a manifold M . Although we shall be primarily concerned with the case where M is a certain polyhedron in a Euclidean space R^n and P is a polynomial with nonnegative coefficients defined on R^n , some of our results are valid in greater generality.

In § 2 we describe the general behavior of a growth transformation of P in the vicinity of a local extremum. These results are of a topological nature and can be thought of as a topological—dynamical description of growth transformations.

In § 3 we turn our attention to a particular class of growth transformation which arise for polynomials with nonnegative coefficients. We shall prove the following result, which is the main theorem of this paper:

THEOREM. *Let $M \cup \partial M$ denote the manifold with boundary given by $x = (x_{ij})$ where*

$$\left\{ x_{ij} : x_{ij} \geq 0 \text{ and } \sum_{j=1}^{q_i} x_{ij} = 1 \right\}$$

where q_1, \dots, q_k is a set of nonnegative integers. Let P be a homogeneous polynomial in the variable $\{x_{ij}\}$, with nonnegative coefficients. Let $\mathcal{T} = \mathcal{T}_P : M \rightarrow M \cup \partial M$ defined by $y = \mathcal{T}_P(x)$ where

$$y_{ij} = x_{ij} \frac{\partial P}{\partial x_{ij}} \left[\sum_{k=1}^{q_i} x_{ik} \frac{\partial P}{\partial x_{ik}} \right]^{-1}.$$

Then

$$(1) \quad P(x) \leq P(t\mathcal{T}_P(x) + (1-t)x), \quad (0 \leq t \leq 1, x \in M).$$

The proof of this is based on a suitable modification of an argument of L. E. Baum and J. A. Eagon, cf., [1].

We also study the problem of extending the mapping \mathcal{T}_P to the boundary ∂M in such a way that it is continuous. These results are stated in Theorem 7. It is a consequence of this that \mathcal{T}_P maps neighborhoods of a local maximum into themselves even if the maximum is on the boundary.

In § 5 we examine other growth transformations that are related

to the mapping \mathcal{F}_P . By using an argument suggested by Professor O. Rothaus we are able to extend the theorem stated above to arbitrary (nonhomogeneous) polynomials with positive coefficients.

2. Growth transformations. In this section we shall investigate the behavior of a growth transformation in the vicinity of an isolated maximum.

DEFINITION. Let P be a continuous function defined on a C^∞ -manifold M . We say that a continuous mapping $\sigma : M \rightarrow M$ is a *growth transformation* (for P) if

$$(2) \quad P(x) \leq P(\sigma(x)), \quad (x \in M).$$

If P is a C^2 -function we say that σ is a *proper growth transformation* (for P) if (2) holds and

$$(3) \quad P(x) = P(\sigma(x)) \text{ implies that } x \text{ is a critical point of } P,$$

which means that $\text{grad } P = 0$ at x . A growth transformation σ is said to *increase P homotopically* if there exists a continuous mapping

$$S_t(x) : [0, 1] \times M \longrightarrow M$$

such that

$$(4) \quad \begin{aligned} & \text{(i) } S_0(x) = x \\ & \text{(ii) } S_1(x) = \sigma(x) \\ & \text{(iii) For each } t, 0 \leq t \leq 1, S_t \text{ is a growth transformation for } P. \end{aligned}$$

A continuous function P is said to have a *local maximum* at q if there is a neighborhood V of q with

$$P(x) \leq P(q)$$

for all x in V . If P is a C^2 -function, the point q is called an *isolated maximum* if it is a local maximum of P and an isolated critical point.

We will describe the asymptotic behavior of the iterates of a growth transformation. Let σ be a growth transformation for P and define the fixed value set $F_{P\sigma}$ by

$$(5) \quad F_{P\sigma} = \{x \in M : P(x) = P(\sigma(x))\}.$$

We say that a set $K \subset M$ is *invariant* if $\sigma^n(x) \in K$ for $n \geq 1$, whenever $x \in K$.

THEOREM 1. *Let P be a real-valued continuous function on a C^∞ -manifold M and let $\sigma : M \rightarrow M$ be a growth transformation for*

P. Let K be the largest invariant subset of the fixed value set $F_{P\sigma}$ given by (5). If a point x in M has the property that $\sigma^n(x)$ remains in a compact for $n \geq 1$, then

$$\sigma^n(x) \longrightarrow K \text{ as } n \longrightarrow \infty .$$

Proof. Let $x \in M$ be a point with the property that $\{x, \sigma(x), \sigma^2(x), \dots\}$ lies in a compact set in M . Since $P(\sigma^n(x))$ is increasing in n , it follows that $\lim_{n \rightarrow \infty} P(\sigma^n(x))$ exists, say that

$$P(\sigma^n(x)) \longrightarrow \alpha \text{ as } n \longrightarrow \infty .$$

Now let $y \in M$ be a limit point of a subsequence $\{\sigma^{n_i}(x)\}$. Then by the continuity of P we have

$$P(\sigma^{n_i}(x)) \longrightarrow P(y) = \alpha .$$

However $\sigma^{n_i+1}(x) \rightarrow \sigma(y)$ and in general $\sigma^{n_i+k}(x) \rightarrow \sigma^k(y)$. From this it follows that $P(\sigma^k(y)) = \alpha, k = 0, 1, \dots$, in other words, $y \in K$, which completes the proof of Theorem 1.

By demanding that the growth transformation σ leave the local maxima of P fixed, we are able to assert something about the behavior of σ in the vicinity of a local maximum.

THEOREM 2. Let P be a real-valued C^2 -function on a C^∞ -manifold M and let $\sigma : M \rightarrow M$ be a growth transformation for P . Assume that every local maximum of P is a fixed point for σ . Then for every local maximum q of P there is a neighborhood V with $\sigma(V) \subset V$. If, in addition, σ is a proper growth transformation and q is an isolated maximum then V can be chosen so that

$$\sigma^n(x) \longrightarrow q \text{ as } n \longrightarrow \infty$$

for every $x \in V$.

Proof. Let q be a local maximum of P and let V be the connected component of

$$\{x : P(x) > P(q) - \eta\}$$

that contains q , where $\eta > 0$ is fixed. Furthermore, we can choose η so that \bar{V} , the closure of V , is compact. Now define

$$\begin{aligned} A &= \{x \in V : \sigma(x) \in \bar{V}\} \\ B &= \{x \in V : \sigma(x) \notin \bar{V}\} . \end{aligned}$$

It is clear that $A \cap B = \emptyset$ and $V = A \cup B$. Furthermore, B is an open set since σ is continuous. The set A can be written as

$$A = \{x \in V : \sigma(x) \in V\}$$

since σ is a growth transformation, that is the set

$$\{x \in V : \sigma(x) \in \bar{V} - V\} = \{x \in V : P(x) = P(q) - \eta\}$$

is empty. It follows then that A is an open set. Since V is connected one of the sets A or B must be empty. However, $\sigma(q) = q$, therefore $B = \emptyset$. Hence $V = A$ and $\sigma(V) \subset V$. (It also follows that $\sigma(\bar{V}) \subset \bar{V}$.)

If q is a maximum, that is an isolated critical point, then η can be chosen so that \bar{V} contains no critical point of P other than q . If K is the largest invariant subset of $F_{P\sigma}$ which is given by (5), then $K \cap V = \{q\}$. Hence by Theorem 1, if $x \in V$, then $\sigma^n(x) \rightarrow q$ as $n \rightarrow \infty$.

A conclusion similar to that of Theorem 2 is possible under a slightly different hypothesis.

THEOREM 3. *Let P be a real-valued continuous function defined on a C^∞ -manifold M and let $\sigma : M \rightarrow M$ be a growth transformation for P . Assume that σ increases P homotopically and let $S_t(x)$ satisfy (4). Then for every local maximum q of P there is a neighborhood V with $S_t(V) \subset V$, $0 \leq t \leq 1$. If, in addition, σ is a proper growth transformation and q is an isolated maximum, then V can be chosen so that for every $x \in V$*

$$\sigma^n(x) \longrightarrow q \quad \text{as } n \longrightarrow \infty,$$

in particular $\sigma(q) = q$.

Proof. Let $q \in M$ be a local maximum of P and let V be the component of $\{x : P(x) > P(q) - \eta\}$ that contains q , where $\eta > 0$ is fixed. We also choose η so that \bar{V} is compact. Now fix x in V . Then

$$(6) \quad P(q) - \eta < P(x) \leq P(S_t(x)), \quad 0 \leq t \leq 1.$$

Let $G = \{S_t(x) : 0 \leq t \leq 1\}$, then G is a connected set since $S_t(x)$ is continuous in t . By (6) we see that the set

$$\{S_t(x) \in \bar{V} - V\} = \{S_t(x) : P(S_t(x)) = P(q) - \eta\}$$

is empty. Thus, $G = (G \cap V) \cup (G \cap (M - \bar{V}))$. Since G is connected one of the sets $G \cap V$ or $G \cap (M - \bar{V})$ is empty. However, $x \in G \cap V$. Hence $G \subset V$, that is $S_t(V) \subset V$ for $0 \leq t \leq 1$. The remainder of the proof follows that of Theorem 2.

In Theorems 2 and 3 one is able to assert that the neighborhood V is a disk provided the function P is a C^2 -function, cf. [4]. Also,

these results are really "local" results, so they are still valid even if σ and P are defined only in a neighborhood of the maximum q . And, finally, they have obvious extensions to manifolds with boundary. Another refinement of Theorem 3, is the following.

THEOREM 4. *Let P be a real-valued continuous function defined on a C^∞ -manifold M and let $\sigma: M \rightarrow M$ be a growth transformation that increases P homotopically. If q is an isolated maximum of P , then $\sigma(q) = q$.*

Proof. Let V_η be the component of $\{x: P(x) > P(q) - \eta\}$ that contains q . It was shown in the proof of Theorem 3 that $S_t(V_\eta) \subseteq V_\eta$. By letting $\eta \rightarrow 0$ we conclude that $\sigma(q) = q$.

The next result seems rather interesting. It asserts that, under appropriate conditions, every growth transformation increases P homotopically in a neighborhood of an isolated maximum.

THEOREM 5. *Let P be a real-valued C^2 -function on a C^∞ -manifold M and let $\sigma: M \rightarrow M$ be a growth transformation. Assume that every isolated maximum of P is a fixed point of σ . Then for every isolated maximum q there is a neighborhood V such that σ increases P homotopically in V .*

Proof. Let $\eta > 0$ be chosen so that the component $V = V_\eta$ of the set $\{x: P(x) > P(q) - \eta\}$ that contains q has the property that \bar{V} is compact and \bar{V} contains no critical points of P other than q . Now consider the differential equation

$$x' = -\text{grad } P$$

in V_η , and let $\varphi(x, t)$ denote the solution that satisfies $\varphi(x, 0) = x$. It is easy to show that for $x \in V_\eta$, $\varphi(x, t) \in V_\eta$, for $t \geq 0$, and $\varphi(x, t) \rightarrow q$ as $t \rightarrow \infty$. Now choose ε , $0 < \varepsilon < \eta$, so that, in the local coordinate system at q , the convex hull of V_ε lies in V_η . We now define a mapping $h: V_\eta \rightarrow V_\varepsilon$ by

$$\begin{aligned} h(x) &= x, & \text{if } x \in V_\varepsilon \\ h(x) &= \varphi(x, T_x), & \text{if } x \in V_\eta - V_\varepsilon, \end{aligned}$$

where $\varphi(x, T_x)$ is the first point at which the trajectory $\varphi(x, t)$ meets V_ε . Since the level surface $P(x) = P(q) - \varepsilon$ is transverse to the flow $\varphi(x, t)$, it follows that h is continuous. Also the mapping

$$g: V_\eta \times [0, 1] \rightarrow V_\eta$$

given by

$$g(x, \tau) = \tau h(\sigma(x)) + (1 - \tau)h(x)$$

is continuous. Now define $S_\tau(x)$, $0 \leq \tau \leq 1$ and $x \in V_\tau$, to be that point in V_τ on the trajectory $\varphi(g(x, \tau), t)$ that satisfies

$$P(S_\tau(x)) = \tau P(\sigma(x)) + (1 - \tau)P(x).$$

It is clear that $S_\tau(x)$ is continuous, and it is easy to verify that $S_\tau(x)$ satisfies (4). This completes the proof of the theorem.

It is apparent from the proofs that if σ is a proper growth transformation then the region of attraction of an isolated maximum q is "large". More precisely, let $\eta > 0$ and define V_η to be the component of $\{x : P(x) > P(q) - \eta\}$ that contains q . We have seen that the region of attraction for q will contain every V_η that has the property that the closure \bar{V}_η is compact and contains only one critical point of P , which must necessarily be the point q . If we let $\eta_0 > 0$ be the first real number for which \bar{V}_{η_0} contains more than one critical point of P then since

$$V_{\eta_0} = \bigcup_{\eta < \eta_0} V_\eta,$$

this implies that the region of attraction always contains V_{η_0} . It should be emphasized that V_{η_0} depends only on the function P and not on the growth transformation σ .

3. Homogeneous polynomials and the transformation \mathcal{F}_p .

Let q_1, \dots, q_k be a set of nonnegative integers with $\sum_i q_i = n$. Let $M \cup \partial M$ denote the set of all vectors

$$\{x = (x_{ij}), i = 1, \dots, k, j = 1, \dots, q_i\}$$

such that

$$x_{ij} \geq 0 \quad \text{and} \quad \sum_{j=1}^{q_i} x_{ij} = 1.$$

The set $M \cup \partial M$ is a polyhedron in R^n . We shall let M denote the interior of the polyhedron, that is

$$M = \{(x_{ij}) \in M \cup \partial M : x_{ij} > 0\}$$

and ∂M is the boundary. The space M is a manifold of dimension $n - k$.

Let $P : R^n \rightarrow R$ be a homogeneous polynomial in the variables (x_{ij}) with positive coefficients. We define a mapping

$$\mathcal{F} = \mathcal{F}_p : M \rightarrow M \cup \partial M$$

by $y = \mathcal{J}_P(x)$ where

$$(7) \quad y_{ij} = \mathcal{J}_P(x)_{ij} = x_{ij} \frac{\partial P}{\partial x_{ij}} \left[\sum_{k=1}^{q_i} x_{ik} \frac{\partial P}{\partial x_{ik}} \right]^{-1}.$$

Note that the range of \mathcal{J} is contained in M unless P does not depend on one of the variables x_{ij} .

In [1] it was shown that $P(x) \leq P(\mathcal{J}(x))$ for all $x \in M$ and equality held if and only if $\mathcal{J}(x) = x$. In other words, the transformation \mathcal{J}_P is a growth transformation for P . We now can assert a stronger result.

THEOREM 6. *The transformation \mathcal{J}_P increases P homotopically. More precisely, if P is a homogeneous polynomial in (x_{ij}) with positive coefficients and $\mathcal{J}_P = \mathcal{J}$ is given by (7), then*

$$(8) \quad P(x) \leq P(t\mathcal{J}(x) + (1-t)x), \quad (x \in M, 0 < t \leq 1).$$

Moreover, equality holds in (8) if and only if $\mathcal{J}(x) = x$.

Note that the transformation \mathcal{J}_P is determined by the first derivatives of P only. In a sense it is similar to moving in the "gradient direction", which also depends on only the first derivatives of P . While moving in the gradient direction will increase the value of P , this is valid only for small steps, and there is no way—without considering second derivatives—for determining the size of the step. On the other hand, the size of the step is completely determined by the first derivatives above for the transformation \mathcal{J}_P .

Proof. One can write $P(x) = \sum_{\alpha} C_{\alpha} m_{\alpha}(x)$ where the coefficients C_{α} are positive and $m_{\alpha}(x)$ is a monomial of degree d , that is

$$m_{\alpha}(x) = \prod_{i,j} x_{ij}^{\alpha_{ij}}$$

where the α_{ij} are nonnegative integers with $\sum_{i,j} \alpha_{ij} = d$. We shall let $\alpha = (\alpha_{ij})$ be the index set for the summation defining P . We note now a few identities which will be needed later.

$$(9) \quad \begin{aligned} P(x) &= \frac{1}{d} \sum_{i,j} x_{ij} \frac{\partial P}{\partial x_{ij}} \\ \sum_{\alpha} \alpha_{ij} C_{\alpha} m_{\alpha}(x) &= x_{ij} \frac{\partial P}{\partial x_{ij}} \\ \frac{m_{\alpha}(x)}{m_{\alpha}(y)} &= m_{\alpha}\left(\frac{x}{y}\right) \end{aligned}$$

where

$$\frac{x}{y} = \left(\frac{x_{ij}}{y_{ij}} \right).$$

By the inequality of geometric and arithmetic means [3, p. 16] we get

$$(10) \quad m_\alpha(x)^{1/d} \leq \frac{1}{d} \sum_{i,j} \alpha_{ij} x_{ij}.$$

Let $t, 0 \leq t \leq 1$, be fixed and let $y = t\mathcal{S}^-(x) + (1-t)x$. Also define Q_α by

$$(11) \quad P(x) = \sum_\alpha \{C_\alpha m_\alpha(y)\}^{1/d+1} Q_\alpha.$$

By applying Hölder's inequality [3, p. 21] to (11) we get

$$(12) \quad P(x) \leq P(y)^{1/d+1} \left(\sum_\alpha Q_\alpha^{d+1/d} \right)^{d/(d+1)}.$$

Now

$$(13) \quad \begin{aligned} \sum_\alpha Q_\alpha^{d+1/d} &= \sum_\alpha C_\alpha m_\alpha(x) \left[\frac{m_\alpha(x)}{m_\alpha(y)} \right]^{1/d} \\ &= \sum_\alpha C_\alpha m_\alpha(x) \left[m_\alpha \left(\frac{x}{y} \right) \right]^{1/d} \\ &\leq \frac{1}{d} \sum_\alpha C_\alpha m_\alpha(x) \sum_{i,j} \alpha_{ij} \left(\frac{x_{ij}}{y_{ij}} \right) \end{aligned}$$

where we apply (10) in the last step.

By substituting for y_{ij} in (13) we get

$$\sum_\alpha Q_\alpha^{(d+1)/d} \leq \frac{1}{d} \sum_\alpha C_\alpha m_\alpha(x) \sum_i \sum_j \alpha_{ij} \left[\frac{\sum_k x_{ik} P_{ik}}{tP_{ij} + (1-t)\sum_k x_{ik} P_{ik}} \right]$$

where

$$P_{ij} = \frac{\partial P}{\partial x_{ij}}.$$

Now by interchanging the order of summation and using (9) we get

$$(14) \quad \sum_\alpha Q_\alpha^{(d+1)/d} \leq \frac{1}{d} \sum_i \left(\sum_k x_{ik} P_{ik} \right) \left[\sum_j \frac{x_{ij} P_{ij}}{tP_{ij} + (1-t)\sum_k x_{ik} P_{ik}} \right].$$

However, by Lemma 1 (see below) with $a_j = x_{ij} P_{ij}$, $b_j = tP_{ij}$ and $c_j = (1-t)\sum_k x_{ik} P_{ik}$, we see that the quantity in the brackets in (14) is bounded by 1 for $0 < t < 1$, and by continuity it is bounded for $0 \leq t \leq 1$. Hence (14) becomes

$$(15) \quad \sum_{\alpha} Q_{\alpha}^{(d+1)/d} \leq \frac{1}{d} \sum_i \sum_k x_{ik} P_{ik} = P(x) .$$

By putting this into (12) we get

$$P(x) \leq P(y)^{1/(d+1)} P(x)^{d/(d+1)} ,$$

which implies that $P(x) \leq P(y)$, and this completes the proof of the theorem.

LEMMA 1. *Let $a_j, b_j, c_j, j = 1, \dots, n$, be positive numbers with $\sum_{j=1}^n a_j/b_j \leq 1/P$ and $\sum_{j=1}^n a_j/c_j \leq 1/Q$, then $\sum_{j=1}^n a_j/(b_j + c_j) \leq 1/(P + Q)$.*

The proof of this is a straightforward induction argument and we will omit the details.

The following consequence of Theorems 3 and 6 asserts that the mapping \mathcal{S}_P cannot leave a ‘‘local hill’’. Furthermore, we are able to conclude something about the region of attraction for an isolated local maximum of P .

COROLLARY. *Let P be a homogeneous polynomial in the variables (x_{ij}) with positive coefficients and let $q \in M$ be an isolated local maximum of P . Then there exists a neighborhood V of q such that $\mathcal{S}(V) \subset V$ and for every $x \in V$*

$$\mathcal{S}^n(x) \longrightarrow q \text{ as } n \longrightarrow \infty .$$

Observe that this corollary can also be obtained from Theorem 2 since \mathcal{S} is a proper growth transformation and $\mathcal{S}(x) = x$ if and only if x is a critical point of P .

The transformation \mathcal{S}_P can, in a limited sense, be extended to the boundary ∂M

$$\partial M = \bigcup_{ij} N_{ij} \cup \partial N_{ij}$$

where each $N_{ij} \cup \partial N_{ij}$ is a polyhedron defined by

$$N_{ij} \cup \partial N_{ij} = \{x \in M \cup \partial M : x_{ij} = 0\} .$$

Following our original convention, we shall let N_{ij} denote the interior of $N_{ij} \cup \partial N_{ij}$ and ∂N_{ij} the boundary.

THEOREM 7. (A) *The transformation \mathcal{S}_P on M can be extended to be continuous, and in fact C^∞ , on*

$$\bigcup_{ij} N_{ij} \cup M .$$

(B) *\mathcal{S}_P can also be continuously extended to any isolated local*

maximum q of P on ∂M by the definition. $\mathcal{F}_P(q) = q$.

(C) The extended transformation \mathcal{F}_P still obeys the inequality

$$P(x) \leq P(t\mathcal{F}_P(x) + (1-t)x), \quad (0 \leq t \leq 1).$$

The proof of this theorem and the following corollaries will be given at the end of the paper.

In an example below we will show that in general \mathcal{F}_P cannot be continuously extended to all of ∂N_{ij} . This occurs when a saddle point of P lies on ∂N_{ij} .

Note that if a local maximum q lies on ∂N_{ij} , then statement (B) above does *not* assert that \mathcal{F} can be extended to a neighborhood of q in $M \cup \partial M$.

COROLLARY 1 (A). *Let P be a homogeneous polynomial in the variables (x_{ij}) with positive coefficients and let $q \in \bigcup_{ij} N_{ij}$ be an isolated maximum of P on the boundary ∂M . Then there exists a neighborhood V of q , such that $\mathcal{F}(V) \subset V$, and for every $x \in V$*

$$\mathcal{F}^n(x) \longrightarrow q \quad \text{as } n \longrightarrow \infty.$$

(B) *If $q \in \bigcup_{ij} \partial N_{ij}$ is an isolated maximum of P , then there is a neighborhood V of q in $M \cup \partial M$ such that for every $x \in V \cap (M \cup \bigcup_{ij} N_{ij})$*

$$\mathcal{F}^n(x) \longrightarrow q \quad \text{as } n \longrightarrow \infty.$$

COROLLARY 2. *Let $q \in N_{ij}$ be an isolated local maximum of P and let $x \in M \cup \partial M$ a point in the domain of attraction of q . Let $y^n = \mathcal{F}_n(x)$ have coordinates (y_{rs}^n) . Then $y_{ij}^n \rightarrow 0$ as $n \rightarrow \infty$. Furthermore, if y_{ij}^n is sufficiently small, say that $y_{ij}^n < \delta$, then we can replace y^n with \tilde{y} where $\tilde{y}_{ij} = 0$, $|\tilde{y}_{rs} - y_{rs}^n| \leq \delta$ and $\mathcal{F}^k(\tilde{y}) \rightarrow q$ as $k \rightarrow \infty$.*

That is we can set the ij component of y^n equal to zero without destroying the convergence property. It should be noted that we do assume q to be an isolated maximum of P in $M \cup \partial M$. If q is only an isolated maximum on ∂M and not on $M \cup \partial M$, then it will still be a critical point of P , but it now will act as a saddle point in the discrete flow induced by \mathcal{F} . A simple example of this phenomenon is given by the polynomial

$$P(x_1, x_2, y_1, y_2) = x_1x_2 + y_1y_2$$

on

$$M \cup \partial M = \{(x_1, x_2, y_1, y_2) : x_i \geq 0, y_i \geq 0, i = 1, 2, x_1 + x_2 = 1, y_1 + y_2 = 1\}.$$

The points $(\frac{1}{2}, \frac{1}{2}, 0, 1)$, $(\frac{1}{2}, \frac{1}{2}, 1, 0)$, $(0, 1, \frac{1}{2}, \frac{1}{2})$ and $(1, 0, \frac{1}{2}, \frac{1}{2})$ are saddle points for \mathcal{F} and they are local extremum for P on ∂M , but are not extremum on $M \cup \partial M$.

4. An illustrative example. Let $M \cup \partial M$ be the subset of R^3 given by

$$\{(x_1, x_2, x_3) : x_i \geq 0 \text{ and } \Sigma x_i = 1\}$$

and let $P(x) = x_1^2 + 4x_2x_3$. Then on M , $\mathcal{F}(x)$ is given by

$$\mathcal{F} : (x_1, x_2, x_3) \longrightarrow \left(\frac{x_1^2}{P}, \frac{2x_2x_3}{P}, \frac{2x_2x_3}{P} \right).$$

The range of \mathcal{F} is then contained in the set

$$\{(x_1, x_2, x_3) \in M \cup \partial M : x_2 = x_3\}$$




Figure 1.

The critical points of P on $M \cup \partial M$ are:

$$(1, 0, 0), (0, 1, 0), (0, 0, 1), (0, \frac{1}{2}, \frac{1}{2}), (\frac{1}{2}, \frac{1}{4}, \frac{1}{4}).$$

The points $(1, 0, 0)$ and $(0, \frac{1}{2}, \frac{1}{2})$ are local maxima, $(0, 1, 0)$ and $(0, 0, 1)$ are local minima, and $(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ is a saddle point. The set of points

$$\{(x_1, x_2, x_3) \in M : x_1^2 = 4x_2x_3\}$$

is mapped onto $(\frac{1}{2}, \frac{1}{4}, \frac{1}{4})$ by \mathcal{F} . This forms a separatrix for the domains of attraction of $(1, 0, 0)$ and $(0, \frac{1}{2}, \frac{1}{2})$. If $x_1^2 > 4x_2x_3$, then $\mathcal{F}^n(x) \rightarrow (0, \frac{1}{2}, \frac{1}{2})$, and if $x_1^2 < 4x_2x_3$ then $\mathcal{F}^n(x) \rightarrow (1, 0, 0)$.

By taking limits from the inside, as indicated in Theorem 7, \mathcal{F} can be extended to all boundary points other than $(0, 1, 0)$ and $(0, 0, 1)$. (\mathcal{F} is defined and continuous even at the corner $(1, 0, 0)$ in this case.)

The corners $(0, 1, 0)$ and $(0, 0, 1)$ are points of discontinuity of \mathcal{J} since there are points arbitrarily close to either that are mapped near $(1, 0, 0)$ and other points that are mapped near $(0, \frac{1}{2}, \frac{1}{2})$.

Finally let us show that the function of t $P(t\mathcal{J}(x) + (1-t)x)$ may fail to be monotone in t . We use this example, which is due to Blakely [2]. Let $x_0 = (4/16, 1/16, 11/16)$

$$\mathcal{J}(4/16, 1/16, 11/16) = (8/30, 11/30, 11/30) .$$

Now

$$\frac{d}{dt}P(t\mathcal{J}(x) + (1-t)x)|_{t=1} = \text{grad } P|_{t=1} \cdot (\mathcal{J}(x) - x) ,$$

and by a direct computation

$$\text{grad } P|_{t=1} \cdot (\mathcal{J}(x_0) - x_0) < 0 .$$

Hence $P(t\mathcal{J}(x_0) + (1-t)x_0)$ is not monotone.

5. **Growth transformations related to \mathcal{J}_P .** Let us, for the moment, consider a special case of the polyhedron $M \cup \partial M$ discussed in § 3, namely where $k = 1$. Then $M \cup \partial M$ is the set of vectors $x = (x_j)$, $j = 1, \dots, n$ such that $x_j \geq 0$ and $\sum_{j=1}^n x_j = 1$. In Theorem 3, when we were studying the behavior of the transformation \mathcal{J}_P generated by a homogeneous polynomial P , we were only interested in the behavior of P on the polyhedron $M \cup \partial M$. Even though P is fixed on $M \cup \partial M$, its extension to R^n is not unique. As a matter of fact, for every integer $m \geq 0$, the polynomial $Q(x) = (\sum x_j)^m P(x)$ is homogeneous and agrees with P on $M \cup \partial M$. However, since the partial derivatives $\partial P/\partial x_i$ and $\partial Q/\partial x_i$ differ, the transformation \mathcal{J}_P and \mathcal{J}_Q generated by P and Q differ. A direct computation shows that the transformation generated by Q is given by

$$(16) \quad \mathcal{J}_Q = \frac{m}{d+m} I + \frac{d}{d+m} \mathcal{J}_P$$

where I denotes the identity and d is the degree of P . Now (16) is valid for every integer $m \geq 0$, and if we set $t = d/(d+m)^{-1}$, (16) becomes

$$\mathcal{J}_Q = t\mathcal{J}_P + (1-t)I .$$

By using the fact that the transformation \mathcal{J}_P is a growth transformation, cf. [1], we can give a partial proof of (8). Indeed,

$$P(x) = Q(x) \leq Q(\mathcal{J}_Q(x)) = P(\mathcal{J}_Q(x)) = P(t\mathcal{J}_P(x) + (1-t)x)$$

provided $t = d/(d+m)^{-1}$ where m is an integer.

It is also interesting to note that when m is large, the transformation \mathcal{S}_Q is a local homeomorphism since the jacobian is (approximately) $m(d + m)^{-1}$. In fact, it is not hard to show that \mathcal{S}_Q is a homeomorphism. This suggests that one may be able to use the topological-dynamical theory of discrete flows in order to study the asymptotic behavior of $\mathcal{S}_Q^n(x)$ when there are nonisolated singularities for P , or Q .

In the general case, where

$$M \cup \partial M = \{x = (x_{ij}) : x_{ij} \geq 0, \sum_{j=1}^{q_i} x_{ij} = 1, i = 1, \dots, k\}$$

the same method yields an improvement of Theorem 6. We are able to conclude not only that

$$P(x_{ij}) \leq P\{(1 - t)(x_{ij}) + t\mathcal{S}_P(x_{ij})_{ij}\}, \quad 0 \leq t \leq 1$$

but also that

$$\begin{aligned} P(x_{ij}) &\leq P\{(1 - t_i)(x_{ij}) + t_i\mathcal{S}_P(x_{ij})_{ij}\}, \\ 0 &\leq t_i \leq 1, \quad i = 1, \dots, k, \end{aligned}$$

that is, not only is the number $P(x)$ smaller than (or equal to) the value of P at any point along the line joining (x) to $\mathcal{S}_P(x)$, but $P(x)$ is also less than (or equal to) the value of P at any point of the k -dimensional rectangle determined by (x) and $\mathcal{S}_P(x)$.

The proof of this fact is obtained by applying the original inequality [1] to the polynomials

$$Q(x_{ij}) = \prod_{i=1}^k \left(\sum_{j=1}^{q_i} x_{ij} \right)^{n_i} P(x_{ij})$$

for integers n_i while noting that P and Q agree on $M \cup \partial M$.

Another interesting consequence of this observation is that we can define proper growth transformations for nonhomogeneous polynomials with positive coefficients.

THEOREM 8. *Let P be a polynomial in the variables (x_{ij}) with positive coefficients. Then P agrees with a homogeneous polynomial Q with positive coefficients on the manifold $M \cup \partial M$ and the transformation \mathcal{S}_Q is a proper growth transformation for P , in fact*

$$P(x) \leq P(t\mathcal{S}_Q(x) + (1 - t)x), \quad (0 < t \leq 1, \quad x \in M),$$

where equality holds if and only if $\mathcal{S}_Q(x) = x$.

Proof. Write P in the form

$$P = H_0 + H_1 + \dots + H_d$$

where H_l is a homogeneous polynomial of degree l . Let

$$Q = H_0\left(\sum_j x_{ij}\right)^d + H_1\left(\sum_j x_{ij}\right)^{d-1} + \cdots + H_d,$$

then $Q = P$ on $M \cup \partial M$. The remainder of the theorem follows from Theorem 6. Observe that Q is not unique since there is great freedom in choosing the multipliers of the H_l .

Finally, the method of proof of this section can be used to extend the basic inequality (10) for homogeneous polynomials with positive coefficients to all polynomials with positive coefficients. We are grateful to Oscar Rothaus for this observation.

THEOREM 9. *Let P be a polynomial in the variables*

$$\{(x_{ij}) : x_{ij} \geq 0, \sum_{j=1}^{q_i} x_{ij} = 1, \quad i = 1, \dots, x\}$$

with nonnegative coefficients. Let

$$\mathcal{F}_P(x)_{ij} = x_{ij} \frac{\partial P}{\partial x_{ij}} \left[\sum_{k=1}^{q_i} x_{ik} \frac{\partial P}{\partial x_{ik}} \right]^{-1}.$$

Then

$$P(x) \leq P((1-t)x + t\mathcal{F}_P(x)), \quad (0 < t \leq 1).$$

Furthermore, equality holds if and only if $\mathcal{F}_P(x) = x$.

Proof. We write $P(x)$ in the form $P(x) = \sum_{l=0}^d H_l(x)$ where H_l is a homogeneous polynomial of degree l . Now we introduce some dummy variables y_1, y_2 and enlarge the domain $M \cup \partial M$. That is, let $N \cup \partial N$ be the domain

$$\{(x_{ij}, y_1, y_2) : (x_{ij}) \in M \cup \partial M, y_1 \geq 0, y_2 \geq 0, y_1 + y_2 = 1\}$$

and consider the polynomial

$$Q(x, y_1, y_2) = \sum_{l=0}^d H_l(x)(y_1 + y_2)^{d-l}.$$

Q is a homogeneous polynomial with nonnegative coefficients and with y_1 and y_2 fixed, Q agrees with P on $M \cup \partial M$. Furthermore, we can apply Theorem 6 of this paper to Q . Since

$$\begin{aligned} \mathcal{F}_Q : x_{ij} &\longrightarrow x_{ij} \frac{\partial Q}{\partial x_{ij}} \left[\sum_{k=1}^{q_i} x_{ik} \frac{\partial Q}{\partial x_{ik}} \right]^{-1} \\ &= x_{ij} \frac{\partial P}{\partial x_{ij}} \left[\sum_{k=1}^{q_i} x_{ik} \frac{\partial P}{\partial x_{ik}} \right]^{-1} \end{aligned}$$

and

$$\mathcal{F}_Q : y_k \longrightarrow y_k \frac{\partial Q}{\partial y_k} \left[\sum_{l=1}^2 y_l \frac{\partial Q}{\partial y_l} \right]^{-1} = y_k$$

it follows that y_1 and y_2 are fixed. Hence

$$\begin{aligned} P(x) &= Q(x, y_1, y_2) \leq Q(t\mathcal{F}_Q(x, y_1, y_2) + (1-t)(x, y_1, y_2)) \\ &= Q(t\mathcal{F}_P(x) + (1-t)x, y_1, y_2) = P(t\mathcal{F}_P(x) + (1-t)x) . \end{aligned}$$

Furthermore, strict inequality holds for $0 < t \leq 1$ unless

$$\mathcal{F}_Q(x, y_1, y_2) = (x, y_1, y_2) ,$$

that is, unless $\mathcal{F}_P(x) = x$.

6. Proof of Theorem 7. For clarity we consider first the special case where there is only one restraint equation. Let

$$M \cup \partial M = \{x = (x_j) : x_j \geq 0 \text{ and } \sum_{j=1}^n x_j = 1\}$$

Let $P_j = \partial P / \partial x_j$. Then

$$\mathcal{F}(x)_j = \frac{x_j P_j}{\sum_k x_k P_k} = \frac{x_j P_j}{dP} ,$$

is C^∞ on the subset of $M \cup \partial M$ where $P \neq 0$, and in particular is continuous on $M \cup \partial M$ at any point where $P \neq 0$ including points on the boundary ∂M . Therefore, $\mathcal{F}(x)$ is well defined on any local maximum of P with respect to $M \cup \partial M$ whether this local maximum is on ∂M or not, since $P \neq 0$ at a local maximum.

Since P has positive coefficients it has no zero in M unless $P \equiv 0$. On the interior of the boundary ∂M , P can vanish only if P is of the form $P = x_j^l Q$ where Q is homogeneous of degree $d - l$ and does not have x_j as a factor. Hence $Q \neq 0$ in the interior of the face $x_j = 0$. In this case $P \equiv 0$ on the face $x_j = 0$ and \mathcal{F} is continuously extended to the interior of the face $x_j = 0$ by

$$\begin{aligned} \mathcal{F}(x)_i &= \frac{x_i Q_i}{dQ} , & i \neq j \\ &= l/d , & i = j . \end{aligned}$$

However, P can vanish on ∂N_{ij} without having a common factor and therefore \mathcal{F} cannot generally be extended to ∂N_{ij} . For example, on the domain

$$M \cup \partial M = \{(x, y, z) : x + y + z = 1, x, y, z \geq 0\}$$

the function $P(x, y, z) = z^2 + 4xy$ vanishes at $z = x = 0, y = 1$ although

P has no common factor and in fact \mathcal{F}_P cannot be continuously extended to $z = 0, x = 0, y = 1$. (See the example of § 4.)

More generally, the transformation \mathcal{F} :

$$\mathcal{F}(x)_{ij} = x_{ij} P_{ij} \left[\sum_k x_{ik} P_{ik} \right]^{-1}$$

(where $P_{ij} = \partial P / \partial x_{ij}$) on the domain

$$M \cup \partial M = \{ (x_{ij}) : x_{ij} \geq 0, \sum_{j=1}^{q_i} x_{ij} = 1, i = 1, \dots, k \}$$

is well defined and C^∞ on the set where no denominator $\sum_k x_{ik} P_{ik}$ vanishes. In particular, this is true in the interior M if we assume P depends on at least one of each set

$$S_i = \{ x_{ij} : j = 1 \dots q_i \} \quad i = 1, 2, \dots, k.$$

(If no term $x_{i_0 j}, j = 1 \dots q_{i_0}$ appears in P then we consider \mathcal{F}_P is acting on the reduced domain without the factor

$$\{ x_{i_0 j} \geq 0, \sum_{j=1}^{q_{i_0}} x_{i_0 j} = 1 \}.)$$

Now $\partial M = \bigcup_{i_0, j_0} N_{i_0 i_0} \cup \partial N_{i_0, j_0}$, where

$$N_{i_0 j_0} = \{ (x_{ij}) : x_{i_0 j_0} = 0, x_{ij} \neq 0 \text{ for } \langle i, j \rangle \neq \langle i_0, j_0 \rangle \}.$$

Let R_{i_1} be the sum of all terms of P that do not involve any factor $x_{i_1 j}, j = 1, \dots, q_{i_1}$. Then for any $\langle i_0, j_0 \rangle$ including $i_0 = i_1, P$ can be expressed as

$$P = x_{i_0 j_0}^l Q(x) + R_{i_1}$$

where Q does not have a common factor $x_{i_0 j_0}$, and hence does not vanish on $N_{i_0 j_0}$. Using this decomposition one can show that \mathcal{F} can be continuously extended to $N_{i_0 j_0}$ as follows:

If $\sum_k x_{ik} P_{ik} \neq 0$ on $N_{i_0 j_0}$, then

$$\mathcal{F} : x_{ij} \longrightarrow x_{ij} P_{ij} [\sum_k x_{ik} P_{ik}]^{-1}.$$

If $\sum_k x_{ik} P_{ik} = 0$ on $N_{i_0 j_0}$ we consider two cases, $i \neq i_0$ and $i = i_0$.

(a) If $i \neq i_0$, then $\sum_k x_{ik} Q_{ik} \neq 0$ and we define

$$\mathcal{F} : x_{ij} \longrightarrow x_{ij} Q_{ij} [\sum_k x_{ik} Q_{ik}]^{-1};$$

(b) If $i = i_0$, and $\sum_k x_{ik} Q_{ik} \neq 0$ we define

$$\mathcal{F} : x_{ij} \longrightarrow x_{ij} Q_{ij} [\sum_k x_{ik} Q_{ik}]^{-1}.$$

(c) If $i = i_0$, and $\sum_k x_{ik} Q_{ik} = 0$, we define

$$\mathcal{F} : x_{i,j} \longrightarrow 0, j \neq j_0, \quad \text{and} \quad \mathcal{F} : x_{i_0, j_0} \longrightarrow 1.$$

At a local maximum on $\bigcup_{ij} \partial N_{ij}$, while $P > 0$, this does not obviously imply that $\sum_{j=1}^{q_i} x_{ij} \partial P / \partial x_{ij} \neq 0$ for each i . Hence a proof that \mathcal{F} can be extended to be continuous at a local maximum of P is most easily made by a route other than that used for the special case where there is a single restraint equation. We proceed as follows: Assume that all variables x_{ij} appear in P . (Otherwise the discussion proceeds in a reduced space.) Then \mathcal{F}_P maps M into M , and for any point $x = (x_{ij})$ in M , the value of P at x is smaller than (or equal to) the value of P at any point along the line joining x to $\mathcal{F}(x)$. If x^0 is an isolated local maximum of P in $M \cup \partial M$, whether on the boundary ∂M or not, then for all sufficiently small $\varepsilon > 0$, the transformation \mathcal{F} maps the connected component of

$$\{(x) \mid P(x) > P(x^0) - \varepsilon\} \cap M$$

surrounding x^0 into itself. Since these sets form a base for the neighborhoods about x^0 , \mathcal{F} can be extended to be continuous at (x^0) by defining $\mathcal{F}(x^0) = x^0$.

Part C of Theorem 7 follows from the known inequalities for points in M and standard limiting arguments.

Corollaries 1 and 2 to Theorem 7 are now direct applications of the above arguments.

We are grateful to the referee for some helpful comments and suggestions.

REFERENCES

1. L. E. Baum, and J. A. Eagon, *An inequality with applications to statistical prediction for functions of Markov processes and to a model for ecology*, Bull. Amer. Math. Soc. **73** (1967), 360-363.
2. G. R. Blakley, *Homogeneous non-negative symmetric quadratic transformations*, Bull. Amer. Math. Soc. **70** (1964), 712-715.
3. G. H. Hardy, J. E. Littlewood, and G. Polya, *Inequalities*, Cambridge University Press, 1959.
4. F. W. Wilson, *The structure of the level surfaces of a Lyapunov function*, J. Differential Equations **3** (1967), 323-329.

Received May 18, 1967. This work was done while the second author was at the Institute for Defense Analyses on leave from the University of Minnesota.

INSTITUTE FOR DEFENSE ANALYSES
PRINCETON, NEW JERSEY

PACIFIC JOURNAL OF MATHEMATICS

EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R. PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
* * *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Pacific Journal of Mathematics

Vol. 27, No. 2

February, 1968

Leonard E. Baum and George Roger Sell, <i>Growth transformations for functions on manifolds</i>	211
Henry Gilbert Bray, <i>A note on CLT groups</i>	229
Paul Robert Chernoff, Richard Anthony Rasala and William Charles Waterhouse, <i>The Stone-Weierstrass theorem for valuable fields</i>	233
Douglas Napier Clark, <i>On matrices associated with generalized interpolation problems</i>	241
Richard Brian Darst and Euline Irwin Green, <i>On a Radon-Nikodym theorem for finitely additive set functions</i>	255
Carl Louis DeVito, <i>A note on Eberlein's theorem</i>	261
P. H. Doyle, III and John Gilbert Hocking, <i>Proving that wild cells exist</i>	265
Leslie C. Glaser, <i>Uncountably many almost polyhedral wild $(k - 2)$-cells in E^k for $k \geq 4$</i>	267
Samuel Irving Goldberg, <i>Totally geodesic hypersurfaces of Kaehler manifolds</i>	275
Donald Goldsmith, <i>On the multiplicative properties of arithmetic functions</i>	283
Jack D. Gray, <i>Local analytic extensions of the resolvent</i>	305
Eugene Carlyle Johnsen, David Lewis Outcalt and Adil Mohamed Yaqub, <i>Commutativity theorems for nonassociative rings with a finite division ring homomorphic image</i>	325
André (Piotrowsky) De Korvin, <i>Normal expectations in von Neumann algebras</i>	333
James Donald Kuelbs, <i>A linear transformation theorem for analytic Feynman integrals</i>	339
W. Kuich, <i>Quasi-block-stochastic matrices</i>	353
Richard G. Levin, <i>On commutative, nonpotent archimedean semigroups</i>	365
James R. McLaughlin, <i>Functions represented by Rademacher series</i>	373
Calvin R. Putnam, <i>Singular integrals and positive kernels</i>	379
Harold G. Rutherford, II, <i>Characterizing primes in some noncommutative rings</i>	387
Benjamin L. Schwartz, <i>On interchange graphs</i>	393
Satish Shirali, <i>On the Jordan structure of complex Banach* algebras</i>	397
Earl J. Taft, <i>A counter-example to a fixed point conjecture</i>	405
J. Roger Teller, <i>On abelian pseudo lattice ordered groups</i>	411