

# Pacific Journal of Mathematics

## **SUMS OF AUTOMORPHISMS OF A PRIMARY ABELIAN GROUP**

FRANK CASTAGNA

# SUMS OF AUTOMORPHISMS OF A PRIMARY ABELIAN GROUP

FRANK CASTAGNA

This paper is concerned with the problem: For which abelian groups  $G$  does the group of automorphisms of  $G$  generate the ring of endomorphisms of  $G$ ? R. S. Pierce has shown that if  $G$  is a 2-primary group, all of whose finite Ulm invariants are equal to one, then the subring of  $E(G)$  generated by the group of automorphisms of  $G$  is properly contained in  $E(G)$ . The groups considered in this paper are  $p$ -primary abelian groups where  $p$  is a fixed prime number greater than two. The paper is divided into four parts. In §1 the following theorem is proved:

**THEOREM.** If  $G$  is a countable reduced  $p$ -primary, ( $p > 2$ ), abelian group, then every endomorphism of  $G$  is a sum of two automorphisms.

The second part gives an extension of this theorem to the case where  $G$  is the direct sum of such groups. In §3 the result of this theorem is established for torsion complete abelian groups, using some known results about their endomorphism rings. Finally an example is given (for an arbitrary prime  $p$ ) of a reduced  $p$ -primary abelian group  $G$  for which there are endomorphisms in  $E(G)$  that are not sums of automorphisms.

**NOTATION.** Throughout this paper  $p$  will represent a fixed prime number greater than two. Mainly the notation will follow that of [3] and [5]. All groups considered are assumed to be  $p$ -primary abelian groups. If  $G$  is such a group, we define for each ordinal  $\beta$  the subgroup  $G_\beta$  as follows:  $G_1 = pG = \{x \in G \mid x = pg, g \in G\}$ ,  $G_\beta = pG_{\beta-1}$  if  $\beta - 1$  exists and  $G_\beta = \bigcap_{\alpha < \beta} G_\alpha$  if  $\beta$  is a limit ordinal. A primary group  $G$  is said to be divisible if  $pG = G$  and is said to be reduced if 0 is its only divisible subgroup. The height function  $h_G$  for a reduced  $p$ -group  $G$  is defined by the conditions

$$\begin{aligned} h_G(x) &= \alpha \text{ if } x \in G_\alpha \text{ and } x \notin G_{\alpha+1} \\ h_G(0) &= \infty \text{ where } \infty > \alpha \text{ for each ordinal } \alpha. \end{aligned}$$

If there is no possibility of confusion, we will write  $h(x)$  for  $h_G(x)$ .

A subgroup  $H$  of  $G$  is pure in  $G$  if  $p^n G \cap H = p^n H$  for each  $n \geq 0$ . For every integer  $e \geq 0$  we define  $G[p^e]$  as

$$G[p^e] = \{x \in G \mid p^e x = 0\}.$$

If  $x \in G$ ,  $o(x)$  is defined to be the order of  $x$ .

Finally for any subset  $X$  of  $G$ , let  $\{X\}$  denote the subgroup of  $G$  which is generated by  $X$ .

**1. Countable groups.** One of the fundamental theorems in abelian groups is Ulm's theorem, [7], which states that any two countable reduced  $p$ -primary abelian groups having the same Ulm invariants are isomorphic. (For a definition of the Ulm invariants see [5], p. 27.) A proof of this theorem is given in [5]. Roughly speaking the concept of the proof is to build an isomorphism between the two groups  $G$  and  $H$ . It is clear that one could by similar methods build an automorphism of the group  $G$ , taking  $G$  and  $H$  to be the same group. Given an endomorphism of  $G$  we might then ask, "Could one not build two automorphisms of  $G$  in such a way that their sum is always the given endomorphism?" This is indeed possible and we get the following theorem.

**THEOREM 1.1.** *If  $G$  is a countable, reduced  $p$ -primary abelian group, then every endomorphism of  $G$  is a sum of two automorphisms.*

The proof of 1.1 is based on several lemmas. We begin with the following definition.

**DEFINITION.** If  $S$  is a subgroup of  $G$ , an element  $x \in G - S$  is said to be proper with respect to  $S$  if  $h(x + s) \leq h(x)$  for all  $s \in S$ .

Let  $S$  be a subgroup of  $G$ , let  $x, y \in G$  with  $h(x) < h(y)$ , and let  $n$  be an integer prime to  $p$ . It is an immediate consequence of this definition that if any one of  $x, x + y$ , or  $nx$  is proper with respect to  $S$ , then so are the other two.

Now suppose that  $S$  and  $T$  are subgroups of  $G$  and that  $\theta$  is a height preserving isomorphism between them. Assume that there exist  $x$  and  $z \in G$ , having height  $\alpha$ , and satisfying

(i)  $px \in S$  and  $\theta(px) = pz$ ,

(ii)  $x$  and  $z$  are proper with respect to  $S$  and  $T$  respectively. Then  $\theta$  can be extended to a height preserving isomorphism  $\theta'$  between  $\{S, x\}$  and  $\{T, z\}$  by defining

$$\theta'(s + nx) = \theta(s) + nz.$$

Some of the lemmas that follow can be found in [5], § 11, and so, mainly, the proofs will be omitted.

**LEMMA 1.2.** *Let  $S$  and  $T$  be subgroups of  $G$  and let  $\theta$  be a height preserving isomorphism between them. Assume that  $x \in G$  satisfies*

- (i)  $x$  is proper with respect to  $S$ ,
- (ii) if  $w \in x + S$  is proper with respect to  $S$  then  $h(pw) \leq h(px)$ ,
- (iii)  $px \in S$  and  $h(px) = h(x) + 1$ .

Then there is a  $z \in G$  with  $pz = \theta(px)$  and  $h(z) = h(x)$ . Furthermore any  $z$  with these two properties will be proper with respect to  $T$ .

If  $S$  is any subgroup of  $G$ , define  $S_\alpha$  by  $S_\alpha = S \cap G_\alpha$ . Define the subgroup  $S_\alpha^*$  of  $S_\alpha$  to be

$$S_\alpha^* = \{x \in S_\alpha \mid px \in G_{\alpha+2}\}.$$

The quotient  $S_\alpha^*/S_{\alpha+1}$  can be considered as a vector space over  $Z_p$ .

**LEMMA 1.3.** *If  $S$  is a Subgroup of  $G$  then there exists a monomorphism  $\zeta^*$  of  $S_\alpha^*/S_{\alpha+1}$  into  $G_\alpha[p]/G_{\alpha+1}[p]$ .*

Since the technique involved in the proof of 1.3 is vital to other parts of this paper the proof will be outlined.

*Proof.* If  $x \in S_\alpha^*$  then  $px \in G_{\alpha+2}$  and there is a  $y \in G_{\alpha+1}$  such that  $px = py$ . The mapping

$$x \rightarrow x - y + G_{\alpha+1}[p]$$

has  $S_{\alpha+1}$  as its kernel.

**LEMMA 1.4.** *Let  $S$  and  $\zeta^*$  be as in 1.3. If  $x \in G_\alpha[p] - G_{\alpha+1}[p]$  then  $x + G_{\alpha+1}[p] \notin \text{Im } \zeta^*$  if and only if  $x$  is proper with respect to  $S$ .*

**LEMMA 1.5.** *Let  $S$  and  $T$  be finite subgroups of  $G$  and let  $\theta$  be a height preserving isomorphism between them. Suppose there is an  $x \in G$  of height  $\alpha$ , which is proper with respect to  $S$ , and such that  $h(px) > \alpha + 1$ . Then there exists  $z \in G_\alpha[p]$  such that  $h(z) = \alpha$  and  $z$  is proper with respect to  $T$ . Furthermore  $z$  can be chosen so that it is also proper with respect to  $S$  and has the following form*

$$z = x + s_0 + w$$

where  $s_0 \in S_\alpha^*$  and  $h(w) > \alpha$ .

*Proof.* Choose  $v \in G_{\alpha+1}$  such that  $x - v \in G_\alpha[p]$ . If  $x$  is proper with respect to  $T$  we let  $s_0 = 0$  and  $w = -v$ . Otherwise let  $\zeta^*$  and  $\xi^*$  be the monomorphisms of  $S_\alpha^*/S_{\alpha+1}$  and  $T_\alpha^*/T_{\alpha+1}$  into  $G_\alpha[p]/G_{\alpha+1}[p]$  respectively. Both  $\text{Im } \xi^*$  and  $\text{Im } \zeta^*$  have the same finite dimension and since, by 1.4,

$$x - v + G_{\alpha+1}[p] \in \text{Im } \xi^* - \text{Im } \zeta^*$$

it follows that there is a  $y \in G_\alpha[p]$  such that

$$y + G_{\alpha+1}[p] \in \text{Im } \zeta^* - \text{Im } \xi^* .$$

From 1.3 we see that  $y$  has the form

$$y = s_0 + u, s_0 \in S_\alpha^* \quad \text{and} \quad h(u) > \alpha .$$

We let  $z = x + s_0 - (u + v)$ .

We can now proceed with a proof of 1.1. Let  $\psi \in E(G)$  and enumerate  $G$  as  $G = \{x_1 = 0, x_2, x_3, \dots\}$ . Suppose that we have finite subsets  $S, T$ , and  $T'$  and height preserving isomorphisms  $\theta$  between  $S$  and  $T$  and  $\theta'$  between  $S$  and  $T'$  such that  $\theta(s) + \theta'(s) = \psi(s)$  for each  $s \in S$ . Assume that

$$x_m \notin S \cap T \cap T' .$$

We must extend  $\theta$  and  $\theta'$  to height preserving isomorphisms  $\bar{\theta}$  and  $\bar{\theta}'$  between finite subgroups  $\bar{S}$  containing  $S$ ,  $\bar{T}$  containing  $T$ , and  $\bar{T}'$  containing  $T'$  such that  $x_m \in \bar{S} \cap \bar{T} \cap \bar{T}'$  and for all  $s \in \bar{S}$ ,  $\bar{\theta}(s) + \bar{\theta}'(s) = \psi(s)$ . The proof is broken into two cases:  $x_m \notin S$  and  $x_m \in T \cap T'$ .

*Case I.*  $x_m \in S$ . We may assume that we have an  $x$  which is proper with respect to  $S$ ,  $px \in S$  and such that if  $y \in x + S$  and  $h(y) = h(x)$  then  $h(py) \leq h(px)$ . Let  $h(x) = \alpha$ ,  $\theta(px) = y$ , and  $\theta'(px) = y'$ .

If  $h(y) = \alpha + 1$ , we can use 1.2 to get a  $z$  which is proper with respect to  $T$  and such that  $h(z) = \alpha$  and  $pz = \theta(px)$ . Define  $z' = \psi(x) - z$ . Then  $h(z') = \alpha$  and  $pz' = \theta'(px)$ . Again using 1.2 we see that  $z'$  is proper with respect to  $T'$ . Letting  $\bar{S} = \{S, x\}$ ,  $\bar{T} = \{T, z\}$ , and  $\bar{T}' = \{T', z'\}$ , the extensions  $\bar{\theta}$  and  $\bar{\theta}'$  of  $\theta$  and  $\theta'$  follow. Since  $z + z' = \psi(x)$  it is clear that  $\bar{\theta}(s) + \bar{\theta}'(s) = \psi(s)$  for all  $s \in \bar{S}$ .

Suppose then that  $h(y) > \alpha + 1$ . By 1.5 there exists  $z_0 \in G_\alpha[p]$ , of height  $\alpha$ , which is proper with respect to  $T$ . We have  $h(pz_0) = h(0) > \alpha + 1$ . Since  $\theta'\theta^{-1}$  is a height preserving isomorphism between  $T$  and  $T'$  we can apply 1.5 again to obtain  $z_1 \in G_\alpha[p]$ , of height  $\alpha$ , which is proper with respect to both  $T$  and  $T'$ . Choose  $u$  and  $u' \in G_{\alpha+1}$  such that  $pu = y$  and  $pu' = y'$ .

If  $h(\psi(x)) > \alpha$  let

$$z = z_1 + u \quad \text{and} \quad z' = \psi(x) - z .$$

A few easy calculations show that with these definitions of  $z$  and  $z'$  the desired extensions follow.

Thus we may suppose that  $h(\psi(x)) = \alpha$ . Letting  $w = u + u'$  we see that  $h(\psi(x) - w) = \alpha$  and  $\psi(x) - w \in G_\alpha[p]$ . Now if  $\psi(x) - w$  is proper with respect to both  $T$  and  $T'$  choose  $v$  such that  $2v = \psi(x) - w$ . Then  $h(v) = \alpha$ ,  $v \in G_\alpha[p]$ , and  $v$  is proper with respect to both  $T$  and

$T'$ . Define

$$z = v + u \quad \text{and} \quad z' = v + u'$$

to get the extensions of  $\theta$  and  $\theta'$ . Therefore we may assume that  $\psi(x) - w$  is not proper with respect to  $T$ . With  $z_1$  as defined above it follows from 1.4 that  $\psi(x) - w - z_1$  is proper with respect to  $T$ . Then with the definitions

$$z = \psi(x) - w - z_1 + u \quad \text{and} \quad z' = z_1 + u'$$

we have completed the proof of Case I.

*Case II.*  $x_m \notin T \cap T'$ . To be explicit, let us say that  $x_m \notin T$ . Then as before, the problem reduces to that of having a  $z$  which is proper with respect to  $T$  and  $pz \in T$ . Let  $h(z) = \alpha$  and  $pz = y$ . The case  $h(y) = \alpha + 1$  is handled just as before. Use 1.2 to get  $x$  and let  $z' = \psi(x) - z$ .

Consider then the situation when  $h(y) > \alpha + 1$ . We use 1.5 to select  $w_0 \in G_\alpha[p]$ , of height  $\alpha$ , such that  $w_0$  is proper with respect to both  $T$  and  $T'$  and has the form

$$w_0 = z + t_0 + u_0,$$

where  $t_0 \in T_\alpha^*$  and  $h(u_0) > \alpha$ .

Let  $\zeta^*$  and  $\gamma^*$  be the monomorphisms of  $S_\alpha^*/S_{\alpha+1}$  and  $T_\alpha'^*/T_{\alpha+1}'$  into  $G_\alpha[p]/G_{\alpha+1}[p]$  respectively as given by 1.3. Let  $S^*$  be a complimentary summand of  $\text{Im } \zeta^*$  in  $G_\alpha[p]/G_{\alpha+1}[p]$ :

$$G_\alpha[p]/G_{\alpha+1}[p] = \text{Im } \zeta^* \oplus S^*.$$

Let  $\psi^*$  be the endomorphism of  $G_\alpha[p]/G_{\alpha+1}[p]$  induced by  $\psi$ . Choose  $x_1 + G_{\alpha+1}[p] \in S^*$  such that  $x_1 \in G_{\alpha+1}[p]$ . Since  $p > 2$  there is a  $k \not\equiv 0 \pmod{p}$  such that

$$k(w_0 + G_{\alpha+1}[p]) + \psi^*(x_1 + G_{\alpha+1}[p]) \notin \text{Im } \gamma^*.$$

Let  $l$  satisfy

$$kl \equiv 1 \pmod{p}.$$

Then

$$(w_0 + \psi(lx_1)) + G_{\alpha+1}[p] \notin \text{Im } \gamma^*.$$

It follows that  $lx_1$  is proper with respect to  $S$  and  $w_0 + \psi(lx_1)$  is proper with respect to  $T'$ . Define

$$w = w_0 - u_0 = z + t_0.$$

Then  $pw \in T$  and  $h(\theta^{-1}(pw)) > \alpha + 1$ . Choose  $v \in G_{\alpha+1}$  such that  $pv = \theta^{-1}(pw)$ . Define  $x$  and  $w'$  by

$$\begin{aligned}x &= -lx_1 + v \\w' &= \psi(x) - w.\end{aligned}$$

It is routine to check that the subgroups  $\{S, x\}$ ,  $\{T, w\}$ , and  $\{T', w'\}$  give rise to the desired extensions of  $\theta$  and  $\theta'$ . This completes the proof of 1.1.

**2. Direct sums of countable groups.** In this section we will be considering subgroups  $G_\lambda, H_\beta$ , etc., of a group  $G$ , where  $\lambda$  and  $\beta$  are members of an index set  $A$ . These are not the subgroups defined in the notation. To avoid confusion, we will denote the subgroups  $G_\alpha$ , defined previously, by  $p^\alpha G$ ,  $p^\alpha H_\beta$ , etc.

The purpose of this section is to extend Theorem 1.1 to reduced  $p$ -primary abelian groups  $G$  that are direct sums of countable groups. In the next lemma and theorem, however, we make no restriction on the summands except that they be countable.

**LEMMA 2.1.** *Let  $G = \sum_{\lambda \in A} G_\lambda$  where  $|G_\lambda| \leq \aleph_0$  for all  $\lambda \in A$  and let  $\psi \in E(G)$ . Then  $A = \bigcup_{\gamma \in A} I_\gamma$  where  $|I_\gamma| \leq \aleph_0$  for each  $\gamma \in A$  and*

$$\psi(\sum_{\lambda \in I_\gamma} G_\lambda) \subseteq \sum_{\lambda \in I_\gamma} G_\lambda.$$

*Proof:* Let  $\gamma$  be a fixed element of  $A$  and define  $\Gamma_0 = \{\gamma\}$ . Having defined the countable subset  $\Gamma_h$  of  $A$  we define  $\Gamma_{h+1}$  as follows. Since  $\sum_{\lambda \in \Gamma_h} G_\lambda$  is a countable set there is a countable subset  $\Gamma_{h+1}$  of  $A$  which contains  $\Gamma_h$  and satisfies

$$\psi(\sum_{\lambda \in \Gamma_h} G_\lambda) \subseteq \sum_{\lambda \in \Gamma_{h+1}} G_\lambda.$$

Define

$$I_\gamma = \bigcup_{h < \omega} \Gamma_h.$$

**THEOREM 2.2.** *Let  $G = \bigoplus \sum_{\lambda \in A} G_\lambda$  where each  $G_\lambda$  is countable and let  $\psi \in E(G)$ . Then  $G$  can be written as  $G = \bigcup_{\beta \in A} H_\beta$  where each  $H_\beta$  has the following properties.*

- (i)  $H_\beta \subseteq H_{\beta+1}$  for all  $\beta \in A$ ,
- (ii)  $H_\beta = \bigcup_{\alpha < \beta} H_\alpha$  if  $\beta$  is a limit cardinal,
- (iii)  $H_\beta = H_{\beta-1} \oplus C_\beta$ , where  $C_\beta$  is countable, if  $\beta$  is not a limit ordinal,
- (iv) for each  $\beta \in A$ ,  $\psi(H_\beta) \subseteq H_\beta$ .

*Proof.* Let  $A = \bigcup_{\gamma \in A} I_\gamma$  as in 2.1. Define  $\Gamma_0 = I_0$  and having defined  $\Gamma_\alpha$  for all  $\alpha < \beta$ , define  $\Gamma_\beta = \Gamma_{\beta-1} \cup I_\beta$  if  $\beta - 1$  exists and  $\Gamma_\beta = \bigcup_{\alpha < \beta} \Gamma_\alpha$  if  $\beta$  is a limit ordinal. Now define  $H_\beta = \bigoplus \sum_{\lambda \in \Gamma_\beta} G_\lambda$ . We note that if  $\beta$  is not a limit ordinal

$$H_\beta = H_{\beta-1} \oplus \sum_{\lambda \in K_\beta} G_\lambda$$

where  $K_\beta = I_\beta - \Gamma_{\beta-1}$ . It is easily seen that the assertions (i)-(iv) are satisfied.

**THEOREM 2.3.** *If  $G$  is a reduced  $p$ -primary abelian group, ( $p > 2$ ), and if  $G$  is a direct sum of countable groups, then every endomorphism of  $G$  is a sum of two automorphisms.*

*Proof.* Let  $G = \bigcup_{\gamma \in A} H_\gamma$  as is given in 2.2 and let  $\psi \in E(G)$ . We define inductively, for each  $\gamma \in A$ , automorphisms  $\theta_\gamma$  and  $\theta'_\gamma$  such that  $\psi|_{H_\gamma} = \theta_\gamma + \theta'_\gamma$  and if  $\alpha < \gamma$ ,  $\theta_\gamma|_{H_\alpha} = \theta_\alpha$  and  $\theta'_\gamma|_{H_\alpha} = \theta'_\alpha$ . For  $\gamma = 0$ , 1.1 gives  $\theta_0$  and  $\theta'_0$ . Assume that  $\theta_\gamma$  and  $\theta'_\gamma$  have been defined for all  $\gamma < \beta$ . If  $\beta$  is a limit ordinal we define  $\theta_\beta = \bigcup_{\gamma < \beta} \theta_\gamma$  and  $\theta'_\beta = \bigcup_{\gamma < \beta} \theta'_\gamma$ . Assume that  $\beta - 1$  exists. Then  $H_\beta = H_{\beta-1} \oplus C_\beta$ . Let  $\pi_1$  and  $\pi_2$  be the projections of  $H_\beta$  onto  $H_{\beta-1}$  and  $C_\beta$  respectively. Then  $\pi_2(\psi|_{C_\beta}) \in E(C_\beta)$  and by 1.1 there exist  $\phi$  and  $\phi'$ , automorphisms of  $C_\beta$ , such that  $\pi_2(\psi|_{C_\beta}) = \phi + \phi'$ . For each  $c \in C$  choose  $v_c \in H_{\beta-1}$  such that  $2v_c = \pi_1\psi(c)$ . Plainly  $v_c$  defines a homomorphism of  $C_\beta$  into  $H_{\beta-1}$ . Now define  $\theta_\beta$  and  $\theta'_\beta$  on  $H_\beta$  by

$$\theta_\beta(x + c) = \theta_{\beta-1}(x) + \phi(c) + v_c$$

$$\theta'_\beta(x + c) = \theta'_{\beta-1}(x) + \phi'(c) + v_c$$

where  $x \in H_{\beta-1}$  and  $c \in C_\beta$ . It is routine to check that  $\theta_\beta$  and  $\theta'_\beta$  satisfy the required conditions.

**3. An application.** In this section we show that if  $G$  is a torsion complete group, (in the  $p$ -adic topology), then each endomorphism of  $G$  is a sum of two automorphisms. This result will follow as a corollary of a theorem in [6].

**NOTATION.**  $B_n = \bigoplus \sum_{i \in I_n} \{b_{in}\}$ , for some index set  $I_n$  and  $o(b_{in}) = p^{n+1}$ .

$$B = \bigoplus \sum_{n < \omega} B_n$$

$$\bar{B} = \text{torsion subgroup of } \prod_{n < \omega} B_n$$

$$C_n = \{p^n \bar{B}, B_n, B_{n+1}, \dots\}.$$

It is not difficult to show that

$$(1) \quad \bar{B} = \bigoplus \sum_{k < n} B_k \oplus C_n.$$

Let  $\pi_n$  be the natural projection of  $\bar{B}$  onto  $C_n$  as determined by (1). Define  $\rho_n = \pi_n - \pi_{n+1}$ . Then  $\rho_n$  is a projection of  $\bar{B}$  onto  $B_n$ . Define  $\lambda_n: E(\bar{B}) \rightarrow E(B_n[p])$  by

$$(\lambda_n \phi)(x) = \rho_n(\phi(x)) \quad \text{for all } x \in B_n[p].$$



LEMMA 3.1.  $\lambda_n$  is a ring homomorphism of  $E(\bar{B})$  into  $E(B_n[p])$ .

*Proof.* If  $\phi \in E(\bar{B})$  then plainly  $\lambda_n \phi$  maps  $B_n[p]$  to  $B_n[p]$ . It is also clear that  $\lambda_n$  is a group homomorphism of  $E(\bar{B})$  to  $E(B_n[p])$ . Note now that  $x \in C_n[p]$  implies  $\phi(x) \in C_n[p]$ . Suppose that  $\phi, \psi \in E(\bar{B})$ . For  $i \in I_n$ , we can write

$$\psi(p^n b_{i_n}) = \sum_{j \in I_n} \alpha_{ij} p^n b_{j_n} + x$$

where  $x \in C_{n+1}[p]$ . Consequently

$$\lambda_n \psi(p^n b_{i_n}) = \sum_{j \in I_n} \alpha_{ij} p^n b_{j_n}$$

so that

$$\psi(p^n b_{i_n}) = \lambda_n \psi(p^n b_{i_n}) + x.$$

Then

$$\lambda_n \phi \psi(p^n b_{i_n}) = \rho_n \phi(\lambda_n \psi(p^n b_{i_n}) + x) = (\lambda_n \phi)(\lambda_n \psi) p^n b_{i_n}.$$

LEMMA 3.2. Define

$$\lambda: E(\bar{B}) \rightarrow \prod_{n < \omega} E(B_n[p])$$

by

$$\lambda: \phi \rightarrow (\lambda_0 \phi, \lambda_1 \phi, \dots, \lambda_n \phi, \dots).$$

Then  $\lambda$  is a ring epimorphism.

*Proof.* Certainly  $\lambda$  is a ring homomorphism. Let

$$(\phi'_0, \phi'_1, \dots, \phi'_n, \dots) \in \pi_{n < \omega} E(B_n[p]),$$

where  $\phi'_n \in (B_n[p])$ . Extend  $\phi'_n$  to  $\phi_n^* \in E(B_n)$ . Now define  $\phi \in E(\bar{B})$  as follows. If  $x \in \bar{B}$ ,  $x = \lim_m \sum_{k < m} \rho_k(x)$ , (the limit being taken in the  $p$ -adic topology) and hence the elements

$$\sum_{k < m} \phi_k^* \rho_k(x), \quad m = 1, 2, \dots$$

form a Cauchy sequence in  $\bar{B}$ , with bounded order. Define  $\phi(x)$  to be the limit of this sequence. If  $x \in B_n[p]$  then  $\phi(x) = \phi'_n(x) \in B_n[p]$  and hence

$$\lambda \phi = (\phi'_0, \phi'_1, \dots, \phi'_n, \dots).$$

LEMMA 3.3. Let  $R$  be a ring and  $J(R)$  the Jacobson radical of  $R$ . If each element of  $R/J(R)$  is a sum of  $n$  units then each element of  $R$  is a sum of  $n$  units.

*Proof.* If  $y + J(R)$  is a unit then it has an inverse  $z + J(R)$  and  $zy + J(R) = 1 + J(R)$ . This implies that  $zy = 1 - r$  for some  $r \in J(R)$

and hence  $zy$  is a unit. Thus  $((zy)^{-1}z)y = 1$  so that  $y$  has a left inverse. Similarly  $y$  has a right inverse and hence  $y$  is a unit. It now follows easily that if  $x + J(R)$  is a sum of  $n$  units in  $R/J(R)$  then  $x$  is a sum of  $n$  units in  $R$ .

**THEOREM 3.4.**  $\prod_{n < \omega} E(B_n[p]) \cong E(\bar{B})/J(E(\bar{B}))$ .

*Proof.* Let  $\lambda$  be as in 3.2. We prove  $\text{Ker } \lambda = J(E(\bar{B}))$ . Suppose  $\lambda(\phi) = 0$ . Then for each  $n, x \in B_n[p]$  implies  $\phi(x) \in C_{n+1}$ . Hence, if  $x \neq 0$ ,  $h(\phi(x)) > h(x)$ . That is,

$$\phi \in H(\bar{B}) = \{\phi \in E(\bar{B}) \mid x \in \bar{B}[p], x \neq 0, \text{ implies } h(\phi(x)) > h(x)\}.$$

Conversely, if  $\phi \in H(\bar{B})$ , then clearly  $\lambda\phi = 0$ . The result now follows from [6], p. 287, which states that  $J(E(\bar{B})) = H(\bar{B})$ .

**COROLLARY 3.5.** *If  $\phi \in E(\bar{B})$  then  $\phi$  is the sum of two automorphisms.*

*Proof.*  $B_n[p]$  is a direct sum of countable groups and therefore, by 2.3, each  $\phi_n \in E(B_n[p])$  is a sum of two automorphisms from which it follows that each element of  $E(\bar{B})/J(E(\bar{B}))$  is a sum of two units. consequently each element of  $E(\bar{B})$  is a sum of two units.

**4. An example.** In this section we exhibit, for an arbitrary prime  $p$ , a reduced  $p$ -primary group  $G$  for which the group of automorphisms does not generate the ring of endomorphisms. The group  $G$  will in fact be without nonzero elements of infinite height and have a countable basic subgroup.

The notation will be as in § 3 except that we require each index set  $I_n$  to be countably infinite.

Define  $\tau$  on  $\bar{B}$  by the condition

$$\tau(b_{m+n}) = b_{m+1+n}.$$

Plainly this property uniquely determines an endomorphism of  $\bar{B}$ . Let  $R$  be the subring of  $E(\bar{B})$  generated by the identity and  $\tau$ .

**LEMMA 4.1.** *If  $\phi \in R - pR$  then  $\phi$  is one-to-one.*

*Proof.* Let  $\phi = \sum_{i=0}^r k_i \tau^i$  and suppose  $\phi(x) = 0$  for some  $x \in \bar{B}[p] - \{0\}$ . We show that  $\phi \in pR$ . Indeed suppose otherwise. Then we can assume that  $p$  does not divide  $k_r$ . Write  $x = \sum_{m < \omega} (\sum_{n < \omega} a_{mn} p^n b_{mn})$  where  $a_{mn} \in \mathbb{Z}$  and, for each  $n$ , almost all  $a_{mn}$  are zero. We have

$$0 = \phi(x) = \sum_{n < \omega} \left( \sum_{m < \omega} a_{mn} p^n \left( \sum_{i=0}^r k_i b_{m+i+n} \right) \right).$$

Since  $x \neq 0$ , we can choose  $n$  so that  $\sum_{m < \omega} a_{mn} p^n b_{mn} \neq 0$  and hence, for this  $n$ , there is a largest  $m = M$  such that  $p$  does not divide  $a_{Mn}$ . It then follows that

$$\sum_{m < \omega} a_{mn} p^n \left( \sum_{i=0}^r k_i b_{m+i, n} \right) = a_{Mn} k_r p^n b_{M+r, n} + \sum_{j < M+r} c_j p^n b_{jn},$$

for some integers  $c_j$ , and from this expression it follows that  $\phi(x) \neq 0$ .

Let  $\mathcal{C}$  be the collection of subrings  $S$  of  $E(\bar{B})$  which satisfy, for all nonnegative integers  $e$ , the condition

$$C_e: \phi \in S \quad \text{and} \quad \phi((p^n \bar{B})[p^e]) = 0 \quad \text{for some } n \text{ implies } \phi \in P^e S.$$

We note that if  $S$  is a subring of  $E(\bar{B})$  which satisfies  $C_1$  then  $S \in \mathcal{C}$ . Clearly our subring  $R \in \mathcal{C}$ .

Let  $\bar{R}$  be the closure of  $R$  in the  $p$ -adic topology on  $E(\bar{B})$ .

LEMMA 4.2. *If  $\phi \in \bar{R} - p\bar{R}$  then  $\phi$  is one-to-one.*

*Proof.* Let  $\{\phi_n\}_{n=1}^\infty$  be a Cauchy sequence in  $R$  which converges to  $\phi$ . We can assume that, for all  $n > 0$ ,  $\phi_n - \phi_{n+1} \in p^n R$ . Now note that if  $\phi_r \in pR$  for some  $r$  then  $\phi_n \in pR$  for all  $n$ . Hence we could write  $\phi_n = p\psi_n$ ,  $\psi_n \in R$ . Then

$$\phi = \lim_n \phi_n = \lim_n p\psi_n = p \lim_n \psi_n = p\psi, \quad \psi \in \bar{R}.$$

Thus we can assume no  $\phi_n \in pR$ .

Let  $x \in \bar{B}[p]$ ,  $x \neq 0$ . We can write  $x = b_m + y$  where  $0 \neq b_m \in B_m$  and  $h(y) > m$ . Since the conditions  $\theta \in R - pR$  and  $0 \neq b_n \in B_n$  always imply  $0 \neq \theta(b_n) \in B_n$  it follows that  $h(\phi_n(x)) \leq m$  for all  $n$ . Hence

$$\phi(x) = \lim \phi_n(x) \neq 0.$$

It is clear that  $\bar{R} \in \mathcal{C}$ . Furthermore  $\bar{R}$  is a closed separable subring of  $E(\bar{B})$ . Thus  $\bar{R}$  satisfies the hypothesis of the following theorem by A. L. S. Corner [1].

THEOREM 4.3. *Let  $C$  be a torsion complete  $p$ -group with unbounded countable basic subgroup  $B$ , and let  $\Phi$  be a separable closed subring of  $E(C)$  such that  $\Phi(B) \subseteq B$  and, for all positive integers  $e$ ,  $\Phi$  satisfies  $C_e$ . Then there is a pure subgroup  $G$  of  $C$  containing  $B$  such that*

$$E(G) = \Phi \oplus E_s(G).$$

Here  $E_s(G)$  is the subring of small endomorphisms of  $G$ . Note, too, that if  $G$  is pure in  $C$  we can consider  $E(G)$  as being embedded in  $E(C)$ .

Now if  $G$  is a pure subgroup of  $\bar{B}$  corresponding to  $\bar{R}$  in this

theorem we can write

$$E(G) = \bar{R} \oplus E_s(G) .$$

We then get the following epimorphism of  $E(G)$

$$\gamma: E(G) \rightarrow \bar{R} \rightarrow \bar{R}/p\bar{R} \simeq R/pR \simeq Z_p[X]$$

where  $X$  is an indeterminate. The only units in  $Z_p[X]$  are the nonzero constant polynomials. Thus, the nonconstant polynomials cannot be written as sums of units and therefore, their pre-images in  $E(G)$  cannot be written as sums of automorphisms.

The author wishes to thank Professor R. S. Pierce for suggesting the problem and for his suggestions and encouragement in solving it.

#### BIBLIOGRAPHY

1. A.L.S. Corner, *On endomorphism rings of primary abelian groups* (to appear)
2. L. Fuchs, *Recent results and problems on abelian groups*, Topics in Abelian Groups, Chicago, 1963.
3. ———, *Abelian groups*, Budapest, 1958.
4. N. Jacobson, *Structure of Rings*, Providence, 1956.
5. I. Kaplansky, *Infinite Abelian Groups*, Ann Arbor, 1954.
6. R. S. Pierce, *Homomorphisms of primary abelian groups*, Topics in Abelian Groups, Chicago, 1963.
7. H. Ulm, *Zur Theorie der abzählbarunendlichen abelschen Gruppen*, Math. Ann. **107** (1933), 774-803.

Received January 15, 1968. This paper is essentially the author's dissertation which was written at the University of Washington.

NEW MEXICO STATE UNIVERSITY

AND

WAYNE STATE UNIVERSITY



# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN

Stanford University  
Stanford, California

J. DUGUNDJI

Department of Mathematics  
University of Southern California  
Los Angeles, California 90007

R. R. PHELPS

University of Washington  
Seattle, Washington 98105

RICHARD ARENS

University of California  
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH

B. H. NEUMANN

F. WOLF

K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA  
CALIFORNIA INSTITUTE OF TECHNOLOGY  
UNIVERSITY OF CALIFORNIA  
MONTANA STATE UNIVERSITY  
UNIVERSITY OF NEVADA  
NEW MEXICO STATE UNIVERSITY  
OREGON STATE UNIVERSITY  
UNIVERSITY OF OREGON  
OSAKA UNIVERSITY  
UNIVERSITY OF SOUTHERN CALIFORNIA

STANFORD UNIVERSITY  
UNIVERSITY OF TOKYO  
UNIVERSITY OF UTAH  
WASHINGTON STATE UNIVERSITY  
UNIVERSITY OF WASHINGTON  
\* \* \*  
AMERICAN MATHEMATICAL SOCIETY  
CHEVRON RESEARCH CORPORATION  
TRW SYSTEMS  
NAVAL WEAPONS CENTER

---

Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California 90024.

Each author of each article receives 50 reprints free of charge; additional copies may be obtained at cost in multiples of 50.

---

The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is \$8.00; single issues, \$3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: \$4.00 per volume; single issues \$1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley 8, California.

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners of publishers and have no responsibility for its content or policies.

Charles A. Akemann, <i>Invariant subspaces of <math>C(G)</math></i> .....	421
Dan Amir and Zvi Ziegler, <i>Generalized convexity cones and their duals</i> ....	425
Raymond Balbes, <i>On <math>(J, M, m)</math>-extensions of order sums of distributive lattices</i> .....	441
Jan-Erik Björk, <i>Extensions of the maximal ideal space of a function algebra</i> .....	453
Frank Castagna, <i>Sums of automorphisms of a primary abelian group</i> .....	463
Theodore Seio Chihara, <i>On determinate Hamburger moment problems</i> .....	475
Zeev Ditzian, <i>Convolution transforms whose inversion function has complex roots in a wide angle</i> .....	485
Myron Goldstein, <i>On a paper of Rao</i> .....	497
Velmer B. Headley and Charles Andrew Swanson, <i>Oscillation criteria for elliptic equations</i> .....	501
John Willard Heidel, <i>Qualitative behavior of solutions of a third order nonlinear differential equation</i> .....	507
Alan Carleton Hindmarsh, <i>Pick's conditions and analyticity</i> .....	527
Bruce Ansgar Jensen and Donald Wright Miller, <i>Commutative semigroups which are almost finite</i> .....	533
Lynn Clifford Kurtz and Don Harrell Tucker, <i>An extended form of the mean-ergodic theorem</i> .....	539
S. P. Lloyd, <i>Feller boundary induced by a transition operator</i> .....	547
Henry B. Mann, Josephine Mitchell and Lowell Schoenfeld, <i>A new proof of the maximum principle for doubly-harmonic functions</i> .....	567
Robert Einsohn Mosher, <i>The product formula for the third obstruction</i> .....	573
Sam Bernard Nadler, Jr., <i>Sequences of contractions and fixed points</i> .....	579
Eric Albert Nordgren, <i>Invariant subspaces of a direct sum of weighted shifts</i> .....	587
Fred Richman, <i>Thin abelian <math>p</math>-groups</i> .....	599
Jordan Tobias Rosenbaum, <i>Simultaneous interpolation in <math>H_2</math>. II</i> .....	607
Charles Thomas Scarborough, <i>Minimal Urysohn spaces</i> .....	611
Malcolm Jay Sherman, <i>Disjoint invariant subspaces</i> .....	619
Joel John Westman, <i>Harmonic analysis on groupoids</i> .....	621
William Jennings Wickless, <i>Quasi-isomorphism and TFM rings</i> .....	633
Minoru Hasegawa, <i>Correction to "On the convergence of resolvents of operators"</i> .....	641