Pacific Journal of Mathematics

# A NOTE ON EXTREMAL LENGTH AND MODULUS

RICHARD EMANUEL KATZ

Vol. 28, No. 2 April 1969

PACIFIC JOURNAL OF MATHEMATICS
Vol. 28, No. 2, 1969

# A NOTE ON EXTREMAL LENGTH AND MODULUS

RICHARD KATZ

**The equality of extremal length and modulus is shown for general annuli in a Riemannian manifold.**

In a recent paper, [2], Ow showed that the modulus and extremal length of an annulus in a Riemannian manifold are equal under the assumption that the set of critical points of the harmonic measure of the annulus has capacity zero. The purpose of this note is to show that this condition can be dispensed with. We refer the reader to [2] for definitions.

THEOREM. *The extremal length* ($\lambda$) *and modulus* ($\mu$) *of an annulus* $(\Omega, \alpha, \beta)$ *are equal.*

*Proof.* The inequality $\lambda \geqq \mu$ was shown in [2].

To show the opposite inequality, let $u$ be the harmonic measure of $(\Omega, \alpha, \beta)$, and $h$ be a function such that

$$h \mid \alpha = 0, h \mid \beta = 1, \mid \nabla h^2 - \nabla u^2 \mid < \varepsilon ,$$

and $h$ has only a finite number of critical points in $\Omega$. For the existence of such a function see Milnor [1]. Let $\Gamma_0$ be the set of integral curves of $\nabla h$ which do not meet a critical point and $P$ the set of admissible densities on $\Omega$. It is immediate that $\mid \nabla h \mid \in P$. Now for $\gamma \in \Gamma_0$, $\rho \in P$

$$\int_\Omega \rho^2 dV = \int_\alpha \left( \int_\gamma \frac{\rho^2}{\nabla h^2} dh \right) * dh \geqq \int_\alpha \left( \int_\gamma \left| \frac{\rho}{\nabla h} \right| dh \right)^2 * dh$$

and

$$\int_\alpha \left( \int_\gamma \left| \frac{\rho}{\nabla h} \right| dh \right)^2 * dh = \int_\alpha \left( \int_\gamma \rho dl \right)^2 * dh \geqq \inf_{\Gamma_0} \left( \int_\gamma \rho dl \right)^2 \int_\alpha * dh .$$

Since $\mid \nabla h^2 - \nabla u^2 \mid < \varepsilon$, it follows that

$$\left| \int_\alpha * dh - \mu^{-1} \right| < \varepsilon V(\Omega)$$

and therefore,

$$\lambda \leqq \sup_P \inf_{\Gamma_0} \frac{\left( \int_\gamma \rho dl \right)^2}{\int_\Omega \rho^2 dV} \leqq \mu + K\varepsilon$$

for a suitable constant $K$. Since $\varepsilon$ was arbitrary, we have $\lambda \leqq \mu$ which completes the proof.

## References

1. J. Milnor, *Lectures on the h-cobordism theorem*, Princeton University Press, Princeton, N. J., 1965.
2. W. Ow, *An extremal length criterion for the parabolicity of Riemannian spaces*, Pacific J. Math. **23** (1967), 585-590.

Received March 4, 1968.

California State College
Los Angeles, California

# PACIFIC JOURNAL OF MATHEMATICS

## EDITORS

H. ROYDEN
Stanford University
Stanford, California

J. DUGUNDJI
Department of Mathematics
University of Southern California
Los Angeles, California 90007

R. R PHELPS
University of Washington
Seattle, Washington 98105

RICHARD ARENS
University of California
Los Angeles, California 90024

## ASSOCIATE EDITORS

E. F. BECKENBACH  B. H. NEUMANN  F. WOLF  K. YOSIDA

## SUPPORTING INSTITUTIONS

UNIVERSITY OF BRITISH COLUMBIA
CALIFORNIA INSTITUTE OF TECHNOLOGY
UNIVERSITY OF CALIFORNIA
MONTANA STATE UNIVERSITY
UNIVERSITY OF NEVADA
NEW MEXICO STATE UNIVERSITY
OREGON STATE UNIVERSITY
UNIVERSITY OF OREGON
OSAKA UNIVERSITY
UNIVERSITY OF SOUTHERN CALIFORNIA
STANFORD UNIVERSITY
UNIVERSITY OF TOKYO
UNIVERSITY OF UTAH
WASHINGTON STATE UNIVERSITY
UNIVERSITY OF WASHINGTON
*  *  *
AMERICAN MATHEMATICAL SOCIETY
CHEVRON RESEARCH CORPORATION
TRW SYSTEMS
NAVAL WEAPONS CENTER

The Supporting Institutions listed above contribute to the cost of publication of this Journal, but they are not owners or publishers and have no responsibility for its content or policies.


Mathematical papers intended for publication in the *Pacific Journal of Mathematics* should be in typed form or offset-reproduced, double spaced with large margins. Underline Greek letters in red, German in green, and script in blue. The first paragraph or two must be capable of being used separately as a synopsis of the entire paper. It should not contain references to the bibliography. Manuscripts, in duplicate if possible, may be sent to any one of the four editors. Please classify according to the scheme of Math. Rev. **36**, 1539-1546. All other communications to the editors should be addressed to the managing editor, Richard Arens, University of California, Los Angeles, California, 90024.

50 reprints are provided free for each article; additional copies may be obtained at cost in multiples of 50.


The *Pacific Journal of Mathematics* is published monthly. Effective with Volume 16 the price per volume (3 numbers) is $8.00; single issues, $3.00. Special price for current issues to individual faculty members of supporting institutions and to individual members of the American Mathematical Society: $4.00 per volume; single issues $1.50. Back numbers are available.

Subscriptions, orders for back numbers, and changes of address should be sent to Pacific Journal of Mathematics, 103 Highland Boulevard, Berkeley, California, 94708.

PUBLISHED BY PACIFIC JOURNAL OF MATHEMATICS, A NON-PROFIT CORPORATION

Printed at Kokusai Bunken Insatsusha (International Academic Printing Co., Ltd.), 7-17, Fujimi 2-chome, Chiyoda-ku, Tokyo, Japan.

**Pacific Journal of Mathematics**
Vol. 28, No. 2 April, 1969