

Pacific Journal of Mathematics

**A QUASI-DECOMPOSABLE ABELIAN GROUP WITHOUT
PROPER ISOMORPHIC QUOTIENT GROUPS AND PROPER
ISOMORPHIC SUBGROUPS**

JOHN MCCORMICK IRWIN AND TAKASHI ITO

A QUASI-DECOMPOSABLE ABELIAN GROUP WITHOUT PROPER ISOMORPHIC QUOTIENT GROUPS AND PROPER ISOMORPHIC SUBGROUPS

JOHN M. IRWIN AND TAKASHI ITO

All of the group in this paper are abelian p -groups without elements of infinite height. A group is said to be quasi-indecomposable if whenever H is a summand of G then either H or G/H is finite. The p -socle of G is the sub-group consisting of all the elements x in G such that $px = 0$.

In this paper it is shown that there are conditions that can be imposed on the socle of G which are sufficient for G to (a) have no proper isomorphic subgroups; (b) have no proper isomorphic quotient groups; and (c) be quasiindecomposable. Furthermore, it is shown that groups which make these results meaningful actually exist.

Let the cardinality of a group G be either \aleph_0 or greater than $c = 2^{\aleph_0}$. Then, as is well known, G has a proper isomorphic subgroup and a proper isomorphic quotient group. However P. Crawley [3] showed that the cardinality c is exceptional. He gave an example G_0 of cardinality c which has a standard basic subgroup and no proper isomorphic subgroups. After Crawley's example appeared, it was clear that a group, of cardinality c and with a standard basic subgroup, supplies examples of groups with strange but interesting properties. In fact R. S. Pierce [7] gave an example G_1 which has no proper isomorphic subgroups and no proper isomorphic quotient groups. And he gave also in [7] an example G_2 which is quasi-indecomposable, that is, every direct summand H of G_2 is either finite or G_2/H is finite.

The relationship between the above three properties (no proper isomorphic subgroups, no proper isomorphic quotient groups and quasi-indecomposability) of a group G with the cardinality c and a standard basic subgroup seems to authors an interesting problem. In this paper we shall give some results about this problem. In our approach the topological structure of the p -socle of the torsion completion of G will be used in an essential way. Theorem 1 tells us that the situation of the p -socle of G in the p -socle of the torsion completion of G gives us sufficient conditions for these three properties of G . In some sense it shows a relationship between the three properties. Theorem 2 shows the existence of a group which has all three properties. Theorem 3 shows the existence of a group which has no proper isomorphic subgroups and no proper isomorphic quotient groups but which is quasi-decomposable.

Now we want to add a simple proof of the following fact which

was mentioned in the opening of this section.

Let G be an infinite reduced p -group with $\text{card } G = \aleph_0$ or $\text{card } G > c$. Then G has a proper isomorphic subgroup and a proper isomorphic quotient group.

Proof. For simplicity we divide the proof into

Case 1; Suppose G is bounded. Then $G = \sum_{k=1}^n B_k$ where B_k is a direct sum of cyclic groups of order p^k , $B_k = \sum C(p^k)$. Now clearly one of these B_k 's is infinite and throwing out a cyclic summand of B_k yields the desired subgroup and quotient group.

Case 2. Suppose $\text{card } G = \aleph_0$ and G is unbounded. Then $G = H \oplus K$ where H is an unbounded direct sum of cyclic groups (Exercise 19 (a), p.143 in [4]). It is easy to find a proper subgroup A of H which is isomorphic to H and a non-zero subgroup B of H such that $H/B \cong H$. Whence we obtain our proper isomorphic subgroup $A \oplus K$ and our proper isomorphic quotient group G/B .

Case 3. Suppose G is unbounded with $\text{card } G > c$, and $B = \sum_{k=1}^{\infty} B_k$ is a basic subgroup where $B_k = \sum C(p^k)$. Then $G = B_1 \oplus B_2 \oplus \cdots \oplus B_n \oplus G_n$ for all n (Theorem 29.3 in [4]). But as is well known ($\text{card } B)^{\aleph_0} \geq \text{card } G > c$ so that some B_n must be infinite. Now throwing out a cyclic summand of B_n yields the result as in Case 1 and the proof is complete.

2. Sufficient conditions for the three properties. Let $p > 1$ be a fixed prime number, $C(p^n)$ be a cyclic group of order p^n , Σ be the direct sum of cyclic groups $C(p^n)$, Π be the direct product of cyclic groups $C(p^n)$ and C be the torsion group of Π , that is, Σ is the standard basic group and C is the torsion completion of Σ .

The p -socle $C[p]$ of C is a vector space over the prime field of characteristic p and can be topologized as a totally disconnected compact topological group, because Π is clearly a totally disconnected compact topological group with respect to the product topology of compact discrete topologies and the p -socle $C[p]$ of C is the closed subgroup $\{x \mid x \in \Pi, px = 0\}$ of Π . Actually $U_n = \{x \mid x \in C[p] \text{ and } h(x) \geq n\} = (p^n C)[p]$ ($n = 1, 2, \dots$) are open compact subgroups of $C[p]$ and $\{U_n\}$ is a fundamental system of 0-neighborhoods in $C[p]$. These two structures on $C[p]$ which are a vector space and a totally disconnected compact group are used in an essential way in this paper.

Every continuous group homomorphism T on $C[p]$ defines compact subgroups $E_q(T) = \{x \mid x \in C[p] \text{ and } Tx = qx\}$ ($0 \leq q < p$) and the compact subgroup $E(T) = E_0(T) \oplus E_1(T) \oplus \cdots \oplus E_{p-1}(T)$. We can define naturally two types of continuous group homomorphism on $C[p]$ as follows. T is a *singular* homomorphism if $E(T)$ is an open compact subgroup of $C[p]$. For instance a continuous projection on $C[p]$ is

singular. T is a *strongly singular* homomorphism if for some q $E_q(T)$ is an open compact subgroup. If a continuous group homomorphism T on $C[p]$ has a dense subgroup which is invariant under T and on which T is one to one, T is called a semi-isomorphism on $C[p]$.

We have the following theorem which is fundamental to the ideas in what follows.

THEOREM 1. *Let G be a pure subgroup of C which contains Σ and $G[p]$ be the p -socle of G .*

(1) *If $G[p]$ is not invariant under any nonsingular onto homomorphism on $C[p]$, then G has no proper isomorphic quotient groups.*

(2) *If $G[p]$ is not invariant under any nonsingular semi-isomorphism on $C[p]$, then G has no proper isomorphic subgroups.*

(3) *If $G[p]$ is not invariant under any nonstrongly singular projection on $C[p]$, then G is quasi-indecomposable.*

Proof. Suppose φ is a homomorphism of G into G . The purity of G in C implies $\varphi(G[p] \cap U_n) \subset U_n$ for all $n = 1, 2, \dots$. This means that the restriction of φ to $G[p]$ is continuous on $G[p]$. since $G[p] \supset \Sigma[p]$ and $\Sigma[p]$ is dense in $C[p]$, $\varphi|_{G[p]}$ has a unique continuous homomorphism extension T on $C[p]$. Clearly $G[p]$ is invariant under T and $T(U_n) \subset U_n$ for all $n = 1, 2, \dots$. If this T is singular, then there exists a positive integer N such that

$$T(U_N) \subset U_N \subset E(T).$$

Then we have the following decomposition of $G[p]$,

$$G[p] = (G[p] \cap U_N) \oplus R_N = (E_0(T) \cap G[p] \cap U_N) \oplus (E_1(T) \cap G[p] \cap U_N) \oplus \dots \oplus (E_{p-1}(T) \cap G[p] \cap U_N) \oplus R_N,$$

where R_N is a finite subgroup of $G[p]$.

Because $C[p]/U_N$ is finite and $G[p]/G[p] \cap U_N$ is isomorphic to a subgroup $C[p]/U_N$, so the dimension of $G[p]/G[p] \cap U_N$ as a vector space over the prime field of characteristic p is finite. Hence there exists a finite subgroup R_N of $G[p]$ such that $G[p] = (G[p] \cap U_N) \oplus R_N$. The decomposition of $G[p] \cap U_N$ can be shown as follows. For each x in $G[p] \cap U_N$ x is the sum of $z_q \in E_q(T)$ ($0 \leq q < p$); $x = \sum_{q=0}^{p-1} z_q$. Then we have $\varphi^\nu(x) = \sum_{q=0}^{p-1} T^\nu z_q = \sum_{q=0}^{p-1} q^\nu z_q$ for $0 \leq \nu \leq p-1$. Since the determinant of Vandermonde's matrix is not zero mod p , each z_q ($0 \leq q \leq p-1$) is a linear combination of $x, \varphi(x), \dots, \varphi^{p-1}(x)$. This means $z_q \in E_q(T) \cap G[p] \cap U_N$ for $0 \leq q \leq p-1$.

Proof of (1). Suppose φ is an onto homomorphism of G . Then

the continuous extension T of $\varphi|_{G[p]}$ is clearly an onto homomorphism of $C[p]$ and $G[p]$ is invariant under T . By our assumption T must be singular, so we have the above decomposition of $G[p]$. Put $Q_N = (E_1(T) \cap G[p] \cap U_N) \oplus (E_2(T) \cap G[p] \cap U_N) \oplus \cdots \oplus (E_{p-1}(T) \cap G[p] \cap U_N)$, clearly $\varphi(Q_N) = Q_N$ and φ is an isomorphism on Q_N , and

$$(E_0(T) \cap G[p] \cap U_N) \oplus R_N \cong G[p]/Q_N = \varphi(G[p])/\varphi(Q_N) \cong \varphi(R_N)$$

but $\dim \varphi(R_N) \leq \dim R_N < +\infty$. This implies that $E_0(T) \cap G[p] \cap U_N = \{0\}$ and R_N is isomorphic to $\varphi(R_N)$ by φ . Therefore $\varphi|_{G[p]}$ is an isomorphism on $G[p]$. Let $0 \neq x \in G$ and the order of $x = p^n > 1$, then $0 \neq \varphi(p^{n-1}x) = p^{n-1}\varphi(x)$, so $\varphi(x) \neq 0$. Thus φ must be an isomorphism on G .

Proof of (2). Suppose φ is an isomorphism of G into G . We have to show $\varphi(G) = G$. The continuous extension T of $\varphi|_{G[p]}$ is a semi-isomorphism and $G[p]$ is invariant under T . By our assumption T must be singular, so we have the same decomposition of $G[p]$ as above. First of all we can see $\varphi(G[p]) = G[p]$. Automatically

$$E_0(T) \cap G[p] \cap U_N = \{0\},$$

because φ is one to one, therefore $G[p] = Q_N \oplus R_N \cong \varphi(Q_N) \oplus \varphi(R_N) = Q_N \oplus \varphi(R_N) \subset G[p]$ but $\dim R_N = \dim \varphi(R_N) < +\infty$, this implies $\varphi(G[p]) = G[p]$. Next we can see $\varphi(G) \supset G[p^2]$. The group $H = \{x | x \in G \text{ and the first } N-1 \text{ coordinates in } \Pi \text{ are zero}\}$ is a direct summand of G and

$$\begin{aligned} H[p] &= G[p] \cap U_N = Q_N \\ &= (E_1(T) \cap Q_N) \oplus (E_2(T) \cap Q_N) \oplus \cdots \oplus (E_{p-1}(T) \cap Q_N). \end{aligned}$$

We can take a finite group L such that $G = H \oplus L$. We have to show first $\varphi(G) \supset H[p^2]$. For arbitrary x in $H[p^2]$ $px = \sum_{q=0}^{p-1} z_q$ for some $z_q \in E_q(T) \cap Q_N$ ($1 \leq q \leq p-1$), then each z_q is a linear combination of $p\varphi(x)$, $p\varphi^2(x)$, \dots , $p\varphi^{p-1}(x)$. This means that there exist $x_q \in G$ ($1 \leq q \leq p-1$) such that $z_q = p\varphi(x_q)$ for $1 \leq q \leq p-1$. Therefore $px = \sum_{q=1}^{p-1} p\varphi(x_q)$, so $x - \varphi(\sum_{q=1}^{p-1} x_q) \in G[p]$, but $G[p] = \varphi(G[p])$ implies $x \in \varphi(G)$. Now $\varphi(G) \supset G[p^2]$ can be shown. For $x \in G[p^2]$ there exists a positive integer M and integers r_i , $0 \leq r_i \leq p-1$ (at least one of them is not zero) such that $\sum_{i=0}^M r_i p\varphi^i(x) \in Q_N = H[p]$, because $G[p]/Q_N$ is finite dimensional. Since $\varphi(Q_N) = Q_N$, we can assume $r_0 = 1$ without loss of generality. Then we find $z \in H[p^2]$ such that $p \sum_{i=0}^M r_i \varphi^i(x) = pz$. But $H[p^2] \subset \varphi(G)$ has been shown, so $z = \varphi(z')$ for some $z' \in G$, therefore $x + \sum_{i=1}^M r_i \varphi^i(x) - \varphi(z') \in G[p] = \varphi(G[p])$, this implies $x \in \varphi(G)$. Now we can see $\varphi(G) \supset G[p^n]$ for all $n = 1, 2, \dots$ by induction. Namely in general $\varphi(G) \supset G[p^n]$ and the special form of φ on Q_N imply $\varphi(G) \supset H[p^{n+1}]$. And $\varphi(G) \supset H[p^{n+1}]$ and the finiteness of L imply $\varphi(G) \supset G[p^{n+1}]$.

Proof of (3). Suppose G is the direct sum of two subgroups G_1 and G_2 and φ is the projection onto G_1 . The continuous extension T of $\varphi|_{G[p]}$ is also a projection defined on $C[p]$, therefore $C[p] = E_0(T) \oplus E_1(T)$ and $G[p] = (E_0(T) \cap G[p]) \oplus (E_1(T) \cap G[p])$. Since $G[p]$ is invariant under T , T must be strongly singular by our assumption about $G[p]$. Suppose $E_1(T)$ is open, then $E_0(T)$ is finite, hence $G_2[p] = E_0(T) \cap G[p]$ is finite. The finiteness of $G_2[p]$ implies the finiteness of G_2 .

The following is a direct corollary of Theorem 1.

COROLLARY. *Let G be a pure subgroup of C which contains Σ . If $G[p]$ is not invariant under any nonstrongly singular homomorphism on $C[p]$, then G has the three properties stated in (1), (2) and (3) in Theorem 1. Namely G has no proper isomorphic quotient group and no proper isomorphic subgroup, and G is quasi-indecomposable.*

3. Existence theorem

THEOREM 2. *There exists a pure subgroup G of C which contains Σ and satisfies three properties;*

- (1) G has no proper isomorphic quotient groups,
- (2) G has no proper isomorphic subgroups,
- (3) G is quasi-indecomposable.

And an arbitrary pure subgroup H of C such that H contains Σ and $H[p] = G[p]$ satisfies above three properties.

This theorem comes from the corollary of Theorem 1 and following two lemmas. Lemma 1 is known as the purification property, so we omit the proof of Lemma 1 (see more general form in [6]).

LEMMA 1. *For an arbitrary subgroup Q between $\Sigma[p]$ and $C[p]$ there exists a pure subgroup G of C such that G contains Σ and $G[p] = Q$.*

LEMMA 2. *For any family $\{T_\lambda | \lambda \in \Lambda\}$ of nonstrongly singular homomorphisms on $C[p]$ there exists a subgroup Q between $\Sigma[p]$ and $C[p]$ such that Q is not invariant under any $T_\lambda (\lambda \in \Lambda)$.*

The existence of such Q can be shown by transfinite induction which is Crawley's idea in [3]. We need following lemma which is also essentially Crawley's.

LEMMA 3. *Suppose T is a nonstrongly singular homomorphism on $C[p]$. Then there exists a one-parameter family $\Delta(T) = \{x_t | 0 \leq t \leq 1\}$ of elements in $C[p]$ such that four elements x_s, x_t, Tx_s and Tx_t are*

linearly independent for arbitrary $s \neq t$.

Proof. The proof can be divided into two cases (a) and (b).

(a) T is singular but not strongly singular. In this case, by Baire's category theorem ($C[p]$ is a complete metric space) there are at least two q and q' such that both $E_q(T)$ and $E_{q'}(T)$ are infinite compact groups, so $\text{card } E_q(T) = \text{card } E_{q'}(T) = c$ (for instance, see [5], p. 31). Therefore $\dim E_q(T) = \dim E_{q'}(T) = c$. Let $\{y_t \mid 0 \leq t \leq 1\}$ be a basis of $E_q(T)$ and $\{y'_t \mid 0 \leq t \leq 1\}$ be a basis of $E_{q'}(T)$. Then $\Delta(T) = \{y_t + y'_t \mid 0 \leq t \leq 1\}$ is the desired family.

(b) T is not singular. In this case, by Baire's category theorem $U_n/E(T) \cap U_n$ are infinite compact groups for all $n = 1, 2, \dots$, so as above $\dim U_n/E(T) \cap U_n = c$. Hence $U_n = (E(T) \cap U_n) \oplus D_n$ with $\dim D_n = c$ for all $n = 1, 2, \dots$. Take $0 \neq x_0 \in D_1$, then x_0 and Tx_0 are linearly independent. Let $\{z_0, z_1, \dots, z_{p^2-1}\}$ be the group generated by x_0 and Tx_0 , then by the continuity of T we can find U_M such that $z_i + U_M + T(U_M)$ ($0 \leq i \leq p^2 - 1$) are mutually disjoint. For this M we take a basis $\{y_t \mid 0 \leq t \leq 1\}$ of D_M . Then $\Delta(T) = \{x_0 + y_t \mid 0 \leq t \leq 1\}$ is the desired system. Because, suppose $n_1(x_0 + y_t) + n_2(Tx_0 + Ty_t) = n'_1(x_0 + y_s) + n'_2(Tx_0 + Ty_s)$ for $s \neq t$ where n_1, n_2, n'_1 and n'_2 are integers, then $n_1x_0 + n_2Tx_0 + n_1y_t + n_2Ty_t = n'_1x_0 + n'_2Tx_0 + n'_1y_s + n'_2Ty_s$, and $n_1x_0 + n_2Tx_0$ must be some z_i and also $n'_1x_0 + n'_2Tx_0$ must be some z_j , but $z_i = z_j$ by our choice of U_M . This implies $n_1 = n'_1 \bmod p$ and $n_2 = n'_2 \bmod p$, therefore we have $n_1y_t + n_2Ty_t = n_1y_s + n_2Ty_s$, whence $n_1(y_t - y_s) = -n_2T(y_t - y_s)$. However $0 \neq y_t - y_s \in D_M$ and $D_M \cap E(T) = \{0\}$, hence $n_1 = n_2 = 0 \bmod p$.

Proof of Lemma 2. $\{T_\lambda \mid \lambda \in A\}$ is given, then $\text{card } A$ is at most c (note that the cardinality of the set of all continuous homomorphisms on $C[p]$ is at most c , because $C[p]$ is a separable compact group). We assume that A is a well ordered set of ordinal numbers which are less than Ω , where Ω is the first ordinal number whose cardinality is c . Choose $e \in C[p]$ but $e \notin \Sigma[p]$, then we can construct a family of subgroups $R_\lambda (\lambda \in A)$ by transfinite induction as follows:

- (a) $\Sigma[p] = R_0 \subset R_\lambda \subset R_\mu \subset C[p]$ if $0 \leq \lambda < \mu$ ($\lambda, \mu \in A$),
- (b) $\text{card } R_\lambda \leq \text{card } \lambda \cdot \aleph_0 < c$ for all $\lambda \in A$,
- (c) $e \notin R_\lambda$ but there exists $x_\lambda \in R_\lambda \cap \Delta(T_\lambda)$ such that $e - T_\lambda x_\lambda \in R_\lambda$.

Suppose R_λ has been constructed for all $\lambda < \mu \in A$. Let $R'_\mu = \bigcup_{\lambda < \mu} R_\lambda$. Then $\text{card } ([e] + R'_\mu) \leq \text{card } \mu \cdot \aleph_0 < c$, where $[e]$ is the group generated by e . The property of $\Delta(T_\mu)$ in Lemma 3 guarantees the existence of $x_{t_0} \in \Delta(T_\mu)$ such that $([e] + R'_\mu) \cap ([x_{t_0}] + [T_\mu x_{t_0}]) = \{0\}$. Then $R_\mu = R'_\mu + [x_{t_0}] + [e - T_\mu x_{t_0}]$ is the desired subgroup. Let $Q = \bigcup_{\lambda \in A} R_\lambda$, then by (a) Q is a subgroup of $C[p]$ which contains $\Sigma[p]$ and by (c) Q is not invariant under any $T_\lambda (\lambda \in A)$.

4. A quasi-decomposable group without proper isomorphic quotient groups and proper isomorphic subgroups.

THEOREM 3. *There exists a pure subgroup G of C which contains Σ and satisfies properties;*

- (1) G has no proper isomorphic quotient groups,
- (2) G has no proper isomorphic subgroups,
- (3) G has a decomposition $G_1 \oplus G_2$ such that G_1 and G_2 are not bounded.

The following lemma is essential for our proof of this theorem.

LEMMA 4. *For any family $\{T_\lambda | \lambda \in \Lambda\}$ of nonsingular homomorphisms on $C[p]$ there exists a subgroup Q between $\Sigma[p]$ and $C[p]$ such that Q is not invariant under any $T_\lambda (\lambda \in \Lambda)$ but invariant under the canonical projection P_e onto even coordinates.*

The outline of the proof of this lemma will be given later.

Proof of Theorem 3. Every element of C has countable coordinates as an element of the product space $\prod_{n=1}^{\infty} C(p^n)$; $x \in C$ is called an even (odd) element if all odd (even) coordinates are zero. For a subset A of C $A^e(A^o)$ means the set of all even (odd) elements in A . Then clearly $C = C^e \oplus C^o$ and $\Sigma = \Sigma^e \oplus \Sigma^o$. By Lemma 4 there exists a subgroup Q between $\Sigma[p]$ and $C[p]$ such that Q is not invariant under any nonsingular homomorphisms on $C[p]$ but is invariant under P_e , therefore $\Sigma^e[p] = \Sigma[p]^e \subset Q^e \subset C[p]^e = C^e[p]$, $\Sigma^o[p] = \Sigma[p]^o \subset Q^o \subset C[p]^o = C^o[p]$ and $Q = Q^e \oplus Q^o$. With exactly the same proof as that of Lemma 1 we can show that there exists a pure subgroup $G_1(G_2)$ of $C^e(C^o)$ which contains $\Sigma^e(\Sigma^o)$ and $G_1[p] = Q^e(G_2[p] = Q^o)$. Clearly G_1 and G_2 are not bounded. Let $G = G_1 \oplus G_2$, then G is a pure subgroup of C which contains Σ and $G[p] = G_1[p] \oplus G_2[p] = Q^e \oplus Q^o = Q$. By Theorem 1 G has the properties (1) and (2) in Theorem 3.

The outline of the proof of Lemma 4. In order to prove Lemma 4 we can apply a similar method to the construction of Q in Lemma 2. However before doing it we have to prepare some reformation of Lemma 3. Precisely our reformation is as follows, hereafter we shall use notations $A^e = P_e(A)(A^o = (I - P_e)(A))$ for a subset A of $C[p]$ and $x^e = P_e x(x^o = x - P_e x)$ for an element x in $C[p]$.

For an arbitrary nonsingular homomorphism T we can find a one-parameter family $\Delta(T) = \{x_t | 0 \leq t \leq 1\}$ of elements in $C[p]$ which has one of the following six properties; $1^o, 2^o, 3^o, 1^e, 2^e$ and 3^e ,

- 1° $x_t, Tx_t \in C[p]^0$ for all $0 \leq t \leq 1$ and four elements x_s, x_t, Tx_s and Tx_t are linearly independent for arbitrary $s \neq t$,
 2° there exists $q, 0 \leq q \leq p-1$ such that $x_t \in C[p]^0$ and

$$Tx_t - qx_t \in C[p]^e$$

for all $0 \leq t \leq 1$ and four elements $x_s, x_t, Tx_s - qx_s$ and $Tx_t - qx_t$ are linearly independent for arbitrary $s \neq t$,

- 3° $x_t \in C[p]^0$ for all $0 \leq t \leq 1$ and six elements $x_s, x_t, (Tx_s)^0, (Tx_s)^e, (Tx_t)^0$ and $(Tx_t)^e$ are linearly independent for arbitrary $s \neq t$.

1°, 2° and 3° are dual properties 1°, 2° and 3° by exchanging odd for even.

In the proof of this we have some difficulty coming from non-commutativity of nonsingular homomorphism and P_e . The proof in our original manuscript needs a long computation, in this paper we omit our detailed computation according to referee's suggestion but authors can supply the detailed proof to interested readers.

Using above $\Delta(T)$ the existence of Q in Lemma 4 can be shown as follows. Let $\{T_\lambda | \lambda \in A\}$ be a given family of nonsingular homomorphisms on $C[p]$. We assume that A is a well ordered set of ordinal numbers which are less than the first ordinal number whose cardinality is c . Choose $c \in C[p]$ but $c^0, c^e \notin \Sigma[p]$. By transfinite induction we can construct the following family of subgroups $R_\lambda (\lambda \in A)$;

- (a) $\Sigma[p] = R_0 \subset R_\lambda \subset R_\mu \subset C[p]$ if $0 \leq \lambda < \mu (\lambda, \mu \in A)$,
- (b) $\text{card } R_\lambda \leq \text{card } \lambda \cdot \aleph_0 < c$ for all $\lambda \in A$,
- (c) R_λ is invariant under P_e for all $\lambda \in A$,
- (d) c^0 and $c^e \notin R_\lambda$ but there exists $x_\lambda \in R_\lambda \cap \Delta(T_\lambda)$ such that $c^0 - T_\lambda x_\lambda$ or $c^e - T_\lambda x_\lambda$ or $c - T_\lambda x_\lambda \in R_\lambda$ for all $\lambda \in A$.

Suppose R_λ has been constructed for all $\lambda < \mu \in A$. Let $R'_\mu = \bigcup_{\lambda < \mu} R_\lambda$. Then $\text{card } R'_\mu \leq \text{card } \mu \cdot \aleph_0 < c$ and R'_μ is invariant under P_e and c^0 and $c^e \notin R'_\mu$. Let $\Delta(T_\mu)$ be one having one of properties $1^0 \sim 3^0$ and $1^e \sim 3^e$. Suppose $\Delta(T_\mu)$ has property 1^0 , then we can find $x_\mu \in \Delta(T_\mu)$ such that $(R'_\mu + [c^0] + [c^e]) \cap ([x_\mu] \oplus [T_\mu x_\mu]) = \{0\}$. Let

$$R_\mu = R'_\mu + [x_\mu] + [c^0 - T_\mu x_\mu],$$

then clearly R_μ satisfies above (a), (b) and (c). And c^0 and $c^e \in R_\mu$ also holds. Suppose $c^0 \in R_\mu$, then $c^0 = x + nx_\mu + m(c^0 - T_\mu x_\mu)$ for some $x \in R'_\mu$ and some integers n and m , so $-x + (1-m)c^0 = nx_\mu - mT_\mu x_\mu$, but by our choice of $x_\mu, nx_\mu - mT_\mu x_\mu = 0$ and $x + (m-1)c^0 = 0$. This implies $n = m = 0 \pmod p$ and $c^0 = x \in R'_\mu$ which is a contradiction. Suppose $c^e \in R_\mu$, then $c^e = x + nx_\mu + m(c^0 - T_\mu x_\mu)$ for some $x \in R'_\mu$ and some integers n and m , but x_μ and $T_\mu x_\mu \in C[p]^0$, so $c^e = x \in R'_\mu$ which

is also a contradiction. Suppose $\Delta(T_\mu)$ has property 2°, then we can find $x_\mu \in \Delta(T_\mu)$ such that $(R'_\mu + [c^0] + [c^e]) \cap ([x_\mu] \oplus [T_\mu x_\mu - qx_\mu]) = \{0\}$. Let $R_\mu = R'_\mu + [x_\mu] + [c^e - T_\mu x_\mu + qx_\mu]$, then clearly R_μ satisfies above (a), (b) and (c). And c^0 and $c^e \notin R_\mu$ also holds. Suppose $c^0 \in R_\mu$, then $c^0 = x + nx_\mu + m(c^e - T_\mu x_\mu + qx_\mu)$ for some $x \in R'_\mu$ and some integers n and m , but $x_\mu \in C[p]^0$ and $T_\mu x_\mu - qx_\mu \in C[p]^e$, hence we have $c^0 = x^0 + nx_\mu$, that is, $-x^0 + c^0 = nx_\mu$. Our choice of x_μ implies $nx_\mu = 0 = -x^0 + c^0$, so we have $c^0 = x^0 \in R'_\mu$ which is a contradiction. Suppose $c^e \in S_\mu$, then $c^e = x + nx_\mu + m(c^e - T_\mu x_\mu + qx_\mu)$ for some $x \in R'_\mu$ and some integers n and m . Hence $-x + (1 - m)c^e = nx_\mu - m(T_\mu x_\mu - qx_\mu)$, but by our choice of x_μ we see $-x + (1 - m)c^e = 0 = nx_\mu - m(T_\mu x_\mu - qx_\mu)$. This implies $n = m = 0 \pmod p$, so $c^e = x \in R'_\mu$ which is also a contradiction. Suppose $\Delta(T_\mu)$ has property 3°, then we can find $x_\mu \in \Delta(T_\mu)$ such that $(R'_\mu + [c^0] + [c^e]) \cap ([x_\mu] \oplus [(T_\mu x_\mu)^0] \oplus [(T_\mu x_\mu)^e]) = \{0\}$. Let

$$R_\mu = R'_\mu + [x_\mu] + [c^0 - (T_\mu x_\mu)^0] + [c^e - (T_\mu x_\mu)^e].$$

Then R_μ clearly satisfies (a), (b) and (c). And c^0 and $c^e \notin R_\mu$ can be seen as follows. Suppose $c^0 = x + nx_\mu + m(c^0 - (T_\mu x_\mu)^0) + m'(c^e - (T_\mu x_\mu)^e)$ for some $x \in R'_\mu$ and integers n, m and m' , then

$$c^0 = x^0 + nx_\mu + m(c^0 - (T_\mu x_\mu)^0),$$

so $-x^0 + (1 - m)c^0 = nx_\mu - m(T_\mu x_\mu)^0$. This implies $nx_\mu - m(T_\mu x_\mu)^0 = 0 = -x^0 + (1 - m)c^0$ by our choice of x_μ . Hence $m = 0$ and $c^0 = x^0 \in R'_\mu$ which is a contradiction. We can see also $c^e \notin R_\mu$ for same reason. And x_μ and $c - T_\mu x_\mu \in R_\mu$ is clear. The construction of R_μ for $\Delta(T_\mu)$ having one of properties 1° ~ 3° is exactly similar by exchanging odd for even.

Let $Q = \bigcup_{\lambda \in A} R_\lambda$. Then the above properties (a) ~ (d) for all R_λ guarantee that Q is a subgroup between $\Sigma[p]$ and $C[p]$ not invariant under any $T_\lambda (\lambda \in A)$ but invariant under P_e .

REFERENCES

1. G. Baumslag, *Hopficity and abelian groups*, Topics in abelian groups, Proceedings of the New Mexico Symposium on Abelian Groups, 1962, 331-335.
2. R. A. Beaumont and R. S. Pierce, *Partly transitive modules and monules with proper isomorphic submodules*, Trans. Amer. Math. Soc. **91** (1959) 209-219.
3. P. Crawley, *An infinite primary abelian group without proper isomorphic subgroups*, Bull. Amer. Math. Soc. **68** (1962) 463-467.
4. L. Fuchs, *Abelian groups*, Pergamon Press, Oxford, 1960.
5. E. Hewitt and K. A. Ross, *Abstract harmonic analysis I*, Academic Press, New York, 1963.
6. J. M. Irwin, F. Richman and E. A. Walker, *Countable direct sums of torsion complete groups*, Proc. Amer. Math. Soc. **17** (1966), 763-766.

7. R. S. Pierce, *The endomorphism rings of primary abelian groups*, Topics in abelian groups, Proceedings of the New Mexico Symposium on Abelian Groups, (1962) 215-310.

Received August 22, 1967.

WAYNE STATE UNIVERSITY AND
HOKKAIDO UNIVERSITY

Jorge Alvarez de Araya, <i>A Radon-Nikodým theorem for vector and operator valued measures</i>	1
Deane Eugene Arganbright, <i>The power-commutator structure of finite p-groups</i>	11
Richard Eugene Barlow, Albert W. Marshall and Frank Proschan, <i>Some inequalities for starshaped and convex functions</i>	19
David Clarence Barnes, <i>Some isoperimetric inequalities for the eigenvalues of vibrating strings</i>	43
David Hilding Carlson, <i>Critical points on rim-compact spaces</i>	63
Allan Matlock Weber Carstens, <i>The lattice of pretopologies on an arbitrary set S</i>	67
S. K. Chatterjea, <i>A bilateral generating function for the ultraspherical polynomials</i>	73
Ronald J. Ensey, <i>Primary Abelian groups modulo finite groups</i>	77
Harley M. Flanders, <i>Relations on minimal hypersurfaces</i>	83
Allen Roy Freedman, <i>On asymptotic density in n-dimensions</i>	95
Kent Ralph Fuller, <i>On indecomposable injectives over artinian rings</i>	115
George Isaac Glauber, <i>Normalizers of p-subgroups in finite groups</i>	137
William James Heinzer, <i>On Krull overrings of an affine ring</i>	145
John McCormick Irwin and Takashi Ito, <i>A quasi-decomposable abelian group without proper isomorphic quotient groups and proper isomorphic subgroups</i>	151
Allan Morton Krall, <i>Boundary value problems with interior point boundary conditions</i>	161
John S. Lowndes, <i>Triple series equations involving Laguerre polynomials</i>	167
Philip Olin, <i>Indefinability in the arithmetic isolic integers</i>	175
Ki-Choul Oum, <i>Bounds for the number of deficient values of entire functions whose zeros have angular densities</i>	187
R. D. Schafer, <i>Standard algebras</i>	203
Wolfgang M. Schmidt, <i>Irregularities of distribution. III</i>	225
Richard Alfred Tapia, <i>An application of a Newton-like method to the Euler-Lagrange equation</i>	235